1,3-Diketones from Acid Chlorides: A Rapid and General One-Pot Synthesis of Pyrazoles

Stephen T. Heller^{P*P} and Swaminathan R. Natarajan

Department of Medicinal Chemistry, Merck Research Laboratories, P.O. Box 2000, Rahway, New Jersey 07065

stephen_heller@merck.com

Supporting Information

Table of Contents

General Procedure			S3
General Experimental Information			S3
Compound Data:			
5, 6a,b			S4
7, 8, 9		S5	
10, 12, 13			S6
14, 15, 16			S7
17, 19, 20			S8
21a,b, 22			S9
23, 24, 25			S10
NMR spectra:			
7	¹ H		S12
7		¹³ C	S13
8	¹ H		S14
8		¹³ C	S15
9	¹ H		S16
9		¹³ C	S17
10	¹ H		S18
10		¹³ C	S19
5	¹ H		S20
5		¹³ C	S21

12	^1H		S22	
12		^{13}C		S23
13	^1H		S24	
13		^{13}C		S25
14	^1H		S26	
14		^{13}C		S27
15	^1H		S28	
15		^{13}C		S29
16	^1H		S30	
16		^{13}C		S31
19	^1H		S32	
19		^{13}C		S33
20	^1H		S34	
20		^{13}C		S35
22	^1H		S36	
22		^{13}C		S37
23	^1H		S38	
23		^{13}C		S39
24	^1H		S40	
24		^{13}C		S41
25	^1H		S42	
25		^{13}C		S43

General Procedure: Ketone (2 mmol) was dissolved in 5 mL dry toluene in a screw cap vial (with septum) and then the solution was cooled to 0°C under nitrogen. LiHMDS (2.1 mL, 1.0 M in THF, 2.1 mmol) was added quickly via syringe with agitation stirring and the formed anion was allowed to sit for approximately 1 minute before the addition *via syringe*¹ of acid chloride (1 mmol) in one portion with stirring. The vial was then removed from the ice bath and allowed to stand for 1 minute, then 2 mL AcOH was added with stirring. 10 mL EtOH and 5 mL THF was added to form a homogeneous mixture, and then hydrazine hydrate (2 mL, 1.1 g, 34.3 mmol) or substituted hydrazine (5 mmol) was added and the mixture was allowed to auto-reflux and was held at that temperature for 5 minutes, when LCMS showed that all diketone had reacted. The resulting solution was added to 1.0 M NaOH solution and extracted with EtOAc. The organic fraction was then washed with brine, dried over Na₂SO₄, and evaporated under reduced pressure. See specific compounds for purification details.

General Comments: All reagents were purchased from Aldrich and were used as received. LiHMDS solutions were rigorously kept under a nitrogen atmosphere after opening. Dry toluene and THF were supplied by Acros. All other solvents were provided by Fisher Scientific. All chemistry was performed under a nitrogen atmosphere using standard techniques. NMR spectra were taken with a Varian Inova (400 MHz or 500 MHz). LC/MS were obtained using a Agilent 1100 series LC and Micromass Zq spectrometer. Column chromatography was performed using a Biotage Horizon system utilizing standard pre-packed silica gel cartridges (40g).

¹ Solid acid chlorides were dissolved in 2 mL toluene and the minimum amount of THF required to make a solution.

Spectroscopic Notes: All spectra calibrated with TMS, or in the case of ^{13}C experiments, the solvent peak ($\text{CDCl}_3 = 77.0$ ppm, $\text{d}_6\text{-DMSO} = 39.5$ ppm). The room temperature ^1H NMR spectra of **11** and **20** showed two distinct tautomers. In both cases, the relevant peaks converged substantially. Reported data is taken from experiments run at 70°C . This effect was also observed in the ^{13}C NMR for **9**, for which the reported data corresponds to the major tautomer. In a few cases quaternary carbons were not detected in ^{13}C NMR experiments.

5-(4-bromophenyl)-3-(4-methoxyphenyl)-1H-pyrazole (5). Crystallized from 2-propanol/water. ^1H NMR (400 MHz, $\text{d}_6\text{-DMSO}$) δ 13.30 (s, 1H), 7.80 (d, $J = 7.6$, 2H), 7.75 (d, $J = 8$ Hz, 2H), 7.64 (d, $J = 7.6$ Hz, 2H), 7.11 (s, 1H), 7.03 (d, $J = 8$ Hz, 2H), 3.80 (s, 3H); ^{13}C NMR (500 MHz, $\text{d}_6\text{-DMSO}$) δ 159.1, 131.7, 127.0, 126.5, 120.7, 114.3, 99.0, 55.1; **HRMS** calcd for $\text{C}_{16}\text{H}_{14}\text{BrN}_2\text{O}$, 331.0271; found, 331.0280.

3-(4-bromophenyl)-5-(4-methoxyphenyl)-1-methylpyrazole (6a). Crystallized from 2-propanol/water. ^1H NMR (400 MHz, CDCl_3) δ 7.73 (d, $J = 8.8$ Hz, 2H), 7.55 (d, $J = 8.4$ Hz, 2H), 7.27 (d, $J = 8.8$ Hz, 2H), 6.92 (d, $J = 8.4$ Hz, 2H), 6.49 (s, 1H), 3.83 (s, 3H), 3.78 (s, 3H); **MS** calcd for $\text{C}_{17}\text{H}_{15}\text{BrN}_2\text{O}$, 343.04; found, 343.06.

5-(4-bromophenyl)-3-(4-methoxyphenyl)-1-methylpyrazole (6b). Crystallized from 2-propanol/water. ^1H NMR (400 MHz, CDCl_3) δ 7.68 (d, $J = 8.8$ Hz, 2H), 7.49 (d, $J = 8.4$ Hz, 2H), 7.34 (d, $J = 8.8$ Hz, 2H), 6.96 (d, $J = 8.4$ Hz, 2H), 6.49 (s, 1H), 3.84 (s, 3H), 3.81 (s, 3H); $\text{C}_{17}\text{H}_{15}\text{BrN}_2\text{O}$, 343.04; found, 343.08.

3-(4-bromophenyl)-5-pentyl-1H-pyrazole (7). Crystallized from 2-propanol/water.

¹H NMR (400 MHz, CDCl₃) δ 7.59 (d, *J* = 8.4 Hz, 2H), 7.47 (d, *J* = 8.4), 6.31 (s, 1H), 2.59-2.53 (m 2H), 1.61-1.58 (m, 2H), 1.30-1.28 (m, 4H), 0.88 (t, 3H); **¹³C NMR** (400 MHz, CDCl₃) δ 131.7, 127.2, 121.6, 100.9, 31.4, 28.8, 26.1, 22.4, 13.9; **HRMS** calcd for C₁₄H₁₈BrN₂, 295.0634; found, 295.0644.

3-(4-bromophenyl)-5-(5-chloropentyl)-1H-pyrazole (8). Crystallized from 2-

propanol/water. **¹H NMR** (400 MHz, CDCl₃) δ 7.59 (d, *J* = 8.4 Hz, 2H), 7.49 (d, *J* = 8.4), 6.33 (s, 1H), 3.51 (t, 2H), 2.63-2.60 (m 2H), 1.80-1.73 (quint, 2H), 1.70-1.62 (quint, 2H), 1.50-1.44 (m, 2H); **¹³C NMR** (500 MHz, CDCl₃) δ 131.8, 127.2, 121.8, 101.1, 44.8, 32.2, 28.4, 26.4, 26.0; **HRMS** calcd for C₁₄H₁₇BrClN₂, 329.0242; found, 329.0240.

5-(4-bromophenyl)-3-(4-cyanophenyl)-1H-pyrazole (9). Recrystallized from 2-

propanol/water. **¹H NMR** (400 MHz, d₆-DMSO) δ 13.71 (s, 1H), 8.04-7.92 (m, 4H), 7.79-7.70 (m, 4H), 7.42 (s, 1H); **¹³C NMR** (500 MHz, d₆-DMSO) δ 149.8, 142.8, 137.9,

132.7, 132.0, 128.2, 127.1, 125.6, 121.4, 119.0, 109.8, 101.1; **HRMS** calcd for $C_{16}H_{11}BrN_3$, 326.0117; found, 326.0107

5-(2-methylphenyl)-3-(4-cyanophenyl)-1H-pyrazole (10). Crystallized from 2-propanol/water. **1H NMR** (400 MHz, d_6 -DMSO, $T = 70^\circ C$) δ 12.97 (s, 1H), 7.88-7.75 (b, 2H), 7.67-7.58 (b, 2H), 7.46 (b, 1H), 7.36-7.20 (b, 3H), (bd, 6.92-6.84, 1H), 2.41 (s, 3H); **^{13}C NMR** (400 MHz, d_6 -DMSO) δ 149.5, 142.9, 135.5, 133.0, 131.9, 131.5, 130.8, 129.4, 128.7, 128.4, 127.1, 126.0, 120.3, 20.5; **HRMS** calcd for $C_{16}H_{14}BrN_2$, 315.0321; found, 315.0324.

3-(4-(N,N-dimethylamino)phenyl)-5-(4-methoxyphenyl)-1H-pyrazole (12). Crystallized from 2-propanol/water. **1H NMR** (400 MHz, d_6 -DMSO) δ 12.92 (s, 1H), 7.75-7.62 (m, 4H), 6.99 (d, 2H), 6.86 (s, 1H), 6.77 (d, $J = 8$ Hz, 2H), 3.79 (s, 3H), 2.94 (s, 6H); **^{13}C NMR** (500 MHz, d_6 -DMSO) δ 158.8, 150.0, 126.4, 126.0, 114.1, 112.3, 97.3, 55.1, 40.0; **HRMS** calcd for $C_{18}H_{20}BrN_3O$, 294.1606; found, 294.1606.

3-(4-cyanophenyl)-5-(4-methoxyphenyl)-1H-pyrazole (13). Recrystallized from toluene. **1H NMR** (400 MHz, d_6 -DMSO) δ 13.47 (s, 1H), 8.04 (d, $J = 7.6$ Hz, 2H), 7.90 (d, $J = 7.6$ Hz, 2H), 7.76 (d, $J = 8.0$ Hz, 2H), 7.25 (s, 1H), 7.05 (d, $J = 8$ Hz, 2H), 3.81 (s,

3H); **¹³C NMR** (500 MHz, d₆-DMSO) δ 159.2, 132.7, 126.6, 125.6, 117.0, 114.3, 109.7, 99.9, 55.1; **HRMS** calcd for C₁₇H₁₄N₃O, 276.1137; found, 276.1161.

5-(4-chlorophenyl)-3-(4-methoxyphenyl)-4-phenyl-1H-pyrazole (14). Crystallized from 2-propanol/water. **¹H NMR** (400 MHz, d₆-DMSO) δ 13.36 (s, 1H), 7.34-7.16 (m, 11H), 6.91-6.83 (m, 2H), 3.74 (s, 3H); **¹³C NMR** (400 MHz, d₆-DMSO) δ 159.0, 147.7, 140.4, 133.7, 131.7, 130.5, 128.9, 128.7, 128.6, 128.1, 127.1, 121.7, 116.2, 114.0, 113.5, 55.1; **HRMS** calcd for C₁₂₂H₁₈ClN₂O, 361.1108; found, 361.1111.

5-(4-chlorophenyl)-3-(4-methoxyphenyl)-4-propyl-1H-pyrazole (15). Crystallized from 2-propanol/water. **¹H NMR** (400 MHz, d₆-DMSO) δ 13.00 (s, 1H), 7.67-7.46 (m, 6H), 7.09-7.03 (m, 2H), 3.81 (s, 3H), 2.64 (t, 2H), 1.41-1.35 (hex, 2H), 0.78 (t, 3H); **¹³C NMR** (400 MHz, d₆-DMSO) δ 159.0, 148.5, 140.6, 133.8, 131.7, 129.4, 129.0, 128.8, 128.4, 122.6, 114.2, 55.1, 25.3, 23.4, 13.9; **HRMS** calcd for C₁₉H₂₀ClN₂O, 327.1264; found, 327.1257.

3-(4-methoxyphenyl)-5-(4-nitrophenyl)-1H-pyrazole (16). Crystallized from 2-propanol/water. **¹H NMR** (400 MHz, d₆-DMSO) δ 13.60 (s, 1H), 8.30 (d, *J* = 8 Hz, 2H),

8.12 (d, $J = 8$ Hz, 2H), 7.76 (d, $J = 8$ Hz, 2H), 7.29 (s, 1H), 7.06 (d, $J = 8$ Hz, 2H), 3.81 (s, 3H); ^{13}C NMR (400 MHz, $\text{d}_6\text{-DMSO}$) δ 159.4, 146.3, 126.7, 125.8, 124.1, 100.0, 55.2; HRMS calcd for $\text{C}_{16}\text{H}_{15}\text{N}_3\text{O}_3$, 296.1035; found, 296.1019.

5-(4-methoxyphenyl)-3-(4-nitrophenyl)-1H-pyrazole (17). Purified by column chromatography (hexanes:ethyl acetate, 19:1). ^1H NMR (400 MHz, CDCl_3) δ 8.30 (d, $J = 7.2$ Hz, 2H), 8.08 (d, $J = 7.2$ Hz, 2H), 7.38 (m, 5H) 7.20 (d, $J = 8.8$ Hz, 2H), 6.87 (d, $J = 8.8$ Hz, 2H), 6.86 (s, 1H), 3.82 (s, 3H); MS calcd for $\text{C}_{22}\text{H}_{17}\text{N}_3\text{O}$, 372.13; found, 372.04. m.p. 176-178°C (Lit. 178-179°C)²

3-(4-pyridyl)-5-(4-methoxyphenyl)-1H-pyrazole (19). Crystallized from ethanol/water. ^1H NMR (400 MHz, $\text{d}_6\text{-DMSO}$) δ 13.53 (s, 1H), 8.63 (d, $J = 4.8$ Hz, 2H), 7.81 (d, $J = 4.8$ Hz, 2H), 7.77 (d, $J = 8$ Hz, 2H), 7.28 (s, 1H), 7.06 (d, $J = 8$ Hz, 2H), 3.81 (s, 3H); ^{13}C NMR (400 MHz, $\text{d}_6\text{-DMSO}$) δ 159.2, 150.2, 126.6, 119.4, 114.4, 99.9, 55.2; MS calcd for $\text{C}_{15}\text{H}_{14}\text{N}_3\text{O}$, 252.11; found, 252.18.

3-(2-thienyl)-5-(4-methoxyphenyl)-1H-pyrazole (20). Crystallized from 2-propanol/water. ^1H NMR (400 MHz, $\text{d}_6\text{-DMSO}$, $T = 70^\circ\text{C}$) δ 12.98 (s, 1H), 7.71 (d, $J = 6$ Hz, 2H), 7.40 (b, 2H), 7.09 (s, 1H), 7.01 (d, $J = 6.8$ Hz, 2H), 6.83 (s, 1H), 3.80 (s, 3H); ^{13}C NMR (400 MHz, $\text{d}_6\text{-DMSO}$) δ 159.2, 146.8, 143.3, 137.1, 127.5, 126.6, 124.5,

² Bishop, B. C. *Synthesis* **2004**, 1, 43.

123.5, 121.7, 114.4, 98.5, 55.2; **HRMS** calcd for C₁₄H₁₃N₂OS, 257.0749; found, 257.0742.

5-(4-methylphenyl)-3-phenyl-1-methylpyrazole (21a). Purified by column chromatography (hexanes:ethyl acetate, 24:1). **¹H NMR** (400 MHz, CDCl₃) δ 7.81 (d, *J* = 7.2 Hz, 2H), 7.41-7.25 (m, 7H), 6.58 (s, 1H), 3.92 (s, 3H), 2.42 (s, 3H);³ **MS** calcd for C₁₇H₁₆N₂, 249.13; found, 249.32.

3-(4-methylphenyl)-5-phenyl-1-methylpyrazole (21b). Purified by column chromatography (hexanes:ethyl acetate, 24:1). **¹H NMR** (400 MHz, CDCl₃) δ 7.73 (d, *J* = 8.4 Hz, 2H), 7.48-7.45 (m, 5H), 7.22 (d, *J* = 8.4 Hz, 2H), 6.58 (s, 1H), 3.92 (s, 3H), 2.37 (s, 3H);³ **MS** calcd for C₁₇H₁₆N₂, 249.13; found, 249.27.

3-(4-methoxyphenyl)-4,5,6,7-tetrahydro-1H-indazole (22). Flash chromatography (2:1, dichloromethane : acetone). Analytical sample crystallized (0°C) from dichloromethane/hexanes. **¹H NMR** (400 MHz, CDCl₃) δ 7.55 (d, *J* = 8.8 Hz, 2H), 6.90 (d, *J* = 8.8 Hz, 2H), 3.82 (s, 3H), 2.68 (t, 2H), 2.63 (t, 2H), 1.82-1.76 (m, 4H); **¹³C NMR** (400 MHz, CDCl₃) δ 158.9, 127.8, 125.5, 114.0, 112.4, 55.2, 28.4, 23.6, 22.3, 22.0; **MS** calcd for C₁₄H₁₇N₂O, 229.13; found, 229.37.

³ Agrees with isomeric assignments in literature; Ahmed, M., S. M.; Kobayashi, K.; Mori, A. *Org. Lett.* **2005**, 7, 4487.

3-pentyl-4,5,6,7-tetrahydro-1H-indazole (23). Flash chromatography (2:1, dichloromethane : acetone). $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 2.64 (t, 2H), 2.55 (t, 2H), 2.43 (t, 2H), 1.76 (quint, 4H), 1.61 (m, 2H), 1.31 (m, 4H), 0.88 (t, 3H); $^{13}\text{C NMR}$ (400 MHz, CDCl_3) δ 144.3, 112.3, 31.6, 28.6, 25.8, 23.4, 23.0, 22.42, 22.4, 20.2, 13.9; **MS** calcd for $\text{C}_{12}\text{H}_{21}\text{N}_2$, 193.17; found, 193.67.

tert-butyl 3-(4-methoxyphenyl)-1,4,6,7-tetrahydro-5H-pyrazolo[4,3-c]pyridine-5-carboxylate (24). Flash chromatography (3:2, dichloromethane : acetone). Analytical sample crystallized from 2-propanol/water. $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.46 (d, J = 8.8 Hz, 2H), 6.90 (bd, 2H), 4.62 (s, 2H), 3.81 (s, 3H), 3.70 (b, 2H), 2.73 (t, 2H), 1.48 (s, 9H); $^{13}\text{C NMR}$ (400 MHz, CDCl_3) δ 159.3, 155.0, 127.5, 114.3, 109.8, 80.1, 55.3, 41.6, 28.4; **MS** calcd for $\text{C}_{18}\text{H}_{24}\text{N}_3\text{O}_3$, 330.18; found, 330.17.

3-(4-methoxyphenyl)-1,4,6,7-tetrahydropyrano[4,3-c]pyrazole (25). Flash chromatography (3:2, dichloromethane : acetone). Analytical sample crystallized from dichloromethane/hexanes. $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.39 (d, J = 8.8 Hz, 2H), 6.95 (d, J = 8.8 Hz, 2H), 4.88 (s, 2H), 3.98 (t, 2H), 3.84 (s, 3H), 2.83 (t, 2H); $^{13}\text{C NMR}$ (400

MHz, CDCl₃) δ 159.5, 127.3, 123.9, 114.4, 111.0, 64.5, 64.2, 55.3, 23.7; **MS** calcd for C₁₃H₁₅N₂O, 231.11; found, 230.96.

3-(4-bromophenyl)-5-(5-chloropentyl)-1H-pyrazole

NMR Spectrum: H1 No: 34
Instrument: Varian Inova400
Solvent: CDCl3 Temp: 25.0
SSID: hellerst Date: Feb 17 2006

cyclohexyl fused n-hexyl pyrazole C13
NMR Spectrum: C13 No: 17
Instrument: Varian Inova400
Solvent: CDCl₃ Temp: 25.0
ISID: hellerst Date: Feb 14 2006