Supplementary Materials

for the manuscript

"3D-Pharmacophore Models for Selective A_{2A} and A_{2B} Adenosine Receptor Antagonists "

Jing Wei,[†] Songqing Wang,[†] Shaofen Gao,[‡] Xuedong Dai[§] and Qingzhi Gao^{*,§}

Table of Contents

- 1, Log file and clustering analysis result for HipHop hypotheses from A_{2A} antagonists ----- p 2-5
- 2, Log file and clustering analysis result for HipHop hypotheses from A_{2B} antagonists ----- p 6-9
- 3, Best fit analysis results of Hypo-A_{2A} towards training set and test set compounds ----- p 10-17
- 4, Best fit analysis results of Hypo-A_{2B} towards training set and test set compounds ----- p 18-26

1, Log file and clustering analysis result for HipHop hypotheses from

A_{2A} antagonists:

logfile:

HipHop / Hypothesis Identification

Version: catHypo 4.10 (330) Friday, December 10, 2004 at 01:58 AM

Confs (cpd): /data1/wei/AR/OutHypo-13579-591Dir/

Spreadsheet: /data1/wei/AR/OutHypo-13579-591Dir/Local-5.spst

Dictionary: /data1/wei/AR/OutHypo-13579-591Dir/dict.chm

Host: YaoXueYuan

Outname: OutHypo-13579

Logname: OutHypo-13579-591.log

Statistics: /home/accelrys/410/conf/catalyst/default.hstat

Summary of parameters:

HyposReported	10
Spacing	300
MinPoints	4
MinSubsetPoints	4
SuperpositionError	1
Misses	1
FeatureMisses	1
CompleteMisses	0.0e+00
ToleranceFactor	1
CheckSuperposition	1
WeightVariation	0.3
MappingCoeff	0.0e+00
Mem	60
IdealHBondGeomOnly	0.0e+00
VariableWeight	0.0e+00
VariableTolerance	0.0e+00

Constraints on numbers of features:

Name:	HBA lipid	HYDROPHOBIC	PosIonizable	RING AROMATIC	Total
Min:	1	0	1	1	2
Max:	5	5	5	5	4

Summary of lead molecules selected:

Name	Principal	MaxOmitFeat	Chirality	Confs	Feats
------	-----------	-------------	-----------	-------	-------

JMC45-115-8h	2	0	74	16.61
JMC45-115-8g	2	0	171	20.42
JMC45-115-11e	2	0	167	25.66
JMC45-115-8k	2	0	89	23.06
JMC45-115-14b	2	0	142	28.27

Summary of feature definitions:

H: HBA lipid

#N,O,S#2: Tolerance= 160

AcceptorPt1: Tolerance= 220

Z: HYDROPHOBIC

Hydrophobic: Tolerance= 160

P: PosIonizable

PosIon-pt: Tolerance= 160

R: RING AROMATIC

centroid: Tolerance= 160

normal: Tolerance= 160

Summary of feature definition hit statistics:

HBA lipid hits/lead: mean=10.50 stddev= 4.52

HYDROPHOBIC hits/lead: mean= 3.10 stddev= 0.11

PosIonizable hits/lead: mean= 1.20 stddev= 0.45

RING AROMATIC hits/lead: mean= 8.00 stddev= 0.00

Summary of feature definition hits by molecule:

	H	Z	P	R
JMC45-115-8h	4.61	3.00	1.00	8.00
JMC45-115-8g	8.18	3.24	1.00	8.00
JMC45-115-11e	13.46	3.20	1.00	8.00
JMC45-115-8k	10.03	3.02	2.00	8.00
JMC45-115-14b	16.23	3.04	1.00	8.00

Summary of ranking coefficients:

Constant: -6.6

Dispersion: 0.0054

H: 2.7

Z: 1.9

P: 3.5

R: 2.7

Processing:

Generating hypos with PH: [c]..... 5560

Generating hypos with RH: [c]..... 98058
Generating hypos with RP: [c]..... 5558
Generating hypos with RPH: [c]..... 66206
Hypos with features PH not desired. Freeing memory.
Hypos with features RH not desired. Freeing memory.
Hypos with features RP not desired. Freeing memory.
Generating hypos with RPHH: [c]..... 75968
Reprocessing hypos with RPHH: 10 remaining.
Generating hypos with RPZH: [c]..... 106872
Reprocessing hypos with RPZH: 10 remaining.
Generating hypos with RRPH: [c]..... 99008
Reprocessing hypos with RRPH: 10 remaining.
Reprocessing hypos with RPH: 10 remaining.
Inspecting hypos with RPHH
Inspecting hypos with RPZH
Inspecting hypos with RRPH

Exporting hypotheses:

1 RPHH	Rank: 68.6099	DH: 11111	PH: 00000
2 RPZH	Rank: 67.8995	DH: 11111	PH: 00000
3 RPZH	Rank: 67.8995	DH: 11111	PH: 00000
4 RPHH	Rank: 67.1473	DH: 11111	PH: 00000
5 RPHH	Rank: 67.1004	DH: 11111	PH: 00000
6 RPHH	Rank: 66.999	DH: 111111	PH: 00000
7 RPHH	Rank: 66.9558	DH: 11111	PH: 00000
8 RRPH	Rank: 66.8051	DH: 11111	PH: 00000
9 RRPH	Rank: 66.8051	DH: 11111	PH: 00000
10 RRPH	Rank: 66.7887	DH: 11111	PH: 00000

Halted normally.

CPU time = 168 s System time = 0 s

Clustering analysis:

This is file: hypoClusterResult_24.txt

The following hypotheses were clustered:

OutHypo-13579.01
OutHypo-13579.02
OutHypo-13579.03
OutHypo-13579.04
OutHypo-13579.05
OutHypo-13579.06
OutHypo-13579.07
OutHypo-13579.08

OutHypo-13579.09

OutHypo-13579.10

using the following clustering method:

Hierarchical Average Linkage method.

In the following table(s), the columns indicate the total number of clusters one wants and the rows indicate the input hypotheses.

The entries in the table indicate which cluster a hypothesis belong to. The hypotheses with the same number belong to the same cluster.

Number of Clusters	2	3	4	5	6	7	8	9
OutHypo-13579.01	1	1	1	1	1	1	1	1
OutHypo-13579.07	1	1	1	2	2	2	2	2
OutHypo-13579.06	1	1	2	3	3	3	3	4
OutHypo-13579.04	1	1	2	3	3	3	3	3
OutHypo-13579.05	1	1	2	3	4	4	4	5
OutHypo-13579.02	2	2	3	4	5	5	5	6
OutHypo-13579.03	2	2	3	4	5	5	5	6
OutHypo-13579.08	2	3	4	5	6	6	6	7
OutHypo-13579.10	2	3	4	5	6	6	7	8
OutHypo-13579.09	2	3	4	5	6	7	8	9

2, Log file and clustering analysis result for HipHop hypotheses from

A_{2B} antagonists:

logfile:

HipHop / Hypothesis Identification

Version: catHypo 4.10 (330) Friday, December 10, 2004 at 01:58 AM

Confs (cpd): /data1/wei/AR/OutHypo-5315-285Dir/

Spreadsheet: /data1/wei/AR/OutHypo-5315-285Dir/Local-4.spst

Dictionary: /data1/wei/AR/OutHypo-5315-285Dir/dict.chm

Host: YaoXueYuan

Outname: OutHypo-5315

Logname: OutHypo-5315-285.log

Statistics: /home/accelrys/410/conf/catalyst/default.hstat

Summary of parameters:

HyposReported 10
Spacing 300
MinPoints 4
MinSubsetPoints 4
SuperpositionError 1
Misses 1
FeatureMisses 1
CompleteMisses 0.0e+00
ToleranceFactor 1
CheckSuperposition 1
WeightVariation 0.3
MappingCoeff 0.0e+00
Mem 60
IdealHBondGeomOnly 0.0e+00
VariableWeight 0.0e+00
VariableTolerance 0.0e+00

Constraints on numbers of features:

Name:	HBA lipid	HYDROPHOB	aliphatRING	AROMATIC	Total
Min:	1	1	1		1
Max:	5	5	5		4

Summary of lead molecules selected:

Name	Principal	MaxOmitFeat	Chirality	Confs	Feats
JMC47-1434-70b-test1	2	0		84	17.49

MRS1706-test1	2	0	70	17.01
MRS1754-test1	2	0	144	15.78
MRE2030F20-test1	2	0	127	18.73
MRE2029F20-test1	2	0	115	17.57

Summary of feature definitions:

H: HBA lipid

#N,O,S#2: Tolerance= 160

AcceptorPt1: Tolerance= 220

Z: HYDROPHOB aliphatic

Hydrophobic1: Tolerance= 160

R: RING AROMATIC

centroid: Tolerance= 160

normal: Tolerance= 160

Summary of feature definition hit statistics:

HBA lipid hits/lead: mean= 8.40 stddev= 0.82

HYDROPHOB aliphatic hits/lead: mean= 2.92 stddev= 1.13

RING AROMATIC hits/lead: mean= 6.00 stddev= 0.00

Summary of feature definition hits by molecule:

	H	Z	R
JMC47-1434-70b-test1	7.36	4.13	6.00
MRS1706-test1	9.01	2.00	6.00
MRS1754-test1	7.78	2.00	6.00
MRE2030F20-test1	8.56	4.17	6.00
MRE2029F20-test1	9.29	2.28	6.00

Summary of ranking coefficients:

Constant: -6.6

Dispersion: 0.0054

H: 2.7

Z: 2.7

R: 2.7

Processing:

Generating hypos with ZH: [c]..... 12755

Generating hypos with RH: [c]..... 46812

Generating hypos with RZ: [c]..... 8896

Generating hypos with RZH: [c]..... 120724

Hypos with features ZH not desired. Freeing memory.

Hypos with features RH not desired. Freeing memory.

Hypos with features RZ not desired. Freeing memory.
Generating hypos with RZHH: [c]..... 284148
Reprocessing hypos with RZHH: 10 remaining.
Generating hypos with RZZH: [c]..... 67874
Reprocessing hypos with RZZH: 10 remaining.
Generating hypos with RRZH: [c]..... 159082
Reprocessing hypos with RRZH: 10 remaining.
Reprocessing hypos with RZH: 10 remaining.
Inspecting hypos with RZHH
Inspecting hypos with RZZH

Exporting hypotheses:

1	RZHH	Rank: 85.43	DH: 11111	PH: 00000
2	RZHH	Rank: 85.43	DH: 11111	PH: 00000
3	RZHH	Rank: 82.582	DH: 11111	PH: 00000
4	RZHH	Rank: 81.7254	DH: 11111	PH: 00000
5	RZHH	Rank: 81.6868	DH: 11111	PH: 00000
6	RZHH	Rank: 81.5595	DH: 11111	PH: 00000
7	RZHH	Rank: 81.285	DH: 11111	PH: 00000
8	RZHH	Rank: 80.4375	DH: 11111	PH: 00000
9	RZHH	Rank: 80.4375	DH: 11111	PH: 00000
10	RZHH	Rank: 80.4036	DH: 11111	PH: 00000

Halted normally.

CPU time = 243 s System time = 0 s

Clustering analysis:

This is file: hypoClusterResult_23.txt

The following hypotheses were clustered:

OutHypo-5315.01
OutHypo-5315.02
OutHypo-5315.03
OutHypo-5315.04
OutHypo-5315.05
OutHypo-5315.10
OutHypo-5315.09
OutHypo-5315.08
OutHypo-5315.07
OutHypo-5315.06

using the following clustering method:

Hierarchical Average Linkage method.

In the following table(s), the columns indicate the total number of clusters one wants and the rows indicate the input hypotheses.

The entries in the table indicate which cluster a hypothesis belong to. The hypotheses with the same number belong to the same cluster.

Number of Clusters	2	3	4	5	6	7	8	9
OutHypo-5315.01	1	1	1	1	1	1	1	1
OutHypo-5315.02	1	1	1	1	1	1	1	1
OutHypo-5315.03	1	1	1	1	1	1	2	2
OutHypo-5315.06	1	1	1	1	2	2	3	3
OutHypo-5315.09	1	1	1	2	3	3	4	4
OutHypo-5315.08	1	2	2	3	4	4	5	5
OutHypo-5315.04	2	3	3	4	5	5	6	6
OutHypo-5315.07	2	3	3	4	5	6	7	8
OutHypo-5315.05	2	3	3	4	5	6	7	7
OutHypo-5315.10	2	3	4	5	6	7	8	9

3, Best fit analysis results of Hypo-A_{2A} towards training set and test set compounds:

Figure 8. (a) Best model Hypo-A_{2A} mapped with selective A_{2A} antagonists No.15, 16, 19 in the A_{2A} test set. All hypothesis features are mapping with the corresponding chemical functional groups on the antagonists. (b) Best model Hypo-A_{2A} mapped with selective A_{2B} antagonists No.6, 7, 8 in the A_{2B} test set. All ligands miss mapping at least one hypothesis feature of the positively ionizable sphere. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid feature; orange represents ring aromatic feature; light blue represents hydrophobic feature; red represents positively ionizable feature. Compound numbers and chemical structures as depicted in Figure 2 and 3.

Figure 9. Best model Hypo-A_{2A} mapped with selective A_{2A} antagonists in the training set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic; red represents positively ionizable

region. Compound numbers and chemical structures as depicted in Figure 1 (No1: yellow; No2: green; No3: blue; No4: red; No5: pink blue).

Figure A. Best model Hypo-A_{2A} mapped with selective A_{2A} antagonists No.1, 2, 5 in the training set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic; red represents positively ionizable region. Compound numbers and chemical structures as depicted in Figure 1.

Figure B. Best model Hypo-A_{2A} mapped with selective A_{2A} antagonists No.15, 16, 18, 19 in the A_{2A} test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic; red represents positively ionizable region. Compound numbers and chemical structures as depicted in Figure 1.

Figure C. Best model Hypo-A_{2A} mapped with selective A_{2A} antagonists No.15, 16, 19 in the A_{2A} test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic; red represents positively ionizable region. Compound numbers and chemical structures as depicted in Figure 1.

Figure D. Best model Hypo-A_{2A} mapped with selective A_{2A} antagonists No.16, 19 in the A_{2A} test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic; red represents positively ionizable region. Compound numbers and chemical structures as depicted in Figure 1.

Figure E. Best model Hypo-A_{2A} mapped with selective A_{2B} antagonists No.6, 7, 8 in the A_{2B} test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic; red represents positively ionizable region. Compound numbers and chemical structures as depicted in Figure 2.

Figure F. Best model Hypo-A_{2A} mapped with selective A_{2B} antagonist No.29 in the A1 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic; red represents positively ionizable region. Compound numbers and chemical structures as depicted in Figure 5.

Figure G. Best model Hypo-A_{2A} mapped with selective A_{2B} antagonists No.32.33 in the A1 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic; red represents positively ionizable region. Compound numbers and chemical structures as depicted in Figure 5.

Figure H. Best model Hypo-A_{2A} mapped with selective A_{2B} antagonist No.33 in the A1 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic; red represents positively ionizable region. Compound numbers and chemical structures as depicted in Figure 5.

Figure I. Best model Hypo-A_{2A} mapped with selective A_{2B} antagonists No.36, 37, 38 in the A3 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic; red represents positively ionizable region. Compound numbers and chemical structures as depicted in Figure 6.

Figure J. Best model Hypo-A_{2A} mapped with selective A_{2B} antagonist No.36 in the A3 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic; red represents positively ionizable region. Compound numbers and chemical structures as depicted in Figure 6.

Figure K. Best model Hypo-A_{2A} mapped with selective A_{2B} antagonist No.37 in the A3 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic; red represents positively ionizable region. Compound numbers and chemical structures as depicted in Figure 6.

Figure L. Best model Hypo-A_{2A} mapped with selective A_{2B} antagonist No.38 in the A3 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic; red represents positively ionizable region. Compound numbers and chemical structures as depicted in Figure 6.

4, Best fit analysis results of Hypo-A_{2B} towards training set and test set compounds:

Figure 12. (a) Best model Hypo-A_{2B} mapped with selective A_{2B} antagonists No.20, 22, 25 in the A_{2B} test set. All hypothesis features are mapping with the corresponding chemical functional groups on the antagonists. (b) Best model Hypo-A_{2B} mapped with selective A_{2A} antagonists No.1, 2, 3 in the A_{2A} test set. All ligands miss mapping at least one hypothesis feature of the hydrogen bond acceptor lipid sphere. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid feature; orange represents ring aromatic feature; light blue represents hydrophobic feature; red represents positively ionizable feature. Compound numbers and chemical structures as depicted in Figure 1 and 4.

Figure 11. Best model Hypo-A_{2B} mapped with selective A_{2B} antagonists in the training set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 2 (No6, green; No7, yellow; No8, blue; No9, red; No10, light blue).

Figure M. Best model Hypo-A_{2B} mapped with selective A_{2B} antagonists No.6, 7, 8 in the training set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 2.

Figure N. Best model Hypo-A_{2B} mapped with selective A_{2A} antagonist No.2 in the A_{2A} test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 1.

Figure O. Best model Hypo-A_{2B} mapped with selective A_{2A} antagonist No.1, 4 in the A_{2A} test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 1.

Figure P. Best model Hypo-A_{2B} mapped with selective A_{2A} antagonist No.28 in the A_{2B} test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 1.

Figure Q. Best model Hypo-A_{2B} mapped with selective A1 antagonists No.29, 30, 31, 32 in the A1 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 5.

Figure R. Best model Hypo-A_{2B} mapped with selective A1 antagonists No.29, 32 in the A1 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 5.

Figure S. Best model Hypo-A_{2B} mapped with selective A1 antagonist No.29 in the A1 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 5.

Figure T. Best model Hypo-A_{2B} mapped with selective A1 antagonists No.30, 31 in the A1 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 5.

Figure U. Best model Hypo-A_{2B} mapped with selective A1 antagonists No.32 in the A1 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 5.

Figure V. Best model Hypo-A_{2B} mapped with selective A3 antagonists No.34 in the A3 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 6.

Figure W. Best model Hypo-A_{2B} mapped with selective A3 antagonists No.35, 36 in the A3 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 6.

Figure X. Best model Hypo-A_{2B} mapped with selective A3 antagonists No.35 in the A3 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 6.

Figure Y. Best model Hypo-A_{2B} mapped with selective A3 antagonists No.37, 38 in the A3 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 6.

Figure Z. Best model Hypo-A_{2B} mapped with selective A3 antagonist No.37 in the A3 test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 6.

Figure α . Best model Hypo-A_{2B} mapped with selective A₃ antagonists No.38 in the A₃ test set. Pharmacophore features are color-coded: green represents hydrogen bond acceptor lipid; orange represents ring aromatic; light blue represents hydrophobic. Compound numbers and chemical structures as depicted in Figure 6.