This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

8000 years of North Atlantic storminess reconstructed from a Scottish peat record: implications for Holocene atmospheric circulation patterns in Western Europe

PLEASE CITE THE PUBLISHED VERSION

<https://doi.org/10.1002/jqs.2983>

PUBLISHER

© Wiley-Blackwell

VERSION

AM (Accepted Manuscript)

PUBLISHER STATEMENT

This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at:
<https://creativecommons.org/licenses/by-nc-nd/4.0/>

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Stewart, Helena, Tom Bradwell, Joanna E. Bullard, S.J. Davies, Nicholas R. Golledge, and Robert McCulloch. 2019. "8000 Years of North Atlantic Storminess Reconstructed from a Scottish Peat Record: Implications for Holocene Atmospheric Circulation Patterns in Western Europe". figshare. <https://hdl.handle.net/2134/25983>.

**8000 years of North Atlantic storminess reconstructed from a Scottish peat record:
implications for Holocene atmospheric circulation patterns in Western Europe**

H. Stewart,¹ T. Bradwell,¹ J. Bullard,² S.J. Davies,³ N. Golledge⁴ and R.D. McCulloch^{1*}

¹ Biological & Environmental Sciences, University of Stirling, Stirling, FK9 4LA, Scotland, UK.

² Department of Geography, Loughborough University, Loughborough, LE11 3TU, England, UK.

³ Department of Geography and Earth Sciences, Aberystwyth University, SY23 3DB, Wales, UK.

⁴ Antarctic Research Centre, Victoria University of Wellington, Wellington 6140, New Zealand.

*Corresponding author: robert.mcculloch@stir.ac.uk

ABSTRACT: North Atlantic storminess can affect human settlements, infrastructure and transport links, all of which strongly impact local, national and global economies. An increase in storm frequency and intensity is predicted over the Northeast Atlantic in the 21st century because of a northward shift in storm tracks and a persistently positive North Atlantic Oscillation (NAO), driven by recent atmospheric warming. Although documentary records of North Atlantic storminess exist, these are generally limited to the last c. 1000-2000 years. This paper presents a continuous high-resolution proxy record of storminess spanning the last 8000 years from a 6 m long core taken from a peat bog in Northern Scotland. Bromine concentrations in the peat, derived from sea spray, are used to reconstruct storm frequency and storm intensity, and mire surface wetness is used as an

indicator of longer-term climate shifts. The results suggest a relationship between positive phases of the NAO and increased North Atlantic storminess. However, subtle differences between bromine concentrations and mire surface wetness suggest that high intensity but perhaps less frequent periods of storminess are not necessarily associated with a wetter climate.

KEYWORDS: Holocene storminess; NAO; micro-XRF; mire surface wetness; Scotland.

Introduction

The location and intensity of mid-latitude storm tracks strongly influence the climate of Europe (Hanna *et al.*, 2008). The most intense and damaging storms affecting Europe originate in the North Atlantic, often causing extensive flooding and damage to infrastructure, and resulting in significant detrimental economic impacts. The highest magnitude storms occur most frequently during winter, when the storm tracks are most intense, and extend in a north-westerly direction from the east coast of North America, to Ireland, Great Britain and Norway (Cheng *et al.*, 2011). North Atlantic storminess has increased over recent decades in association with warming air temperatures over the same period (Alexander and Tett, 2005, Allan *et al.*, 2009, Wang *et al.*, 2009). Predictions suggest that over the next 100 years North Atlantic storm tracks will shift northwards and storm frequency will increase in the British Isles due to an intensified jet stream (Pinto *et al.*, 2009; Orme *et al.*, 2015).

Records of past storminess have been reconstructed through both observational and sedimentary (or palaeoenvironmental) records. Observational records tend to span the past few hundred years and are based on air temperature (Dawson *et al.*, 2003), sea surface temperature (Hurrell, 1995) and wind speed (Clarke and Rendell, 2009; Dawson *et al.*, 2010). In Europe and the North East Atlantic, proxy measures for increased wind strength include aeolian sand influx (de la Vega-Leinert *et al.*, 2000; Clarke *et al.*, 2002; Sommerville *et al.*, 2003; de Jong *et al.*, 2006; Clarke and Rendell, 2009; Tisdall *et al.*, 2013), over-wash deposits in coastal lagoons (Sabatier *et al.*, 2012), cliff-top storm deposits left by extreme waves (Hansom and Hall, 2009), marine records reflecting wind-blown current strength and storm deposits (Hass, 1996; Andresen *et al.*, 2005). and Na⁺ from the Greenland ice cores (Dawson *et al.*, 2003). Paleoenvironmental reconstructions of North Atlantic storminess tend to span from the mid-Holocene to the present and so high-resolution records on a millennial timescale (10³ a) are an important goal in order to gauge longer-term trends and better understand Holocene climate variability.

The position and strength of the polar front jet stream in the Northern Hemisphere strongly determines the process of cyclogenesis between ~40 and 65°N, and, therefore, the number and frequency of high-energy storms in the North Atlantic region (Fig. 1). The development of a vigorous jet stream in winter, enabled by strong temperature contrasts between mild moist mid-latitude and cold polar air, accounts for the strength and trajectory of dominant westerly winds as well as the frequency of extra-tropical storms and their tracks over north-western Europe (Hurrell 1995, Hurrell *et al.*, 2003; Athanasiadis *et al.*, 2010). Secular changes in the North Atlantic Oscillation (NAO), defined as the sea-level pressure difference between the main North Atlantic pressure dipole measured in Iceland and the Azores, are strongly

associated with changes in the polar-front jet stream (Hurrell 1995; Woollings *et al.*, 2008, 2010). An enhanced NAO (positive) mode typically results from undisturbed strong zonal (east to west) jet stream flow between 50 and 60°N; whilst a negative NAO mode results from disturbed, meridional flow with a large north-south component leading to blocking highs and cut-low pressure systems. During a negative NAO, the dominant westerly winds often follow a more southern trajectory owing to the development of a large quasi-stationary high-pressure system over Greenland (Woollings *et al.*, 2008, 2010). Both strongly positive and negative NAO modes represent end members and require relatively vigorous jet-stream-driven atmospheric circulation in the North Atlantic sector. These conditions are normally optimised during European winter, hence the strong positive correlation between NAO mode and winter precipitation in NW Britain and western Norway (Hurrell, 1995; Hurrell *et al.*, 2003). However, other more subdued long-term synoptic situations can occur. These include a neutral NAO phase, where the pressure index is neither strongly positive nor strongly negative, normally associated with a weakening of the main Iceland cyclonic and Azores anticyclonic pressure systems. In this situation, the development of large persistent Scandinavian high-pressure systems can block the path of westerly winds into NW Europe, further reducing cyclonic activity and dampening the NAO index into a more neutral state (Mauri *et al.*, 2014).

This paper presents chemical variability and palaeo-moisture indices from a six metre-long peat core in maritime Northern Scotland. Analyses of down-core variations in bromine concentrations as an indicator of storm intensity are coupled with mire surface wetness (MSW) as an indicator of longer-term storm-track position and climate shifts in the

Northeast Atlantic. Finally, we compare our record to other proxies for shifts in the NAO during the Holocene.

Bromine as an indicator of storminess

The generation of sea-salt aerosol is the principal global source of atmospheric Br, producing ~6.2 Tg/a (Sanders *et al.*, 2003). The bursting of air bubbles and the direct formation of droplets by wave crests injects sea-salt aerosol into the atmosphere and so the flux of Br is dependent on wind speed (Moldanová and Ljungström, 2001). Marine aerosols may be transported long distances (10s-100s km) and be dry deposited or scavenged from the atmosphere by rainfall (Gustafsson and Franzén, 2000). Other sources of inorganic Br are dust, biomass-burning and fossil fuels but these are of an order of magnitude less than marine sources. Crustal sources are estimated to be 4% of the global Br flux and its deposition is geographically restricted to the equatorial Atlantic and North Pacific Oceans (Sanders *et al.*, 2003). Ombrotrophic peat bogs receive only atmospheric inputs and so provide excellent archives of climatic change. It has been argued that Br concentrations are determined by climate-controlled biogenic processes (Biester *et al.*, 2004; Moreno *et al.*, 2015) and are not stratigraphically retained. However, other studies have demonstrated that Br is stably retained in the humic acid content of peat and down-core variations are conserved (Zaccone *et al.*, 2008; Orme *et al.*, 2015; Turner *et al.*, 2015). Br concentrations in peat bogs around the maritime fringes of the North Atlantic, such as in the British Isles, are very likely to be derived from sea spray; hence higher Br levels suggest an increase in sea turbulence, accentuated wave action, and increased windiness during storms. Mire surface wetness (MSW), an additional measure, indicates longer-term changes in precipitation

patterns with increased MSW signalling a more persistent wetter climate (Charman *et al.*, 2006; Turner *et al.*, 2014).

Materials and methods

Site description and field sampling

A continuous high-resolution peat core (to 6.08 m depth) was obtained from the central part of a large mire at Shebster, northern Caithness (58°33'06.6" N, 003°42'39.0" W; 82 m asl; 4.8km from the North Atlantic coast) (Fig. 2). The mire surface is bordered by the Burn of Shebster that drains northwards. The bedrock underlying the site is Middle Devonian sandstone of the Bighouse Formation (Auton *et al.*, 2005). The core was sampled from within the deepest part of the Shebster peat bog using a 1-metre long (75 mm diameter) Russian D-section sampler (with 10 cm overlapping sections). Recovered cores were transferred into plastic guttering, sealed in lay-flat tubing, and stored at the University of Stirling at a constant temperature of 4°C.

Minerogenic analysis

The core was sub-sampled in contiguous 2cm³ sections for acid digestion to remove organic matter (c.f. Dugmore *et al.*, 1995a). The mineral residue was then scanned using light microscopy to identify tephra glass shards to supplement the radiocarbon chronology with tephrochronology.

Organic content

To provide a record of bio-productivity and organic content, contiguous samples of 2 cm depth were combusted in a muffle furnace at 550°C for 4 hours to enable the calculation of the percentage loss-on-ignition (LOI₅₅₀).

Mire surface wetness

The degree of peat humification, as a proxy of MSW, was estimated using the colorimetric alkali extract method modified from Blackford (1993). Under drier conditions peat is more oxidised, the accumulation rate is slower and there is an increase in humic material. The greater the humic content the darker the extract solution and the lower the transmitted light values. Therefore, lower percentage transmission values indicate drier peat accumulation conditions whereas higher percentage transmission values indicate wetter conditions. Contiguous sub-samples of 2 cm³ were taken from the 608 cm core. Samples were oven dried at 80°C for 24 hrs and then ground using a small rotating blade grinder. Sub-samples of 0.2 g were placed into 50 ml falcon centrifuge tubes and 50 ml of NaOH 8% w/v was added to each sample and the tubes placed in a boiling water bath for 60 minutes and intermittently stirred. The samples were then centrifuged at 3000 rpm for 5 minutes and a 0.5 ml aliquot pipetted into a 10 mm cuvette and 2.5 ml of distilled water added. The cuvettes were analysed in a Thermo Scientific Genesys 20 spectrophotometer and the percentage transmitted light measured at 540 nm. This method allowed for batches of 20 samples to be analysed in under 30 min. from removal from the water bath and so minimise any fading of the solution (cf. Blackford, 1993).

Micro X-Ray Fluorescence Geochemistry

The core was analysed using an Itrax® X-Ray Fluorescence core scanner at Aberystwyth University. Non-destructive elemental analysis, including Br was performed at 2mm intervals using a molybdenum (Mo) anode X-ray tube (settings: 30 kV, 50 mA, count time 10 seconds). The results are expressed as a ratio of the coherent+incoherent (coh/incoh) values, which are used as proxies for organic and moisture content. The coh/incoh ratio minimises any influence from these proxies on the geochemical data. Density and colour information was further obtained using X-radiography and digital RGB optical imagery.

Chronology

Five samples of wood material were AMS radiocarbon dated to enable the construction of a Bayesian age-depth model. All samples were pre-treated with an acid/alkali/acid treatment. To augment the radiocarbon chronology, a cryptotephra layer (346 cm depth) was analysed through the geochemical fingerprinting of individual shards by the SX100 Cameca Electron microprobe at the University of Edinburgh using methods established by Hunt and Hill (1993) and Hayward (2013).

The cryptotephra at 346 cm is geochemically correlated to the eruption of Hekla 4 (TephraBase; Newton, 1996) and provides an isochronic marker dated to 3826 ± 12 ¹⁴C a BP (Dugmore *et al.* 1995b). The AMS radiocarbon and tephra ages were calibrated using Calib ver.7.10 (Stuiver and Reimer, 1993) and IntCal13 (Reimer *et al.*, 2013)(Table 1). The Bayesian program, BACON v2.2 (Blaauw and Christen, 2011) was used to construct an age-depth model to constrain the stratigraphic results (Fig. 3). The age-depth model indicates a mean accumulation rate of 13 yrs/cm and a uniform rate of peat accumulation during the Holocene.

186

187 **Results**

188 From 612 to 608 cm depth greyish-brown organic mud grades to a dark brown moderately
189 humified peat (with wood fragments at 404 – 406 cm depth) that continues to ~370 cm.
190 From ~ 370 to 271 cm, the core comprises very dark brown well-humified amorphous peat.
191 At 271 cm, there is a marked transition to a dark brown peat with abundant sedge
192 fragments. The sedge-rich peat continues to 27 cm and is moderately well humified and
193 with occasional wood fragments. From 27 cm to the surface root mat, the core comprises a
194 dark brown fibrous sphagnum peat.

195

196 To aid the interpretation, the Shebster stratigraphic data is divided into 8 major zones based
197 on major changes in the MSW as this is a site-specific proxy (Fig. 3).

198

199 SH-1 8210 – 7400 cal a BP: MSW was not measured in the lowest ~10 cm of the peat to
200 avoid any potential influence from the underlying lake sediments; measurements
201 commenced at 590 cm. In Zone SH-1 the MSW curve is characterised by two peaks at c.
202 7960 and 7610 cal a BP, separated by a nadir at c. 7740 cal a BP. The Br ratios appear to be
203 in antiphase with the largest peak of the entire record occurring at c. 7800 cal a BP, before
204 both MSW and bromine decline to a low at the upper zone boundary at c. 7400 cal a BP.

205

206 SH-2 7400 - 5270 cal a BP: At the start of this zone there is a marked step up to higher MSW
207 values that fluctuate (between ~20 and 30 %T) through the first half of the zone. Br values
208 commence from a peak between c. 7400 and 7300 cal a BP and then decline to a broad low

209 also during the first half of the zone, between c. 7300 and 6200 cal a BP. At c. 6145 cal a BP
210 there is a rapid increase in Br to a sustained peak until a fall at c. 5700 to lower but
211 fluctuating values and then increasing towards the upper zone boundary. The MSW values
212 in the second half of the zone also continue to fluctuate (between ~25 and 35 %T) but at
213 higher values than previously.

214

215 SH-3 5200 - 4000 cal a BP: Br increase to a peak between c. 5200 and 4900 cal a BP followed
216 by a sustained declining trend towards the upper zone boundary at c. 4000 cal a BP. The
217 decline in Br is reflected in a similar profile in the MSW, excepting a brief minor peak in
218 MSW at c. 4540 cal a BP.

219

220 SH-4 4000 - 3300 cal a BP: The Br ratios reach their minima at c. 3850 cal a BP and remain
221 relatively stable before gradually increasing from c. 3400 cal a BP to a peak at the upper
222 zone boundary. During this zone the stratigraphy changes to well-humified peat and this is
223 reflected in the sharp fall of MSW values where they reach their lowest values of the entire
224 record before rising again at the top of the zone.

225

226 SH-5 3300 – 2400 cal a BP: MSW values rise to a peak at c. 3160 cal a BP (25.6 %T) and
227 continue to rise to ~30%, punctuated by a brief decline to 17.5 %T at c. 3005 cal a BP. Br
228 values remain relatively stable, with small fluctuations occurring throughout this period.

229

230 SH-6 2400 - 1400 cal a BP: MSW values fluctuate between ~33 and 20 %T during this zone
231 and Br values continue to be relatively stable.

232

SH-7 1400 – 600 cal a BP: Br ratios peak between c. 1225 and 1130 cal a BP and between c. 855 and 750 cal a BP before declining to a low at c. 600 cal a BP. MSW values increase to a peak at c. 1365 cal a BP (39.1 %T) then decline to a low at c. 1115 cal a BP (18.0 %T) followed by peaks at c. 950 (34.1 %T) and 675 cal a BP (34.8 %T).

SH-8 600-0 cal a BP: MSW values decline to a low at c. 300 cal a BP (18.8 %T) before rising to a peak of 43.3 %T at c. 140 cal a BP. In contrast, Br values rise rapidly to a sustained peak between c. 380 and 200 cal a BP.

Discussion

The down-core variations in Br and MSW indicate that changes in Br concentrations likely reflect longer-term changes in storminess in the NE Atlantic rather than biogenic processes within the peat bog. Therefore, our ~8000 year palaeo-wetness and storminess record from Shebster, Northern Scotland, can be interpreted alongside other key proxy records from around the North Atlantic to place the inferred palaeoenvironmental trends in a wider context of Holocene atmospheric circulation changes. These proxy records include: a 5000 year glacier record from Folgefanna in Southern Norway (Bakke *et al.*, 2008); a reconstructed 5000 year NAO index based on a lake-sediment core in SW Greenland (Olsen *et al.*, 2012); a Scottish speleothem record (Baker *et al.*, 2015); and an Iberian speleothem record (Walkzak *et al.*, 2015) (Fig. 4). Variations in speleothem laminae thickness provide an annual growth-rate record that can be used as a proxy for past climate and environmental change. Growth rates are determined by changes in precipitation and higher growth rates suggest warmer and wetter conditions (positive NAO state) whereas low growth rates are

associated with colder and drier conditions (negative NAO state) (Baker *et al.*, 2015). The NAO index based on a south-west Greenland lake-sediment core is reconstructed from deep-water anoxia data (Olsen *et al.*, 2012). A negative NAO is associated with above average temperatures and below average winter precipitation, leading to earlier ice melt and allows stronger vertical mixing of the water column. This weakens thermal stratification and hence increases oxygen transfer into the deep-water zone, with associated implications for redox processes. A positive NAO, associated with colder conditions, leads to later ice melt, which coincides with maximum solar radiation and results in limited water column mixing. This leads to the rapid onset of thermal stratification, and hence maintenance of hypoxic conditions. Deep-water anoxia can alter cycling of redox-sensitive elements. The Mn/Fe ratio reflects the strength of seasonal thermal stratification and is therefore a proxy for dominant NAO circulation patterns. A higher Mn/Fe ratio and carbonate concentration reflects predominantly weaker stratification and is associated with negative NAO conditions, whereas a lower Mn/Fe ratio and carbonate concentration reflects stronger stratification and is associated with positive NAO conditions.

SH-1 8210 – 7400 cal a BP: To avoid the influence of any mineral material from the underlying lacustrine sediments the degree of peat humification was not analysed at the base of the peat (608 cm). The lowest part of the available record is characterised by several high magnitude fluctuations in wetness and storminess. Although the base of the analysed peat record starts at c. 8200 cal a BP, neither the Br record nor the MSW data completely capture the 8.2ka event, expressed as a marked rapid negative temperature anomaly across much of the Northern Hemisphere (Larsen *et al.*, 2012; Tipping *et al.*, 2012). However, SH-1 is dominated by a large Br positive anomaly centred at c. 7800 cal a BP, suggesting a period of enhanced

storminess lasting c. 300-400 years. By contrast, the humification record at this time suggests conditions of below-average wetness. This anti-phase relationship between MSW and Br appears paradoxical, but may suggest that these storms were cold, moisture-starved Polar vortex systems rather than moisture-bearing westerly winds. This is supported by the Iberian speleothem record, which shows a stable, positive record indicating warmer and wetter conditions. Rainfall in southern Iberia was more evenly distributed throughout the year, typical of a more temperate climate lacking a clear dry season (Walczak *et al.*, 2015) and may indicate a more southerly position of the jet stream and, therefore, a persistent negative NAO phase (phase B1, Fig. 1).

SH-2 7400-5270 cal a BP: After c. 7400 cal a BP there was a marked increase in MSW at Shebster and although the values fluctuate during the period between c. 7400 and 6200 cal a BP, they remain high suggesting the persistence of wetter conditions. However, Br values decrease from the initial peak in zone SH-1 which may suggest that although this zone reflects wetter conditions storminess was less intense than in SH-1. This period probably indicates a northerly migration of the jet-stream-driven westerly winds accompanied by a movement of the North Atlantic storm tracks to a position located over Northern Scotland. This atmospheric circulation pattern is best described by the positive mode of the NAO (phase A, Fig. 1).

In the second half of SH-2, from c. 6200 to 5270 cal a BP, MSW continues to reflect increased and more sustained wetter conditions. At the same time, Br values remain high indicating higher levels of storminess throughout this period. A pronounced peak in storminess occurs in zone SH-2 between c. 6145 and 5700 cal a BP. It is not certain what might have caused this ~400-year window of increased storminess, during a period of relatively wetter but more

stable climatic conditions. The Iberian speleothem record is characterised by a decline in growth rates between 6100 and 5300 cal a BP suggesting decreased moisture availability (Walczak *et al.*, 2015) and this is consistent with a stronger polar-front jet stream bringing an increase in the number and intensity of storms tracking over Northern Scotland (i.e. a strongly positive NAO mode).

SH-3 5270-4000 cal a BP: Between c. 5300 and 4000 cal a BP, MSW and Br values reflect a synchronised period of gradual and near-continuous decline in both precipitation and storminess over Northern Scotland. Several proxy records reflect a mixed climate signal at this time. The glacier-ELA-reconstructed winter precipitation record from Bakke *et al.* (2005) indicates a comparable near-continuous decrease in wetness over most of this period (c. 5000-4000 cal a BP) and is consistent with the Iberian speleothem record which also shows a shift to drier conditions with the exception of an increase in wetness at c. 4200 cal a BP. This climate period is also captured by the earliest part of the SW Greenland lake-sediment-reconstructed NAO index, which shows a sustained positive NAO phase between c. 5200 and 4400 cal a BP (Olsen *et al.*, 2012). These climatic conditions are all compatible with a period of geographically unstable jet stream position and/or declining jet stream strength bringing generally warmer, drier summer conditions to NW Europe accompanied by a decline in cyclogenesis with fewer storms tracking across northern Scotland. After c. 4400 cal a BP, the reconstructed NAO index (Olsen *et al.*, 2012) enters a relatively neutral phase consistent with a decrease in jet stream vigour, at a time when the Shebster peat record indicates steadily decreasing North Atlantic storm activity.

SH-4 4000-3300 cal a BP: This period marks the most striking departure in the Shebster peat record when MSW and Br values are at their lowest for the entire ~ 8000-year record. We relate these values to a sustained period of relatively drier climate and greatly reduced storminess following on from the decline in storminess seen in the preceding millennium (c. 5300-4000 cal a BP; SH-3). A marked decrease in North Atlantic storminess in Northern Scotland could be associated with two different atmospheric circulation scenarios: (1) westerly wind and storm-track migration to a more southerly latitude (ca. 40°N) equating to a strongly negative winter NAO phase; (2) reduced jet stream strength and a low-value or neutral NAO phase. This period of unusual drier and calmer climate identified in the Shebster peat record is not restricted to Caithness, but is probably the expression of a pan-European/North Atlantic event seen widely in other Holocene palaeo-climate proxies. Peat surface-wetness records from a composite of 12 sites in Northern Britain (Charman *et al.*, 2012) show a period of considerably decreased wetness from c. 3900 to 3400 cal a BP, the most pronounced in the mid to Late Holocene (interrupted by a brief increase in wetness at c. 3750 cal a BP). In southern Europe, often in antiphase with the climate of northern Britain, Mediterranean records from south-eastern Italy to south-western France record a period of relatively drier conditions between c. 4000 and 3400 cal a BP (e.g. Di Rita and Magri, 2009; Genty *et al.*, 2006; Walczak *et al.*, 2015), and this can also be seen in the Iberian speleothem record. Further afield, in continental North America and Mexico declining monsoon strength are recorded in a number of geographically diverse proxies from c. 4200 to 3500 cal a BP (Booth *et al.*, 2005; Metcalfe *et al.*, 2015). This is also coincident with the marked southerly migration of the ITCZ (c. 4000 cal a BP) seen in a number of low-latitude records, including the high-resolution Cariaco Basin (Haug *et al.*, 2001; Metcalfe *et al.*, 2015). Closer to Scotland, the same SH-3 time interval (c. 4000-3300 cal a BP) sees lower than present precipitation in

the Norwegian glacier record (Bakke *et al.*, 2005); whilst temperatures in south-west Ireland inferred from speleothems show a broad thermal minimum c. 3800 to 3400 cal a BP (McDermott *et al.*, 2001). It is notable that this period is also characterised by a marked negative departure in chlorine in the GISP2 record, inferred as a weakening of the Icelandic Low pressure system between c. 4500 and 3600 cal a BP (O'Brien *et al.*, 1995; Mayewski *et al.*, 2004).

The collected multi-proxy evidence from both sides of the North Atlantic points towards jet stream weakening during SH-4, leading to a decrease in cyclonic activity which reaches a minimum at c. 3800 cal a BP. Negative NAO conditions normally result in increased rainfall and storminess over southern Europe (e.g. Hurrell *et al.*, 2003; Pinto *et al.*, 2009), something that is not seen in most of the proxy records between c. 4000 and 3300 cal a BP (see above). Secondly, negative NAO phases have been strongly linked with meridional airflow and strong temperature contrasts causing enhanced but intermittent cyclogenesis in north-west Europe (Trouet *et al.*, 2012; Vliet-Lanoë *et al.*, 2014). Again, this is not evident in the Shebster record during SH-4, with storm-driven Br values reaching their 8000-year minimum within this time window. However, these observations, in combination with a prolonged neutral phase of the reconstructed NAO index (Olsen *et al.*, 2012), are entirely consistent with a decrease in jet stream vigour during SH-4. We would expect that this period experienced considerably reduced westerly (zonal) airflow at 50-60°N, accompanied by a higher incidence of quasi-stationary high-pressure systems over Northern Europe. There is evidence from the Shebster Br and MSW data that this c. 700-year period of subdued westerly winds (reduced jet stream vigour) and cyclogenesis over northern Scotland is the ultimate expression of a declining trend in storminess and wetness that started at c. 4500 cal a BP (in SH-3), coincident with the switch

from positive to low-value or neutral NAO values (<1.0) in the Greenland sediment record (Olsen *et al.*, 2012).

SH-5 3300 – 2400 cal a BP: A marked broadly synchronous increase in MSW and Br values at c. 3200-3300 cal a BP indicates the return to wet and stormy conditions in northern Scotland. This period (SH-5) is characterised by generally increasing MSW levels throughout (c. 3300-2400 cal a BP) and a relatively high but fluctuating storminess index. Supporting proxy data suggest more vigorous cyclogenesis, increased precipitation and raised water tables in north-west Europe at this time (Hughes *et al.*, 2000; Charman, 2010; Swindles *et al.*, 2007; Oldfield *et al.*, 2010), although the reconstructed NAO index displays a strong fluctuation from initially positive (c. 3300-3000 cal BP) to strongly negative values (c. 3000-2400 cal a BP) (Olsen *et al.*, 2010). A marked concomitant rise in air temperatures and winter precipitation, seen in the Irish speleothem, Norwegian glacier and Iberian speleothem proxy-records (Fig 4) between c. 3300 and 2700 cal a BP would also suggest a return to more dynamic atmospheric circulation patterns over north-west Europe with strongly zonal moisture-bearing winds and more moderate levels of cyclogenesis.

The most sustained period of negative NAO in the Greenland lake-sediment record is synchronous with an increase in MSW at Shebster (c. 2800-2400 cal a BP). This probably relates to a strengthening of the westerly winds (after the quiescent SH-4 phase) and a mean storm track positioned to the south of Scotland, consistent with the relative decrease in storminess at this time. The annually resolved north-west Scotland speleothem record also starts during this time (Baker *et al.*, 2015). Although no overall trend in the composite speleothem climate-index is apparent during the first ~500 years, relatively high-magnitude

peaks in speleothem growth rates at c. 2900 and 2600 cal a BP probably reflect decadal to centennial periods of higher precipitation in northern Scotland (Baker *et al.*, 2015).

At c. 3000 cal a BP glaciers become permanently established at some marginal sites in Iceland, Norway and southern Greenland for the first time since their complete disappearance in the early Holocene (c. 8000-7000 cal a BP) (Andresen & Bjork, 2005; Balascio *et al.*, 2015; Larsen *et al.*, 2012). Numerous other studies have linked this renewed ice growth, or Neoglaciation, with a shift towards wetter and/or cooler climate in Northern Hemisphere higher latitudes after c. 4200 cal a BP (Blaauw *et al.*, 2004; Swindles *et al.*, 2007; Wang *et al.*, 2012).

SH-6 2400-1400 cal a BP: During this period MSW at Shebster remains relatively high whilst Br levels are somewhat subdued (close to, but slightly below the 8000-year mean), continuing the long-term trend established in SH-5. The north-west Scotland speleothem record exhibits high growth rates suggesting moist but more stable climatic conditions. Elsewhere around Europe this millennium is synonymous with the 'Roman Warm Period' (2500-1600 cal a BP) (Wang *et al.*, 2012) and is characterised by a predominantly positive (>1.0) NAO index in Greenland (Olsen *et al.*, 2012). This strong pressure dipole, but relatively stable low-storm index state, suggests a poleward shift of the westerly storm tracks to a position between Iceland and Scotland, as seen during positive NAO summers. However, the low storm index suggests a more complex relationship, possibly with an increased polar front latitudinal range in winter with storms tracking to the south of northern Scotland. Support for this hypothesis comes from a peat record from Cors Fochno, mid Wales where sustained higher Br values between c. 2200 and 1600 cal a BP suggest that although the

422 Roman Warm period was comparatively dry at Shebster in Northern Scotland, North
423 Atlantic storms still tracked across central and southern Britain with relatively high
424 frequency (Orme *et al.*, 2015).

425

426 SH-7 1400 – 600 cal a BP: This period includes the Medieval Climate Anomaly (MCA: 700-1100
427 cal a BP; Mann and Jones, 2003) and is characterised at Shebster by higher but variable MSW
428 alongside relatively higher and variable Br levels in the peat record. Together they suggest a
429 wetter and stormier period in Northern Scotland than the previous millennium (SH-6) which
430 is consistent with the unusually long and unbroken, strongly positive, NAO phase (Trouet *et*
431 *al.*, 2009). This is seen in the Greenland lake-sediment record from c. 1400 to 600 cal a BP
432 (Olsen *et al.*, 2012). A positive NAO mode is normally associated with a vigorous jet stream
433 and a North Atlantic winter storm track focused between 55-60°N (at the latitude of northern
434 mainland Scotland) (Hurrell *et al.*, 2003; Woollings *et al.*, 2008, 2010). This circulation pattern
435 is supported by several other palaeoenvironmental proxies from the British Isles and adjacent
436 areas. Firstly, the composite British peat-surface wetness record compiled by Charman (2010)
437 shows a continuous phase of elevated water tables spanning the entire 800-year period with
438 a peak c. 1100-1200 cal a BP. Secondly, the Irish speleothem record shows several centuries
439 of increasing above-average (inferred) temperatures, with a peak c. 700-900 cal a BP
440 (McDermott *et al.*, 2001). Thirdly, the winter moisture index from Norwegian glaciers shows
441 well above-average precipitation (120-140% present day) in this time interval (Bakke *et al.*,
442 2008). Fourthly, the occurrence of outsized wave-transported boulders (cliff-top storm
443 deposits) 15-60 m above sea level in Shetland, northern Scotland, dated to between c. 1300
444 and 800 cal a BP (Hansom and Hall, 2009), indicate enhanced storminess at ~60°N. Finally,
445 lower values of Br at Cors Fochno peat bog, relative to the preceding period (Orme *et al.*,

2015), suggest that the main westerly storm tracks were not focused at the latitude of mid Wales (52°N) but further north over Scotland. However, more complexity is introduced when comparing these proxy records with growth rates from the north-west Scotland speleothem record. Baker *et al.* (2015) reconstruct strongly negative NAO-like conditions from c. 1400 to 1100 cal a BP, at which point the trend is reversed and their reconstruction shows a strongly positive phase throughout the MCA, similar to the Olsen *et al.* (2012) NAO record. Therefore, we interpret the MCA period to be one of a strong polar-front jet stream and enhanced cyclogenesis, bringing westerly storms tracking across northern Scotland (57-60°N). Although the variable antiphase relationship between MSW and Br records at Shebster perhaps suggest, at times, decreased storm frequency but higher storm intensity across the northern British Isles, consistent with the generation of high-energy storm deposits around northern Scotland's coasts (Hansom and Hall, 2009).

SH-8 600-0 cal a BP: The most recent period captured in the Shebster peat record spans from c. 600 to 100 cal a BP and almost exactly corresponds to the Little Ice Age (LIA: c. 150-700 cal a BP; Mann and Jones, 2003). This period is characterised at Shebster by high but variable MSW and generally high Br values, indicating increased wetness and storminess for much of this 500-year window. A notable exception is the period between c. 100 and 200 cal a BP when Br (i.e. storminess) is subdued with levels equivalent to SH-6. However, the cause of the LIA cooling (and/or any associated storminess) has been the source of considerable research and debate (Lamb, 1995; Orme *et al.*, 2016; Trouet *et al.*, 2012). A clear LIA signal is seen in the proxy-reconstructed NAO indices of Olsen *et al.* (2012) and Baker *et al.* (2015), where an abrupt shift from strongly positive to negative NAO occurs at c. 600 cal a BP in both records. The shift is larger and more sustained in the reconstruction provided by Baker *et al.*

(2015). The record suggests that this dominantly negative NAO phase was associated with a vigorous jet stream, a higher incidence of moisture-bearing winds and a higher frequency of storms generally tracking across the latitude of Northern Britain (55-60°N) for much of the LIA (c.100-600 cal a BP). The normal negative NAO configuration involves a significant southward shift in dominant westerly winds and storm tracks, to the latitude of southern France, northern Iberia and the western Alps (40-45°N) (Woollings *et al.*, 2008, 2010). However, other records from around Scotland show with a high level of certainty that the LIA period (esp. from 400-100 BP) was one of periodically enhanced storminess, increased sea state and wave activity, and generally disrupted weather patterns (Sommerville *et al.*, 2003; McIlvenny *et al.*, 2013; Orme *et al.*, 2016). These features are the hallmarks of an unusually turbulent period of atmospheric circulation, typically associated with disturbed jet stream strength and an unstable location (switching from zonal to meridional flow pattern), consistent with variable but high levels of Br-inferred storminess at Shebster (this study) and to a lesser degree at Cors Fochno (Orme *et al.*, 2015) during the second half of the LIA. However, the speleothem and MSW records reflect a shift to relatively drier although perhaps less stable conditions. Again the contrast between the records of storminess and local wetness is probably due to the LIA being dominated by overall colder and drier conditions but affected by lower-frequency higher-intensity storm events (supporting the findings of Trouet *et al.*, 2012).

Conclusions

The Shebster climate record provides insights into the timing and nature of North Atlantic climate changes and is a significant advance to the existing records in that it spans much of the Holocene. The combined Br and MSW records highlight the millennial to centennial scale changes in the position of the polar jet stream – a significant driver of environmental

change in northern Scotland and the wider North Atlantic region. The Shebster climate record is consistent with the Norwegian glacier record, Greenland sediment-inferred NAO index, Scottish speleothem record and Iberian speleothem record but most importantly advances our understanding of the development and fluctuations of the NAO from the early Holocene. We infer from the data that periods of high Br and MSW levels probably relate to a jet stream position over Northern Scotland and, therefore, increased storminess and a positive NAO mode. Periods of reduced Br and mire wetness levels probably relate to a more southerly position of the jet stream and, therefore, a decline in storminess and a negative NAO. Between c. 4000 and 3300 cal a BP there are very low levels of Br and mire wetness consistent with a drier period across much of the northern hemisphere which may relate to a neutral NAO state and a weaker jet stream. However, subtle differences between these two proxies suggest that single indicators of storminess may not be sufficient to reconstruct changes in jet stream movement and NAO index. These differences also suggest that higher intensity but perhaps less frequent periods of storminess are not necessarily associated with a wetter climate, which may be exemplified during the Little Ice Age. This work shows that important high-resolution palaeoenvironmental information can be gleaned by XRF-analysis of peat accumulations in cold-temperate climates. Furthermore, these analyses in combination with other established techniques offer a novel and under-used way to examine the climate record of the recent past on a decadal to millennial scale.

Acknowledgements The authors wish to thank the NERC-BGS opportunities fund and support from Dounreay Site Restoration Ltd, which led to the development of this project. HS was supported by a NERC-BUFI – University of Stirling joint studentship (NE/K501156/1). We would like to thank Dr Andrew Finlayson, Dr Mary McCulloch, Natasha Rolph and James

Blaikie for invaluable assistance in the field. We are also grateful to Dr Chris Hayward for his guidance and support for our use of the electron microprobe at the School of GeoSciences, The University of Edinburgh. Two anonymous reviewers are thanked for their comments.

References

Alexander LV, Tett SFB. 2005. Recent observed changes in severe storms over the United Kingdom and Iceland. *Geophysical Research Letters* **32** : 1-4.

Allan R, Tett S, Alexander L. 2009. Fluctuations in autumn-winter severe storms over the British Isles: 1920 to present. *International Journal of Climatology* **29** : 357-371.

Andresen CS, Björk S. 2005. Holocene climate variability in the Denmark Strait region. *Geografiska Annalar* **1**: 159-174.

Andresen CS, Bond G, Kuijpers A, Knutz PC, Björck S. 2005. Holocene climate variability at multidecadal time scales detected by sedimentological indicators in a shelf core NW off Iceland. *Marine Geology* **214**: 323-338.

Athanasiadis PJ, Wallace JM, Wettstein JJ. 2010. Patterns of jet stream wintertime variability and their relationship to the storm tracks. *Journal of Atmospheric Sciences* **67**: 1361-1381.

Auton CA, Gillespie MR, Lott G.K, McKervey JA, Milodowski AE, Stephenson MH. 2005. Palynological and petrological constraints on correlation of Devonian strata across the

541 Dounreay district, BGS Commissioned Report CR/05/009. British Geological Survey,
542 Edinburgh.

543

544 Bakke J, Lie Ø, Nesje A, Dahl SO, Paasche Ø. 2005. Utilizing physical sediment variability in
545 glacier-fed lakes for continuous glacier reconstructions during the Holocene, northern
546 Folgefonna, western Norway. *Journal of Quaternary Science* **15**: 161-176.

547

548 Bakke J, Lie Ø, Dahl SO, Nesje A, Bjune AE. 2008. Strength and spatial patterns of the
549 Holocene wintertime westerlies in the NE Atlantic region. *Global and Planetary Change* **60**:
550 28-41.

551

552 Baker A, Hellstrom J, Kelly B, Mariethoz G, Trouet V. 2015. A composite annual-resolution
553 stalagmite record of North Atlantic climate over the last three millennia. *Scientific Reports* **5**:
554 doi:10.1038/srep10307.

555

556 Balascio NL, D'Andrea WJ, Bradley RS. 2015. Glacier response to North Atlantic climate
557 variability during the Holocene. *Climate of the Past* **11**: 1587-1598.

558

559 Biester H, Keppler F, Putschew A, Martinez-Cortizas A, Petri M. 2004. Halogen retention,
560 organohalogens, and the role of organic matter decomposition on halogen enrichment in
561 two Chilean peat bogs. *Environmental Science and Technology* **38**: 1984– 1991.

562

563 Blaauw M, Van Geel B, Van der Plicht J. 2004. Solar forcing of climatic changes during the
564 mid-Holocene: indications from raised bogs in the Netherlands. *The Holocene* **14**: 1-35.

565

566 Blaauw M, Christen JA. 2011. Flexible paleoclimate age-depth models using an
567 autoregressive gamma process. *Bayesian Analysis* **6**: 457-474.

568

569 Blackford JJ, 1993. Peat bogs as sources of proxy climatic data: past approaches and future
570 research. In *Climatic change and human impact on the landscape. Studies in palaeoecology*
571 *and environmental archaeology*, Chambers FM (ed.). Chapman & Hall: London; 47-56.

572

573 Booth RK, Jackson SL, Forman JE, Kutzbach EA, Bettis III, Kreig J, Wright DK. 2005. A severe
574 centennial-scale drought in continental North America 4200 years ago and apparent global
575 linkages. *The Holocene* **15**: 321-328.

576

577 Charman DJ, Blundell A, Chiverrell RC, Hendon D, Langdon PG. 2006. Compilation of non-
578 annually resolved Holocene proxy climate records: stacked Holocene peatland palaeo-water
579 table reconstructions from northern Britain. *Quaternary Science Reviews* **25**: 336–350.

580

581 Charman DJ, Barber KE, Blaauw M, Langdon PG, Mauquoy D, Daley TJ, Hughes PDM,
582 Karofeld E. 2009. Climate drivers for peatland paleoclimate reconstructions. *The Holocene*
583 **17**: 217-227.

584

585 Charman DJ. 2010. Centennial climate variability in the British Isles during the mid-late
586 Holocene. *Quaternary Science Reviews* **29**: 1539-1554.

587

588 Cheng X, Xie S-P, Tokinaga H, Du Y. 2011. Interannual Variability of High-Wind Occurrence
589 over the North Atlantic. *Journal of Climate* **24**: 6515-6527.

590

591 Clarke M, Rendell H, Tastet J-P. 2002. Late-Holocene sand invasion and North Atlantic
592 storminess along the Aquitaine coast, southwest France. *The Holocene* **12**: 231–238.

593

594 Clarke ML, Rendell HM. 2009. The impact of North Atlantic storminess on western European
595 coasts: A review. *Quaternary International* **195**: 31-41.

596

597 Dawson AG, Elliot L, Mayewski P, Lockett P, Noone S, Hickey K, Holt T, Wadhams P, Foster I.
598 2003. Late Holocene North Atlantic climate ‘seesaws’, storminess changes and Greenland
599 ice sheet (GISP2) paleoclimates. *The Holocene* **13**: 381-392.

600

601 Dawson AG, McIlveny J, Warren J. 2010. Winter Gale Day Frequency in Shetland and
602 Faeroes, AD 1866–1905: Links to Sea Ice History and the North Atlantic Oscillation. *Scottish*
603 *Geographical Journal*, **126**: 141 - 152.

604

605 De Jong R, Björck S, Björckman L, Clemmensen LB. 2006. Storminess variation during the last
606 6500 years as reconstructed from an ombotrophic peat bog in Halland, Southwest Sweden.
607 *Journal of Quaternary Science* **21**: 905-919.

608

609 De la Vega-Leinert AC, Keen DH, Jones R L, Wells JM, Smith DE. 2000, Mid-Holocene
610 environmental changes in the Bay of Skail, Mainland Orkney, Scotland: an integrated

611 geomorphological, sedimentological and stratigraphical study. *Journal of Quaternary*
612 *Science*, 15: 509–528.

613

614 Di Rita F, Magri D. 2009. Holocene drought, deforestation and evergreen vegetation
615 development in the central Mediterranean: A 5500 year record from Lago Alimini, Piccolo,
616 Apulia, southeast Italy. *The Holocene* **19**: 295-306.

617

618 Dugmore AJ, Larsen G, Newton, AJ. 1995a. Seven tephra isochrones in Scotland. *The*
619 *Holocene* **5**: 257-266.

620

621 Dugmore AJ, Cook GT, Shore JS, Newton AJ, Edwards KJ, Larsen G. 1995b. Radiocarbon
622 Dating Tephra Layers in Britain and Iceland. *Radiocarbon* **37**: 379-388.

623

624 Genty D, Blamart D, Ghaleb B, Plagnes V, Causse C, Bakalowicz M, Melières MA, Zouari K,
625 Chkir N. 2006. Timing and Dynamics of the Last Deglaciation from European and North
626 African $\delta^{13}\text{C}$ stalagmite profiles – Comparison with South-Hemisphere stalagmite records.
627 *Quaternary Science Review* **25**: 2118-2142.

628

629 Hanna E, Cappelen J, Allan R, Jónsson T, Le Blanc F, Lillington T, Hickey K. 2008. New insights
630 into North European and North Atlantic surface pressure variability, storminess, and related
631 climate change since 1830. *Journal of Climate* **21**: 6739-6766.

632

633 Hansom JD, Hall AM. 2009. Magnitude and frequency of extra-tropical North Atlantic
634 cyclones: A chronology from cliff-top storm deposits. *Quaternary International* **195**: 42-52.

635

636 Haug GH, Hughen KA, Sigman DM, Petersen LC, Röhl U. 2001. Southward migration of the
637 Intertropical Convergence Zone through the Holocene. *Science* **293**: 1304-1308.

638

639 Hayward C. 2013. High spatial resolution electron probe microanalysis of tephras and melt
640 inclusions without beam-induced chemical modification. *The Holocene* **22**: 119–125.

641

642 Hughes PDM, Mauquoy D, Barber KE, Langdon PG. 2000. Mire-development pathways and
643 paleoclimatic records from a full Holocene peat archive at Walton Moss, Cumbria, England.
644 *The Holocene* **10**: 465-479.

645

646 Hunt JB, Hill PG. 1993. Tephra geochemistry: a discussion of some persistent analytical
647 problems. *The Holocene* **3**: 271-278.

648

649 Hurrell JW. 1995. Decadal Trends in the North Atlantic Oscillation: Regional Temperatures
650 and Precipitation. *Science* **269** : 676-679.

651

652 Hurrell J W, Kushnir Y, Ottersen G, Visbeck M. 2003. An Overview of the North Atlantic
653 Oscillation. In *The North Atlantic Oscillation: Climatic Significance and Environmental Impact*
654 Hurrell JW, Kushnir Y, Ottersen G, Visbeck M. (eds.). American Geophysical Union:
655 Washington, D. C.; 1-35.

656

657 Larsen DJ, Miller GH, Geirsdóttir Á, Ólafsdóttir S. 2012. Non-linear Holocene climate
658 evolution in the North Atlantic: a high-resolution, multi-proxy record of glacier activity and

659 environmental change from Hvítárvatn, central Iceland. *Quaternary Science Reviews* **39**: 14-
660 25.

661

662 McDermott F, Matthey DP, Hawkesworth C. 2001. Centennial-scale Holocene climate
663 variability revealed by a high-resolution speleothem delta O-18 record from SW Ireland.
664 *Science* **294**: 1328–1331.

665

666 McIlvenny JD, Muller FLL, Dawson A. 2013. A 7600-year sedimentary record of climatic
667 instability in Dunnet Bay, North Scotland. *Marine Geology* **335**: 100-113.

668

669 Mann ME, Jones PD. 2003. Global surface temperatures over the past two millennia.
670 *Geophysical Research Letter*, **30**: 10.1029/2003GL017814.

671

672 Mauri A, Davis BAS, Collins PM, Kaplan JO. 2014. The influence of atmospheric circulation on
673 the mid-Holocene climate of Europe: a data-model comparison. *Climates of the Past* **10**:
674 1925–1938.

675

676 Mayewski PA, Rohling EE, Stager JC, Karlen W, Maasch KA, Meeker LD, Meyerson EA, Gasse
677 F, Van Kreveld S, Holmgren K, Lee-Thorp J, Rosqvist G, Rack F, Staubwasser M, Schneider RR,
678 Steig EJ. 2004. Holocene climate variability. *Quaternary Research* **62**: 243–255.

679

680 Metcalfe SE, Barron JA, Davies SJ. 2015. The Holocene history of the North American
681 Monsoon: known knowns and known unknowns in understanding its spatial and temporal
682 complexity. *Quaternary Science Reviews* **120**: 1-27.

683

684 Moreno j, Fatela F, Leorri E, Araújo MF, Moreno F, De la Rosa J, Freitas MC, Valente T,
685 Corbett R. 2015. Bromine enrichment in marsh sediments as a marker of environmental
686 changes driven by grand solar minima and anthropogenic activity (Caminha, NW of Portugal)
687 *Science of the Total Environment* **506–507**: 554–566.

688

689 Newton AJ. 1996. Tephabase: A Tephrochronological Database. *Quaternary Newsletter* **78**:
690 8-13.

691

692 O'Brien SR, Mayewski PA, Meeker LD, Meese DA, Twickler MS, Whitlow SI. 1995. Complexity
693 of Holocene climate as reconstructed from a Greenland ice core. *Science* **270**: 1962–1964.

694

695 Oldfield F, Batterbee RW, Boyle JF, Cameron NG, Davis B, Evershed RP, McGovern AD, Jones
696 V, Thompson R. 2010. Terrestrial and aquatic ecosystem responses to the late Holocene
697 climate change recorded in the sediments of Lochan Uaine, Cairngorms, Scotland.
698 *Quaternary Science Reviews* **29**: 1040-1054.

699

700 Olsen J, Anderson NJ, Knudsen MF. 2012. Variability of the North Atlantic Oscillation over
701 the past 5200 years. *Nature Geoscience* **5**: 808–812.

702

703 Orme LC, Davies SJ, Duller GAT. 2015. Reconstructed centennial variability of Late Holocene
704 storminess from Cors Fochno, Wales, UK. *Journal of Quaternary Science* **30**: 478-488.

705

706 Orme LC, Reinhardt L, Jones RT, Charman DJ, Barkwith A, Ellis MA. 2016. Aeolian sediment
 707 reconstructions from the Scottish Outer Hebrides: Late Holocene storminess and the role of
 708 the North Atlantic Oscillation. *Quaternary Science Reviews* **132**: 15-25.
 709
 710 Pinto JG, Zacharias S, Fink AH, Leckebusch GC, Ulbrich U. 2009. Factors contributing to the
 711 development of extreme North Atlantic cyclones and their relationship with the NAO.
 712 *Climate Dynamics* **32**: 711-737.
 713
 714 Reimer PJ, Bard E, Bayliss AJ, Beck W, Blackwell PG, Bronk Ramsey C, Buck CE, Cheng H,
 715 Edwards RL, Friedrich M, Grootes PM, Guilderson TP, Haflidason H, Hajdas I, Hatté C, Heaton
 716 TJ, Hoffmann DL, Hogg AG, Hughen KA, Kaiser KF, Kromer B, Manning SW, Niu M, Reimer
 717 RW, Richards DA, Marian Scott EM, Southon JR, Staff RA, Turney CSM, Van der Plicht J. 2013.
 718 INTCAL13 and MARINE13 Radiocarbon Age Calibration Curves 0–50,000 years cal. BP.
 719 *Radiocarbon* **55**: 1869–1887.
 720
 721 Sabatier P, Dezileau L, Colin C. 2012. 7000 years of paleostorm activity in the NW
 722 Mediterranean Sea in response to Holocene climate events. *Quaternary Research* **77**: 1–11.
 723
 724 Sander R, Keene WC, Pszenny AAP, Arimoto R, Ayers GP, Baboukas E, Cainey JM, Crutzen PJ,
 725 Duce RA, Hönninger G, Huebert BJ, Maenhaut W, Mihalopoulos N, Turekian VC, Van
 726 Dingenen R. 2003. Inorganic bromine in the marine boundary layer: a critical review.
 727 *Atmospheric Chemistry and Physics* **3**: 1301-1336
 728

729 Stuiver M, Reimer PJ. 1993. Extended 14C database and revised CALIB radiocarbon
730 calibration program. *Radiocarbon* **35**: 215-230.
731
732 Swindles G, Plunkett G, Roe HM. 2007. A delayed response to solar forcing at 2800 cal. BP:
733 multiproxy evidence from three Irish peatlands. *The Holocene* **17**: 177-182.
734
735 Sommerville AA, Hansom JD, Sanderson DCW, Housley RA. 2003. Optically stimulated
736 luminescence dating of large storm events in Northern Scotland. *Quaternary Science*
737 *Reviews* **22**: 1085-1092.
738
739 Strong C, Davis RE. 2008. Variability in the position and strength of winter jet stream cores
740 related to Northern Hemisphere teleconnections. *Journal of Climatology* **21**: 584–592.
741
742 Tipping R, Bradley R, Sanders J, McCulloch R, Wilson R. 2012. Moments of crisis: climate
743 change in Scottish prehistory. *Society of Antiquaries of Scotland* **142**: 9-25.
744
745 Tisdall E, McCulloch R, Sanderson D, Simpson I, Woodward N. 2013. Living with sand: A
746 record of landscape change and storminess during the Bronze and Iron Ages Orkney,
747 Scotland. *Quaternary International* **308-309**: 205-215.
748
749 Trouet V, Esper J, Graham NE, Baker A, Scourse JD, Frank, DC. 2009. Persistent Positive
750 North Atlantic Oscillation Mode Dominated the Medieval Climate Anomaly. *Science* **324**: 78-
751 80
752

753 Trouet V, Scourse JD, Raible CC. 2012. North Atlantic storminess and Atlantic Meridional
754 Overturning Circulation during the last millennium: Reconciling contradictory proxy records
755 of NAO variability. *Global and Planetary Change* **84-85**: 48-55.

756

757 Turner TE, Swindles GT, Roucoux KH. 2014. Late Holocene ecohydrological and carbon
758 dynamics of a UK raised bog: impact of human activity and climate change. *Quaternary*
759 *Science Reviews* **84**: 65-85.

760

761 Vliet-Lanoë B, Penaud AI, Hénaff A, Delacourt C, Fernane A, Goslin JRM, Hallégouët B, Le
762 Cornec E. 2014. Middle-to-late Holocene storminess in Brittany (NW France): Part II – the
763 chronology of events and climate forcing, *Holocene* **24**: 296-310.

764

765 Walczak IW, Baldini JUL, Baldini LM, McDermott F, Marsden S, Standish CD, Richards DA,
766 Andreo B, Slater J. 2015. Reconstructing high-resolution climate using CT scanning of
767 unsectioned stalagmites: A case study identifying the mid-Holocene onset of the
768 Mediterranean climate in southern Iberia. *Quaternary Science Reviews* **127**: 117-128.

769

770 Wang XL, Zwiers FW, Swail VR, Feng Y. 2009. Trends and variability of storminess in the
771 Northeast Atlantic region. 1874-2007. *Climate Dynamics* **33**: 1179-1195.

772

773 Wang T, Surge D, Mithen S. 2012. Seasonal temperature variability of the Neoglacial (3300–
774 2500 BP) and Roman Warm Period (2500–1600 BP) reconstructed from oxygen isotope
775 ratios of limpet shells (*Patella vulgata*), Northwest Scotland. *Palaeogeography,*
776 *Palaeoclimatology, Palaeoecology* **317–318**: 104–113.

777

778 Woollings T, Hoskins B, Blackburn M, Berrisford P. 2008. A new Rossby wave-breaking
779 interpretation of the North Atlantic Oscillation. *Journal of the Atmospheric Sciences* **65**: 609-
780 626.

781

782 Woollings T, Hannachi A, Hoskins B. 2010. Variability of the North Atlantic eddy-driven jet
783 stream. *Quarterly journal of the Royal Meteorological Society* **136**: 856-868.

784

785 Zaccone C, Coccozza C, Shotyk W, Miano TM. 2008. Humic acids role in Br accumulation along
786 two ombrotrophic peat bog profiles. *Geoderma*, **146**: 26-31.

787

788

789 **List of Figures**

790 Figure 1: The preferred positions of the Polar Front Jet Stream and the corresponding
791 phases of the NAO). White lines - Polar Front (i.e. average southern winter limit of Polar air
792 masses). Grey long-dashed line - mean position of jet stream core between 1958 and 2006
793 (from Strong & Davis, 2008). Coloured arrows -- approximate position of Polar vortex winds
794 (colours match storms). Mode A dominates during strongly positive NAO phases; mode B₁ or
795 B₂ dominates during strongly negative NAO. Neutral NAO phase (neither positive or
796 negative) equates to a weak jet stream, with much-reduced storm frequency probably along
797 negative NAO tracks (B₁, B₂).

798

799 Figure 2: A) Location map showing the study site in Northern Scotland. B) Detailed map of
800 peat coring site near the Burn of Shebster. [Grid ticks are in British National Grid.] C)

Average climatological conditions (1981-2010) at the nearest long-running weather station (Wick Airport; 58.454 N, 3.089 W; 36 m asl). Mean monthly air temperature (maximum and minimum) and normal wind speed envelope (at 10 m) plotted on the same axis. Mean monthly precipitation shown as blue bars. [Data from metoffice.gov.uk]. Note the marked seasonality in average wind strength, peaking in winter (Dec-Mar).

Figure 3: Shebster stratigraphy, LOI₅₅₀, Mire Surface Wetness, Bromine (ratio of Br / Inc+Coh) and BACON age/depth model.

Figure 4: Shebster Mire Surface Wetness, Bromine, 5000-yr glacier record from Folgefanna in Southern Norway (Bakke *et al.*, 2008), reconstructed 5000-yr NAO index based on a lake-sediment core in SW Greenland (Olsen *et al.*, 2012), Scottish speleothem record (Baker *et al.*, 2015) and Iberian speleothem record (Walkzak *et al.*, 2015). The Hekla 4 tephra layer is indicated by a grey dotted line.

List of Tables

Table 1: Radiocarbon dates and ages for the H4 tephra layer correlated to the Shebster record. ¹⁴C dates have been calibrated using CALIB Rev. 7.10 (Stuiver and Reimer, 1993) and IntCal 13.14c. (Reimer *et al.*, 2013).

Figure 1

Figure 2

Figure 3

Figure 4

Table 1: Radiocarbon dates and ages for the H4 tephra layer correlated to the Shebster record. ¹⁴C dates have been calibrated using CALIB Rev. 7.10 (Stuiver and Reimer, 1993) and IntCal 13.14c. (Reimer *et al.*, 2013).

Depth (cm)	¹⁴ C a BP (1σ)	δ ¹³ C _{V-PDB} ‰	cal a BP (1σ) ²	Weighted Mean Age (cal a BP) ³	Laboratory Code
150	1740 ± 40	-28.3	1606 – 1706	1679	Beta 251972
271	3010 ± 40	-28.0	3083 – 3322	3215	Beta 251973
346	3826 ± 12 ¹	n/a	4159 – 4241	4199	N/A
405	4110 ± 40	-29.4	4532 – 4802	4841	Beta 251974
500	5730 ± 40	-25.6	6454 – 6601	6507	Beta 251975
612	7530 ± 50	-24.7	8320 – 8403	8286	Beta 251976

¹ ¹⁴C age for Hekla 4 cryptotephra layer (Dugmore *et al.*, 1995b)

² Calibrated ages produced using Calib Ver.7.1 (Stuiver and Reimer, 1993) and IntCal13 (Reimer *et al.*, 2013)

³ Weighted mean ages produced using BACON Bayesian age-depth program (Blaauw and Christen, 2011)