A 2 : 1 Coupling Reaction of Arynes with Aldehydes via *o*-Quinone Methides: Straightforward Synthesis of 9-Arylxanthenes

Hiroto Yoshida,* Masahiko Watanabe, Hiroyuki Fukushima, Joji Ohshita and
Atsutaka Kunai*

*Department of Applied Chemistry, Graduate School of Engineering, Hiroshima University,
Higashi-Hiroshima 739-8527, Japan.*

E-mail: yhiroto@hiroshima-u.ac.jp, akunai@hiroshima-u.ac.jp

Supporting Information

General Remarks. All manipulations of oxygen- and moisture-sensitive materials were conducted with a standard Schlenk technique under a purified argon atmosphere. Melting points were determined using a Yanaco micro melting point apparatus. Nuclear magnetic resonance spectra were taken on a JEOL EX-270 (^1H , 270 MHz; ^{13}C , 67.8 MHz) spectrometer or a JEOL Lambda-400 (^1H , 400 MHz; ^{13}C , 99.5 MHz) spectrometer using residual chloroform (^1H) or CDCl_3 (^{13}C) as an internal standard. High-resolution mass spectra were obtained with a Hitachi M-80B spectrometer or a JEOL JMS-HX110A spectrometer. The preparative recycling gel permeation chromatography was performed with GL Science PU 614 equipped with Shodex GPC H-2001L and -2002L columns (chloroform as an eluent). Column chromatography was carried out using Merck Kieselgel 60. Unless otherwise noted, commercially available reagents were used without purification. 18-Crown-6 was recrystallized from distilled MeCN. KF (spray-dried) was vacuum dried at 100 °C for 12 h. THF was distilled from sodium/benzophenone ketyl. MeCN was distilled from phosphorus pentoxide.

Aryne Precursors. 2-(Trimethylsilyl)phenyl triflate (**1a**),¹ 4,5-dimethyl-2-(trimethylsilyl)phenyl triflate (**1b**),² 6-(trimethylsilyl)-5-indanyl triflate (**1c**),² 3,6-dimethoxy-2-(trimethylsilyl)phenyl triflate (**1d**)² and 3-methoxy-2-(trimethylsilyl)phenyl triflate (**1e**)³ were prepared according to literature procedures.

Aldehydes. 2-Benzoyloxy-1-naphthaldehyde (**2a**)⁴ and 2,6-dibenzoyloxybenzaldehyde (**2e**)⁵ were synthesized according to literature methods. 3,3'-Diformyl-4,4'-dimethoxydiphenyl ether (**2o**) was prepared from 3,3'-dibromo-4,4'-dimethoxydiphenyl ether in a similar manner as reported.⁶ 3,3'-Dibromo-4,4'-dimethoxydiphenyl ether was synthesized by the methylation of 3,3'-dibromo-4,4'-dihydroxydiphenyl ether⁷ with methyl iodide (8 eq) and K_2CO_3 (8 eq) in refluxing acetone.

Reaction of Arynes with Aldehydes. A General Procedure. To a THF solution (1.0 mL) of an aryne precursor (0.45 mmol), an aldehyde (0.15 mmol) and 18-crown-6 (0.238 g, 0.90 mmol) was added KF (0.052 g, 0.90 mmol), and the resulting mixture was stirred at 0 °C. After the time specified in Table 1 or Scheme 2, the mixture was diluted with ethyl acetate, filtered through a Celite plug, and concentrated. Silicagel column chromatography (hexane/ethyl acetate as an eluent) followed by gel permeation chromatography (chloroform as an eluent) gave the corresponding product.

9-(2-Benzyloxy-1-naphthyl)xanthene (3aa). (A 65 : 35 mixture of two conformers)

Isolated in 70% yield as a white solid: m.p. 167–168 °C; ¹H NMR (CDCl₃) δ 4.74 (s, minor, -CH₂Ph), 5.37 (s, major, -CH₂Ph), 6.37 (s, minor, methine H), 6.65 (d, *J* = 7.5 Hz), 6.69–6.88 (m), 6.95 (d, *J* = 8.0 Hz), 7.06 (t, *J* = 7.0 Hz), 7.09–7.25 (m), 7.29–7.41 (m), 7.43–7.54 (m), 7.56 (d, *J* = 8.7 Hz), 7.65 (t, *J* = 7.5 Hz), 7.75 (d, *J* = 8.7 Hz), 7.83–7.91 (m), 8.46 (d, *J* = 8.9 Hz); ¹³C NMR (CDCl₃) δ 33.4, 34.4, 70.0, 72.0, 114.7, 115.3, 115.8, 116.1, 122.0, 122.3, 123.1, 123.3, 124.1, 124.7, 125.7, 125.9, 126.8, 127.1, 127.3, 127.46, 127.52, 127.58, 127.62, 128.0, 128.1, 128.4, 128.5, 128.6, 128.7, 128.77, 128.82, 129.0, 129.7, 130.7, 131.7, 133.5, 136.8, 137.0, 150.9, 151.3, 154.1, 154.9; Anal. Calcd for C₃₀H₂₂O₂ C, 86.93; H, 5.35. Found: C, 86.63; H, 5.33.

¹H NMR integration ratio

4.74 (s) : 5.37 (s) : 6.37 (s) : 6.65 (d) : 6.69–6.88 (m) : 6.95 (d) : 7.06 (t) : 7.09–7.25 (m) : 7.29–7.41 (m) : 7.43–7.54 (m) : 7.56 (d) : 7.65 (t) : 7.75 (d) : 7.83–7.91 (m) : 8.46 (d) = 2.11 : 3.94 : 1.05 : 3.89 : 12.34 : 2.41 : 2.03 : 16.15 : 6.05 : 6.92 : 2.16 : 1.18 : 3.05 : 3.06 : 1.00.

9-(2-Methoxy-1-naphthyl)xanthene (3ab). (A 65 : 35 mixture of two conformers)

Isolated in 66 % yield as a white solid: m.p. 219–221 °C; ¹H NMR (CDCl₃) δ 3.32 (s, minor, -OCH₃), 4.06 (s, major, -OCH₃), 6.29 (s, minor, methine H), 6.64–6.67 (m), 6.77–6.83 (m), 7.02–7.19 (m), 7.43–7.48 (m), 7.52 (d, *J* = 4.1 Hz), 7.65 (t, *J* = 8.0 Hz), 7.75 (d, *J* = 8.2 Hz), 7.80 (d, *J* = 8.9 Hz), 7.88 (d, *J* = 8.9 Hz), 8.46 (d, *J* = 8.7 Hz); ¹³C NMR (CDCl₃) δ 33.2, 34.4, 56.6, 57.2, 113.0, 115.6, 116.1, 116.2, 122.2, 122.3, 123.0, 123.2, 123.5, 123.6, 124.2, 124.8, 125.6, 125.9, 127.3, 127.4, 127.5, 128.4, 128.49, 128.52, 128.8, 129.1, 129.2, 129.8, 130.4, 131.7, 133.2, 151.0, 151.2, 155.7, 156.0; HRMS Calcd for C₂₄H₁₈O₂: M⁺, 338.1307. Found: *m/z* 338.1313.

¹H NMR integration ratio

3.32 (s) : 4.06 (s) : 6.29 (s) : 6.64–6.67 (m) : 6.77–6.83 (m) : 7.02–7.19 (m) : 7.43–7.48 (m) : 7.52 (d) : 7.65 (t) : 7.75 (d) : 7.80 (d) : 7.88 (d) : 8.46 (d) = 3.00 : 5.68 : 0.96 : 5.61 : 8.08 : 16.46 : 2.90 : 2.02 : 1.09 : 1.77 : 1.17 : 2.95 : 0.97.

9-(1-Naphthyl)xanthene (3ac). Isolated in 17 % yield as a white solid: m.p. 159–161 °C; ¹H NMR (CDCl₃) δ 6.02 (s, 1 H), 6.82–6.90 (m, 4 H), 7.14–7.22 (m, 4 H), 7.35–7.46 (m, 4 H), 7.79 (d, *J* = 8.0 Hz, 1 H), 7.86 (d, *J* = 8.6 Hz, 1 H), 8.08 (s, 1 H); ¹³C NMR (CDCl₃) δ 29.7, 116.4, 123.1, 123.6, 124.5, 124.8, 125.5, 125.6, 126.1, 127.9, 128.5, 129.0, 129.1, 129.2, 131.1, 134.5, 150.7; HRMS Calcd for C₂₃H₁₆O: M⁺, 308.1201. Found: *m/z* 308.1235.

9-(2,6-Dimethoxyphenyl)xanthene (3ad). Isolated in 61 % yield as a white solid: m.p. 158–159 °C; ¹H NMR (CDCl₃) δ 3.33 (brs, 3 H), 3.96 (brs, 3 H), 6.07 (s, 1 H), 6.55 (brs, 2 H), 6.87 (t, *J* = 7.4 Hz, 2 H), 6.96 (d, *J* = 7.0 Hz, 2 H), 7.03 (d, *J* = 8.2 Hz, 2 H), 7.11–7.20 (m, 3 H); ¹³C NMR (CDCl₃) δ 32.0, 55.9, 115.4, 122.3, 123.6, 124.3, 126.9, 128.1, 128.6, 151.5; HRMS Calcd for C₂₁H₁₈O₃: M⁺, 318.1256. Found: *m/z* 318.1179.

9-(2,6-Dibenzyloxyphenyl)xanthene (3ae). Isolated in 60% yield as a white solid: m.p. 142–145 °C; ¹H NMR (CDCl₃) δ 4.67 (s, 2 H), 5.28 (s, 2 H), 6.27 (s, 1 H), 6.40 (brs, 1 H), 6.68 (brs, 2 H), 6.75 (brs, 1 H), 6.84–7.02 (m, 6 H), 7.02–7.29 (m, 6 H), 7.29–7.48 (m, 3 H), 7.54 (brs, 2 H); ¹³C NMR (CDCl₃) δ 32.0, 69.5, 70.7, 104.9, 106.2, 115.6, 122.2 (brs), 123.3, 124.1, 127.1 (brs), 128.1 (brs), 128.6 (brs), 136.7, 137.1, 151.4, 156.5, 157.4; HRMS Calcd for C₂₆H₁₉O₃: M⁺-CH₂Ph, 379.1334. Found: *m/z* 379.1331.

9-(2-Methoxyphenyl)xanthene (3af). Isolated in 54 % yield as a white solid: m.p. 143–145 °C; ¹H NMR (CDCl₃) δ 3.86 (s, 3 H), 5.82 (s, 1 H), 6.82 (t, *J* = 7.5 Hz, 1 H), 6.90 (d, *J* = 8.2 Hz, 1 H), 6.93–6.97 (m, 3 H), 7.09–7.21 (m, 7 H); ¹³C NMR (CDCl₃) δ 36.9, 55.6, 111.0, 116.2, 121.1, 123.0, 124.8, 127.5, 127.7, 129.5, 130.3, 135.3, 151.5, 156.2; HRMS Calcd for C₂₀H₁₆O₂: M⁺, 288.1150. Found: *m/z* 288.1159.

9-(2,4-Dimethoxyphenyl)xanthene (3ag). Isolated in 53 % yield as a white solid: m.p. 89–91 °C; ¹H NMR (CDCl₃) δ 3.75 (s, 3 H), 3.84 (s, 3 H), 5.72 (s, 1 H), 6.36 (dd, *J* = 8.4, 2.4 Hz, 1 H), 6.48 (d, *J* = 2.4 Hz, 1 H), 6.84 (d, *J* = 8.4 Hz, 1 H), 6.96 (t, *J* = 7.5 Hz, 2 H), 7.09–7.21 (m, 6 H); ¹³C NMR (CDCl₃) δ 36.3, 55.2, 55.5, 98.6, 104.7, 116.1, 123.0, 125.0, 127.4, 128.0, 129.4, 130.7, 151.4, 157.2, 159.3; HRMS Calcd for C₂₀H₁₅O₃: M⁺-Me, 303.1021. Found: *m/z* 303.0989.

9-(2,4,6-Trimethoxyphenyl)xanthene (3ah). Isolated in 44 % yield as a white solid: m.p. 162–163 °C; ¹H NMR (CDCl₃) δ 3.29 (brs, 3 H), 3.78 (s, 3 H), 3.96 (brs, 3 H), 5.94 (s, 1 H), 6.10 (brs, 2 H), 6.87 (t, *J* = 7.3 Hz, 2 H), 6.96 (d, *J* = 7.3 Hz, 2 H), 7.02 (d, *J* = 8.0 Hz, 2 H), 7.12 (t, *J* = 7.5 Hz, 2 H); ¹³C NMR (CDCl₃) δ 31.7, 55.2, 55.9, 115.4, 116.5, 122.2, 124.7, 126.8, 128.56, 151.5, 160.0; HRMS Calcd for C₂₂H₂₀O₄: M⁺, 348.1362. Found: *m/z* 348.1393.

9-(4-Methoxyphenyl)xanthene (3ai). Isolated in 42% yield as a white solid: m.p. 110–112 °C; ^1H NMR (CDCl_3) δ 3.76 (s, 3 H), 5.22 (s, 1 H), 6.82 (d, $J = 9.1$ Hz, 2 H), 7.06 (d, $J = 6.8$ Hz, 2 H), 6.98 (t, $J = 7.0$ Hz, 2 H), 7.11–7.14 (m, 4 H), 7.21 (td, $J = 7.6, 1.7$ Hz, 2 H); ^{13}C NMR (CDCl_3) δ 43.5, 55.2, 114.1, 116.5, 123.2, 124.7, 127.8, 129.3, 129.7, 138.8, 151.0, 158.3; HRMS Calcd for $\text{C}_{20}\text{H}_{16}\text{O}_2$: M^+ , 288.1150. Found: m/z 288.1183.

9-(2,4,6-Trimethylphenyl)xanthene (3aj).⁸ Isolated in 58 % yield as a white solid: ^1H NMR (CDCl_3) δ 1.65 (s, 3 H), 2.31 (s, 3 H), 2.55 (s, 3 H), 5.90 (s, 1 H), 6.70–6.84 (m, 3 H), 6.90 (td, $J = 7.2, 1.0$ Hz, 2 H), 7.00 (s, 1 H), 7.06 (d, $J = 7.6$ Hz, 2 H), 7.18 (t, $J = 7.3$ Hz, 2 H); ^{13}C NMR (CDCl_3) δ 20.6, 20.9, 21.3, 37.1, 116.0, 123.0, 123.7, 127.6, 128.1, 128.6, 131.4, 136.5, 137.0, 137.4, 137.7, 150.8.

9-(2,4-Dimethylphenyl)xanthene (3ak). Isolated in 41 % yield as a white solid: m.p. 113–115 °C; ^1H NMR (CDCl_3) δ 2.18 (s, 3 H), 2.31 (s, 3 H), 5.52 (s, 1 H), 6.88–6.99 (m, 6 H), 7.05–7.11 (m, 3 H), 7.19 (td, $J = 7.4, 1.7$ Hz, 2 H); ^{13}C NMR (CDCl_3) δ 20.0, 21.0, 41.0, 116.2, 123.1, 124.3, 127.1, 127.7, 129.2, 131.0, 132.0, 135.7, 136.4, 140.6, 150.9; HRMS Calcd for $\text{C}_{21}\text{H}_{18}\text{O}$: M^+ , 286.1358. Found: m/z 286.1349.

9-(2,6-Dimethylphenyl)xanthene (3al).⁸ Isolated in 38 % yield as a white solid: ^1H NMR (CDCl_3) δ 1.69 (s, 3 H), 2.59 (s, 3 H), 5.94 (s, 1 H), 6.74 (d, $J = 7.5$ Hz, 2 H), 6.88–6.95 (m, 3 H), 7.07 (d, $J = 8.2$ Hz, 2 H), 7.12–7.21 (m, 4 H); ^{13}C NMR (CDCl_3) δ 21.7, 21.3, 37.4, 116.1, 123.1, 123.5, 127.1, 127.7, 127.9, 128.0, 130.7, 137.2, 138.0, 140.3, 150.9.

9-Phenylxanthene (3am). Isolated in 22% yield as a white solid: m.p. 136–137 °C; ^1H NMR (CDCl_3) δ 5.26 (s, 1 H), 6.98 (t, $J = 7.0$ Hz, 2 H), 7.06 (d, $J = 7.5$ Hz, 2 H), 7.13 (d, $J = 8.2$ Hz, 2 H), 7.17–7.29 (m, 7 H); ^{13}C NMR (CDCl_3) δ 44.4, 116.5, 123.2, 124.4, 126.6, 127.9, 128.4, 128.7, 129.7, 146.5, 151.0; HRMS Calcd for $\text{C}_{19}\text{H}_{14}\text{O}$: M^+ , 258.1045. Found: m/z 258.1068.

9-(2-Benzyloxy-1-naphthyl)-2,3,6,7-tetramethylxanthene (3ba). (A 70 : 30 mixture of two conformers) Isolated in 51 % yield as a green solid; m.p. 219–222 °C; ^1H NMR (CDCl_3) δ 1.94 (s), 2.01 (s), 2.16–2.27 (m), 4.78 (s, minor, $-\text{CH}_2\text{Ph}$), 5.39 (s, major, $-\text{CH}_2\text{Ph}$), 6.24 (s, minor, methine H), 6.40 (s), 6.56 (s), 6.64 (s), 6.69 (d, $J = 7.2$ Hz), 6.74 (s), 6.90 (s), 7.02–7.25 (m), 7.29–7.61 (m), 7.62–7.79 (m), 7.79–7.93 (m), 8.49 (d, $J = 8.9$ Hz); ^{13}C NMR (CDCl_3) δ 18.7, 18.8, 18.9, 19.0, 19.1, 19.4, 19.5, 32.6, 32.7, 33.8, 71.9, 114.7, 114.8, 115.6, 115.7, 116.2, 116.5, 116.6, 116.8, 116.9, 121.1, 121.5, 122.3, 123.3 (brs), 125.5, 125.7, 125.9 (brs), 126.1, 126.9, 127.0, 127.2, 127.6, 127.7, 127.9, 128.1, 128.2, 128.3, 128.5, 128.6, 128.8,

128.9, 129.08, 129.13, 129.2, 129.4, 130.0, 130.65, 130.69, 132.0, 133.6, 135.3, 135.7, 137.1; HRMS Calcd for $C_{27}H_{23}O_2$: $M^+ - CH_2Ph$, 379.1698. Found: m/z 379.1690.

1H NMR integration ratio

1.94 (s) : 2.01 (s) : 2.16-2.27 (m) : 4.78 (s) : 5.39 (s) : 6.24 (s) : 6.40 (s) : 6.56 (s) : 6.64 (s) : 6.69 (d) : 6.74 (s) : 6.90 (s) : 7.02-7.25 (m) : 7.29-7.61 (m) : 7.62-7.79 (m) : 7.79-7.93 (m) : 8.49 (d) = 13.75 : 6.99 : 22.25 : 2.00 : 4.64 : 1.01 : 4.64 : 2.03 : 2.20 : 2.25 : 2.62 : 4.77 : 9.16 : 15.55 : 6.77 : 3.48 : 1.00.

9-(2-Benzyloxy-1-naphthyl)-dicyclopenta[*b,i*]xanthene (3ca). (A 70 : 30 mixture of two conformers) Isolated in 50 % yield as a dark brown solid: m.p. decomposed; 1H NMR ($CDCl_3$) δ 1.87-2.21 (m), 2.47-2.76 (m), 2.76-3.02 (m), 4.81 (s, minor, $-CH_2Ph$), 5.37 (s, major, $-CH_2Ph$), 6.31 (s, minor, methine *H*), 6.45 (s), 6.61-6.74 (m), 6.74-6.86 (m), 6.93-7.03 (m), 7.03-7.22 (m), 7.28-7.39 (m), 7.39-7.57 (m), 7.57-7.78 (m), 7.78-7.92 (m), 8.48 (d, $J = 8.7$ Hz); ^{13}C NMR ($CDCl_3$) δ 25.55, 25.64, 25.8, 32.0, 32.68, 32.72, 32.8, 33.7, 33.8, 34.9, 69.9, 72.0, 72.4, 100.5, 111.3, 111.4, 111.5, 111.6, 111.8, 114.85, 114.91, 115.0, 115.6, 115.7, 121.6, 122.1, 122.3, 123.0, 123.3, 123.55, 123.63, 123.9, 124.4, 125.8, 125.9, 126.1, 126.2, 126.8, 126.9, 127.2, 127.7, 127.8, 127.9, 128.0, 128.3, 128.4, 128.5, 128.6, 128.8, 129.20, 129.24, 129.4, 130.7, 132.0, 133.6, 137.0, 137.1, 137.6, 138.4, 143.2, 143.5, 149.8, 150.3, 153.9, 154.5; HRMS Calcd for $C_{29}H_{23}O_2$: $M^+ - CH_2Ph$, 403.1698. Found: m/z 403.1684.

1H NMR integration ratio

1.87-2.21 (m) : 2.47-2.76 (m) : 2.76-3.02 (m) : 4.81 (s) : 5.37 (s) : 6.31 (s) : 6.45 (s) : 6.61-6.74 (m) : 6.74-6.86 (m) : 6.93-7.03 (m) : 7.03-7.22 (m) : 7.28-7.39 (m) : 7.39-7.57 (m) : 7.57-7.78 (m) : 7.78-7.92 (m) : 8.48 (d) = 18.78 : 17.86 : 20.10 : 2.21 : 5.37 : 1.12 : 5.58 : 8.21 : 2.80 : 5.73 : 12.12 : 9.39 : 10.10 : 8.55 : 4.91 : 1.00.

9-(4-Methoxyphenyl)-1,4,5,8-tetramethoxyxanthene (3di). Isolated in 45 % yield as a brown solid: m.p. 142–145 °C; 1H NMR ($CDCl_3$) δ 3.43 (s, 3 H), 3.79 (s, 3 H), 3.81 (s, 3 H), 3.83 (s, 3 H), 3.88 (s, 3 H), 6.81-6.92 (m, 6 H), 7.36 (s, 1 H), 7.72 (d, $J = 8.9$ Hz, 2 H); ^{13}C NMR ($CDCl_3$) δ 54.2, 55.2, 56.0, 57.0, 58.0, 84.8, 89.7, 112.8, 114.0, 114.3, 126.0, 126.6, 127.3, 129.8, 130.5, 131.2, 133.9, 134.7, 149.0, 150.5, 159.9, 190.7; HRMS Calcd for $C_{24}H_{24}O_6$: M^+ , 408.1573. Found: m/z 408.1570.

1,5-Dimethoxy-9-(2,4,6-trimethylphenyl)xanthene (3ej). (A 91 : 9 mixture of two conformers)⁹ Isolated in 48 % yield as a white solid; m.p. 125–131 °C; 1H NMR ($CDCl_3$) δ 1.92 (s, minor, $-CH_3$), 1.98 (s, major, $-CH_3$), 2.12 (s, major, $-CH_3$), 2.15 (s, minor, $-CH_3$), 2.30 (s, major, $-CH_3$), 3.08 (s, minor, $-OCH_3$), 3.24 (s, major, $-OCH_3$), 3.72-3.83 (m), 3.86 (s, minor, $-OCH_3$), 3.92 (s), 3.97-4.07 (m), 4.53 (dd, $J = 6.3, 1.7$ Hz, major, methine *H*), 5.08 (dd, $J = 6.4, 1.7$ Hz, minor, methine *H*), 6.46-6.54 (m), 6.61-6.73 (m), 6.73-6.83 (m), 6.86 (d, $J = 8.2$ Hz),

6.92 (dd, $J = 8.0, 1.7$ Hz), 6.96 (dd, $J = 7.4, 0.7$ Hz), 7.10 (d, $J = 7.7$ Hz), 7.13-7.34 (m); ^{13}C NMR (CDCl_3) δ 19.9, 20.5, 20.8, 21.00, 21.03, 50.7, 54.0, 54.6, 55.5, 56.2, 58.2, 67.4, 100.9, 106.2, 106.6, 108.9, 112.5, 113.9, 117.8, 126.8, 127.0, 127.1, 127.4, 127.7, 127.9, 128.2, 128.7, 129.1, 129.3, 129.8, 130.3, 132.3, 132.8, 132.9, 134.9, 135.2, 135.6, 135.8, 136.0, 136.3, 138.2, 155.2, 191.7; HRMS Calcd for $\text{C}_{24}\text{H}_{24}\text{O}_3$: M^+ , 360.1725. Found: m/z 360.1726.

^1H NMR integration ratio

1.93 (s) : 1.99 (s) : 2.13 (s) : 2.15 (s) : 2.30 (s) : 3.08 (s) : 3.24 (s) : 3.72-3.83 (m) : 3.86 (s) : 3.92 (s) : 3.97-4.07 (m) : 4.53 (dd) : 5.08 (dd) : 6.46-6.54 (m) : 6.61-6.73 (m) : 6.73-6.83 (m) : 6.86 (d) : 6.92 (dd) : 6.96 (dd) : 7.10 (d) : 7.13-7.34 (m) = 0.44 : 3.21 : 2.58 : 0.85 : 3.52 : 0.36 : 3.09 : 3.34 : 0.57 : 0.16 : 0.30 : 1.00 : 0.10 : 0.46 : 1.33 : 2.22 : 1.40 : 1.20 : 1.32 : 0.21 : 2.78.

4,5-Dimethoxy-9-(2,4,6-trimethylphenyl)xanthene (3'ej**).** Isolated in 16 % yield as a white solid; m.p. 269–276 °C; ^1H NMR (CDCl_3) δ 1.90 (brs, 3 H), 2.19 (s, 3 H), 2.75 (brs, 3 H), 3.59 (s, 6 H), 5.80 (s, 1 H), 6.46 (dd, $J = 8.0, 1.0$ Hz, 2 H), 6.57 (brs, 1 H), 6.67 (dd, $J = 8.2, 1.0$ Hz, 2 H), 6.83 (brs, 1 H), 7.11 (t, $J = 8.2$ Hz, 2 H); ^{13}C NMR (CDCl_3) δ 20.7, 30.9, 55.4, 105.4, 108.3, 112.5, 127.6, 134.6, 137.7, 152.3, 158.2; HRMS Calcd for $\text{C}_{24}\text{H}_{24}\text{O}_3$: M^+ , 360.1725. Found: m/z 360.1723.

Structure Determination of **3'ej**

The structure of **3'ej** was determined by NOE in ^1H NMR. As shown below, irradiation of the methine proton of **3'ej** did not enhance the MeO moiety, however, the absence of NOE between the methine proton and H_a as well as the observed NOE in **3ej** would support the assignment of the stereochemistry.

Reaction of **2o with benzyne.** To a THF solution (2.0 mL) of **1a** (0.269 g, 0.90 mmol), **2o** (0.043 g, 0.15 mmol) and 18-crown-6 (0.476 g, 1.80 mmol) was added KF (0.105 g, 1.80 mmol), and the resulting mixture was stirred at 0 °C for 23 h before dilution with ethyl acetate. Then the mixture was filtered through a Celite plug, and concentrated. Silicagel column chromatography (hexane/ethyl acetate as an eluent) followed by gel permeation chromatography (chloroform as an eluent) gave 4,4'-dimethoxy-3,3'-di(9-xanthyl)diphenyl ether (**3ao**) (25% yield) as a waxy greenish yellow solid: ^1H NMR (CDCl_3) δ 3.74 (s, 6 H), 5.66 (s, 2 H), 6.59-

6.67 (m, 4 H), 6.75 (d, $J = 9.4$ Hz, 2 H), 6.95 (td, $J = 7.5, 1.2$ Hz, 4 H), 7.07 (d, $J = 8.2$ Hz, 8 H), 7.17 (td, $J = 7.2, 1.4$ Hz, 4 H); ^{13}C NMR (CDCl_3) δ 37.7, 56.2, 112.3, 116.2, 117.0, 120.7, 123.0, 124.2, 127.6, 129.3, 136.5, 151.1, 151.4, 152.3; HRMS Calcd for $\text{C}_{40}\text{H}_{30}\text{O}_5$: M^+ , 590.2093. Found: m/z 590.2099.

References

- (1) Himeshima, Y.; Sonoda, T.; Kobayashi, H. *Chem. Lett.* **1983**, 1211–1214.
- (2) Yoshida, H.; Sugiura, S.; Kunai, A. *Org. Lett.* **2002**, 4, 2767–2769.
- (3) Peña, D.; Pérez, D.; Guitián, E.; Castedo, L. *J. Am. Chem. Soc.* **1999**, 121, 5827–5828.
- (4) Quideau, S.; Pouyse'gu, L.; Oxoby, M.; Looney, M. A. *Tetrahedron* **2001**, 57, 319–329.
- (5) Smith, J. R. L.; O'Brien, P.; Reginato, G. *Tetrahedron: Asymmetry* **1997**, 8, 3415–3420.
- (6) Kuhnert, N.; Rossignolo, G. M.; Lopez-Periago, A. *Org. Biomol. Chem.* **2003**, 1, 1157–1170.
- (7) Smith, D. W. Jr.; Babb, D. A.; Snelgrove, R. V.; Townsend, P. H. III; Martin, S. J. *J. Am. Chem. Soc.* **1998**, 120, 9078–9079.
- (8) McKinley, S. V.; Grieco, P. A.; Young, A. E.; Freedman, H. H. *J. Am. Chem. Soc.* **1970**, 92, 5900–5907.
- (9) Although **3ej** has a symmetrical 9-aryl moiety (2,4,6-trimethylphenyl) in contrast to other xanthenes that form a mixture of conformers, the asymmetric center at C-9 as well as a high rotational barrier about 9-aryl $\text{sp}^2\text{--sp}^3$ bond should make **3ej** a mixture of conformers unless the 9-aryl moiety is perpendicular (or parallel) to the xanthyl moiety. For rotational barriers in 9-arylxanthenes, see ref. 8.