Supporting Information

A Cascade Reaction Consisting of Pictet-Spengler-type Cyclization and Smiles Rearrangement: Application to the Synthesis of Novel Pyrrole-fused Dihydropteridines

*Jinbao Xiang, Lianyou Zheng, Feng Chen, Qun Dang, and Xu Bai**

The Center for Combinatorial Chemistry and Drug Discovery, Jilin University, 75
Haiwai Street, Changchun, Jilin 130012, P. R. China.

xbai@jlu.edu.cn

Table of Contents

Experimental Section-General Considerations	S2
Experimental Details and Spectroscopic Data	S2
Figure Legends	S9
Spectra	S14

* Tel: +86-431-85188955; Fax: +86-431-85188900; E-mail: xbai@jlu.edu.cn

Experimental Section

General considerations: Dichloromethane (DCM) was dried with P₂O₅ and distilled. All other commercial reagents were used as received without additional purification. Melting point was uncorrected. Mass spectra and HPLC data was recorded on an 1100 LC/MS system (Agilent Technology Corporation) with Alltech ELSD 2000. The ¹H and ¹³C NMR data were obtained on a Varian Mercury (300 MHz) or Varian INOVA (500 MHz) NMR spectrometer with TMS as the internal standard and CDCl₃ as solvent. Multiplicities are indicated as the following: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; dd, doubled doublet; br, broad. Coupling constants (*J* values) where noted are quoted in Hertz.

4,6-Dichloro-5-pyrrol-1-yl-pyrimidine 1. To a stirred solution of 4,6-dichloro-5-aminopyrimidine (15.0 g, 92.02 mmol) in acetic acid (150 mL), 2,5-dimethoxytetrahydrofuran (11.9 mL, 92.02 mmol) was added. The resulting solution was stirred for 1 h at reflux. Then acetic acid was evaporated and the residue was diluted with DCM and washed with water. The organic layers were dried over MgSO₄, filtered, and concentrated in vacuo. Purification by flash column chromatography (Petroleum ether/EtOAc 20:1, v/v) afforded 15.3g (78%) of **1** as a white solid. mp: 120-122 °C; ¹H NMR: δ 8.80 (s, 1H), 6.74 (t, 2H, *J* = 2.4), 6.45 (t, 2H, *J* = 2.1); MS (ESI): *m/z* 214.0 [M+H⁺]

6-[N-Benzyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine 2a: To a stirred solution of 4,6-dichloro-5-pyrrol-1-yl-pyrimidine (4.0 g, 18.78 mmol) and *N*-benzylglycinol (4.088 g, 27.05 mmol) in *n*-BuOH (100 mL) was added TEA (4.05 mL, 29.06 mmol). The resulting solution was stirred for 3.5 h at reflux. Then *n*-BuOH was evaporated and the residue was diluted with DCM and washed with water. The organic layers were dried over MgSO₄, filtered, and concentrated in vacuo. Purification by flash column chromatography (Petroleum ether/EtOAc 2:1, v/v) afforded 5.667g (92%) of **2a** as a white solid. mp: 71-73 °C; ¹H NMR: δ 8.39 (s, 1H), 7.31-7.24 (m, 3H), 7.06 (d, 2H, *J* = 8.1), 6.58 (t, 2H, *J* = 2.1), 6.28 (t, 2H, *J* = 2.1), 4.33 (s, 2H), 3.63 (q, 2H, *J*₁ = 5.4, *J*₂ = 5.1), 3.33 (t, 2H, *J* = 5.4) 2.44 (br, 1H); ¹³C NMR: δ 160.4, 160.0, 155.5, 137.2, 128.5, 127.2, 126.6, 122.6, 117.0, 110.5, 60.8, 52.9, 51.8; MS (ESI): *m/z* 329.1 [M+H⁺].

6-[N-Methyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine 2b: This compound was obtained by following the above procedure of **2a**. 92%; white solid; mp: 109-110 °C; ¹H NMR: δ 8.31 (s, 1H), 6.70 (d, 2H, *J* = 1.8), 6.35 (s, 2H), 3.81 (t, 2H, *J* = 4.8), 3.60 (t, 2H, *J* = 4.8), 3.28 (s, 1H), 2.61 (s, 3H); ¹³C NMR: δ 159.9, 155.3, 123.5, 116.8, 110.2, 109.6, 60.4, 54.6, 36.1; MS (ESI): *m/z* 253.1 [M+H⁺].

General procedure for the synthesis of pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4 and 5,6-dihydropyrrolo[1,2-*f*]pteridine 5: 6-[N-Benzyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine **2a** (164 mg, 0.5 mmol) in 6 mL DMSO:DCM (1:1) was treated with TEA (0.703 mL, 5.0 mmol). A solution of SO₃-Pyridine (481mg, 3.0 mmol) in 6 mL DMSO:DCM (1:1) was added, and the mixture was stirred for 35 min at ambient temperature. The reaction solution was then diluted with DCM (30 mL), washed twice with 10% citric acid and brine, dried over MgSO₄, and filtered to give crude aldehyde **3a** in DCM. To the above solution was added the amine (aromatic amine 0.6 mmol or aliphatic amine 3.0 mmol) and TFA (68 μL) (For aliphatic amines, additional 68 μL of TFA was added after stirring 3 h). The mixture was stirred for corresponding time at ambient temperature or 40 °C, then washed with saturated aqueous NaHCO₃, dried over MgSO₄, and concentrated in vacuo. Purification by flash chromatography (Petroleum ether/EtOAc = 4:1 or 2:1, v/v) afforded the desired products.

1-Chloro-5,6-dihydro-5-methyl-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine

5.1a: 30%. mp: 105-106 °C; ¹H NMR: δ 8.18 (s, 1H), 7.94 (t, 1H, *J* = 1.5), 7.30-7.27 (m, 2H), 7.23 (t, 1H, *J* = 7.5), 7.13 (d, 2H, *J* = 7.5), 6.39 (d, 1H, *J* = 1.0), 6.11 (t, 1H, *J* = 1.5), 4.70 (t, 1H, *J* = 4.5), 3.68 (A of AB, 1H, *J* = 13.5), 3.65 (B of AB, 1H, *J* = 13.0), 3.21 (s, 3H), 2.94-2.80 (m, 2H); ¹³C NMR: δ 154.4, 152.1, 141.5, 139.6, 128.4, 127.8, 127.0, 126.4, 119.6, 116.4, 111.3, 105.8, 58.1, 53.5, 51.3, 35.8; MS (ESI): *m/z* 340.1 [M+H⁺].

5-(*n*-Butyl)-1-chloro-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine

5.2a: 32%. Syrup; ¹H NMR: δ 8.14 (s, 1H), 7.91 (dd, 1H, *J*₁ = 3.0. *J*₂ = 1.5), 7.32-7.20 (m, 3H), 7.15 (d, 1H, *J* = 7.2), 6.38 (t, 1H, *J* = 3.3), 6.11 (dd, 1H, *J*₁ = 3.3. *J*₂ = 1.2), 4.72 (dd, 1H, *J*₁ = 7.5. *J*₂ = 4.2), 4.30-4.21 (m, 1H), 3.68 (s, 2H), 3.12-3.03 (m, 1H), 2.90-2.71 (m, 2H), 1.65-1.54 (m, 2H), 1.36-1.26 (m, 2H), 0.91 (t, 3H, *J* = 7.2); ¹³C NMR: δ 153.9,

152.1, 141.8, 139.6, 128.4, 127.8, 127.0, 126.4, 119.7, 116.1, 111.3, 105.7, 56.0, 53.5, 51.7, 47.3, 29.6, 20.0, 13.8; MS (ESI): m/z 382.2 $[M+H]^+$.

5-Benzyl-1-chloro-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.3a:

34%. syrup. 1H NMR: δ 8.19 (s, 1H), 7.94 (dd, 1H, $J_1 = 3.3$, $J_2 = 1.2$), 7.34-7.20 (m, 8H), 7.12 (d, 2H, $J = 8.1$), 6.37 (t, 1H, $J = 3.3$), 6.02 (dd, 1H, $J_1 = 3.6$, $J_2 = 1.5$), 5.57 (d, 1H, $J = 15.3$), 4.68 (dd, 1H, $J_1 = 6.6$, $J_2 = 2.1$), 4.35 (d, 1H, $J = 15.3$), 3.62 (s, 1H), 2.85-2.72 (m, 2H); ^{13}C NMR: δ 154.6, 152.3, 142.4, 140.0, 136.6, 129.0, 128.7, 128.13, 128.07, 128.01, 127.3, 126.9, 119.9, 116.7, 111.6, 106.1, 55.1, 53.7, 51.7, 50.4; MS (ESI): m/z 416.1 $[M+H]^+$.

1-Chloro-5-cyclohexyl-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine

5.4a: 35%. syrup; 1H NMR: δ 8.17 (s, 1H), 7.85 (dd, 1H, $J_1 = 3.3$, $J_1 = 1.5$), 7.32-7.20 (m, 3H), 7.16 (d, 2H, $J = 8.1$), 6.38 (t, 1H, $J = 3.3$), 6.09 (dd, 1H, $J_1 = 3.3$, $J_1 = 1.5$), 4.83 (dd, 1H, $J_1 = 9.0$, $J_2 = 3.9$), 4.53-4.50 (m, 1H), 3.69 (s, 1H), 2.78-2.61 (m, 2H), 1.98 (d, 1H, $J = 6.3$), 1.83 (br, 2H), 1.72-1.47 (m, 4H), 1.47-1.37 (m, 4H); ^{13}C NMR: δ 153.9, 152.0, 142.4, 139.7, 128.3, 127.7, 127.4, 127.0, 119.6, 116.3, 111.3, 105.1, 57.1, 53.3, 53.1, 50.8, 31.7, 30.7, 25.93, 25.86, 25.4; MS (ESI): m/z 408.0 $[M+H]^+$.

1-Chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(iso-propyl)pyrrolo[1,2-*f*]pteridine

5.5a: 42%. mp: 62-63 °C. 1H NMR: δ 8.17 (s, 1H), 7.86 (d, 1H, $J = 1.5$), 7.30-7.21 (m, 3H), 7.15 (d, 2H, $J = 7.0$), 6.38 (d, 1H, $J = 3.5$), 6.11 (s, 1H), 4.88-4.83 (m, 2H), 3.68 (s, 2H), 2.80-2.64 (m, 2H), 1.32 (d, 3H, $J = 7.0$), 1.28 (d, 3H, $J = 6.5$); ^{13}C NMR: δ 153.0, 151.0, 141.4, 138.6, 127.4, 126.8, 126.3, 126.1, 118.6, 115.4, 110.4, 104.3, 52.3, 52.1, 49.6, 48.1, 20.3, 19.4; MS (ESI): m/z 368.0 $[M+H]^+$.

1-Chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(2-methoxyphenyl)pyrrolo[1,2-*f*]

pteridine 5.6a: 61%. syrup. 1H NMR: δ 8.05 (s, 1H), 8.02 (dd, 1H, $J_1 = 3.0$, $J_2 = 1.2$), 7.37 (t, 1H, $J = 7.5$), 7.28-7.22 (m, 4H), 7.10 (d, 2H, $J = 6.3$), 7.02 (d, 2H, $J = 8.1$), 6.43 (t, 1H, $J = 3.3$), 6.12 (d, 1H, $J = 2.4$), 5.00 (t, 1H, $J = 4.8$), 3.74 (s, 3H), 3.62 (s, 2H), 2.88 (d, 2H, $J = 5.4$); ^{13}C NMR: δ 151.83, 151.79, 142.2, 139.9, 131.1, 129.5, 128.3, 127.8, 126.9, 126.8, 121.0, 119.64, 119.61, 116.0, 112.4, 111.4, 105.8, 57.0, 55.5, 53.5, 52.1; MS (ESI): m/z 432.0 $[M+H]^+$.

1-Chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-methoxyphenyl)pyrrolo[1,2-*f*]pteridine 5.7a: 40%. mp: 111-113 °C. ¹H NMR: δ 8.10 (s, 1H), 7.99 (dd, 1H, *J*₁ = 3.0, *J*₂ = 1.5), 7.30-7.23 (m, 3H), 7.17-7.10 (m, 4H), 6.96-6.92 (m, 2H), 6.44 (t, 1H, *J* = 3.3), 6.15 (d, 1H, *J* = 2.1), 5.01 (dd, 1H, *J*₁ = 6.6, *J*₂ = 3.6), 3.83 (s, 3H), 3.67 (A of AB, 1H, *J* = 13.2), 3.61 (B of AB, 1H, *J* = 13.2), 3.01-2.85 (m, 2H); ¹³C NMR: δ 158.6, 154.3, 152.0, 142.6, 139.7, 134.0, 128.8, 128.4, 127.9, 127.0, 126.6, 119.8, 116.4, 114.7, 111.6, 106.0, 59.3, 55.4, 53.5, 52.0; MS (ESI): *m/z* 432.1 [M+H⁺].

7-(2-Methylphenylamino)-5-benzyl-1-chloro-6,7-dihydropyrrolo[2,1-*d*][1,5]

benzodiazepine 4.8a: 86%. mp: 127-128 °C. ¹H NMR: δ 8.35 (s, 1H), 7.29-7.21 (m, 5H), 7.02-6.98 (m, 2H), 6.91 (t, 1H, *J* = 6.9), 6.65 (t, 1H, *J* = 7.2), 6.37 (d, 1H, *J* = 8.1), 6.27 (t, 1H, *J* = 3.3), 6.16 (t, 1H, *J* = 1.5), 4.90 (br, 2H), 4.79 (t, 1H, *J* = 5.1), 3.83-3.63 (m, 3H), 1.99 (s, 3H); ¹³C NMR: δ 157.1, 153.7, 153.4, 143.4, 136.6, 133.6, 130.3, 128.8, 127.9, 127.7, 127.1, 124.3, 122.7, 118.3, 116.8, 110.8, 109.1, 105.4, 57.7, 55.0, 48.9, 17.3; MS (ESI): *m/z* 416.1 [M+H⁺].

1-Chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(3-methylphenyl)pyrrolo[1,2-*f*]pteridine 5.9a:

31%. mp: 103-105 °C. ¹H NMR: δ 8.11 (s, 1H), 7.98 (dd, 1H, *J*₁ = 3.0, *J*₂ = 1.5), 7.33-7.22 (m, 4H), 7.14-7.05 (m, 5H), 6.43 (t, 1H, *J* = 3.3), 6.14 (dd, 1H, *J*₁ = 3.3, *J*₂ = 0.9), 5.04 (dd, 1H, *J*₁ = 6.6, *J*₂ = 3.6), 3.68 (A of AB, 1H, *J* = 13.5), 3.62 (B of AB, 1H, *J* = 13.5), 3.01-2.85 (m, 2H), 2.35 (s, 3H); ¹³C NMR: δ 153.9, 152.0, 142.8, 141.4, 139.6, 139.5, 129.2, 128.34, 128.25, 127.9, 127.5, 127.0, 126.6, 124.1, 119.7, 116.6, 111.6, 106.0, 58.9, 53.4, 51.9, 21.4; MS (ESI): *m/z* 416.2 [M+H⁺].

7-(4-Methylphenylamino)-5-benzyl-1-chloro-6,7-dihydropyrrolo[2,1-*d*][1,5]

pyrimidodiazepine 4.10a: 26%. mp: 175-176 °C. ¹H NMR: δ 8.34 (s, 1H), 7.31-7.22 (m, 5H), 6.98 (dd, 1H, *J*₁ = 3.0, *J*₂ = 1.5), 6.88 (d, 2H, *J* = 7.5), 6.40 (d, 2H, *J* = 8.4), 6.25 (t, 1H, *J* = 3.3), 6.12 (dd, 1H, *J*₁ = 3.6, *J*₂ = 0.9), 4.96 (br, 1H), 4.83 (br, 1H), 4.69 (br, 1H), 3.76 (dd, 1H, *J*₁ = 12.0, *J*₂ = 2.1), 3.63 (br, 2H), 2.20 (s, 3H); ¹³C NMR: δ 156.9, 153.7, 143.4, 136.5, 133.9, 129.7, 128.7, 128.0, 127.8, 127.7, 124.1, 114.0, 109.71, 109.68, 109.0, 105.2, 58.3, 55.0, 49.6, 20.3; MS (ESI): *m/z* 416.1 [M+H⁺].

1-Chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-methylphenyl)pyrrolo[1,2-*f*]pteridine 5.10a: 58%. mp: 113-114 °C. ¹H NMR: δ 8.10 (s, 1H), 7.98 (dd, 1H, *J*₁ = 3.3, *J*₂ = 1.5), 7.31-7.21 (m, 5H), 7.15-7.10 (m, 4H), 6.43 (t, 1H, *J* = 3.3), 6.13 (dd, 1H, *J*₁ = 3.6, *J*₂ = 1.5), 5.02 (dd, 1H, *J*₁ = 6.6, *J*₂ = 4.2), 3.67 (A of AB, 1H, *J* = 13.2), 3.61 (B of AB, 1H, *J* = 13.5), 3.01-2.86 (m, 2H), 2.38 (s, 3H); ¹³C NMR: δ 154.0, 151.8, 142.6, 139.6, 138.6, 137.3, 130.0, 128.3, 127.8, 127.0, 126.9, 126.5, 119.6, 116.4, 111.5, 105.9, 59.0, 53.3, 51.9, 21.0; MS (ESI): *m/z* 416.0 [M+H⁺].

1-Chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-phenylpyrrolo[1,2-*f*]pteridine 5.11a: 55%. mp: 91-92 °C. ¹H NMR: δ 8.10 (s, 1H), 7.97 (d, 1H, *J* = 1.5), 7.41 (t, 2H, *J* = 7.5), 7.33-7.19 (m, 6H), 7.10 (d, 2H, *J* = 6.9), 6.42 (t, 1H, *J* = 3.3), 6.13 (d, 1H, *J* = 2.7), 5.05 (dd, 1H, *J*₁ = 6.3, *J*₂ = 3.6), 3.66 (A of AB, 1H, *J* = 13.5), 3.60 (B of AB, 1H, *J* = 13.5), 3.00-2.85 (m, 2H); ¹³C NMR: δ 153.8, 151.8, 147.2, 142.9, 141.4, 139.6, 129.4, 128.3, 127.8, 127.3, 127.0, 126.6, 119.7, 116.7, 111.5, 106.0, 58.9, 53.4, 51.9; MS (ESI): *m/z* 402.2 [M+H⁺].

7-(2-Chlorophenylamino)-5-benzyl-1-chloro-6,7-dihydropyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.12a: 64%. mp: 120-121 °C. ¹H NMR: δ 8.36 (s, 1H), 7.31-7.27 (m, 5H), 7.21 (d, 1H, *J* = 7.5), 6.99 (s, 1H), 6.93 (t, 1H, *J* = 7.5), 6.63 (t, 1H, *J* = 7.5), 6.45 (br, 1H), 6.27 (d, 1H, *J* = 2.5), 6.19 (s, 1H), 4.92 (br, 2H), 4.78 (t, 1H, *J* = 6.5), 4.51 (s, 1H), 3.81-3.72 (m, 2H); ¹³C NMR: δ 156.9, 153.6, 153.4, 141.3, 136.4, 132.6, 129.1, 128.7, 127.8, 127.7, 127.6, 124.3, 119.7, 118.5, 116.5, 112.3, 109.0, 105.8, 57.6, 54.9, 48.8; MS (ESI): *m/z* 436.0 [M+H⁺].

1-Chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(3-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.13a: 38%. mp: 108-109 °C. ¹H NMR: δ 8.15 (s, 1H), 7.97 (dd, 1H, *J*₁ = 3.3, *J*₂ = 1.5), 7.36-7.23 (m, 6H), 7.19-7.11 (m, 3H), 6.44 (t, 1H, *J* = 3.3), 6.14 (dd, 1H, *J*₁ = 3.3, *J*₂ = 1.2), 5.05 (dd, 1H, *J*₁ = 6.0, *J*₂ = 3.9), 3.69 (A of AB, 1H, *J* = 13.5), 3.63 (B of AB, 1H, *J* = 13.2), 3.00-2.84 (m, 2H); ¹³C NMR: δ 153.5, 151.8, 143.3, 142.7, 139.5, 134.7, 130.1, 128.4, 127.8, 127.2, 127.1, 126.9, 126.5, 124.9, 119.7, 117.0, 111.6, 106.1, 58.8, 53.4, 51.9; MS (ESI): *m/z* 436.1 [M+H⁺].

1-Chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.14a: 62%. mp: 139-140 °C. ¹H NMR: δ 8.12 (s, 1H), 8.00-7.98 (m, 1H),

7.38-7.34 (m, 2H), 7.31-7.24 (m, 3H), 7.20-7.17 (m, 2H), 7.12-7.09 (m, 2H), 6.45 (t, 1H, $J = 3.3$), 6.14 (d, 1H, $J = 3.0$), 5.02 (dd, 1H, $J_1 = 6.0$, $J_2 = 3.9$), 3.68 (A of AB, 1H, $J = 13.2$), 3.60 (B of AB, 1H, $J = 13.5$), 2.98-2.83 (m, 2H); ^{13}C NMR: δ 153.7, 151.7, 143.1, 139.9, 139.6, 132.7, 129.5, 128.3, 128.3, 127.9, 127.0, 126.4, 119.7, 116.9, 111.6, 106.1, 58.9, 53.3, 51.8; MS (ESI): m/z 436.0 $[\text{M}+\text{H}^+]$.

1-Chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-florophenyl)pyrrolo[1,2-*f*]

pteridine 5.15a: 50%. mp: 136-138 °C. ^1H NMR: δ 8.10 (s, 1H), 8.00 (dd, 1H, $J_1 = 3.0$, $J_2 = 1.5$), 7.30-7.18 (m, 5H), 7.12-7.06 (m, 4H), 6.45 (t, 1H, $J = 3.3$), 6.14 (dd, 1H, $J_1 = 3.6$, $J_2 = 1.5$), 5.00 (dd, 1H, $J_1 = 6.0$, $J_2 = 3.9$), 3.68 (A of AB, 1H, $J = 13.5$), 3.60 (B of AB, 1H, $J = 13.2$), 2.99-2.84 (m, 2H); ^{13}C NMR: δ 162.9, 159.6, 154.0, 151.7, 142.8, 139.6, 137.2, 129.4, 129.2, 128.3, 128.0, 127.9, 127.0, 126.4, 119.7, 116.4, 116.1, 111.6, 106.0, 59.2, 53.4, 51.8; MS (ESI): m/z 420.1 $[\text{M}+\text{H}^+]$.

7-(2-Nitrophenylamino)-5-benzyl-1-chloro-6,7-dihydropyrrolo[2,1-*d*][1,5]

pyrimidodiazepine 4.16a: 32%. mp: 167-168 °C. ^1H NMR: δ 8.39 (s, 1H), 8.27 (br, 1H), 8.13 (d, 1H, $J = 8.4$), 7.31-7.20 (m, 5H), 7.00 (s, 1H), 6.66 (t, 1H, $J = 7.5$), 6.58 (br, 1H), 6.28-6.26 (m, 2H), 5.04 (br, 2H), 4.91 (s, 1H), 3.79 (s, 2H); ^{13}C NMR: δ 156.8, 153.89, 153.86, 142.9, 136.4, 136.1, 132.8, 131.7, 128.8, 128.1, 127.9, 126.8, 124.6, 116.6, 116.3, 114.1, 109.3, 106.2, 57.4, 55.0, 48.3; MS (ESI): m/z 447.0 $[\text{M}+\text{H}^+]$.

1-Chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-nitrophenyl)pyrrolo[1,2-*f*]

pteridine 5.17a: 43%. mp: 186-188 °C. ^1H NMR: δ 8.25 (s, 1H), 8.22 (d, 2H, $J = 9.0$), 7.95 (d, 1H, $J = 1.8$), 7.47 (d, 2H, $J = 9.3$), 7.34-7.26 (m, 3H), 7.16 (d, 2H, $J = 7.5$), 6.45 (t, 2H, $J = 3.6$), 6.18 (d, 1H, $J = 2.7$), 5.17 (t, 1H, $J = 6.6$), 3.70 (d, 1H, $J = 3.6$), 3.02-2.85 (m, 2H); ^{13}C NMR: δ 152.3, 151.4, 146.2, 144.8, 130.1, 129.8, 129.6, 129.4, 129.3, 124.9, 123.4, 122.8, 121.4, 118.1, 112.5, 109.3, 53.4, 51.5, 46.4; MS (ESI): m/z 447.0 $[\text{M}+\text{H}^+]$.

1-Chloro-5,6-dihydro-5-[benzylhydroxyl]-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]

pteridine 5.18a: 21%. mp: 90-91 °C. ^1H NMR: δ 8.37 (s, 1H), 7.92 (dd, 1H, $J_1 = 3.3$, $J_2 = 1.2$), 7.48-7.45 (m, 2H), 7.38-7.36 (m, 3H), 7.27-7.16 (m, 3H), 7.06 (d, 2H, $J = 6.3$), 6.32 (t, 1H, $J = 3.3$), 6.10 (d, 1H, $J = 2.4$), 5.12 (d, 1H, $J = 10.8$), 4.99 (d, 1H, $J = 10.8$), 4.77 (t, 1H, $J = 4.2$), 3.61 (A of AB, 1H, $J = 13.5$), 3.58 (B of AB, 1H, $J = 13.5$), 3.13-

3.00 (m, 1H); ^{13}C NMR: δ 156.2, 152.5, 142.0, 139.5, 135.1, 129.4, 128.8, 128.6, 128.3, 127.7, 127.0, 126.9, 119.8, 116.8, 111.0, 107.1, 77.2, 59.1, 53.6, 50.2; MS (ESI): m/z 432.1 $[\text{M}+\text{H}^+]$.

1-Chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-methylphenyl)pyrrolo[1,2-*f*]pteridine 5.10b: 51%. mp: 122-124 °C. ^1H NMR: δ 8.11 (s, 1H), 7.98-7.97 (m, 1H), 7.28 (d, 1H, $J = 8.7$), 7.24 (d, 1H, $J = 8.4$), 6.44 (t, 1H, $J = 3.3$), 6.15 (d, 1H, $J = 3.0$), 5.02 (dd, 1H, $J_1 = 7.1$, $J_2 = 4.2$), 2.94-2.77 (m, 2H), 2.39 (s, 3H), 2.33 (s, 3H); ^{13}C NMR: δ 154.0, 152.4, 143.3, 139.1, 137.8, 130.5, 127.2, 126.6, 120.1, 116.5, 111.9, 106.5, 59.0, 54.7, 36.6, 21.4; MS (ESI): m/z 340.1 $[\text{M}+\text{H}^+]$.

1-Chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.14b: 36%. mp: 86-87 °C. ^1H NMR: δ 8.13 (s, 1H), 7.97 (d, 1H, $J = 1.8$), 7.43 (d, 1H, $J = 8.7$), 7.33 (d, 1H, $J = 8.7$), 6.44 (t, 1H, $J = 3.3$), 6.16 (d, 1H, $J = 2.7$), 5.03 (dd, 1H, $J_1 = 7.8$, $J_2 = 4.5$), 2.88-2.78 (m, 2H), 2.33 (s, 3H); ^{13}C NMR: δ 153.7, 152.2, 143.7, 140.3, 133.1, 129.9, 128.5, 126.5, 120.2, 117.0, 112.0, 106.7, 58.9, 54.7, 36.6; MS (ESI): m/z 360.0 $[\text{M}+\text{H}^+]$.

5-Benzyl-1-chloro-6,7-dihydro-7-hydroxypyrrolo[2,1-*d*][1,5]pyrimido- diazepine 6a: 6-[*N*-Benzyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine **2a** (164 mg, 0.5 mmol) in 6 mL DMSO:DCM (1:1) was treated with TEA (0.703 mL, 5.0 mmol). A solution of SO_3 -Pyridine (481 mg, 3.0 mmol) in 6 mL DMSO:DCM (1:1) was added, and the mixture was stirred for 35 min at ambient temperature. The reaction solution was diluted with DCM (30 mL), washed twice with 10% citric acid and brine, dried over MgSO_4 , filtered, and concentrated in vacuo to give crude product **3a**. Crude product **3a** was dissolved in DCM (0.5 mL) and mixed with silica gel (0.6 g). The mixture was stirred overnight in air. Purification by flash chromatography (Petroleum ether/EtOAc= 4:1, v/v) afforded 74mg (45%) of **6a** as a white solid. mp: 171-173 °C. ^1H NMR: δ 8.26 (s, 1H), 7.36-7.24 (m, 5H), 7.01 (dd, 1H, $J_1 = 3.0$, $J_2 = 1.8$), 6.23 (t, 1H, $J = 3.0$), 6.16 (dd, 1H, $J_1 = 3.6$, $J_2 = 1.8$), 5.10 (d, 1H, $J = 15.3$), 5.00 (dd, 1H, $J_1 = 5.1$, $J_2 = 2.1$), 4.69 (d, 1H, $J = 15.3$), 3.81-3.67 (m, 2H); ^{13}C NMR: δ 157.8, 153.1, 153.0, 152.0, 136.4, 133.1, 128.6, 127.5, 127.3, 124.8, 108.7, 106.9, 64.3, 59.2, 55.0; MS (ESI): m/z 327.2 $[\text{M}+\text{H}^+]$.

Figure Legends:

Fig. S-1: LC-MS-ELSD of 6-[*N*-benzyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine 2a.

Fig. S-2: ¹H Spectra of 6-[*N*-benzyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine 2a.

Fig. S-3: ¹³C Spectra of 6-[*N*-benzyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine 2a.

Fig. S-4: LC-MS-ELSD of 6-[*N*-methyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine 2b.

Fig. S-5: ¹H Spectra of 6-[*N*-methyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine 2b.

Fig. S-6: ¹³C Spectra of 6-[*N*-methyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine 2b.

Fig. S-7: LC-MS-ELSD of 1-chloro-5,6-dihydro-5-methyl-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.1a.

Fig. S-8: ¹H Spectra of 1-chloro-5,6-dihydro-5-methyl-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.1a.

Fig. S-9: ¹³C Spectra of 1-chloro-5,6-dihydro-5-methyl-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.1a.

Fig. S-10: LC-MS-ELSD of 5-(*n*-butyl)-1-chloro-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.2a.

Fig. S-11: ¹H Spectra of 5-(*n*-butyl)-1-chloro-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.2a..

Fig. S-12: ¹³C Spectra of 5-(*n*-butyl)-1-chloro-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.2a..

Fig. S-13: LC-MS-ELSD of 5-benzyl-1-chloro-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.3a.

Fig. S-14: ¹H Spectra of 5-benzyl-1-chloro-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.3a.

Fig. S-15: ¹³C Spectra of 5-benzyl-1-chloro-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.3a.

Fig. S-16: LC-MS-ELSD of 1-chloro-5-cyclohexyl-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.4a.

Fig. S-17: ¹H Spectra of 1-chloro-5-cyclohexyl-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.4a.

Fig. S-18: ¹³C Spectra of 1-chloro-5-cyclohexyl-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.4a.

Fig. S-19: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(isopropyl)pyrrolo[1,2-*f*]pteridine 5.5a.

Fig. S-20: ¹H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(isopropyl)pyrrolo[1,2-*f*]pteridine 5.5a.

Fig. S-21: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(isopropyl)pyrrolo[1,2-*f*]pteridine 5.5a.

Fig. S-22: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(2-methoxyphenyl)pyrrolo[1,2-*f*]pteridine 5.6a.

Fig. S-23: ¹H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(2-methoxyphenyl)pyrrolo[1,2-*f*]pteridine 5.6a.

Fig. S-24: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(2-methoxyphenyl)pyrrolo[1,2-*f*]pteridine 5.6a.

Fig. S-25: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-methoxyphenyl)pyrrolo[1,2-*f*]pteridine 5.7a.

Fig. S-26: ¹H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-methoxyphenyl)pyrrolo[1,2-*f*]pteridine 5.7a.

Fig. S-27: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-methoxyphenyl)pyrrolo[1,2-*f*]pteridine 5.7a.

Fig. S-28: LC-MS-ELSD of 7-(2-methylphenylamino)-5-benzyl-1-chloro-6,7-dihydropyrrolo[2,1-*d*][1,5]benzodiazepine 4.8a.

Fig. S-29: ¹H Spectra of 7-(2-methylphenylamino)-5-benzyl-1-chloro-6,7-dihydropyrrolo[2,1-*d*][1,5]benzodiazepine 4.8a.

Fig. S-30: ¹³C Spectra of 7-(2-methylphenylamino)-5-benzyl-1-chloro-6,7-dihydropyrrolo[2,1-*d*][1,5]benzodiazepine 4.8a.

Fig. S-31: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(3-methylphenyl)pyrrolo[1,2-*f*]pteridine 5.9a.

Fig. S-32: ¹H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(3-methylphenyl)pyrrolo[1,2-*f*]pteridine 5.9a.

Fig.S-33: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(3-methylphenyl)pyrrolo[1,2-*f*]pteridine 5.9a.

Fig. S-34: LC-MS-ELSD of 7-(4-methylphenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.10a

Fig. S-35: ¹H Spectra of 7-(4-methylphenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.10a.

Fig. S-36: ¹³C Spectra of 7-(4-methylphenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.10a.

Fig. S-37: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-methylphenyl)pyrrolo[1,2-*f*]pteridine 5.10a

Fig. S-38: ¹H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-methylphenyl)pyrrolo[1,2-*f*]pteridine 5.10a.

Fig. S-39: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-methylphenyl)pyrrolo[1,2-*f*]pteridine 5.10a.

Fig. S-40: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-phenyl pyrrolo[1,2-*f*]pteridine 5.11a.

Fig. S-41: ¹H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-phenyl pyrrolo[1,2-*f*]pteridine 5.11a.

Fig. S-42: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-phenyl pyrrolo[1,2-*f*]pteridine 5.11a.

Fig. S-43: LC-MS-ELSD of 7-(2-chlorophenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.12a.

Fig. S-44: ¹H Spectra of 7-(2-chlorophenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.12a.

Fig. S-45: ¹³C Spectra of 7-(2-chlorophenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.12a.

Fig. S-46: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(3-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.13a.

Fig. S-47: ¹H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(3-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.13a.

Fig. S-48: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(3-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.13a.

Fig. S-49: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.14a.

Fig. S-50: ¹H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.14a.

Fig. S-51: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.14a.

Fig. S-52: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-fluorophenyl)pyrrolo[1,2-*f*]pteridine 5.15a.

Fig. S-53: ¹H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-fluorophenyl)pyrrolo[1,2-*f*]pteridine 5.15a.

Fig. S-54: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-fluorophenyl)pyrrolo[1,2-*f*]pteridine 5.15a.

Fig. S-55: LC-MS-ELSD of 7-(2-nitrophenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.16a.

Fig. S-56: ¹H Spectra of 7-(2-nitrophenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.16a.

Fig. S-57: ¹³C Spectra of 7-(2-nitrophenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.16a.

Fig. S-58: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-nitrophenyl)pyrrolo[1,2-*f*]pteridine 5.17a.

Fig. S-59: ¹H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-nitrophenyl)pyrrolo[1,2-*f*]pteridine 5.17a.

Fig. S-60: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-nitrophenyl)pyrrolo[1,2-*f*]pteridine 5.17a.

Fig. S-61: LC-MS-ELSD of 1-chloro-5,6-dihydro-5-[benzylhydroxyl]-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.18a.

Fig. S-62: ^1H Spectra of 1-chloro-5,6-dihydro-5-[benzylhydroxyl]-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.18a.

Fig. S-63: ^{13}C Spectra of 1-chloro-5,6-dihydro-5-[benzylhydroxyl]-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.18a.

Fig. S-64: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-methylphenyl)pyrrolo[1,2-*f*]pteridine 5.10b.

Fig. S-65: ^1H Spectra of 1-chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-methylphenyl)pyrrolo[1,2-*f*]pteridine 5.10b.

Fig. S-66: ^{13}C Spectra of 1-chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-methylphenyl)pyrrolo[1,2-*f*]pteridine 5.10b.

Fig. S-67: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.14b.

Fig. S-68: ^1H Spectra of 1-chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.14b.

Fig. S-69: ^{13}C Spectra of 1-chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.14b.

Fig. S-70: LC-MS-ELSD of 5-benzyl-1-chloro-6,7-dihydro-7-hydroxylpyrrolo[2,1-*d*][1,5]pyrimidodiazepine 6a.

Fig. S-71: ^1H Spectra of 5-benzyl-1-chloro-6,7-dihydro-7-hydroxylpyrrolo[2,1-*d*][1,5]pyrimidodiazepine 6a.

Fig. S-72: ^{13}C Spectra of 5-benzyl-1-chloro-6,7-dihydro-7-hydroxylpyrrolo[2,1-*d*][1,5]pyrimidodiazepine 6a.

2a
 $C_{17}H_{17}ClN_4O$
 Exact Mass: 328.11
 Mol. Wt.: 328.8

Fig. S-1: LC-MS-ELSD of 6-[N-benzyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine 2a.

Fig. S-2: ^1H Spectra of 6-[N-benzyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine 2a.

$C_{11}H_{13}ClN_4O$
 Exact Mass: 252.08
 Mol. Wt.: 252.7

2b

Fig. S-4: LC-MS-ELSD of 6-[N-methyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine 2b.

Fig. S-5: ^1H Spectra of 6-[N-methyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine 2b.

Fig. S-6: ^{13}C Spectra of 6-[*N*-methyl(2-ethanol)amino]-4-chloro-5-pyrrol-1-yl-pyrimidine 2b.

$C_{18}H_{18}ClN_5$
Exact Mass: 339.13
Mol. Wt.: 339.82

5.1a

Fig. S-7: LC-MS-ELSD of 1-chloro-5,6-dihydro-5-methyl-6-[(benzylamino)methyl]pyrrolo[1,2-f]pteridine 5.1a.

Fig. S-9: ¹³C Spectra of 1-chloro-5,6-dihydro-5-methyl-6-[(benzylamino)methyl]pyrrolo[1,2-f]pteridine 5.1a.

$C_{21}H_{24}ClN_5$
Exact Mass: 381.17
Mol. Wt.: 381.9

5.2a

Fig. S-10: LC-MS-ELSD of 5-(*n*-butyl)-1-chloro-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-*f*]pteridine 5.2a.

$C_{24}H_{22}ClN_5$
Exact Mass: 415.16
Mol. Wt.: 415.92

5.3a

Fig. S-13: LC-MS-ELSD of 5-benzyl-1-chloro-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-f]pteridine 5.3a.

Fig. S-14: ^1H Spectra of 5-benzyl-1-chloro-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-f]pteridine 5.3a.

$C_{23}H_{26}ClN_5$
Exact Mass: 407.19
Mol. Wt.: 407.94

5.4a

Fig. S-16: LC-MS-ELSD of 1-chloro-5-cyclohexyl-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-f]pteridine 5.4a.

Fig. S-18: ^{13}C Spectra of 1-chloro-5-cyclohexyl-5,6-dihydro-6-[(benzylamino)methyl]pyrrolo[1,2-f]pteridine 5.4a.

$C_{20}H_{22}ClN_5$
Exact Mass: 367.16
Mol. Wt.: 367.88

5.5a

Fig. S-19: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(isopropyl)pyrrolo[1,2-*f*]pteridine 5.5a.

Fig. S-21: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(isopropyl)pyrrolo[1,2-f]pteridine 5.5a.

$C_{24}H_{22}ClN_5O$
Exact Mass: 431.15
Mol. Wt.: 431.92

5.6a

Fig. S-22: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(2-methoxyphenyl)pyrrolo[1,2-f]pteridine 5.6a.

Fig. S-24: ^{13}C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(2-methoxyphenyl)pyrrolo[1,2-f]pteridine 5.6a.

$C_{24}H_{22}ClN_5O$
Exact Mass: 431.15
Mol. Wt.: 431.92

5.7a

Fig. S-25: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-methoxyphenyl)pyrrolo[1,2-f]pteridine 5.7a.

Fig. S-26: ^1H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-methoxyphenyl)pyrrolo[1,2-*f*]pteridine 5.7a.

Fig. S-27: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-methoxyphenyl)pyrrolo[1,2-*f*]pteridine 5.7a.

$C_{24}H_{22}ClN_5$
Exact Mass: 415.16
Mol. Wt.: 415.92

4.8a

Fig. S-28: LC-MS-ELSD of 7-(2-methylphenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-d][1,5]pyrimidodiazepine 4.8a.

Fig. S-30: ¹³C Spectra of 7-(2-methylphenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine.

Fig.S-33: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(3-methyl phenyl)pyrrolo[1,2-f]pteridine 5.9a.

$C_{24}H_{22}ClN_5$
 Exact Mass: 415.16
 Mol. Wt.: 415.92

4.10a

Fig. S-34: LC-MS-ELSD of 7-(4-methylphenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.10a.

Fig. S-35: ^1H Spectra of 7-(4-methylphenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.10a.

$C_{24}H_{22}ClN_5$
Exact Mass: 415.16
Mol. Wt.: 415.92

5.10a

Fig. S-37: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-methylphenyl)pyrrolo[1,2-f]pteridine 5.10a.

Fig. S-39: ^{13}C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-methylphenyl)pyrrolo[1,2-f]pteridine 5.10a.

$C_{23}H_{20}ClN_5$
Exact Mass: 401.14
Mol. Wt.: 401.89

5.11a

Fig. S-40: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-phenyl pyrrolo[1,2-f]pteridine 5.11a.

Fig. S-41: ^1H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-phenyl pyrrolo[1,2-f]pteridine 5.11a.

$C_{23}H_{19}Cl_2N_5$
Exact Mass: 435.1
Mol. Wt.: 436.34

4.12a

$C_{23}H_{19}Cl_2N_5$
Exact Mass: 435.1
Mol. Wt.: 436.34

5.13a

Fig. S-46: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(3-chlorophenyl)pyrrolo[1,2-f]pteridine 5.13a.

Fig. S-47: ¹H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(3-chlorophenyl)pyrrolo[1,2-f]pteridine 5.13a.

$C_{23}H_{19}Cl_2N_5$
Exact Mass: 435.1
Mol. Wt.: 436.34

5.14a

Fig. S-49: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-f]pteridine 5.14a.

Fig. S-50: ^1H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-f]pteridine 5.14a.

Fig. S-51: ^{13}C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-f]pteridine 5.14a.

$C_{23}H_{19}ClFN_5$
Exact Mass: 419.13
Mol. Wt.: 419.88

5.15a

Fig. S-52: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-fluorophenyl)pyrrolo[1,2-f]pteridine 5.15a.

Fig. S-53: ¹H Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-floro phenyl)pyrrolo[1,2-f]pteridine 5.15a.

Fig. S-54: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-floro phenyl)pyrrolo[1,2-f]pteridine 5.15a.

$C_{23}H_{19}ClN_6O_2$
Exact Mass: 446.13
Mol. Wt.: 446.89

4.16a

Fig. S-55: LC-MS-ELSD of 7-(2-nitrophenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.16a.

Fig. S-56: ^1H Spectra of 7-(2-nitrophenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.16a.

Fig. S-57: ¹³C Spectra of 7-(2-nitrophenylamino)-5-benzyl-1-chloro-6,7-dihydro pyrrolo[2,1-*d*][1,5]pyrimidodiazepine 4.16a.

$C_{23}H_{19}ClN_6O_2$
Exact Mass: 446.13
Mol. Wt.: 446.89

5.17a

Fig. S-58: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-nitrophenyl)pyrrolo[1,2-f]pteridine 5.17a.

Fig. S-60: ¹³C Spectra of 1-chloro-5,6-dihydro-6-[(benzylamino)methyl]-5-(4-nitrophenyl)pyrrolo[1,2-f]pteridine 5.17a.

$C_{24}H_{22}ClN_5O$
Exact Mass: 431.15
Mol. Wt.: 431.92

5.18a

Fig. S-61: LC-MS-ELSD of 1-chloro-5,6-dihydro-5-[benzylhydroxyl]-6-[(benzylamino)methyl]pyrrolo[1,2-f]pteridine 5.18a.

Fig. S-62: ^1H Spectra of 1-chloro-5,6-dihydro-5-[benzylhydroxyl]-6-[(benzylamino)methyl]pyrrolo[1,2-f]pteridine 5.18a.

Fig. S-63: ^{13}C Spectra of 1-chloro-5,6-dihydro-5-[benzylhydroxyl]-6-[(benzylamino)methyl]pyrrolo[1,2-f]pteridine 5.18a.

$C_{18}H_{18}ClN_5$
Exact Mass: 339.13
Mol. Wt.: 339.82

5.10b

Fig. S-64: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-methylphenyl)pyrrolo[1,2-f]pteridine 5.10b.

Fig. S-65: ^1H Spectra of 1-chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-methylphenyl)pyrrolo[1,2-f]pteridine 5.10b.

Fig. S-66: ^{13}C Spectra of 1-chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-methylphenyl)pyrrolo[1,2-f]pteridine 5.10b.

$C_{17}H_{15}Cl_2N_5$
Exact Mass: 359.07
Mol. Wt.: 360.24

5.14b

Fig. S-67: LC-MS-ELSD of 1-chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-f]pteridine 5.14b.

Fig. S-68: ^1H Spectra of 1-chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.14b.

Fig. S-69: ^{13}C Spectra of 1-chloro-5,6-dihydro-6-[(methylamino)methyl]-5-(4-chlorophenyl)pyrrolo[1,2-*f*]pteridine 5.14b.

6a
 $C_{17}H_{15}ClN_4O$
 Exact Mass: 326.09
 Mol. Wt.: 326.78

Fig. S-70: LC-MS-ELSD of 5-benzyl-1-chloro-6,7-dihydro-7-hydroxypyrrolo[2,1-d][1,5]pyrimidodiazepine 6a.

Fig. S-72: ¹³C Spectra of 5-benzyl-1-chloro-6,7-dihydro-7-hydroxylpyrrolo[2,1-*d*][1,5]pyrimidodiazepine 6a.