SUPPORTING INFORMATION

A Comprehensive Approach to the Synthesis of Sulfate Esters

Levi S. Simpson and Theodore S. Widlanski*

Contribution from the Department of Chemistry, Indiana University, Bloomington, Indiana

Contents	page
Synthesis of compounds	3-18
Details of optical rotation measurements	19
General procedures for direct assessment of protecting group stability	19-20
Procedure for reactivity study of tetrahydropyran-2-methyl- and isobutyl-protected sulfate monoesters	21
Line graph of rate study results	21
Spectroscopic characterization of compounds	22-67
References	68

Synthesis of compounds:

Neopentyl chlorosulfate.¹ A solution of freshly distilled sulfuryl chloride (10 mL, 0.124 mol) in 10 mL Et₂O was cooled to -75 °C. A solution of neopentyl alcohol (10.97 g, 0.124 mol) and pyridine (10 mL, 0.124 mol) in 25 mL Et₂O was added dropwise (over 1 h) to the cooled solution. The mixture was allowed to warm to room temperature and was stirred for an additional 2 h. The resulting white precipitate was filtered off and the filtrate was concentrated *in vacuo*. The resulting clear liquid was distilled under vacuum to yield neopentyl chlorosulfate (19.4 g, 84%): ¹H NMR (400 MHz, CDCl₃) δ 4.17 (s, 2H), 1.04 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ 85.59, 31.99, 25.94; IR (neat) 2967, 2908, 2874, 1478, 1410, 1371, 1189, 950, 856; HRMS (ESI) m/z calcd for C₅H₁₁O₃ClS (M⁺) 186.0117, found 186.0123.

Isobutyl chlorosulfate. A solution of freshly distilled sulfuryl chloride (15 mL, 0.187 mol) in 5 mL of THF was cooled to -75 °C. A solution of isobutyl alcohol (18 mL, 0.195 mol) in 5 mL of THF was added dropwise (over 1 h) to the cooled solution. The mixture was allowed to warm to room temperature, stirred for an additional 2 h, and then concentrated *in vacuo*. The resulting clear liquid was distilled under vacuum to yield isobutyl chlorosulfate (23.5 g, 73%): ¹H NMR (400 MHz, CDCl₃) δ 4.30 (d, *J* = 6.4 Hz, 2H), 2.16 (m, 1H), 1.05 (d, *J* = 6.8 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 82.51, 27.96, 18.59; IR (neat) 2971, 2880, 1471, 1410, 1192, 943, 849; HRMS (CI) m/z calcd for C₄H₁₀O₃ClS (M+H⁺) 173.0039, found 173.0038.

Phenyl neopentyl sulfate (1). Phenol (0.3043 g, 3.23 mmol) was dissolved in 4 mL of THF and 3 mL of DMPU, and the resulting solution was cooled to -75 °C. Sodium bis(trimethylsilyl)amide (0.93 M solution in THF, 3.8 mL, 3.53 mmol, 1.09 eq) was added dropwise to the cooled solution and stirred for 10 min. Neat neopentyl chlorosulfate (0.56 mL, 3.52 mmol, 1.09 eq) was added quickly to the reaction mixture. After 10 min, the reaction mixture was allowed to warm to room temperature. Upon completion (TLC), ethyl acetate and saturated aqueous NaHCO₃ were added. The organic layer was separated and washed with saturated aqueous NaHCO₃, water and brine, then dried over MgSO₄ and concentrated *in vacuo*. Purification by silica gel chromatography (10% ethyl acetate/hexanes) gave **1** as clear liquid (0.750 g, 95%): ¹H NMR (400 MHz, CDCl₃) δ 7.44-7.30 (m, 5H), 4.09 (s, 2H), 1.00 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ 150.42, 130.10, 127.42, 121.17, 83.54, 32.05, 26.07; IR (neat) 2964, 2873, 1587, 1489, 1410, 1370, 1206, 1175, 1149, 962, 880; HRMS (EI) m/z calcd for C₁₁H₁₆O₄S (M⁺) 244.0769, found 244.0776.

Estrone neopentyl sulfate (2). Estrone (0.3244 g, 1.10 mmol) was dissolved in 3.5 mL of THF and 2 mL of DMPU, and the resulting solution was cooled to -75 °C. Sodium bis(trimethylsilyl)amide (0.93 M solution in THF, 1.3 mL, 1.21 mmol, 1.1 eq) was added dropwise to the cooled solution and stirred for 10 min. Neat neopentyl chlorosulfate (0.2 mL, 1.26 mmol, 1.1 eq) was added quickly to the reaction mixture. After 10 min, the reaction mixture was allowed to warm to room temperature. Upon completion (TLC), ethyl acetate and saturated aqueous NaHCO₃ were added. The organic layer was separated and washed with saturated aqueous NaHCO₃, water and brine, then dried over

MgSO₄ and concentrated *in vacuo*. Purification by silica gel chromatography (20% ethyl acetate/hexanes) gave **2** as a white solid (0.453 g, 98%): ¹H NMR (400 MHz, CDCl₃) δ 7.30 (d, *J* = 8.4 Hz, 1H), 7.06-7.01 (m, 2H), 4.07 (s, 2H), 2.91-2.88 (m, 2H), 2.51-1.90 (m, 7H), 1.66-1.38 (m, 6H), 0.99 (s, 9H), 0.89 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 148.18, 139.02, 138.72, 126.81, 121.07, 118.12, 99.44, 83.23, 50.39, 47.87, 44.09, 37.87, 35.80, 31.92, 31.49, 29.37, 26.18, 25.98, 25.71, 21.56, 13.79; IR (film) 2933, 1739, 1490, 1408, 1193, 1143, 964, 920, 868; HRMS (EI) *m/z* calc for C₂₃H₃₂O₅S (M⁺) 420.1970, found 420.1961.

Benzyl N-carbobenzyloxy-L-tyrosine neopentyl sulfate (3). Benzyl *N*-carbobenzyloxy-L-tyrosine² (0.5001 g, 1.23 mmol) was dissolved in 3 mL of THF and 3 mL of DMPU, and the resulting solution was cooled to -75 °C. Sodium bis(trimethylsilyl)amide (0.93 M solution in THF, 2.0 mL, 1.86 mmol, 1.5 eq) was added dropwise to the cooled solution and stirred for 10 min. Neat neopentyl chlorosulfate (0.22 mL, 1.38 mmol, 1.12 eq) was added quickly to the reaction mixture. After 10 min, the reaction mixture allowed to warm to room temperature. Upon completion (TLC), ethyl acetate and saturate aqueous NaHCO₃ were added. The organic layer was separated and washed with saturated aqueous NaHCO₃, water and brine, then dried over MgSO₄ and concentrated *in vacuo*. Purification by silica gel chromatography (20% ethyl acetate/hexanes) gave **3** as viscous oil (0.675 g, 99%): ¹H NMR (400 MHz, CDCl₃) δ 7.41-7.32 (m, 10H), 7.17 (d, *J* = 8.8 Hz, 2H), 7.06 (d, *J* = 8.4 Hz, 2H) 5.58 (d, *J* = 8 Hz, 1H), 5.23-5.08 (m, 4H), 4.72 (dd, *J* = 6.4, 14 Hz, 1H), 4.10 (s, 2H), 3.20-3.06 (m, 2H), 1.03 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ 170.85, 155.44, 148.97, 135.93, 134.89,

134.71, 130.59, 128.44, 128.20, 127.98, 127.85, 127.18, 126.67, 120.77, 83.18, 67.12, 66.74, 54.50, 37.08, 31.62, 25.65; IR (film) 3034, 2962, 1725, 1504, 1399, 1208, 1182, 1152, 962, 884; HRMS (EI) m/z calcd for $C_{29}H_{34}NO_8S$ ($M+H^+$) 556.2005, found 556.2024.

1,2:5,6-Diisopropylidene- α -D-glucofuranose 3-neopentyl sulfate (4). Diacetone D-glucose (0.7051 g, 2.71 mmol) was dissolved in 4 mL of THF and 3 mL of DMPU, and the resulting solution was cooled to $-75\text{ }^{\circ}\text{C}$. Sodium bis(trimethylsilyl)amide (0.88 M solution in THF, 3.4 mL, 3.00 mmol, 1.1 eq) was added dropwise to the cooled solution and stirred for 10 min. Neat neopentyl chlorosulfate (0.47 mL, 2.95 mmol, 1.09 eq) was added quickly to the reaction mixture. After 10 min, the reaction mixture was allowed to warm to room temperature. Upon completion (TLC), ethyl acetate and saturated aqueous NaHCO_3 were added. The organic layer was separated and washed with saturated aqueous NaHCO_3 , water and brine, then dried over MgSO_4 and concentrated *in vacuo*. Purification by silica gel chromatography (20% ethyl acetate/hexanes) gave **4** as a white solid (1.051 g, 95%): ^1H NMR (400 MHz, CDCl_3) δ 5.93 (d, $J = 4.0$ Hz, 1H), 4.97 (d, $J = 4.0$ Hz, 1H), 4.91 (d, $J = 2.4$ Hz, 1H), 4.31–4.26 (m, 1H), 4.16–4.11 (m, 3H), 4.06–4.01 (m, 2H), 1.513 (s, 3H), 1.43 (s, 3H), 1.34 (s, 3H), 1.33 (s, 3H), 1.01 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ 112.64, 109.76, 104.93, 84.11, 82.71, 82.65, 79.81, 71.63, 67.45, 31.83, 27.11, 26.65, 26.23, 25.94, 25.38; IR (film) 2988, 1400, 1200, 1076, 1025; HRMS (EI) m/z calcd for $C_{16}H_{27}O_9S$ ($M-\text{CH}_3^+$) 395.1375, found 395.1373.

Phenyl isobutyl sulfate (5). Phenol (0.2016 g, 2.14 mmol) was dissolved in 40 mL of THF, and the resulting solution was cooled to -15 °C. Sodium bis(trimethylsilyl)amide (1.0 M solution in THF, 2.4 mL, 2.4 mmol, 1.1 eq) was added dropwise to the cooled solution and stirred for 10 min. Neat isobutyl chlorosulfate (1.5 mL, 10.58 mmol, 4.9 eq) was added quickly to the reaction mixture. After 10 min, the reaction mixture was allowed to warm to room temperature. Upon completion (TLC), ethyl acetate and saturated aqueous NaHCO₃ were added. The organic layer was separated and washed with saturated aqueous NaHCO₃, water and brine, then dried over MgSO₄ and concentrated in vacuo. Purification by silica gel chromatography (10% ethyl acetate/hexanes) gave **5** as a clear liquid (0.392 g, 80%): ¹H NMR (400 MHz, CDCl₃) δ 7.44-7.30 (m, 5H), 4.20 (d, *J* = 6.8 Hz, 2H), 2.09 (m, 1H), 0.99 (d, *J* = 6.8 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 150.41, 130.06, 127.41, 121.15, 80.38, 28.19, 18.61; IR (neat) 2968, 2878, 1587, 1489, 1408, 1207, 1149, 973, 879; HRMS (CI) *m/z* calc for C₁₀H₁₅O₄S (M+H⁺) 231.0686, found 231.0687.

Estrone isobutyl sulfate (6). Estrone (0.100 g, 0.37 mmol) was dissolved in 10 mL of THF, and the resulting solution was cooled to -15 °C. Sodium bis(trimethylsilyl)amide (0.99 M solution in THF, 0.41 mL, 0.406 mmol, 1.1 eq) was added dropwise to the cooled solution and stirred for 10 min. Neat isobutyl chlorosulfate (0.52 mL, 3.67 mmol, 9.9 eq) was added quickly to the reaction mixture. After 10 min, the reaction mixture was allowed to warm to room temperature. Upon completion (TLC), ethyl acetate and saturated aqueous NaHCO₃ were added. The organic layer was separated and washed with saturated aqueous NaHCO₃, water and brine, then dried over MgSO₄ and

concentrated in vacuo. Purification by silica gel chromatography (10 → 20% ethyl acetate/hexanes) gave **6** as a viscous oil (0.123 g, 82%): ¹H NMR (400 MHz, CDCl₃) δ 7.30 (d, *J* = 8.4 Hz, 1H), 7.06-7.02 (m, 2H), 4.20 (d, *J* = 6.4 Hz, 2H), 2.94-2.91 (m, 2H), 2.54-1.95 (m, 8H), 1.66-1.42 (m, 6H), 1.01 (d, *J* = 6.8 Hz, 6H), 0.91 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 220.57, 148.23, 139.12, 138.82, 126.89, 121.09, 118.15, 80.15, 77.48, 77.16, 76.84, 50.42, 47.92, 44.13, 37.91, 35.85, 31.55, 29.44, 28.13, 26.22, 25.77, 21.61, 18.60, 13.79; IR (film) 2931, 1738, 1489, 1404, 1194, 920; HRMS (EI) *m/z* calc for C₂₂H₃₁O₅S (M+H⁺) 407.1887, found 407.1878.

1,2:5,6-Diisopropylidene- α -D-glucofuranose 3-isobutyl sulfate (7). Diacetone D-glucose (0.5006 g, 1.92 mmol) was dissolved in 40 mL of THF and the resulting solution was cooled to -75 °C. Sodium bis(trimethylsilyl)amide (0.98 M solution in THF, 2.4 mL, 2.35 mmol, 1.22 eq) was added dropwise to the cooled solution and stirred for 10 min. Neat isobutyl chlorosulfate (1.1 mL, 7.76 mmol, 4.04 eq) was added quickly to the reaction mixture. After 1 h, the reaction mixture was allowed to warm to room temperature. Upon completion (TLC), ethyl acetate and saturated aqueous NaHCO₃ were added. The organic layer was separated and washed with saturated aqueous NaHCO₃, water and brine, then dried over MgSO₄ and concentrated *in vacuo*. Purification by silica gel chromatography (15% ethyl acetate/hexanes) gave **7** as white solid (0.726 g, 95%): ¹H NMR (400 MHz, CDCl₃) δ 5.92 (d, *J* = 3.6 Hz, 1H), 4.95 (d, *J* = 3.6 Hz, 1H), 4.92 (d, *J* = 2.8 Hz, 1H), 4.30–4.02 (m, 6H), 2.07 (m, 1H), 1.51 (s, 3H), 1.43 (s, 3H), 1.33 (s, 3H), 1.32 (s, 3H), 1.00 (d, *J* = 6.8 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 112.75, 109.82, 105.05, 84.24, 82.77, 79.89, 79.73, 71.80, 67.60, 28.10, 27.16, 26.75, 26.34, 25.41,

18.61; IR (film) 2985, 1406, 1384, 1199, 1077, 1025, 972, 949, 909, 879, 844; HRMS (EI) m/z calcd for $C_{16}H_{28}O_9S$ (M^+) 396.1454, found 396.1467.

1,2-Isopropylidene-3,5-dibenzyl- α -D-glucofuranose 3-isobutyl sulfate (8). 1,2-Isopropylidene-3,5-dibenzyl- α -D-glucofuranose³ (0.555 g, 1.39 mmol) was dissolved in 35 mL of THF, and the resulting solution was cooled to -75 °C. Sodium bis(trimethylsilyl)amide (0.98 M solution in THF, 1.7 mL, 1.67 mmol, 1.2 eq) was added dropwise to the cooled solution and stirred for 10 min. Neat isobutyl chlorosulfate (1.0 mL, 7.05 mmol, 5.1 eq) was added quickly to the reaction mixture. After 1 h, the reaction mixture was allowed to warm to room temperature. Upon completion (TLC), ethyl acetate and saturated aqueous $NaHCO_3$ were added. The organic layer was separated and washed with saturated aqueous $NaHCO_3$, water and brine, then dried over $MgSO_4$ and concentrated *in vacuo*. Purification by silica gel chromatography (10 $\rightarrow$ 20% ethyl acetate/hexanes) gave **8** as a viscous oil (0.637 g, 86%): 1H NMR (400 MHz, $CDCl_3$) δ 7.33-7.24 (m, 10H), 5.89 (d, J = 4.0 Hz, 1H), 4.73-4.62 (m, 4H), 4.48 (d, J = 2.8 Hz, 1H), 4.46 (d, J = 2.8 Hz, 1H), 4.36 (dd, J = 5.2, 10.8 Hz, 1H), 4.28 (dd, J = 3.2, 8.8 Hz, 1H), 4.14-3.99 (m, 2 H), 4.12 (d, J = 3.2 Hz, 2H) 2.01 (m, 1H), 1.47 (s, 3H), 1.31 (s, 3H), 0.94 (dd, J = 4.0, 6.8 Hz, 6H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 137.80, 137.32, 128.71, 128.58, 128.21, 127.99, 127.80, 127.73, 112.18, 105.25, 81.81, 81.61, 79.09, 78.52, 73.90, 72.85, 72.29, 72.15, 28.09, 26.98, 26.44, 18.61; IR (film) 2965, 2934, 1455, 1408, 1893, 1197, 1075, 1025, 943, 916, 867; HRMS (CI) m/z calcd for $C_{27}H_{36}O_9S$ (M^+) 536.2075, found 536.2087.

***N*-(9-fluorenylmethoxycarbonyl) L-tyrosine neopentyl sulfate (9).**⁴ Benzyl *N*-carbobenzyloxy-L-tyrosine neopentyl sulfate **3** (0.584 g, 1.05 mmol) was dissolved in 10 mL of methanol. To the resulting solution was added 10% Pd on activated carbon (0.0522 g). The flask was fitted with an H₂ balloon and the reaction was stirred 1.5 h. Upon completion (TLC), the reaction was filtered and concentrated *in vacuo*. The crude product was dissolved in 4 mL of dioxane and 8 mL of 10% aqueous Na₂CO₃. 9-Fluorenylmethyl chloroformate (0.2997 g, 1.159 mmol) was added and the reaction was stirred overnight. The reaction mixture was acidified (pH 3) and the product was extracted with ethyl acetate, dried over MgSO₄ and concentrated *in vacuo*. Purification by silica gel chromatography (10% methanol/CH₂Cl₂) gave **9** as white solid (0.317 g, 98%): ¹H NMR (400 MHz, MeOH-d₄) δ 7.73 (d, *J* = 7.6 Hz, 2H), 7.52 (m, 2H), 7.35-7.14 (m, 8H), 4.30 (m, 2H), 4.13-4.05 (m, 2H), 4.02 (s, 2H), 3.25-2.93 (m, 2H), 0.90 (s, 9H); ¹³C NMR (100 MHz, DMSO-d₆) δ 155.73, 148.15, 143.79, 140.67, 138.23, 130.88, 127.58, 127.02, 125.25, 125.17, 120.73, 120.07, 83.25, 65.47, 55.83, 46.61, 36.06, 31.55, 25.43; IR (film) 2961, 1690, 1575, 1504, 1406, 1206, 1150, 1049, 960, 883; HRMS (ESI) *m/z* calcd for C₂₉H₃₂O₈NS (M+H⁺) 554.1849, found 544.1858.

***N*-(*tert*-Butoxycarbonyl) L-tyrosine neopentyl sulfate (10).**⁵ Benzyl *N*-carbobenzyloxy-L-tyrosine neopentyl sulfate **3** (0.181 g, 0.326 mmol) was dissolved in 5 mL of methanol. To the resulting solution was added 10% Pd on activated carbon (0.020 g). The flask was fitted with an H₂ balloon and the reaction was stirred 2 h. Upon completion (TLC), the reaction was filtered and concentrated *in vacuo*. The crude product was dissolved in 3.5 mL of DMF and 0.5 mL of triethylamine. Di-*tert*-butyl

dicarbonate (0.1538 g, 0.705 mmol, 2.16 eq) was added with vigorous stirring. The solution was heated to 50 °C for 20 min and then stirred at room temperature for 1 h. The solvents were removed *in vacuo* and the residue was stirred with dilute HCl. The product was extracted with ethyl acetate, dried over MgSO₄ and concentrated *in vacuo*. Purification by silica gel chromatography (10% methanol/CH₂Cl₂) gave **10** as white solid (0.069 g, 88%): ¹H NMR (400 MHz, DMSO-d₆) δ 7.25-7.24 (m, 4H), 4.18 (s, 2H), 3.90 (m, 1H), 3.13-2.92 (m, 2H), 1.31 (s, 9H), 0.93 (s, 9H); ¹³C NMR (100 MHz, DMSO-d₆) δ 155.13, 148.15, 138.46, 130.91, 129.24, 127.98, 126.08, 120.71, 83.25, 77.80, 55.55, 55.40, 36.89, 36.30, 31.61, 28.19, 27.86, 25.49; IR (solid) 2969, 1692, 1504, 1394, 1368, 1206, 1151, 961, 882; HRMS (ESI) m/z calcd for C₁₉H₂₉O₈NNaS (M+Na⁺) 454.1506, found 454.1493.

D-Glucose 3-neopentyl sulfate (11). 1,2:5,6-Diisopropylidene- α -D-glucofuranose 3-neopentyl sulfate **4** (0.0448 g, 0.109 mmol) was dissolved in 2 mL of THF. A 50% (v/v) aqueous solution of H₂SO₄ (0.5 mL) was added and the solution was stirred vigorously and heated to 60 °C in an oil bath. After 2 h the entire solution was loaded onto a silica gel column and the product was eluted with 10% methanol/ethyl acetate to give **11** as a hygroscopic solid (0.034 g, 94%): ¹H NMR (400 MHz, MeOH-d₄) δ (α -anomer, 45%): 5.11 (d, *J* = 3.6 Hz, 1H), 4.72 (t, *J* = 9.2 Hz, 1H), (β -anomer, 55%): 4.50 (d, *J* = 8 Hz, 1H), 4.43 (t, *J* = 9.2 Hz, 1H), (both anomers): 4.07 (s, 2H), 3.84-3.27 (m, 5H), 0.96 (s, 9H); ¹³C NMR (100 MHz, MeOH-d₄) δ 98.03, 94.18, 90.92, 89.65, 83.11, 77.41, 74.49, 72.97, 71.98, 69.73, 69.63, 62.35, 62.17, 32.53, 26.31; IR (film) 3236, 2963, 2936, 1389,

1367 1190, 1091, 973, 884; HRMS (ESI) m/z calcd for $C_{11}H_{22}O_9NaS$ ($M+Na^+$) 353.0877, found 353.0891.

D-Glucose 3-isobutyl sulfate (12). 1,2:5,6-Diisopropylidene- α -D-glucofuranose 3-isobutyl sulfate **7** (0.0546 g, 0.138 mmol) was dissolved in 2 mL of THF. A 50% (v/v) aqueous solution of H_2SO_4 (0.4 mL) was added and the solution was stirred vigorously and heated to 60 °C in an oil bath. After 1.5 h the entire solution was loaded onto a silica gel column and the product was eluted with 10% methanol/ethyl acetate to give **12** as a hygroscopic solid (0.040 g, 92%): 1H NMR (400 MHz, $MeOH-d_4$) δ (α -anomer, 60%): 5.14 (d, $J = 3.6$ Hz, 1H), 4.73 (t, $J = 9.2$ Hz, 1H), (β -anomer, 40%): 4.53 (d, $J = 8$ Hz, 1H), 4.45 (t, $J = 9.2$ Hz, 1H), (both anomers): 4.18 (d, $J = 6.8$ Hz, 2H), 3.87-3.30 (m, 5H), 2.04 (m, 1H), 0.98 (d, $J = 6.8$ Hz, 6H); ^{13}C NMR (100 MHz, $MeOH-d_4$) δ 98.02, 94.18, 90.87, 89.56, 80.14, 80.10, 77.40, 74.47, 72.97, 71.97, 69.72, 69.63, 62.36, 62.18, 29.18, 18.85; IR (film) 3371, 2965, 1370, 1192, 1045, 946, 880; HRMS (CI) m/z calcd for $C_{10}H_{19}O_8S$ ($M-OH^+$) 299.0795, found 299.0793.

D-Glucose 6-isobutyl sulfate (13). 1,2-Isopropylidene-3,4-dibenzyl- α -D-glucofuranose 3-isobutyl sulfate **8** (0.1098 g, 0.205 mmol) was dissolved in 1.5 mL of THF. To the resulting solution was added 10% Pd on activated carbon (0.0248 g). The flask was fitted with an H_2 balloon and the reaction was stirred overnight. Upon completion (TLC), the reaction was filtered, concentrated *in vacuo*, and the crude product was dissolved in 3 mL of THF. A 50% (v/v) aqueous solution of H_2SO_4 (0.5 mL) was added and the solution was stirred vigorously and heated to 60 °C in an oil bath. After 0.5 h the entire solution

was loaded onto a silica gel column and the product was eluted with 10% methanol/ethyl acetate to give **13** as a hygroscopic solid (0.062 g, 96%): ^1H NMR (400 MHz, MeOH-d_4) δ (α -anomer, 65%): 5.13 (d, $J = 3.6$ Hz, 1H), 4.04 (m, 1H), 3.69 (t, $J = 9.6$ Hz, 1H), (β -anomer, 35%): 4.45 (d, $J = 2.0$ Hz, 1H) 3.57 (m, 1H), 3.16 (m, 1H), (both anomers): 4.53-4.33 (m, 2H), 4.13-4.09 (m, 2H), 3.40-3.27 (m, 2H), 2.06 (m, 1H), 1.01 (d, $J = 6.8$ Hz, 6H); ^{13}C NMR (100 MHz, MeOH-d_4) δ 98.23, 94.01, 80.21, 80.07, 77.85, 76.04, 74.82, 74.72, 73.62, 73.11, 72.78, 71.32, 71.06, 70.40, 29.17, 18.83; IR (film) 3357, 2967, 1389, 1195, 1061, 934, 875; HRMS (ESI) m/z calcd for $\text{C}_{10}\text{H}_{20}\text{O}_9\text{NaS}$ ($\text{M}+\text{Na}^+$) 339.0720, found 339.0711.

Phenyl sulfate (14). In a flask fitted with a reflux condenser, phenyl neopentyl sulfate **1** (0.0545 g, 0.223 mmol) was dissolved in 1 mL of DMF. Sodium azide (0.0199 g, 0.306 mmol, 1.3 eq) was added and the solution was stirred and heated to 70 $^\circ\text{C}$ in an oil bath overnight. Removal of the solvent *in vacuo* and purification of the crude product by silica gel chromatography (20% ethanol/ CH_2Cl_2) gave **14** as the sodium salt (0.041 g, 96%): ^1H NMR (400 MHz, D_2O) δ 7.34-7.17 (m, 5H); ^{13}C NMR (100 MHz, D_2O) δ 151.08, 129.89, 126.42, 121.58; IR (film) 2924, 2820, 1419, 1390, 1264, 1229, 1051, 870; HRMS (FAB) m/z calcd for $\text{C}_6\text{H}_5\text{O}_4\text{S}$ (M^+) 172.9909, found 172.9904.

Benzyl N-carbobenzyloxy-L-tyrosine sulfate (15). In a flask fitted with a reflux condenser, benzyl *N*-carbobenzyloxy-L-tyrosine neopentyl sulfate **3** (0.0730 g, 0.131 mmol) was dissolved in 1 mL of DMF. Sodium azide (0.0117 g, 0.180 mmol, 1.4 eq) was added and the solution was stirred and heated to 70 $^\circ\text{C}$ in an oil bath overnight. The

solvent was removed *in vacuo* and the crude product was dissolved in CHCl_3 and filtered to give **15** as the sodium salt (0.064 g, 96%): ^1H NMR (400 MHz, MeOH-d_4) δ 7.33-7.27 (m, 10H), 7.19 (d, $J = 8.4$ Hz, 2H), 7.12 (d, $J = 8.8$, 2H), 5.13 (s, 2H), 5.03 (s, 2H), 4.47-4.43 (m, 1H), 3.14-2.91 (m, 2H); ^{13}C NMR (100 MHz, MeOH-d_4) δ 173.21, 158.37, 152.84, 138.06, 137.01, 134.59, 130.97, 129.55, 129.43, 129.35, 129.30, 128.94, 128.66, 122.46, 68.00, 67.57, 57.08, 37.72; IR (solid) 3039, 2972, 1720, 1512, 1281, 1180, 942; HRMS (ESI) m/z calcd for $\text{C}_{24}\text{H}_{22}\text{O}_8\text{NNa}_2\text{S}$ ($\text{M}+2\text{Na}^+$) 530.0856, found 530.0860.

Estrone sulfate (16). In a flask fitted with a reflux condenser, estrone neopentyl sulfate **2** (0.1011 g, 0.240 mmol) was dissolved in 1 mL of DMF. Sodium azide (0.0213 g, 0.328 mmol, 1.4 eq) was added and the solution was stirred and heated to 70 °C in an oil bath overnight. Removal of the solvent *in vacuo* and purification of the crude product by silica gel chromatography (20% ethanol/ CH_2Cl_2) gave **16** as the sodium salt (0.088 g, 98%): ^1H NMR (400 MHz, MeOH-d_4) δ 7.24 (d, $J = 8.4$ Hz, 1H), 7.06-7.03 (m, 2H), 2.89-2.87 (m, 2H), 2.52-1.88 (m, 7H), 1.66-1.40 (m, 6H), 0.91 (s, 3H); ^{13}C NMR (100 MHz, MeOH-d_4) δ 151.76, 138.71, 137.51, 126.93, 122.48, 119.77, 45.46, 45.33, 39.59, 36.72, 32.79, 30.42, 27.52, 27.04, 22.52, 14.30, 14.21; IR (solid) 3453, 2930, 2867, 1724, 1493, 1227, 1049, 933; HRMS (ESI) m/z calcd for $\text{C}_{18}\text{H}_{21}\text{O}_5\text{Na}_2\text{S}$ ($\text{M}+2\text{Na}^+$) 395.0900, found 395.0909.

1,2:5,6-Diisopropylidene- α -D-glucofuranose 3-sulfate (17). In a flask fitted with a reflux condenser, 1,2:5,6-diisopropylidene- α -D-glucofuranose 3-neopentyl sulfate **4** (0.0829 g, 0.202 mmol) was dissolved in 1 mL of DMF. Sodium azide (0.0234 g, 0.360

mmol, 1.8 eq) was added and the solution was stirred and heated to 70 °C in an oil bath overnight. Removal of the solvent *in vacuo* and purification of the crude product by silica gel chromatography (20% ethanol/CH₂Cl₂) gave **17** as the sodium salt (0.072 g, 98%): ¹H NMR (400 MHz, D₂O) δ 6.09 (d, *J* = 3.6 Hz, 1H), 4.98 (d, *J* = 3.6 Hz, 1H), 4.84 (d, *J* = 1.6 Hz, 1H), 4.54–4.24 (m, 2H), 4.17–4.04 (m, 2H), 1.54 (s, 3H), 1.47 (s, 3H), 1.39 (s, 3H), 1.38 (s, 3H); ¹³C NMR (100 MHz, D₂O) δ 113.42, 109.47, 104.95, 82.85, 80.07, 79.71, 73.10, 64.99, 25.81, 25.49, 25.38, 23.90; IR (solid) 3496, 3451, 2986, 1401, 1238, 1221, 1032; HRMS (ESI) *m/z* calcd for C₁₂H₁₉O₉Na₂S (M+2Na⁺) 385.0540, found 385.0533.

3-azido-3-deoxy D-glucose (18). In a flask fitted with a reflux condenser, D-glucose 3-neopentyl sulfate **11** (0.0336 g, 0.102 mmol) was dissolved in 0.5 mL of DMF. Sodium azide (0.0102 g, 0.157 mmol, 1.5 eq) was added and the solution was stirred and heated to 70 °C in an oil bath overnight. The solvent was removed *in vacuo* for characterization as the crude material: ¹H NMR (400 MHz, D₂O) δ 4.81 (d, *J* = 8.0 Hz, 1H), 4.24 (t, *J* = 3.6 Hz, 1H), 3.86–3.81 (m, 2H), 3.73–3.65 (m, 2H), 3.57 (dd, *J* = 3.6, 8.4 Hz, 1H); IR (film) 3548, 2050 (N≡N⁺ stretch), 1653, 1223, 994; HRMS (FAB) *m/z* calcd for C₁₂H₁₉O₉Na₂S (M+H⁺) 206.0766, found 206.0748.

Neopentyl sulfate (19). Obtained as a secondary product of the previous reaction. ¹H NMR (400 MHz, D₂O) δ 3.71 (s, 2H), 0.94 (s, 9H); HRMS (FAB) *m/z* calcd for C₅H₁₁O₄Na₂S (M+2Na⁺) 213.0168, found 213.0171.

D-Glucose 3-sulfate (20). In a flask fitted with a reflux condenser, D-glucose 3-isobutyl sulfate **12** (0.038 g, 0.120 mmol) was dissolved in 4 mL of acetone. Sodium iodide (0.0544 g, 0.363 mmol, 3.0 eq) was added and the solution was stirred and heated to 55 °C in an oil bath for 5 h. The precipitated product was washed several times with cold acetone to give **20** as a hygroscopic solid (0.033 g, 97%): ¹H NMR (400 MHz, D₂O) δ (α-anomer, 45%): 5.30 (d, *J* = 4.0 Hz, 1H), 4.51 (t, *J* = 9.6 Hz, 1H), 3.45 (m, 1H), (β-anomer, 55%): 4.77-4.74 (m, 1H), 4.33 (t, *J* = 9.2 Hz, 1H) 3.58-3.54 (m, 1H), (both anomers): 3.95-3.72 (m, 3H), 3.64 (t, *J* = 9.6 Hz, 1H); ¹³C NMR (100 MHz, D₂O) δ 96.45, 92.84, 85.11, 83.05, 76.22, 73.68, 71.99, 71.00, 69.13, 69.06, 61.48, 61.31; IR (solid) 3362, 1637, 1215, 1055, 999, 934, 816; HRMS (ESI) *m/z* calcd for C₆H₁₁O₉Na₂S (M+2Na⁺) 304.9914, found 304.9910.

D-Glucose 6-sulfate (21). In a flask fitted with a reflux condenser, D-glucose 6-isobutyl sulfate **13** (0.057 g, 0.180 mmol) was dissolved in 4 mL of acetone. Sodium iodide (0.0423 g, 0.282 mmol, 1.57 eq) was added and the solution was stirred and heated to 55 °C in an oil bath for 5 h. The precipitated product was washed several times with cold acetone to give **21** as a hygroscopic solid (0.055 g, 96%): ¹H NMR (400 MHz, D₂O) δ (α-anomer, 40%): 5.26 (d, *J* = 3.6 Hz, 1H), 3.75 (t, *J* = 9.6 Hz, 1H), 3.60-3.57 (m, 1H), (β-anomer, 60%): 4.70 (d, *J* = 7.6 Hz, 1H), 4.07-4.04 (m, 1H), 3.30 (t, *J* = 8.4 Hz, 1H), (both anomers): 4.37-4.21 (m, 3H), 3.73-3.69 (m, 1H); ¹³C NMR (100 MHz, D₂O) δ 96.83, 93.01, 76.40, 74.82, 74.59, 73.47, 72.18, 70.33, 70.13, 67.97; IR (solid) 3375, 1637, 1215, 1055, 999, 934, 816; HRMS (ESI) *m/z* calcd for C₆H₁₁O₉Na₂S (M+2Na⁺) 304.9914, found 304.9917.

Estrone sulfate (22). In a flask fitted with a reflux condenser, estrone isobutyl sulfate **6** (0.0420 g, 0.103 mmol) was dissolved in 1 mL of acetone. Sodium thiocyanate (0.0177 g, 0.218 mmol, 2.1 eq) and triethylamine (0.04 mL) were added and the solution was stirred and heated to 55 °C in an oil bath for 2 h. The precipitated product was washed several times with cold acetone to give **22** as a white solid (0.037 g, 96%): ¹H NMR, ¹³C NMR, IR, and HRMS identical to estrone sulfate **16**.

Tetrahydropyran-2-methyl chlorosulfate (23). A solution of freshly distilled sulfuryl chloride (1.7 mL, 21.16 mmol) in 5 mL of THF was cooled to -75 °C. A solution of tetrahydropyran-2-methanol (2.0 mL, 17.68 mmol) in 10 mL of THF was added dropwise (over 0.5 h) to the cooled solution. The mixture was allowed to warm to room temperature, stirred for an additional 0.5 h and the solvent was removed *in vacuo*. Purification by silica gel chromatography (10% ethyl acetate/hexanes) gave **23** as a clear liquid (3.683 g, 81%): ¹H NMR (400 MHz, CDCl₃) δ; 4.40 (d, *J* = 4.8 Hz, 2H), 4.04-4.00 (m, 1H), 3.72-3.66 (m, 1H), 3.48-3.42 (m, 1H), 1.93-1.89 (m, 1H), 1.62-1.52 (m, 4H), 1.49-1.34 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ; 78.22, 74.14, 68.52, 27.29, 25.45, 22.78; IR (neat) 2944, 2854, 1404, 1189, 1103, 1076, 1049, 960, 845; HRMS (CI) *m/z* calcd for C₆H₁₂O₄ClS (M+H⁺) 215.0139, found 215.0135.

1,2:5,6-diisopropylidene- α -D-glucofuranose 3-tetra hydropyran-2-methyl sulfate (24). Diacetone D-glucose (0.2500 g, 0.961 mmol) was dissolved in 15 mL of THF, and the resulting solution was cooled to -15 °C. Sodium bis(trimethylsilyl)amide (1.0 M

solution in THF, 1.0 mL, 1.00 mmol, 1.04 eq) was added dropwise to the cooled solution and stirred for 10 min. Neat tetrahydropyran-2-methyl chlorosulfate (0.214 g, 0.988 mmol, 1.03 eq) was added quickly to the reaction mixture. After 0.5 h, the reaction mixture was allowed to warm to room temperature. Upon completion (TLC), ethyl acetate and saturated aqueous NaHCO₃ were added. The organic layer was separated and washed with saturated aqueous NaHCO₃, water and brine, then dried over MgSO₄ and concentrated in vacuo. Purification by silica gel chromatography (20% ethyl acetate/hexanes) gave **24** as a white solid (diastereomers, 0.361 g, 86%): IR (film) 2931, 1738, 1489, 1404, 1194, 920; HRMS (EI) m/z calc for C₁₈H₃₁O₁₀S (M+H⁺) 439.1632, found 439.1639.

Details of optical rotation measurements:

Tyrosine neopentyl sulfate derivative **3** was desulfated by removal of the neopentyl protecting group with NaN₃/DMF and hydrolysis of the resulting sulfate monoester in aqueous acid to give benzyl *N*-carbobenzyloxy-L-tyrosine ($[\alpha]^{22}_{\text{D}} -14.17^\circ$). This optical rotation was compared with the optical rotation of pure benzyl *N*-carbobenzyloxy-L-tyrosine ($[\alpha]^{22}_{\text{D}} -14.26^\circ$) to determine that **3** had undergone less than 1% racemization.

General procedures for direct assessment of protecting group stability:

Assays in 50% TFA. A stock solution was prepared by dissolving the model compound (0.5 mmol) in CDCl₃ (1 ml). An aliquot of the stock solution (250 μL) was transferred to an NMR tube. Distilled TFA (250 μL) was added and the tube was capped and sealed. ¹H NMR spectra were obtained periodically to follow the degradation progress.

Assays in 10% TFA. A stock solution was prepared by dissolving the model compound (0.5 mmol) in CDCl₃ (1 ml). An aliquot of the stock solution (100 μL) was transferred to an NMR tube. CDCl₃ (350 μL) and distilled TFA (50 μL) were added and the tube was capped and sealed. ¹H NMR spectra were obtained periodically to follow the degradation progress.

Assays in 20% piperidine. A stock solution was prepared by dissolving the model compound (0.5 mmol) in CDCl₃ (1 ml). An aliquot of the stock solution (300 μL) was

transferred to an NMR tube. CDCl_3 (100 μL) and distilled piperidine (100 μL) were added and the tube was capped and sealed. ^1H NMR spectra were obtained periodically to follow the degradation progress.

Assays in 6% piperidine. A stock solution was prepared by dissolving the model compound (0.5 mmol) in CDCl_3 (1 ml). An aliquot of the stock solution (100 μL) was transferred to an NMR tube. CDCl_3 (370 μL) and distilled piperidine (30 μL) were added and the tube was capped and sealed. ^1H NMR spectra were obtained periodically to follow the degradation progress.

Procedure for reactivity study of tetrahydropyran-2-methyl- and isobutyl-protected sulfate monoesters:

Each model compound (0.1 mmol) was dissolved in acetone- d_6 (1 mL) and placed in separate NMR tubes. Sodium iodide (0.2 mmol) was added to each NMR tube and the tubes were capped and sealed. ^1H NMR spectra were obtained periodically to follow the reaction progress.

Line graph of rate study results:

Figure 1. ^1H NMR spectrum of neopentyl chlorosulfate

Figure 2. ^{13}C NMR spectrum of neopentyl chlorosulfate

Figure 3. ^1H NMR spectrum of isobutyl chlorosulfate

Figure 4. ^{13}C NMR spectrum of isobutyl chlorosulfate

Figure 5. ^1H NMR spectrum of phenyl neopentyl sulfate (1)

Figure 6. ^{13}C NMR spectrum of phenyl neopentyl sulfate (1)

Figure 7. ^1H NMR spectrum of estrone neopentyl sulfate (2)

Figure 8. ^{13}C NMR spectrum of estrone neopentyl sulfate (2)

Figure 9. ^1H NMR spectrum of benzyl *N*-carbobenzyloxy-L-tyrosine neopentyl sulfate (3)

Figure 10. ^{13}C NMR spectrum of benzyl *N*-carbobenzyloxy-L-tyrosine neopentyl sulfate
(3)

Figure 11. ^1H NMR spectrum of 1,2:5,6-diisopropylidene- α -D-glucofuranose 3-neopentyl sulfate (4)

Figure 12. ^{13}C NMR spectrum of 1,2:5,6-diisopropylidene- α -D-glucofuranose 3-neopentyl sulfate (4)

Figure 13. ^1H NMR spectrum of phenyl isobutyl sulfate (5)

Figure 14. ^{13}C NMR spectrum of phenyl isobutyl sulfate (5)

Figure 15. ^1H NMR spectrum of estrone isobutyl sulfate (6)

Figure 16. ^{13}C NMR spectrum of estrone isobutyl sulfate (6)

Figure 17. ^1H NMR spectrum of 1,2:5,6-diisopropylidene- α -D-glucofuranose 3-isobutyl sulfate (7)

Figure 18. ^{13}C NMR spectrum of 1,2:5,6-diisopropylidene- α -D-glucofuranose 3-isobutyl sulfate (7)

Figure 19. ^1H NMR spectrum of 1,2-isopropylidene-3,5-dibenzyl- α -D-glucufuranose 3-isobutyl sulfate (8)

Figure 20. ^{13}C NMR spectrum of 1,2-isopropylidene-3,5-dibenzyl- α -D-glucofuranose 3-isobutyl sulfate (8)

Figure 21. ^1H NMR spectrum of *N*-(9-fluorenylmethoxycarbonyl) L-tyrosine neopentyl sulfate (9)

Figure 22. ^{13}C NMR spectrum of *N*-(9-fluorenylmethoxycarbonyl) L-tyrosine neopentyl sulfate (9)

^{13}C OBSERVE
 Archive directory: /1400/dob/vnmrsys/data
 Sample directory:
 File: CARBON
 Pulse Sequence: s2pul
 Date: Aug 18 2005
 1400
 PULSE SEQUENCE: standard
 OBSERVE C13
 DECOUPLE H1

Figure 23. ^1H NMR spectrum of *N*-(tert-butoxycarbonyl) L-tyrosine neopentyl sulfate (10)

Figure 24. ^{13}C NMR spectrum of *N*-(tert-butoxycarbonyl) L-tyrosine neopentyl sulfate (10)

Figure 25. ^1H NMR spectrum of D-glucose 3-neopentyl sulfate (11)

Figure 26. ^{13}C NMR spectrum of D-glucose 3-neopentyl sulfate (11)

Figure 27. ^1H NMR spectrum of D-glucose 3-isobutyl sulfate (12)

Figure 28. ^{13}C NMR spectrum of D-glucose 3-isobutyl sulfate (12)

Figure 29. ^1H NMR spectrum of D-glucose 6-isobutyl sulfate (13)

Figure 30. ^{13}C NMR spectrum of D-glucose 6-isobutyl sulfate (13)

Figure 31. ^1H NMR spectrum of benzyl *N*-carbobenzyloxy-L-tyrosine sulfate (15)

Figure 32. ^{13}C NMR spectrum of benzyl *N*-carbobenzyloxy-L-tyrosine sulfate (15)

Figure 33. ^1H NMR spectrum of estrone sulfate (16)

Figure 34. ^{13}C NMR spectrum of estrone sulfate (16)

Figure 35. ^1H NMR spectrum of 1,2:5,6-diisopropylidene- α -D-glucofuranose 3-sulfate (17)

Figure 36. ^{13}C NMR spectrum of 1,2:5,6-diisopropylidene- α -D-glucofuranose 3-sulfate (17)

Figure 37. ^1H NMR spectrum of 3-azido-3-deoxy D-glucose (18) and neopentyl sulfate (19)

Figure 38. IR spectrum of 3-azido-3-deoxy D-glucose (18) and neopentyl sulfate (19)

Figure 39. ^1H NMR spectrum of D-glucose 3-sulfate (20)

Figure 40. ^{13}C NMR spectrum of D-glucose 3-sulfate (20)

Figure 41. ^1H NMR spectrum of D-glucose 6-sulfate (21)

Figure 42. ^{13}C NMR spectrum of D-glucose 6-sulfate (21)

Figure 43. ^1H NMR spectrum of tetrahydropyran-2-methyl chlorosulfate (23)

Figure 44. ^{13}C NMR spectrum of tetrahydropyran-2-methyl chlorosulfate (23)

Figure 45. ^1H NMR spectrum of 1,2:5,6-diisopropylidene- α -D-glucofuranose 3-tetrahydropyran-2-methyl sulfate (24)

Figure 46. Mass spectrum of 1,2:5,6-diisopropylidene- α -D-glucofuranose 3-tetrahydropyran-2-methyl sulfate (24)

References:

- (1) Buncel, E.; Millington, J. P., *Can. J. Chem.* **1965**, 43, 556-564.
- (2) Bodanszky, M.; Bodanszky, A., *The Practice of Peptide Synthesis*. Springer-Verlag: Berlin, 1984; p 37-38.
- (3) Dax, K.; Wolflehner, W.; Weidmann, H., *Carbohydr. Res.* **1978**, 65, 132-138.
- (4) Carpino, L. A.; Han, G. Y., *J. Org. Chem.* **1972**, 37, 3404-3409.
- (5) Ponnusamy, E.; Fotadar, U.; Spisni, A.; Fiat, D., *Synthesis* **1986**, 48-49.