A Concise Synthesis of Butylcycloheptylprodigiosin

Jonathan T. Reeves*

Department of Chemical Development, Boehringer Ingelheim Pharmaceuticals, Inc., 900 Old Ridgebury Road/P.O. Box 368, Ridgefield, CT 06877

jreeves@rdg.boehringer-ingelheim.com

Supporting Information

Table of Contents

General -----	1
Procedures and data for 1, 3-5, 8 and 10 -----	2-6
¹H and ¹³C NMR spectra for 1, 3-5, 8 and 10 -----	7-20

General. Reagents and solvents were used as received from commercial sources. Reactions were conducted under an atmosphere of dry N₂. Reaction temperatures refer to internal temperatures unless noted otherwise. HPLC analyses were performed on a 4.6 mm x 10 cm column with particle size of 2 micron, wavelength 220 nm and 254 nm, at 25 °C. Mobile phases are A: water with 0.05% TFA, B: MeCN with 0.05% TFA. Gradient conditions are 90% A to 10% A in 15 min, hold 5 min, back to 90% A followed by a 3-min post run. ¹H and ¹³C NMR spectra were recorded on a Bruker DPX-400 NMR spectrometer at 400 and 100 MHz, respectively, in CDCl₃ unless indicated otherwise. COSY, NOESY and ROESY NMR spectra were recorded on a Bruker 600 MHz NMR spectrometer. Melting points were measured using a Mel-Temp 3.0 melting point apparatus and are uncorrected. IR spectra were recorded neat using an Omnic IR spectrometer.

Aldol 5. To a suspension of CuI (179 mg, 0.94 mmol) in THF (15 mL) at $-40\text{ }^{\circ}\text{C}$ was added *n*-BuMgCl (5.2 mL, 10.4 mmol, 2.0 M/THF). The reaction mixture was stirred for 10 min. A solution of 2-cyclononen-1-one (**6**) (1.30 g, 9.4 mmol) in THF (5 mL) was added dropwise over 20 min at $-40\text{ }^{\circ}\text{C}$. After 30 min at $-40\text{ }^{\circ}\text{C}$, a solution of **7** (1.00 g, 10.4 mmol) in THF (10 mL) was added over 2 min. The cooling bath was removed, and after 30 min the reaction mixture was quenched with saturated aq. NH_4Cl , extracted with EtOAc, and the organic layer concentrated. ^1H NMR analysis showed a single diastereomer. The residue was chromatographed on SiO_2 (80:20 hexanes/EtOAc) to give 2.15 g of **5** as a white solid (78%). Mp (hexanes/EtOAc) $87\text{--}88\text{ }^{\circ}\text{C}$; IR 3427, 2931, 1685, 1506, 1431, 1105, 1056 cm^{-1} ; ^1H NMR δ 7.79 (d, 1 H, $J = 0.9\text{ Hz}$), 7.60 (t, 1 H, $J = 1.0\text{ Hz}$), 4.96 (d, 1 H, $J = 2.8\text{ Hz}$), 3.74 (br s, 1 H), 3.19 (dd, 1 H, $J = 4.1, 8.6\text{ Hz}$), 2.62 (ddd, 1 H, $J = 4.1, 8.1, 10.5\text{ Hz}$), 2.30 (ddd, 1 H, $J = 4.6, 8.6, 10.1\text{ Hz}$), 2.04–1.95 (m, 1 H), 1.83–1.68 (m, 2 H), 1.59–0.97 (m, 14 H), 0.75 (t, 3 H, $J = 6.9\text{ Hz}$); ^{13}C NMR δ 219.5, 150.6, 141.1, 135.7, 68.1, 58.9, 45.4, 36.4, 32.6, 28.6, 25.9, 25.8, 24.3, 22.7, 22.6, 21.6, 13.8; HRMS calcd for $\text{C}_{17}\text{H}_{28}\text{NO}_3$ $[\text{M}+\text{H}]^+$: 294.2063; found: 294.2083.

Pyrrole 4. To a solution of aldol **5** (1.00 g, 3.41 mmol), Et_3N (2.4 mL, 17.1 mmol) and THF (6.8 mL) at $-10\text{ }^{\circ}\text{C}$ was added dropwise MsCl (0.32 mL, 4.09 mmol). The reaction mixture was warmed to rt over 1

h. Aqueous 2 N NaOH (10.0 mL) was added, the reaction mixture was heated at 70 °C for 4 h, cooled to rt, diluted with 20 mL saturated aqueous NaHCO₃, and extracted with CH₂Cl₂. The organic layer was dried (MgSO₄), concentrated, and the residue was chromatographed on SiO₂ (95:5→90:10 hexanes/EtOAc) to give 571 mg of **4** as an orange oil (68%). IR 3247, 2926, 2857, 1712, 1463, 1346, 1087 cm⁻¹; ¹H NMR δ 9.47 (br s, 1 H), 9.28 (s, 1 H), 6.70 (d, 1 H, *J* = 2.6 Hz), 2.79-2.61 (m, 3 H), 1.97-1.90 (m, 1 H), 1.79-1.17 (m, 14 H), 1.12-0.98 (m, 1 H), 0.87 (t, 3 H, *J* = 6.8 Hz); ¹³C NMR δ 177.5, 141.9, 130.9, 128.3, 120.3, 37.1, 36.6, 35.9, 30.3, 27.1 (3 C), 26.7, 23.4, 22.8, 14.1; HRMS calcd for C₁₆H₂₆NO [M+H]⁺: 248.2008; found: 248.2023.

Enone 8. To a solution of **5** (1.35 g, 4.60 mmol), Et₃N (3.2 mL, 23.0 mmol) and CH₂Cl₂ (15 mL) at 0 °C was added MsCl (0.43 mL, 5.52 mmol). The reaction mixture was stirred at rt for 2 h, at reflux for 20 h, and then cooled to rt, washed with saturated aq. NH₄Cl, and the organic phase dried (MgSO₄) and concentrated. ¹H NMR of the crude product showed a 6:1 *E*:*Z* mixture of olefin isomers. The residue was chromatographed on SiO₂ (90:10 hexanes/EtOAc) to give 1.10 g of **8** as a colorless oil (87%). Pure isomers were obtained by further chromatography on SiO₂ (96:4 hexanes/EtOAc). *E*-**8**: colorless oil; IR 2927, 2858, 1712, 1676, 1517, 1467, 1099 cm⁻¹; ¹H NMR δ 7.91 (s, 1 H), 7.77 (s, 1 H), 6.71 (s, 1 H), 3.18-3.10 (m, 1 H), 3.05-2.99 (m, 1 H), 2.40-2.30 (m, 1 H), 1.86-1.55 (m, 6 H), 1.48-1.16 (m, 10 H), 0.86 (t, 3 H, *J* = 6.9 Hz); ¹³C NMR δ 209.8, 150.7, 148.0, 137.6, 136.6, 122.3, 42.9, 42.0, 35.0, 30.1, 27.8, 26.3, 23.04, 22.96, 22.6, 22.5, 14.1; HRMS calcd for C₁₇H₂₆NO₂ [M+H]⁺: 276.1958; found: 276.1977. *Z*-**8**: colorless oil; IR 2929, 2858, 1722, 1695, 1467, 1309, 1099 cm⁻¹; ¹H NMR δ 7.80 (s, 1

H), 7.60 (s, 1 H), 6.23 (s, 1 H), 2.75-2.60 (m, 2 H), 2.48-2.41 (m, 1 H), 1.82-1.77 (m, 2 H), 1.73-1.25 (m, 11 H), 0.88 (t, 3 H, $J = 6.8$ Hz); ^{13}C NMR δ 214.4, 150.8, 150.3, 136.6, 136.4, 115.4, 44.6, 43.5, 36.3, 33.5, 29.8, 27.8, 25.6, 24.8, 24.2, 22.8, 14.1; HRMS calcd for $\text{C}_{17}\text{H}_{26}\text{NO}_2$ $[\text{M}+\text{H}]^+$: 276.1958; found: 276.1984.

Individual hydrolysis of *E*-8 and *Z*-8. A solution of either *E*-8 or *Z*-8 (120 mg, 0.436 mmol) in THF (3.0 mL) was treated with aqueous 2N NaOH (3.0 mL) and the reaction mixture was heated at 70 °C until HPLC analysis indicated consumption of starting material (*E*-8: 6.0 h; *Z*-8: 9.5 h). The reaction mixture was diluted with saturated aqueous NaHCO_3 , extracted with CH_2Cl_2 , and the organic layer was dried (MgSO_4), concentrated, and the residue chromatographed on SiO_2 (95:5→90:10 hexanes/EtOAc) to give **4** as an orange oil. From *E*-8: 92 mg (85% yield). From *Z*-8: 80 mg (74% yield).

Lactam 3. A solution of **4** (550 mg, 2.22 mmol), **9** (503 mg, 4.45 mmol) and DMSO (7.8 mL) was treated with aq. 2 N NaOH (6.2 mL) and the reaction mixture was heated at 60 °C for 8 h. After cooling to rt, the reaction mixture was poured into EtOAc (50 mL) and the layers cut. The organic layer was washed with 50% brine, dried (MgSO_4) and concentrated to give crude **3**. Trituration with hexanes/MTBE gave 569 mg of **3** as a bright yellow solid (75%). Mp (hexanes/MTBE) 233-236 °C; IR 3351, 2922, 2855, 1665, 1593, 1579, 1441, 1361, 1225 cm^{-1} ; ^1H NMR ($\text{DMSO}-d_6$) δ 10.49 (s, 1 H), 9.37

(s, 1 H), 6.47 (d, 1 H, $J = 2.5$ Hz), 6.03 (s, 1 H), 5.17 (d, 1 H, $J = 1.3$ Hz), 3.83 (s, 3 H), 2.70-2.54 (m, 2 H), 1.91-1.79 (m, 1 H), 1.66-1.13 (m, 15 H), 1.03-0.90 (m, 1 H), 0.84 (t, 3 H, $J = 6.9$ Hz); ^{13}C NMR (DMSO- d_6) δ 170.6, 166.8, 134.2, 125.6, 124.2, 123.7, 110.7, 98.5, 90.8, 58.2, 36.9, 36.0, 35.3, 30.0, 27.3, 26.8, 26.4, 26.0, 23.2, 22.3, 14.0; HRMS calcd for $\text{C}_{21}\text{H}_{31}\text{N}_2\text{O}_2$ $[\text{M}+\text{H}]^+$: 343.2380; found: 343.2400.

Triflate 10. A suspension of **3** (310 mg, 0.905 mmol) in CH_2Cl_2 was treated dropwise at 0°C with Tf_2O (0.18 mL, 1.09 mmol). The reaction mixture was stirred for 2 h at 0°C , washed with 2% aq. NaHCO_3 solution, and the organic phase dried (MgSO_4) and concentrated. The residue was filtered through a short (2 cm) SiO_2 pad using 95:5 hexanes/MTBE and the filtrate concentrated to give 361 mg of **10** as a brown oil (84%). IR 3356, 2923, 2853, 1626, 1569, 1552, 1418, 1370, 1332, 1196, 1129 cm^{-1} ; ^1H NMR δ 10.62 (br s, 1 H), 6.97 (s, 1 H), 6.44 (s, 1 H), 5.39 (s, 1 H), 3.82 (s, 3 H), 2.78-2.62 (m, 3 H), 2.02-1.92 (m, 1 H), 1.71-1.18 (m, 14 H), 1.16-1.02 (m, 1 H), 0.86 (t, 3 H, $J = 6.8$ Hz); ^{13}C NMR δ 167.7, 160.5, 144.5, 131.6, 129.3, 127.9, 121.9, 120.9, 118.8 (q, $J = 319$ Hz, CF_3), 86.7, 58.6, 37.2, 36.7, 36.0, 30.3, 27.6, 27.3, 27.2, 26.8, 23.5, 22.8, 14.1; HRMS calcd for $\text{C}_{22}\text{H}_{30}\text{N}_2\text{O}_4\text{F}_3\text{S}$ $[\text{M}+\text{H}]^+$: 475.1872; found: 475.1902.

Butylcycloheptylprodigosin (1). To a nitrogen-purged flask containing **10** (380 mg, 0.80 mmol), **11** (338 mg, 1.60 mmol), Pd(PPh₃)₄ (74 mg, 0.064 mmol) and LiCl (102 mg, 2.40 mmol) was added DME (23.0 mL) followed by 2 M aq. Na₂CO₃ (1.7 mL). The reaction mixture was heated at 80 °C for 10 h, cooled to rt, diluted with saturated aq. NaHCO₃ and CH₂Cl₂, and the layers were separated. The organic layer was dried (Na₂SO₄), concentrated and the residue was chromatographed on SiO₂ (95:5→80:20 hexanes/EtOAc with 2% Et₃N) to give 220 mg of **1** as an amorphous dark red solid (70%). IR 3318, 3105, 2921, 2852, 1619, 1578, 1552, 1449, 1361, 1333, 1284, 1223, 1149, 1116, 1086, 1057, 997, 955, 729 cm⁻¹; ¹H NMR (CD₂Cl₂) δ 12.8-11.7 (br, 1 H), 10.7-9.2 (br, 1 H), 6.86 (s, 1 H), 6.70 (dd, 1 H *J* = 1.2, 3.6 Hz), 6.61 (dd, 1 H, *J* = 1.3, 2.4 Hz), 6.34 (s, 1 H), 6.15-6.14 (m, 1 H), 6.14 (s, 1 H), 3.98 (s, 3 H), 2.58-2.50 (m, 1 H), 2.22-2.07 (m, 2 H), 1.61-1.45 (m, 5 H), 1.35-1.06 (m, 10 H), 0.84 (t, 3 H, *J* = 6.9 Hz), 0.79-0.65 (m, 1 H); ¹³C NMR (CD₂Cl₂) δ 169.5, 160.2, 143.7, 138.5, 129.1, 128.2, 122.7, 119.2, 116.6, 112.9, 110.5, 96.0, 58.9, 38.2, 37.1, 36.0, 30.6, 28.4, 28.1, 27.6, 27.5, 23.3, 23.1, 14.3; HRMS calcd for C₂₅H₃₄N₃O [M+H]⁺: 392.2696; found: 392.2722.