SUPPORTING INFORMATION

A Concise Synthesis of Spirotryprostatin A

Fumiko Y. Miyake, Kenichi Yakushijin, and David A. Horne*

Department of Chemistry, Oregon State University, Corvallis, Oregon 97331

Experimental procedures, ^1H and ^{13}C NMR spectra for compounds **1**, **5**, **6**, **8-12**.

Experimental

General. Materials were obtained from commercial suppliers and used without further purification unless otherwise specified. Silica gel (particle size 32-63 μ) was used for flash chromatography.

2,6-Dibromotryptophan methyl ester (5) and 2,5-dibromotryptophan methyl ester (6). To a stirred solution of tryptophan methyl ester hydrobromide (**4**) (3.0 g, 10 mmol) in 50 mL of AcOH-HCO₂H (4:1) was added NBS (3.8 g, 21 mmol) at 23 °C. After 20 m, the reaction mixture was concentrated under reduced pressure ($T < 40$ °C) to afford a residue. Ether was added and the resulting organic layer was washed with sat. NaHCO₃, brine, dried (MgSO₄), and filtered. Concentration of the filtrate and flash chromatography of the resulting residue using CH₂Cl₂:EtOAc (1:1) as eluent afforded **5** (0.9 g, 24%) and **6** (1.2 g, 32%) as a mixture. Small amounts of pure **5** and **6** were also obtained from the chromatography procedure for characterization purposes. **5**·HBr: $[\alpha]_D^{23} +25.9$ (c 0.08, MeOH); ¹H NMR (DMSO-*d*₆, 300 MHz). δ 12.09 (1H, bs), 8.50 (3H, bs), 7.51 (1H, d, $J = 8.5$ Hz), 7.48 (1H, d, $J = 1.8$ Hz), 7.21 (1H, dd, $J = 8.5, 1.8$ Hz), 4.08 (1H, bs), 3.58 (3H, s), 3.23 (1H, dd, $J = 14.5, 5.7$ Hz), 3.12 (1H, dd, $J = 14.5, 9.0$ Hz); ¹³C NMR (DMSO-*d*₆, 100 MHz) δ 170.2 (s), 137.7 (s), 126.9 (s), 123.4 (d), 120.3 (d), 115.5 (s), 114.3 (d), 112.5 (s), 108.2 (s), 53.8 (q), 52.9 (d), 26.9 (t). **6**·HBr: $[\alpha]_D^{23} +23.6$ (c 0.11, MeOH); ¹H NMR (DMSO-*d*₆, 300 MHz). δ 12.12 (1H, bs), 8.49 (3H, bs), 7.79 (1H, d, $J = 1.7$ Hz), 7.28 (1H, d, $J = 8.6$ Hz), 7.23 (1H, dd, $J = 8.6, 1.7$ Hz), 4.10 (1H, dd, $J = 9.1, 5.6$ Hz), 3.61 (3H, s), 3.22 (1H, dd, $J = 14.6, 5.6$ Hz), 3.10 (1H, dd, $J = 14.6, 9.1$ Hz); ¹³C NMR (DMSO-*d*₆, 100 MHz) δ 170.2 (s), 135.7 (s), 129.5 (s), 125.2 (d), 120.9 (d), 113.8 (d), 113.19 (s), 113.18 (s), 107.6 (s), 53.8 (q), 52.9 (d), 26.9 (t).

6-Bromodemethoxyspirotryprostatin A (8) and 5-bromodemethoxyspirotryprostatin A (9). To a stirred solution of a 3:2 mixture **5** and **6** (1.4 g, 3.74 mmol), respectively, in 50 mL of CH₂Cl₂ was added 3-methyl-2-butenal (0.44 mL, 4.5 mmol) at 23 °C. After 2 h, *N*-2,2,2-trichloroethoxycarbonyl-L-prolinyl chloride (1.4 g, 4.5 mmol) and triethylamine (0.63 mL, 4.5 mmol) was added and the whole was stirred an additional 3 h. Trifluoroacetic acid (1.7 mL, 22.5 mmol) was then added to the reaction mixture and the whole was stirred for an additional 2 h. The reaction mixture was diluted with 100 mL of CH₂Cl₂ and the organic layer was washed with sat. Na₂CO₃, brine, dried (MgSO₄), and filtered. Concentration of the filtrate afforded a residue that was used directly in the next step. The residue was dissolved in 50 mL of MeOH to which Zn powder (2.5 g, 37.4 mmol) was added. The reaction mixture was refluxed for 2 h, filtered over celite, followed by hot MeOH (3x) washes. Concentration of the filtrate afforded a residue that was purified by flash chromatography using an eluent mixture of hexane:EtOAc:MeOH (sat. NH₃) (5:4.25:0.75) to afford **8** (0.26 g, 26%), 3-*epi*-6-bromodemethoxyspirotryprostatin A (**8a**) (40 mg, 4%) and **9** (0.2 g, 30%). **8**: $[\alpha]_D^{23} -101$ (c 0.3, CHCl₃) ¹H NMR (CDCl₃, 400 MHz) δ 8.26 (1H, bs), 7.13 (1H, dd, $J = 8.1, 1.7$ Hz), 7.05 (1H, d, $J = 1.7$ Hz), 6.91 (1H, d, $J = 8.1$ Hz), 5.03 (1H, dm, $J = 9.1$ Hz), 5.01 (1H, dd, $J = 10.7, 7.0$ Hz), 4.82 (1H, d, $J = 9.1$ Hz), 4.32 (1H, bt, $J = 8.0$ Hz), 3.64-3.57 (2H, m), 2.63 (1H, dd, $J = 13.4, 10.7$ Hz), 2.42 (1H, dd, $J = 13.4, 7.0$ Hz), 2.37-2.22 (2H, m), 2.07-1.96 (2H, m), 1.66 (3H, s), 1.19 (3H, s); ¹³C NMR (CDCl₃, 75 MHz) δ 180.6 (s), 167.4 (s), 167.3 (s), 142.5 (s), 139.3 (s), 128.1 (d), 126.3 (s), 125.4 (d), 122.8 (s), 121.4 (d), 113.4 (d), 61.5 (d), 60.5 (d), 58.9 (d), 56.1 (s), 45.6 (t), 34.6 (t), 27.8 (t), 25.9 (q), 24.1 (t), 18.5 (q). **8a**: ¹H NMR (CDCl₃, 300 MHz) δ 8.13

(1H, bs), 7.18 (1H, dd, $J = 8.0, 1.7$ Hz), 7.08 (1H, d, $J = 1.7$ Hz), 7.03 (1H, d, $J = 8.0$ Hz), 5.23 (1H, dm, $J = 9.8$ Hz), 4.76 (1H, d, $J = 9.8$ Hz), 4.55 (1H, bt, $J = 8.1$ Hz), 4.24 (1H, bt, $J = 8.0$ Hz), 3.68-3.57 (2H, m), 3.05 (1H, dd, $J = 13.6, 7.7$ Hz), 2.34 (1H, dd, $J = 13.6, 8.7$ Hz), 2.39-2.23 (2H, m), 2.11-1.93 (2H, m), 1.66 (3H, s), 1.43 (3H, s); ^{13}C NMR (CDCl_3 , 75 MHz) δ 176.7 (s), 167.4 (s), 166.2 (s), 142.2 (s), 137.2 (s), 130.7 (s), 125.9 (d), 124.5 (d), 122.8 (s), 119.1 (d), 113.8 (d), 64.4 (d), 61.5 (d), 59.6 (d), 56.2 (s), 45.7 (t), 34.8 (t), 28.1 (t), 26.3 (q), 24.0 (t), 18.6 (q). **9**: $[\alpha]_D^{23}$ -28 (c 0.17, CHCl_3) ^1H NMR (CDCl_3 , 400 MHz) δ 8.20 (1H, bs), 7.36 (1H, dd, $J = 8.3, 2.0$ Hz), 7.18 (1H, d, $J = 2.0$ Hz), 6.75 (1H, d, $J = 8.3$ Hz), 5.04 (1H, dm, $J = 9.0$ Hz), 5.01 (1H, dd, $J = 10.7, 7.0$ Hz), 4.82 (1H, d, $J = 8.9$ Hz), 4.30 (1H, bt, $J = 7.8$ Hz), 3.65-3.57 (2H, m), 2.62 (1H, dd, $J = 13.4, 10.7$ Hz), 2.44 (1H, dd, $J = 13.4, 7.0$ Hz), 2.35-2.26 (2H, m), 2.07-1.96 (2H, m), 1.70 (3H, s), 1.12 (3H, s); ^{13}C NMR (CDCl_3 , 100 MHz) δ 180.4 (s), 167.4 (s), 167.2 (s), 140.2 (s), 139.7 (s), 131.9 (s), 130.5 (d), 129.4 (s), 121.5 (d), 115.0 (s), 111.2 (d), 61.5 (d), 60.8 (d), 59.0 (d), 56.6 (s), 45.7 (t), 34.6 (t), 27.9 (t), 25.8 (q), 24.1 (t), 18.5 (q).

Spirotryprostatin A (1) and 9,12-bis-epi-spirotryprostatin A (10). To a stirred solution of 1.5 mL of MeOH under argon was added Na (43 mg, 1.8 mmol) at 23 °C followed by the addition of DMF (1 mL). After 20 m, CuI (68 mg, 0.36 mmol), **8** (80 mg, 0.18 mmol), and DMF (1 mL) were added to the reaction mixture and the whole was refluxed for 6 h. Upon cooling, the reaction mixture was filtered and the collected solid was rinsed with generous amounts of EtOAc. The filtrate was washed with 2% NaOH and brine and the organic layer dried (MgSO_4) and filtered. Concentration of the filtrate afforded a residue that was purified by preparative TLC (SiO_2 , Hex:EtOAc:MeOH (sat. NH_3) 5:4:1) to yield **1** (8 mg, 11%) and **10** (12 mg, 17%). A solution of **10** (8 mg) in 1 mL of MeOH, Na (5 mg), and DMF (1.5 mL) was refluxed (oil bath T = 120 °C) for 6 h. After cooling, EtOAc was added and the organics were washed with 2% NaOH, brine, and dried (MgSO_4). Concentration of the filtrate and preparative TLC of the resulting residue afforded **1** (2.6 mg) and recovered **10** (4.0 mg). **1**: $[\alpha]_D^{23}$ -94.3 (c 0.14, CHCl_3); ^1H NMR (acetone- d_6 , 400 MHz) δ 9.42 (1H, bs), 7.11 (1H, d, $J = 8.3$ Hz), 6.53 (1H, dd, $J = 8.3, 2.4$ Hz), 6.50 (1H, d, $J = 2.4$ Hz), 5.14 (1H, dm, $J = 9.3$ Hz), 4.90 (1H, dd, $J = 10.4, 7.0$ Hz), 4.66 (1H, d, $J = 9.3$ Hz), 4.34 (1H, t, $J = 8.0$ Hz), 3.78 (3H, s), 3.53-3.43 (2H, m), 2.54 (1H, dd, $J = 13.2, 10.4$ Hz), 2.27 (1H, dd, $J = 13.2, 7.0$ Hz), 2.23 (1H, m), 2.13 (1H, m), 1.95 (2H, m), 1.62 (3H, s), 1.19 (3H, s); ^{13}C NMR (acetone- d_6 , 75 MHz) δ 180.9 (s), 167.1 (s), 166.8 (s), 160.9 (s), 143.7 (s), 136.3 (s), 127.6 (d), 123.1 (d), 119.7 (s), 106.5 (d), 96.6 (d), 61.0 (d), 60.4 (d), 58.7 (d), 55.7 (s), 55.2 (q), 45.0 (t), 34.7 (t), 27.8 (t), 25.0 (q), 23.7 (t), 17.5 (q). **1**: ^1H NMR (CDCl_3 , 400 MHz) δ 7.64 (1H, bs), 6.94 (1H, d, $J = 8.4$ Hz), 6.51 (1H, dd, $J = 8.4, 2.3$ Hz), 6.44 (1H, d, $J = 2.3$ Hz), 5.04 (1H, dm, $J = 9.1$ Hz), 5.01 (1H, dd, $J = 10.8, 6.9$ Hz), 4.79 (1H, d, $J = 9.1$ Hz), 4.30 (1H, t, $J = 7.9$ Hz), 3.81 (3H, s), 3.64-3.57 (2H, m), 2.62 (1H, dd, $J = 13.4, 10.8$ Hz), 2.40 (1H, dd, $J = 13.4, 6.9$ Hz), 2.39-2.55 (2H, m), 2.06-1.96 (2H, m), 1.66 (3H, s), 1.18 (3H, s); ^{13}C NMR (CDCl_3 , 75 MHz) δ 181.2 (s), 167.5 (s), 167.4 (s), 160.8 (s), 142.1 (s), 138.8 (s), 127.7 (d), 121.8 (d), 119.1 (s), 107.2 (d), 97.1 (d), 61.5 (d), 60.6 (d), 58.9 (d), 55.9 (q and s), 45.6 (t), 34.8 (t), 27.8 (t), 25.9 (q), 24.1 (t), 18.4 (q). **10**: $[\alpha]_D^{23}$ +45.4 (c 0.18, CHCl_3); ^1H NMR (CDCl_3 , 400 MHz) δ 8.23 (1H, bs), 7.02 (1H, d, $J = 8.3$ Hz), 6.51 (1H, dd, $J = 8.3, 2.3$ Hz), 6.47 (1H, d, $J = 2.3$ Hz), 5.12 (2H, s), 4.63 (1H, dd, $J = 9.2, 5.3$ Hz), 4.25 (1H, t, $J = 7.6$ Hz), 3.80 (3H, s), 3.71-3.56 (2H, m), 2.94 (1H, dd, $J = 13.6, 5.3$ Hz), 2.51 (1H, dd, $J = 13.6, 9.2$ Hz), 2.34-2.28 (2H, m), 2.05 (1H, m), 1.93 (1H, m), 1.67 (3H, s), 1.43 (3H, s); ^{13}C NMR (CDCl_3 , 75 MHz) δ 180.4 (s), 166.5 (s), 166.0 (s), 160.7 (s), 142.6 (s), 138.8 (s), 126.1 (d), 119.9 (s), 119.4 (d), 107.3 (d),

97.5 (d), 62.0 (d), 60.4 (d), 59.3 (d), 55.8 (q), 54.7 (s), 46.1 (t), 36.3 (t), 28.2 (t), 26.2 (q), 23.7 (t), 18.5 (q).

5-Methoxydemethoxyspirotryprostatin A (11) and 9,12-bis-epi-5-methoxydemethoxyspirotryprostatin A (12). A similar protocol as described above for **1** and **10** was followed wherein **9** (80 mg, 0.18 mmol) yielded **11** (10 mg, 14%) and **12** (15 mg, 21%). Similarly, **12** was epimerized to **11**. **11**: $[\alpha]_D^{23}$ -40 (c 0.075, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ 7.47 (1H, bs), 6.79 (1H, dd, J = 8.4, 2.3 Hz), 6.75 (1H, d, J = 8.4, Hz), 6.67 (1H, d, J = 2.3 Hz), 5.06 (1H, dm, J = 9.1 Hz), 5.03 (1H, dd, J = 10.7, 7.0 Hz), 4.84 (1H, d, J = 9.1 Hz), 4.31 (1H, t, J = 7.9 Hz), 3.74 (3H, s), 3.65-3.56 (2H, m), 2.64 (1H, dd, J = 13.4, 10.7 Hz), 2.44 (1H, dd, J = 13.4, 7.0 Hz), 2.32 (2H, m), 2.03 (2H, m), 1.68 (3H, s), 1.18 (3H, s); ¹³C NMR (CDCl₃, 100 MHz) δ 180.6 (s), 167.5 (s), 167.3 (s), 155.8 (s), 138.9 (s), 134.3 (s), 128.6 (s), 121.6 (d), 114.6 (d), 113.5 (d), 110.1 (d), 61.5 (d), 60.8 (d), 59.0 (d), 56.6 (s), 56.2 (q), 45.6 (t), 34.8 (t), 27.8 (t), 26.0 (q), 24.1 (t), 18.4 (q). **12**: $[\alpha]_D^{23}$ +68.8 (c 0.16, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ 7.77 (1H, bs), 6.78-6.76 (3H, m), 5.17 (1H, d, J = 9.5, Hz), 5.12 (1H, dm, J = 9.5 Hz), 4.66 (1H, dd, J = 9.1, 5.6 Hz), 4.26 (1H, t, J = 7.6 Hz), 3.78 (3H, s), 3.72-3.56 (2H, m), 2.98 (1H, dd, J = 13.6, 5.6 Hz), 2.53 (1H, dd, J = 13.6, 9.1 Hz), 2.34-2.28 (2H, m), 2.06 (1H, m), 1.95 (1H, m), 1.68 (3H, s), 1.45 (3H, s); ¹³C NMR (CDCl₃, 100 MHz) δ 179.4 (s), 166.6 (s), 165.9 (s), 155.8 (s), 139.0 (s), 134.7 (s), 129.7 (s), 119.3 (d), 113.4 (d), 113.0 (d), 110.4 (d), 62.2 (d), 60.5 (d), 59.4 (d), 56.3 (q), 55.5 (s), 46.2 (t), 36.4 (t), 28.2 (t), 26.2 (q), 23.7 (t), 18.6 (q).

spirotryprostatin A (**1**)

^1H NMR CDCl_3

spirotryprostatin A (**1**)
 ^1H NMR acetone- d_6

^1H NMR DMSO- d_6

6

¹H NMR DMSO-d₆

8

^1H NMR CDCl_3

8a

^1H NMR CDCl_3

9

^1H NMR CDCl_3

10

^1H NMR CDCl_3

11

 ^1H NMR CDCl_3

12

^1H NMR CDCl_3