A Concise Total Synthesis of Naamidine A

Nicholas S. Aberle, Guillaume Lessene and Keith G. Watson*

*Medicinal Chemistry Group, The Walter and Eliza Hall Institute
of Medical Research, Melbourne, Australia*

SUPPORTING INFORMATION

General Experimental Procedures.

All non-aqueous reactions were performed in oven-dried glassware under an atmosphere of dry nitrogen, unless otherwise specified. THF was freshly distilled from sodium/benzophenone under N₂; CH₂Cl₂ and toluene were freshly distilled from CaH₂ under N₂. Anhydrous acetonitrile (99.8%) was purchased from Aldrich and used as supplied. All other solvents were reagent grade. Petroleum ether describes a mixture of hexanes in the bp range 40-60 °C. Analytical thin-layer chromatography was performed on Merck silica gel 60F₂₅₄ aluminium-backed plates. Flash chromatography was performed with silica gel 60 (particle size 0.040-0.063mm) following the guidelines of Still et al.¹ All NMR spectra were recorded on a Bruker Avance DRX 300 with the solvents indicated (¹H NMR at 300 MHz; ¹³C NMR at 75 MHz). Chemical shifts are reported in ppm on the δ scale, referenced to the appropriate solvent peak. Infrared spectra were recorded on a Bruker Tensor27 FT-IR spectrometer. Mass spectrometry was performed on a Finnigan LCQ Advantage MAX. High resolution mass spectra were measured at the University of Canterbury on a Kratos MS80 RFA mass spectrometer. Elemental analyses were carried out for carbon, nitrogen and hydrogen at the University of Otago microanalytical laboratory.

¹ Still, W.C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, *43*, 2923.

2-(*tert*-Butoxycarbonyl-methyl-amino)-3-(4-benzyloxyphenyl)-propionic acid (7).

Iodomethane (5.00 mL, 80.32 mmol, 8.0 eq.) was added to a solution of Boc-Tyr(Bzl)-OH (**6**) (3.721 g, 10.02 mmol) in THF (30 mL) and cooled to 0 °C. Sodium hydride (60% dispersion in mineral oil; 1.32 g, 33.00 mmol, 3.3 eq) was added in small portions over five minutes, during which time hydrogen gas evolved. The solution became a light brown suspension and was removed from the ice-bath after 20 minutes then stirred at room temperature under N₂ for 24 hours. Diluting the reaction with EtOAc (50 mL) and quenching by dropwise addition of H₂O (50 mL) was followed by removal of the solvent *in vacuo*. The residue was partitioned between ether (30 mL) and water (100 mL), and the organic layer was washed with saturated aqueous sodium bicarbonate (50 mL). The combined aqueous layer was acidified to pH 3 with 5% citric acid and extracted with EtOAc (3 x 100 mL). The organic extracts were washed with H₂O (2 x 75 mL), 5% Na₂S₂O₃ (2 x 50 mL) and H₂O (50 mL) then dried (MgSO₄), filtered and concentrated to give **7** as a pale yellow oil (3.717 g, 96%) which partially crystallised on standing.

¹H NMR (CDCl₃, 300 MHz, exists as a 1:1 mixture of two rotamers) δ 7.27-7.42 (m, 5 H), 7.07-7.12 (m, 2 H), 6.89 (d, *J* = 8.4 Hz, 2 H), 5.03 (s, 2 H), 4.67 and 4.55 (two m, 1 H), 3.22 (m, 1 H), 2.93-3.11 (m, 1 H), 2.73 and 2.66 (two s, 3 H), 1.40 and 1.33 (two s, 9 H).

IR (ATR, cm⁻¹) 2800-3300 (br), 2978, 2935, 1738, 1629, 1369, 1142.

MS (ESI) 384 (M - H⁺), 769 (2M - H⁺).

EA C₂₂H₂₇NO₅ (385.45): Calcd C 68.55, H 7.06, N 3.63; Found C 68.27, H 7.24, N 3.63.

***tert*-Butyl 1-(*N*-methoxy-*N*-methylcarbamoyl)-2-(4-benzyloxy)phenyl)ethyl-methylcarbamate (**8**).**

A solution of **7** (3.516 g, 9.12 mmol) in CH₂Cl₂ (40 mL) at 0 °C was treated with DIEA (2.40 mL, 13.78 mmol) then DAST (0.77 mL, 10.89 mmol). Stirring continued at 0 °C for 15 minutes before addition of a second solution prepared thus: *N,O*-dimethylhydroxylamine (1.340 g, 13.737 mmol) in CH₂Cl₂ (20 mL) at 0 °C was treated with DIEA (3.54 mL, 20.954 mmol) by dropwise addition and stirred for 15 minutes. The combined solution was then stirred for a further 15 minutes at 0 °C followed by 3 h at room temperature, whereupon it was washed successively with 40 mL of saturated aqueous solutions of sodium bicarbonate, ammonium chloride and sodium chloride. The organic layer was dried (MgSO₄), filtered and concentrated *in vacuo* to a crude orange oil. Purification by flash chromatography (100% CH₂Cl₂ to 90% CH₂Cl₂ in methanol) afforded **8** as a colourless oil (3.36 g, 86%).

¹H NMR (DMSO-*d*₆, 300 MHz, 360 K; exists as a complex mixture of four rotamers at room temperature) δ 7.25–7.40 (m, 5 H), 7.08 (d, *J* = 8.6 Hz, 2 H), 6.89 (d, *J* = 8.6 Hz, 2 H), 5.13 (bs, 1 H), 5.04 (s, 2 H), 3.59 (s, 3 H), 3.08 (s, 3 H), 2.75–2.95 (m, 2 H), 2.71 (s, 3 H), 1.25 (s, 9 H).

¹³C NMR (DMSO-*d*₆, 300 MHz, 360 K) δ 170.6, 157.3, 154.7, 137.5, 130.3, 130.1, 128.4, 127.7, 127.4, 115.0, 79.0, 69.8, 61.2, 56.2, 33.7, 32.4, 29.9, 28.0.

IR (ATR, cm⁻¹) 2976, 2931, 1674, 1665, 1511, 1391, 1240, 1173, 696, 495.

HRMS (ESI) Calcd for (M + H⁺) C₂₄H₃₃N₂O₅⁺ 429.2389, found 429.2382.

***tert*-Butyl 1-(4-(benzyloxy)phenyl)-4-(4-methoxyphenyl)-3-oxobutan-2-yl methyl carbamate (**9**).**

A solution of Weinreb amide **8** (2.911 g, 6.790 mmol) in THF (27 mL) at 0 °C was treated dropwise over five minutes with 4-methoxybenzylmagnesium chloride (0.20 M in THF; 50 mL; 10 mmol). Stirring continued at 0 °C for 10 min then for 3 h at room temperature. Quenching the reaction with saturated aqueous ammonium chloride (30 mL) was tempered by an ice bath. Resulting solids were dissolved with water (20 mL), the phases separated and the aqueous layer was extracted with diethyl ether (3 x 50 mL). The combined organic layers were washed with brine (75 mL) then dried (Na₂SO₄), filtered and concentrated to give a clear residue which was purified by flash chromatography (4:1 petroleum ether:EtOAc). **9** was obtained as a clear oil which crystallised on standing (2.071 g, 62%).

¹H NMR (CDCl₃, 300 MHz, exists as a 1:1 mixture of 2 rotamers) δ 7.28-7.41 (m, 5 H), 6.97-7.08 (m, 4 H), 6.80-6.85 (m, 4 H), 5.00 (s, 2 H), 4.72 and 4.30 (two m, 1 H), 3.77 (s, 3 H), 3.66 (s, 2 H), 3.09 (m, 1 H), 2.79 (m, 1 H), 2.57 and 2.51 (two s, 3 H), 1.41 and 1.33 (two s, 9 H).

¹³C NMR (CDCl₃, 300 MHz, exists as a 1:1 mixture of 2 rotamers) δ 205.7 and 205.4, 158.7 and 158.5, 157.5 and 157.3, 155.7, 154.8, 137.0, 130.3, 130.2, 130.0, 128.5, 127.8, 127.3, 125.7 and 125.6, 114.9 and 114.7, 114.1 and 114.0, 80.8 and 80.2, 70.0, 66.5, 64.5, 55.2, 46.1 and 45.7, 32.8, 32.4 and 32.1, 28.3 and 28.1.

IR (ATR, cm⁻¹) 2969, 2935, 1726, 1672, 1510, 1239, 1162.

MS (ESI) 512 (M + Na⁺).

EA C₃₀H₃₅NO₅ (489.60): Calcd C 73.59, H 7.21, N 2.86; Found C 73.40, H 7.40, N 2.89.

Naamine A (10).

Aminoketone **9** (695 mg, 1.42 mmol) was dissolved in 4 M HCl in diethyl ether (28 mL) and allowed to stand in a stoppered flask for 1 hour. Evaporation of the solvent followed by suspension in ether resulted in formation of a white powder which was collected by filtration. The filtrate was concentrated to dryness and suspended again in ether to produce further solids. The combined precipitate was washed with ether and dried *in vacuo*.

The HCl salt was suspended in H₂O (30 mL). Cyanamide was added in one portion and the reaction stirred at 90 °C for 1 h, becoming a homogeneous pale yellow solution upon heating. An off-white precipitate formed as the flask cooled, and the mixture was made basic to pH 10 with 0.3 M aqueous NaOH. The solids were collected by filtration and dried *in vacuo*, affording the desired tri-substituted 2-aminoimidazole **O-Bn-10** (488 mg, 86%).

¹H NMR (CDCl₃, 300 MHz) δ 7.28-7.43 (m, 5 H), 7.14 (d, J = 8.6 Hz, 2 H), 6.96 (d, J = 8.6 Hz, 2 H), 6.85 (d, J = 8.6 Hz, 2 H), 6.77 (d, J = 8.6 Hz, 2 H), 5.01 (s, 2 H), 3.84 (s, 2 H), 3.74 (s, 5 H), 3.05 (s, 3 H).

¹³C NMR (CDCl₃, 300 MHz) δ 157.6, 157.1, 146.5, 136.6, 132.2, 131.1, 130.4, 129.1, 128.5, 128.2, 127.6, 127.1, 120.7, 114.7, 113.5, 69.7, 54.9, 31.7, 29.0, 28.2.

IR (ATR, cm⁻¹) 3382, 3117, 2904, 1644, 1611, 1509, 1247, 810, 737.

MS (ESI) 414 (M + H⁺), 827 (2M + H⁺).

EA C₂₆H₂₇N₃O₂ (413.51): Calcd C 75.52, H 6.58, N 10.16; Found C 75.33, H 6.46, N 10.39.

A flask containing a suspension of benzyl ether **O-Bn-10** as obtained above (68 mg, 0.16 mmol) and Pd/C (10 wt. %; 9 mg) in methanol (4 mL) was stirred under H₂ until all starting material was consumed (monitored by MS). The suspension was

filtered through a plug of celite and washed with methanol. Evaporation of the filtrate afforded naamine A (**10**) (53 mg, 100%) as a partially crystalline solid.

¹H NMR (MeOD, 300 MHz) δ 7.07 (d, J = 8.6 Hz, 2 H), 6.83 (d, J = 8.5 Hz, 2 H), 6.75 (d, J = 8.6 Hz, 2 H), 6.63 (d, J = 8.5 Hz, 2 H), 3.75 (s, 2 H), 3.71 (s, 3 H), 3.68 (s, 2 H), 3.07 (s, 3 H).

¹³C NMR (DMSO- d_6 , 300 MHz) δ 157.2, 155.5, 148.3, 133.8, 131.6, 129.9, 129.5, 128.9, 119.8, 115.1, 113.4, 55.0, 32.0, 28.8, 27.9.

IR (ATR, cm^{-1}) 3322, 2913, 2362, 1633, 1609, 1508, 1242, 813.

HRMS (ESI) Calcd for ($M + H^+$) $\text{C}_{19}\text{H}_{22}\text{N}_3\text{O}_2^+$ 324.1712, found 324.1713.

Naamidine A (**1**).

1-methylparabanic acid **11** (105 mg, 0.820 mmol) in acetonitrile (750 μ L) was treated with *N,O*-bis(trimethylsilyl)acetamide (253 μ L, 1.02 mmol) and stirred at reflux for 45 min. The yellow solution was concentrated to dryness under vacuum, opened to dry N_2 to minimise exposure to air of the silylated product **12**. The residue was taken in toluene (1.30 mL) and added to **10** (53 mg, 0.16 mmol). The mixture was stirred overnight at reflux then diluted with EtOAc (15 mL) and washed with H_2O . The organic layer was dried over Na_2SO_4 and concentrated to dryness. Purification by flash chromatography (85:15 toluene:methanol with 1% NEt_3) provided naamidine A (**1**) as a yellow powder (57 mg, 80%).

1H NMR ($CDCl_3$, 300 MHz) δ 7.09 (d, J = 8.6 Hz, 2 H), 6.78-6.82 (m, 4 H), 6.72 (d, J = 8.6 Hz, 2 H), 3.86 (s, 2 H), 3.85 (s, 2 H), 3.76 (s, 3 H), 3.35 (s, 3 H), 3.15 (s, 3 H).

^{13}C NMR ($CDCl_3$, 300 MHz) δ 162.3, 158.0, 155.8, 154.9, 145.9, 144.5, 133.3, 130.2, 128.7, 127.8, 127.4, 126.8, 115.5, 113.8, 54.9, 31.0, 29.6, 28.2, 24.5.

IR (ATR, cm^{-1}) 3308, 2949, 2907, 2832, 1786, 1707, 1563, 1440, 1242, 1143, 606.

MS (ESI) 434 ($M + H^+$).

EA $C_{23}H_{23}N_5O_4$ (433.46): Calcd C 63.73, H 5.35, N 16.16; Found C 63.63, H 5.31, N 16.03.