A General Synthesis of Iminosugars

Ciaran McDonnell, Linda Cronin, Julie L. O' Brien and Paul V. Murphy*

Centre for Synthesis and Chemical Biology, Chemistry Department, Conway Institute of Biomolecular and Biomedical Research, University College Dublin, Belfield, Dublin 4, Ireland

Email: Paul.V.Murphy@ucd.ie

Index to Supporting Information

General Experimental Conditions.....	2
Methyl 2,3,4-tri- <i>O</i> -acetyl-6-deoxy-6-iodo- α -D-mannopyranoside 9a	2
2,3,4-Tri- <i>O</i> -acetyl-6-deoxy-6-iodo- α -D-mannopyranosyl azide	3
2,3,4-Tri- <i>O</i> -acetyl-6-deoxy- α -D- <i>lyxo</i> -hex-5-enopyranosyl azide 10a	3
Epoxides 11a	4
2,3,6-Tri- <i>O</i> -acetyl-5- <i>C</i> -methoxy- β -L-gulopyranoside 13	4
5- <i>C</i> -Methoxy- β -L-gulopyranosyl azide 14	5
1-Deoxymannojirimycin 2	5
Methyl 2,3,4-tri- <i>O</i> -acetyl-6-deoxy-6-iodo- α -D-glucopyranoside 9b	5
1,2,3,4-Tetra- <i>O</i> -acetyl-6-deoxy-6-iodo- α -D-glucopyranose	6
2,3,4-Tri- <i>O</i> -acetyl-6-deoxy-6-iodo- β -D-glucopyranosyl azide	6
2,3,4-Tri- <i>O</i> -acetyl-6-deoxy- β -D- <i>xylo</i> -hex-5-enopyranosyl azide 10b	7
Epoxides 11b	7
Acyclic compound 19	8
1,2:3,4-Di- <i>O</i> -isopropylidene-6-deoxy-6-iodo-D-galactopyranoside	8
1,2,3,4-Tetra- <i>O</i> -acetyl-6-deoxy-6-iodo-D-galactopyranose	9
2,3,4-Tri- <i>O</i> -acetyl-6-deoxy-6-iodo- β -D-galactopyranosyl azide	9
2,3,4-Tri- <i>O</i> -acetyl-6-deoxy- β -D- <i>arabino</i> -hex-5-enopyranosyl azide 20	10
5- <i>C</i> -Methoxy- β -D-glucopyranosyl and α -L-idopyranosyl azide	10
NMR data for 1-deoxynojirimycin 1 (salt)	10
NMR data for 1-deoxygalactostatin 3	11
NMR data for 1,6-dideoxygalactostatin 4	11
NMR Spectra	12-38

General Experimental Conditions

Optical rotations were determined at the sodium D line at 23 °C. Chemical shifts in ^1H -NMR spectra are reported relative to internal Me_4Si in CDCl_3 (δ 0.0) or HOD for D_2O (δ 4.63) or CD_2HOD (δ 3.36) for ^1H and CDCl_3 (δ 77.0) or CD_3OD (δ 47.7) for ^{13}C . ^1H NMR signals were assigned with the aid of COSY. ^{13}C -NMR signals were assigned with the aid of DEPT. Coupling constants are reported in hertz. The IR spectra were recorded using either a thin film between NaCl plates or KBr discs, as specified. Thin layer chromatography (TLC) was performed on aluminium sheets precoated with silica gel 60 (HF_{254} , E. Merck) and spots visualized by UV and charring with H_2SO_4 -EtOH (1:20). Flash column chromatography was carried out with silica gel 60 (0.040-0.630 mm, E. Merck) and using a stepwise solvent polarity gradient (EtOAc-petroleum ether unless otherwise stated) correlated with TLC mobility. Chromatography solvents used were EtOAc and MeOH (Riedel-deHaen) and petroleum ether (b.p. 40-60°C, BDH laboratory supplies). Toluene (Aldrich) and CH_2Cl_2 (Riedel-deHaen) were freshly distilled from calcium hydride.

Methyl 2,3,4-tri-*O*-acetyl-6-deoxy-6-iodo- α -D-mannopyranoside 9a

To a solution of methyl α -D-mannopyranoside (9 g, 46.3 mmol) in toluene (200 mL) at 80 °C was added triphenylphosphine (13.9 g, 53.3 mmol) and imidazole (10.9 g, 160 mmol). The solution was stirred for 10 min and then iodine (17.2 g, 66.7 mmol) was added portionwise over 0.5 h. After a further 2 h heating the solution was cooled and the intermediate extracted into water (3 x 200 mL). The combined water layers were then washed with toluene, the water removed *in vacuo* and the residue dissolved in acetic anhydride and pyridine (1:1, 200 mL). The solution was left to stand for 12 h after which time pyridine and acetic anhydride were removed *in vacuo* and xylene was evaporated from residue (2 x 50 mL). The residue (a dark syrup) was reacted without further purification in the next step. A small portion of the material was purified further by chromatography to obtain an analytical sample of the title compound¹ and the remainder used in the next reaction without further purification; $[\alpha]_{\text{D}} +42.4^\circ$ (*c* 2.5, CHCl_3); ^1H NMR (300 MHz, CDCl_3): δ 5.31 (dd, 1H, J = 3.5 and 10.0), 5.22 (dd, 1H, J = 1.5 and 10.0), 5.11 (apparent t, 1H, J = 9.5), 4.73 (d, 1H, J = 1.5), 3.80 (apparent dt, 1H, J = 2.5 and 9.5) 3.48 (s, 3H, OCH_3), 3.31 (dd, 1H, J = 10.5 and 2.5), 3.18 (dd, 1H, J = 10.5 and 9.5), 2.15, 2.07, 1.99 (each s, each CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 170.2, 170.1, 170.0 (each s, each $\text{C}=\text{O}$), 98.8 (d, C-1), 70.4, 69.9,

¹ Garegg, P. J.; Samuelsson, B. *J. Chem. Soc. Perkin Trans. 1* **1980**, 2866.

69.0, (each d), 55.8 (q, OCH₃), 21.1, 21.0, 20.9 (each q, each CH₃), 4.1 (t, C-6); IR (KBr): 1748 (C=O), 1376, 1245, 1135, 1042, 972 cm⁻¹. Anal. Calcd. for C₁₃H₁₉O₈I: C, 36.28; H, 4.42; I, 29.54. Found: C, 36.24; H, 4.48; I, 29.62.

2,3,4-Tri-*O*-acetyl-6-deoxy-6-iodo- α -D-mannopyranosyl azide

The iodide **9a** was dissolved in acetic anhydride:sulphuric acid (50:1, 250 mL) and the mixture allowed to stir for 12 h at rt. The solution was then concentrated *in vacuo* and the residue separated between EtOAc and water. The organic layer was washed several times with aq. NaHCO₃ (5 x 500 mL), then water (2 x 500 mL) and the removal of solvent *in vacuo* gave 1,2,3,4-tetra-*O*-acetyl-6-deoxy-6-iodo- α -D-mannopyranose as a pale yellow oil which was used in the next reaction without further purification; [α]_D +39.7° (*c* 1.5, CHCl₃); ¹H NMR (300 MHz, CDCl₃): δ 6.10 (d, 1H, *J* = 1.5), 5.34 (dd, 1H, *J* = 3.5 and 10.0), 5.24 (overlapping signals 2H), 3.80 (apparent dt, 1H, *J* = 2.5, 7.0 and 9.0), 3.32 (dd, 1H, *J* = 11.0 and 2.5), 3.17 (dd, 1H, *J* = 11.0 and 7.0), 2.17, 2.08, 2.01 (each s, 9H); ¹³C NMR (125 MHz, CDCl₃): δ 170.1, 169.9, 169.7, 168.1 (each s, each C=O), 90.6 (d, C-1), 72.0, 69.8, 68.7, 68.6 (each d), 21.0, 20.9, 20.9, 20.7 (each q, each CH₃), 3.7 (t, C-6); IR (KBr): 1748 (C=O), 1371, 1220, 1151, 1058 cm⁻¹. CI-HRMS: Found 476.0422, required 476.0418 (M+NH₄)⁺. To a solution of this intermediate in CH₂Cl₂ (250 mL) under an inert atmosphere was added first SnCl₄ (3 mL, 25 mmol), then TMS-N₃ (10 mL, 76 mmol). The resulting solution was stirred at room temp for 10 min, then diluted with CH₂Cl₂ (300 mL), washed with aq. NaHCO₃ (4 x 500 mL), water (2 x 500 mL), dried over magnesium sulphate; the solvent was removed to give the title compound as a pale oil (quantitative yield) which was used in the next reaction without further purification; [α]_D +86.5° (*c* 1.1, CHCl₃); ¹H NMR (300 MHz, CDCl₃): δ 5.41 (d, 1H, *J* = 2.0), 5.24 (dd, 1H, *J* = 3.5 and 9.5), 5.16 (apparent t, 1H, *J* = 9.5), 5.12 (dd, 1H, *J* = 2.0 and 3.5), 3.92 (ddd, 1H, *J* = 3.0 and 8.0), 3.35 (dd, 1H, *J* = 11.0 and 3.0), 3.18 (dd, 1H, *J* = 11.0 and 8.0), 2.16, 2.08, 1.99 (each s, 9H); ¹³C NMR (125 MHz, CDCl₃): δ 170.0, 169.9, 169.8 (each s, each C=O), 87.6 (d, C-1), 72.1, 69.8, 69.5, 69.2 (each d), 21.0, 20.9, 20.8 (each q, each CH₃), 3.5 (t, C-6); IR (thin film): 2121 (N₃), 1753 (C=O), 1372, 1235, 1126, 1055, 949 cm⁻¹. CI-HRMS: Found 459.0372, required 459.0377 (M+NH₄)⁺.

2,3,4-Tri-*O*-acetyl-6-deoxy- α -D-lyxo-hex-5-enopyranosyl azide **10a**

2,3,4-Tri-*O*-acetyl-6-deoxy-6-iodo- α -D-mannopyranosyl azide from the previous reaction and DBU (14.4 mL, 96.4 mmol) in toluene (150 mL) was stirred and heated at reflux for 1.5 h. The mixture was

then washed with water, concentrated and the residue chromatographed to give **10a** as a clear oil (7.64 g, 50 % from methyl α -D-mannopyranoside); $[\alpha]_D +129.6^\circ$ (*c* 0.5, CHCl_3); ^1H NMR (300 MHz, CDCl_3): δ 5.67 (apparent dt, 1H, $J = 2.0$ and 9.5), 5.36 (d, 1H, $J = 3.0$), 5.28 (dd, 1H, $J = 3.0$ and 9.5), 5.19 (apparent t, 1H, $J = 9.5$), 4.95 (apparent t, 1H, $J = 2.0$), 3.18 (apparent t, 1H, $J = 2.0$), 2.16, 2.15, 2.03 (each s, 9H, each CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 169.9, 169.7, 169.6 (each s, each $\text{C}=\text{O}$), 151.3 (s, C-5), 99.4 (t, C-6), 87.5 (d, C-1), 69.0, 68.2, 67.2 (each d), 20.9, 20.9, 20.8 (each q, each CH_3); IR (KBr): 2118 (N_3) 1751 ($\text{C}=\text{O}$), 1663 ($\text{C}=\text{C}$), 1432, 1373, 1210 cm^{-1} . CI-HRMS: Found 331.1249, required 331.1254 ($\text{M}+\text{NH}_4$) $^+$.

2,3,4-Tri-*O*-acetyl-5,6-anhydro-5-*C*-hydroxy- α -D-mannopyranosyl azide and 2,3,4-tri-*O*-acetyl-5,6-anhydro-5-*C*-hydroxy- β -L-gulopyranosyl azide 11a

To a cold (0 $^\circ\text{C}$) solution of **10a** (0.5 g, 1.6 mmol) in acetonitrile (10 mL) was added aqueous sodium EDTA (9 mL, 0.4 M), then 1,1,1 trifluoroacetone (1.2 mL, 12.8 mmol) via a precooled syringe. To this was added sodium hydrogen carbonate (0.8 g, 9.0 mmol) and oxone (3.5 g, 6.5 mmol), portionwise over 1 h. The resulting suspension was allowed to warm to room temp and after a further 0.5 h the reaction was poured onto ice (25 g). The resulting solution was extracted into CH_2Cl_2 (3 x 50 mL), washed several times with water and concentrated *in vacuo* to provide a mixture of epoxides **11a** (97 %), which was used in the next reaction without further purification; CI-HRMS: Found 347.1202, required 347.1203 ($\text{M}+\text{NH}_4$) $^+$; ^1H NMR (Isomer 1, 300 MHz, CDCl_3): δ 5.79 (d, 1H, $J = 10.5$), 5.53 (dd, 1H, $J = 3.5$ and 10.5), 5.39 (d, 1H, $J = 2.0$), 5.27 (dd, 1H, $J = 2.0$ and 10.5), 2.97 (d, 1H, $J = 4.0$), 2.65 (d, 1H, $J = 4.0$), 2.18, 2.07, 2.01 (each s, 9H); ^{13}C NMR (125 MHz, CDCl_3): δ 170.2, 169.9, 169.6 (each s, each $\text{C}=\text{O}$), 87.7 (d, C-1), 80.8 (s, C-5), 69.5, 66.7, 64.1 (each d), 47.3 (t, C-6), 20.9, 20.8, 20.7 (each q, each CH_3); ^1H NMR (Isomer 2, 300 MHz, CDCl_3): δ 5.39 (d, 1H, $J = 6.5$), 5.37 (dd, 1H, $J = 3.0$ and 6.5), 5.39 (dd, 1H, $J = 2.0$ and 6.5), 5.27 (d, $J = 2.0$), 2.90 (d, 1H, $J = 4.5$), 2.96 (d, 1H, $J = 4.5$), 2.14, 2.12, 2.11 (each s, 9H).

2,3,6-Tri-*O*-acetyl-5-*C*-methoxy- β -L-gulopyranoside 13

The mixture of epoxides **11a** (0.16 g, 0.46 mmol) were dissolved in (5 mL) and heated at reflux overnight. The solvent was removed *in vacuo* and chromatography gave **13** (0.17 g, 98 %); ^1H NMR (300 MHz, CDCl_3): δ 5.25 (apparent t, 1H, $J = 3.5$), 5.19 (dd, 1H, $J = 9.0$ and 3.5), 5.05 (d, 1H, $J = 9.0$), 4.42 (d, 1H, $J = 12.5$), 2.97 (d, 1H, $J = 12.5$), 3.88 (apparent t, 1H, $J = 3.5$), 3.79 (d, 1H, $J = 3.5$), 3.37 (s,

3H) 2.12, 2.10, 2.09 (each s, 9H); ^{13}C NMR (125 MHz, CDCl_3): δ 171.5, 170.7, 169.7 (each s, each C=O), 101.7 (s, C-5), 81.4 (d, C-1), 70.8, 66.9, 66.8 (each d), 59.9 (t, C-6), 49.0 (q, OCH_3), 21.1, 21.0, 20.9 (each q, each CH_3); CI-HRMS: Found 379.1468, required 379.1465 ($\text{M}+\text{NH}_4$) $^+$.

5-C-Methoxy- β -L-gulopyranosyl azide **14**

Compound **13** (0.12 g, 0.32 mmol) was dissolved in MeOH (5 mL) and elemental sodium (10 mg) was added. The solution was stirred for 4 h and then amberlite IR-120 (plus) ion exchange resin was added and stirring continued. After 10 min. the solution was filtered and concentrated to give **14** (quantitative) as a pale brown oil. Attempts to purify by chromatography led to substantial loss of **14** by decomposition and so the residue from the reaction was used without purification in the next step; $[\alpha]_{\text{D}}^{25} +83.6^\circ$ (*c* 0.75, MeOH); ^1H NMR (300 MHz, D_2O): δ 5.02 (d, 1H, $J = 9.0$), 4.02 (br s, 1H), 3.91 (br s, 1H), 3.76-3.64 (overlapping signals, 3H), 3.34 (s, 3H); ^{13}C NMR (125 MHz): δ 103.3 (d, C-5), 84.0 (d, C-1), 71.6, 68.0, 67.2 (each d), 57.7 (t, C-6), 27.7 (q, OCH_3); IR (KBr): 3414 (OH), 2123 (N_3), 1247, 1073 cm^{-1} . CI-HRMS: Found 253.1145, required 253.1148 ($\text{M}+\text{NH}_4$) $^+$.

1-Deoxymannojirimycin **2**

Compound **14** (0.40 g, 1.7 mmol) was dissolved in ethanol (20 mL) and shaken in a Parr Hydrogenator under an atmosphere of hydrogen (40 p.s.i) in the presence of Pd-C (0.1 g) for 12 h. The suspension was filtered and the solvent was removed *in vacuo* to give the title compound (0.23 g). Some of the residue (0.15 g) was dissolved in MeOH (20 mL) and HCl (1 M in Et_2O , 1.5 mL) was added. The solvents were evaporated and the salt was chromatographed (eluting with -chloroform-triethylamine, 100:100:1). This gave the free amine (64 mg, 23 %); treatment with HCl as above gave the hydrochloride salt. The NMR data of the salt agreed with those of an authentic sample purchased from Sigma; ^1H NMR (300 MHz, D_2O , HCl salt): δ = 4.09 (apparent dt, 1H, $J = 2.7$, 2.7 and 1.2), 3.82 (dd, 1H, $J = 12.0$ and 3.0), 3.69 (apparent t, 1H, $J = 9.6$), 3.64 (dd, $J = 5.2$ and 12.0), 3.53 (dd, 1H, $J = 2.7$ and 9.6), 3.24 (dd, 1H, $J = 2.7$ and 14.4), 3.06 (dd, 1H, $J = 1.2$ and 14.4), 2.96 (ddd, 1H, $J = 5.2$, 3.3 and 9.6); ^{13}C NMR (75MHz, D_2O , free base) δ 48.2 (C-1), 60.3 (C-6), 60.6 (C-5), 67.9, 68.6, 74.2 (C-2, C-3, C-4).

Methyl 2,3,4-tri-*O*-acetyl-6-deoxy-6-iodo- α -D-glucopyranoside **9b**

Methyl α -D-glucopyranoside (9 g, 46.3 mmol) was treated as described above for methyl α -D-mannopyranoside and gave the title compound¹ which was used without purification in the next

reaction; +114.6 ° (*c* 0.56, acetone); ^1H NMR (300 MHz, CDCl_3): δ 5.46 (dd, 1H, J = 9.4 and 10.1), 4.96 (d, 1H, J = 3.7), 4.88 (dd, 1H, J = 3.7 and 10.1), 4.87 (apparent t, 1H, J = 10.1), 3.79 (ddd, 1H, J = 9.4, 2.5 and 8.3), 3.48 (s, 3H), 3.30 (dd, 1H, J = 10.9 and 2.5), 3.13 (dd, 1H, J = 10.9 and 8.3), 2.07, 2.05, 2.0 (each s, 9H); ^{13}C NMR (125 MHz, CDCl_3): δ 199.5, 170.0, 169.9 (each s, each C=O), 96.6 (d, C-1), 72.4, 70.8, 69.0, 68.6 (each d), 55.7 (q, OCH_3), 20.6, 20.6, 20.5 (each q, each CH_3), 3.79 (t, C-6); IR (KBr): 1748 (C=O), 1376, 1245, 1135, 1042, 972 cm^{-1} . Anal. Calcd. for $\text{C}_{13}\text{H}_{19}\text{O}_8\text{I}$: C, 36.28; H, 4.42; I, 29.54. Found: C, 36.24; H, 4.48; I, 29.62. CI-HRMS: Found 448.0470, required 448.0468 ($\text{M}+\text{NH}_4$) $^+$.

1,2,3,4-Tetra-*O*-acetyl-6-deoxy-6-iodo- α -D-glucopyranose

All of the unpurified iodide **9a** obtained as described above was dissolved in acetic anhydride and sulphuric acid (50:1, 250 mL) and the mixture was stirred for 12 h at rt. The solution was then concentrated *in vacuo* and residue was separated between EtOAc and water. The organic layer was washed with aq. NaHCO_3 (5 x 500 mL), water (2 x 500 mL) and dried and the title compound was obtained as a pale yellow oil after the removal of solvent. A small portion was further purified by chromatography (EtOAc: petroleum ether gradient commencing with 1:4) so as to obtain an analytical sample of the title compound; $[\alpha]_D$ +86.5° (*c* 0.2, CHCl_3); ^1H NMR (300 MHz, CDCl_3): δ 6.34 (d, 1H, J = 3.7), 5.47 (apparent t, 1H, J = 9.8), 5.08 (dd, 1H, J = 3.7 and 9.8), 4.98 (apparent t, 1H, J = 9.8), 3.83 (ddd, 1H, J = 2.9 and 6.3), 3.31 (dd, 1H, J = 11.2 and 2.9), 3.14 (dd, 1H, J = 11.2 and 6.3), 2.18, 2.07, 2.02, 2.01 (each s, each CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 170.0, 169.8, 169.5, 167.1 (each s, each C=O), 90.8 (d, C-1), 72.1, 69.9, 68.6, 68.5 (each d), 21.0, 20.9, 20.8, 20.6 (each q, each CH_3), 3.7 (t, C-6); IR (KBr): 1748, 1371, 1220, 1151, 1058 cm^{-1} . CI-HRMS: Found 476.0416, required 476.0418 ($\text{M}+\text{NH}_4$) $^+$. Anal. Calcd. for $\text{C}_{14}\text{H}_{19}\text{O}_9\text{I}$: C, 36.70; H, 4.18; I, 27.7. Found: C, 36.79; H, 4.01; I, 27.51.

2,3,4-Tri-*O*-acetyl-6-deoxy-6-iodo- β -D-glucopyranosyl azide

To a solution of 1,2,3,4-tetra-*O*-acetyl-6-deoxy-6-iodo- α -D-glucopyranose, used as obtained in the above reaction without purification, in CH_2Cl_2 (250 mL) under an inert atmosphere was added first SnCl_4 (3.0 mL, 25 mmol), then azidotrimethylsilane (10.0 mL, 76 mmol). The resulting solution was stirred at room temp for 10 min, then diluted with CH_2Cl_2 (300 mL), washed with aq. NaHCO_3 (4 x 500 mL), water (2 x 500 mL) and dried over magnesium sulphate. The solvent was removed *in vacuo* to give the title compound as a pale oil; $[\alpha]_D$ +86.5° (*c* 0.2, CHCl_3); ^1H NMR (300 MHz, CDCl_3): δ 5.23

(apparent t, 1H, $J = 9.4$), 4.95 (2 x overlapping apparent t, 2H, $J = 9.4$), 4.70 (d, 1H, $J = 9.4$), 3.57 (ddd, 1H, $J = 2.6$, 7.2 and 9.4), 3.34 (dd, 1H, $J = 11.1$ and 2.6), 3.17 (dd, 1H, $J = 11.1$ and 7.2), 2.07, 2.07, 2.06, 2.06 (each s, 12H); ^{13}C NMR (125 MHz, CDCl_3): δ 170.0, 169.9, 169.8, (each s, each C=O), 87.6 (d, C-1), 72.2, 69.7, 69.6, 68.2 (each d), 20.0, 19.0, 18.7, (each q, each CH_3), 3.7 (t, C-6); IR (thin film): 2120 (N_3), 1742, 1376, 1321, 1126, 1090, 820 cm^{-1} . CI-HRMS: Found 459.0373, required 459.0377 ($\text{M}+\text{NH}_4$) $^+$.

2,3,4-Tri-*O*-acetyl-6-deoxy- β -D-xylo-hex-5-enopyranosyl azide **10b**

A solution of 2,3,4-tri-*O*-acetyl-6-deoxy-6-iodo- β -D-glucopyranosyl azide and DBU (14.4 mL, 96.4 mmol) in toluene (150 mL) was stirred and heated at reflux for 1.5 h, after which time the UV active iodide starting material had been consumed. The solution was washed with water, the solvent was removed and the residue chromatographed (EtOAc: petroleum ether gradient, commencing with 1:6, as eluant) to give the title compound² as a clear oil (7.53 g, 52% from methyl α -D-glucopyranoside); $[\alpha]_{\text{D}}^{40}$ $^\circ$ (c 0.08, acetone); ^1H NMR (300 MHz, CDCl_3): δ 5.52 (apparent dt, 1H, $J = 1.6$ and 7.5), 5.16 (apparent t, 1H, $J = 7.5$), 4.98 (apparent t, 1H, $J = 7.5$), 4.93-5.04 (overlapping d ($J = 7.5$) and apparent t ($J = 1.6$), 2H), 4.65 (apparent t, 1H, $J = 1.6$), 2.12, 2.09, 2.05 (each s, each CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 169.9, 169.3, 169.2 (each s, each C=O), 150.7 (s, C-5), 98.3 (t, C-6), 88.2 (d, C-1), 71.6, 70.9, 68.6 (each d), 20.9, 20.8, 20.77 (each q, each CH_3); IR (KBr): 2105 (N_3), 1749 (C=O), 1635, 1432, 1375, 1048 cm^{-1} . Anal. Calcd. for $\text{C}_{12}\text{H}_{15}\text{O}_7\text{N}_3$: C, 46.12; H, 4.81; N, 13.25. Found: C, 46.01; H, 4.83; N, 13.41.

2,3,4-Tri-*O*-acetyl-5,6-anhydro-5-*C*-hydroxy- β -D-glucopyranosyl azide and 2,3,4-tri-*O*-acetyl-5,6-anhydro-5-*C*-hydroxy- α -L-idopyranosyl azide **11b**

To a cold (0 $^\circ\text{C}$) solution of **10b** (0.5 g, 1.6 mmol) in acetonitrile (10 mL) was added aqueous sodium EDTA (9 mL, 0.4 M), then 1,1,1 trifluoroacetone (1.2 mL, 12.8 mmol) *via* a precooled syringe. To this was added sodium hydrogen carbonate (0.8 g, 9.0 mmol) and oxone (3.5 g, 6.5 mmol), portionwise over 1 h. The resultant suspension was allowed to warm to room temp and after a further 0.5 h, was poured onto ice (25 g). The products were extracted into CH_2Cl_2 (3 x 50 mL), washed several times with water and the solvent then removed *in vacuo* (95 %) to give the title compounds as a 1:1.7 mixture. Anal. Calcd. for $\text{C}_{12}\text{H}_{15}\text{O}_8\text{N}_3$ (mixture): C, 43.77; H, 4.59; N, 12.76. Found: C, 43.86; H, 4.75; N, 12.79.

NMR data for isomer 1: ^1H NMR (300 MHz, CDCl_3): δ 5.53 (d, 1H, J 9.5), 5.43 (apparent t, 1H, J = 9.5), 5.1 (apparent t, 1H, J = 9.5), 4.97 (d, 1H, J = 9.5), 3.15 (d, 1H, J = 4.0), 2.78 (d, 1H, J = 4.0), 2.09, 2.04, 2.02 (each s, 9H); ^{13}C NMR (125 MHz, CDCl_3), δ 169.8, 169.7, 169.4 (each s, each C=O), 86.1 (d, C-1), 80.7, (s, C-5), 70.8, 70.7, 66.1 (each d), 49.8 (t, C-6), 20.7, 20.6 (each q, each CH_3).

NMR data for isomer 2: ^1H -NMR (300 MHz, CDCl_3): δ 5.34 (d, 1H, J = 7.5), 5.28 (apparent t, 1H, J = 7.5), 5.09 (apparent t, 1H, J = 7.5), 4.95 (d, 1H, J = 7.5), 2.97 (d, 1H, J = 4.9), 2.95 (d, 1H, J = 4.9), 2.11, 2.06, 2.05 (each s, 9H); ^{13}C NMR (125 MHz, CDCl_3), δ 170.0, 169.3, 169.0 (each s, each C=O), 87.1 (d, C-1), 80.8, (s, C-5), 70.9, 70.3, 67.5 (each d), 49.0 (t, C-6), 20.8 and 20.5 (each q, each CH_3).

Acyclic compound 19

Epoxides **11b** (0.75 g, 2.0 mmol) were dissolved in (23 mL) and the mixture was heated at reflux overnight. The solution was allowed to cool and pyridine (7.0 mL) and acetic anhydride (7.0 mL) were added. The solution was stirred for 30 min after which time xylene was added and the volatile substances were removed *in vacuo*. The residue was purified by chromatography (EtOAc: petroleum ether gradient, commencing with 1:3, as eluant) which gave **19** as a white solid (0.45 g, 58%); $[\alpha]_D^{+42.7^\circ}$ (c 0.61, CHCl_3); ^1H NMR (300 MHz, CDCl_3): δ 5.80 (overlapping dd and d, 2H, J = 2.8 and 6.5), 5.61 (d, 1H, J = 3.4), 5.34 (dd, 1H, J = 6.5 and 2.8), 4.12 (2 x d, 2H, J = 17.0), 3.46, 3.42 (each s, 6H), 2.18, 2.13, 2.08, 2.04 (each s, 12H); ^{13}C NMR (125 MHz, CDCl_3): δ 201.6, 170.4, 170.2, 170.0, 169.8 (each s, each C=O), 95.5 (d), 76.5 (t), 74.3, 70.8, 68.3 (each d), 59.7, 57.7 (each q, each OCH_3), 21.0, 20.7, 20.7, 20.5 (each q, each CH_3); IR (KBr): 2941, 2124 (N_3), 1750 (C=O), 1432, 1376, 1213; CI-HRMS: Found 410.1665, required 410.1662 ($\text{M}+\text{NH}_4$) $^+$. Anal. Calcd. for $\text{C}_{16}\text{H}_{24}\text{O}_{11}$: C, 48.98; H, 6.17; N. Found: C, 48.97; H, 6.13.

1,2:3,4-Di-*O*-isopropylidene-6-deoxy-6- α -iodo-D-galactopyranoside

To 1,2:3,4-di-*O*-propylidene- α -D-galactopyranoside (25 g, 96.1 mmol) in toluene (400 mL) at 80 $^\circ\text{C}$, triphenylphosphine (15.4 g, 115.3 mmol), imidazole (31.4 g, 461.2 mmol) and iodine (48.8 g, 192.2 mmol) were added successively in small portions. The mixture was allowed to stir for 10 h, cooled to room temp and washed with 10% aqueous sodium thiosulphate (2 x 400 mL), water (2 x 400 mL), dried (Na_2SO_4), filtered, and the solvent removed *in vacuo*. Chromatography (6:1 petroleum ether-EtOAc) gave the title compound as a white solid (33.4 g, 94 %); R_F 0.76 (3:1 petroleum ether-EtOAc); m.p. 64-66 $^\circ\text{C}$ $[\alpha]_D^{-44.2^\circ}$ (c 0.5 CHCl_3); ^1H NMR (CDCl_3 , 300MHz); δ 5.55 (d, 1H, J 5.0), 4.63 (dd, 1H, J =

7.9 and 2.5), 4.42 (dd, 1H, $J = 7.9$ and 1.9), 4.31 (dd, 1H, $J = 5.0$ and 2.5), 3.96 (apparent dt, 1H, $J = 7.0$ and 1.9), 3.33 (dd, 1H, $J = 10.0$ and 7.0), 3.22 (dd, 1H, $J = 10.0$ and 7.0), 1.56, 1.46, 1.37, 1.35 (each s, 12H); ^{13}C NMR (CDCl_3 , 125MHz); δ 109.7, 109.0 (each s, 2 x $\text{C}(\text{CH}_3)_2$), 96.9 (d, C-1), 71.7, 71.3, 70.8, 69.2 (each d, C-2-5), 26.2, 26.1, 25.0, 24.6 (each q, 4 x $\text{C}(\text{CH}_3)_2$), 2.4 (t, C-6); IR (KBr) ν 2989, 1383, 1066 cm^{-1} . Anal. Calcd. for $\text{C}_{12}\text{H}_{19}\text{IO}_5$ requires C, 38.93; H, 5.17; I, 34.28; Found C, 38.85; H, 5.06; I, 34.48; HRMS-Cl: Found 388.0621, required 388.0622 $[\text{M}+\text{NH}_4]^+$.

1,2,3,4-Tetra-*O*-acetyl-6-deoxy-6-iodo-D-galactopyranose

To 1,2,3,4-Di-*O*-isopropylidene-6-deoxy-6-iodo-D-galactopyranoside (15 g, 40.5 mmol) stirring in acetic acid (300 mL) at 0 °C, acetic anhydride (75 mL) and H_2SO_4 (40 mL) was added slowly over 20 min. The solution was allowed to warm to 35 °C and was stirred at this temperature for a further 2 days. The mixture was poured onto ice water (300 mL) and extracted with CH_2Cl_2 (2 x 300 mL). The combined organic layers were washed with water (2 x 300 mL), dried (Na_2SO_4), filtered and the solvent was removed *in vacuo* to give the title compound ($\alpha:\beta$, 5:1) as a yellow oil (11.6 g, 62 %); R_F 0.40 (2:1 petroleum ether-EtOAc); HRMS-Cl: Found 476.0414 required 476.0418 $[\text{M}+\text{NH}_4]^+$.

2,3,4-Tri-*O*-acetyl-6-deoxy-6-iodo- β -D-galactopyranosyl azide

To 1,2,3,4-tetra-*O*-acetyl-6-deoxy-6-iodo-D-galactopyranose (6.33 g, 13.8 mmol) in dry CH_2Cl_2 (90 mL) under a N_2 atmosphere at 0 °C, SnCl_4 (0.81 mL, 6.9 mmol) was added dropwise and mixture was stirred for 20 min. Azidotrimethylsilane (3.66 mL, 27.6 mmol) was then added dropwise and the resulting solution was allowed warm to and stirred at room temp for a further 16 h. The mixture was diluted with CH_2Cl_2 (50 mL) and washed successively with aq. NaHCO_3 (150 mL) and brine (150 mL). The organic phase was dried (Na_2SO_4), filtered and the solvent removed *in vacuo*. Chromatography (4:1 petroleum ether-EtOAc) gave the title compound² as a white solid (6.08 g, 91 %); R_F 0.55 (1:1 petroleum ether-EtOAc); m.p. 70-72 °C; $[\alpha]_D +39.6^\circ$ (c 0.5, CHCl_3); ^1H NMR (CDCl_3 , 300 MHz); δ 5.61 (dd, 1H, $J_{4,3}$ 3.2, 1.0, H-4), 5.13 (dd, 1H, $J_{2,3}$ 10.4, $J_{2,1}$ 8.5, H-2), 5.04 (dd, 1H, $J_{3,2}$ 10.4, $J_{3,4}$ 3.2, H-3), 4.64 (d, 1H, $J_{1,2}$ 8.5, H-1), 4.0 (apparent t, $J_{5,6a}$ 7.0, $J_{5,6b}$ 7.0, 1H, H-5), 3.25 (dd, 1H, $J_{6b,6a}$ 10.5, $J_{6b,5}$ 7.0, H-6b), 3.16 (dd, 1H, $J_{6a,6b}$ 10.5, $J_{6a,5}$ 7.0, H-6a), 2.19, 2.09, 2.05 (each s, 9H, each CH_3); ^{13}C NMR (CDCl_3 , 125MHz); δ 170.1, 170.1, 169.5 (each s, each C=O), 88.3 (d, C-1), 76.0, 70.9, 68.1, 67.9 (each d, C-

² Györgydeák, Z.; Szilagyi, L. *Liebigs Ann. Chem.* **1987**, 235.

2-5), 20.8, 20.7, 20.6 (each q, each CH₃), -1.1 (t, C-6); IR (KBr) ν 2946, 2119 (N₃), 1753 (C=O), 1371, 1221 cm⁻¹. Anal. Calcd. for C₁₂H₁₆IN₃O₅ requires C, 32.67; H, 3.66; I, 28.77; N, 9.52; Found C, 32.63; H, 3.52; I, 28.60; N, 9.34. HRMS-Cl: Found 459.0377 required 459.0377 [M+NH₄]⁺.

2,3,4-Tri-*O*-acetyl-6-deoxy- β -D-arabino-hex-5-enopyranosyl azide **20**

To 2,3,4-tri-*O*-acetyl-6-deoxy-6-iodo- β -D-galactopyranosyl azide (5.30 g, 12.0 mmol) in dry CH₂Cl₂ (80 mL) under a N₂ atmosphere at 35 °C, DBU (9.05 mL, 60.0 mmol) was added. The solution was stirred for 2 days, cooled and was directly added to a column of silica gel and chromatographed (6:1 petroleum ether-EtOAc) to give **20**² as a white solid (2.96 g, 78 %); R_F 0.55 (1:1 petroleum ether-EtOAc); m.p. 71-72 °C; [α]_D -52.0 ° (*c* 0.5 CHCl₃); ¹H NMR (300 MHz, CDCl₃); δ 5.72 (d, 1H, *J*_{1,2} 3.5, H-1), 5.25 (dd, 1H, *J*_{3,2} 8.8, *J*_{3,4} 6.7, H-3), 5.11 (dd, 1H *J*_{2,3} 8.8, *J*_{2,1} 3.5, H-2), 5.01 (d, 1H, *J*_{6,6a} 1.6, H-6), 4.85 (d, 1H, *J*_{4,3} 6.7, H-4), 4.84 (d, 1H, *J*_{6a,6} 1.6, H-6a), 2.16, 2.14, 2.07 (each s, 9H, each CH₃); ¹³C NMR (125 MHz, CDCl₃); δ 169.9, 169.8, 169.4 (each s, each C=O), 150.3 (s, C-5), 102.8 (t, C-6), 88.6 (d, C-1), 69.3, 68.2, 67.8 (each d, C-2-C-4), 20.9, 20.7, 20.6 (each q, each C=O); IR (KBr) ν 2938, 2117 (N₃), 1752 (C=O), 1368, 1240, 1071 cm⁻¹. Anal. Calcd. for C₁₂H₁₅N₃O₇ required: C, 46.01; H, 4.83, N, 13.41; Found C, 45.85; H, 4.75; N, 13.36; HRMS-Cl: Found 331.1254, required 331.1257 [M+NH₄]⁺.

5-*C*-Methoxy- β -D-glucopyranosyl azide and 5-*C*-methoxy- α -L-idopyranosyl azide

These two isomers were obtained by deacetylation of **12**. Analytical data for first isomer: [α]_D -59.4° (*c* 0.5, MeOH); ¹H NMR (300 MHz, D₂O): δ 4.92 (d, 1H, *J* = 9.0), 3.91 and 3.78 (each d, each 1H, *J* = 12.9 and 12.9), 3.85 (d, 1H, *J* = 9.0), 3.70 (apparent t, 1H, *J* = 9.0), 3.46 (apparent t, 1H, *J* = 9.0), 3.26 (s, 3H); ¹³C NMR (125 MHz): δ 101.6 (s, C-5), 86.9 (d, C-1), 72.9, 72.8, 70.5 (each d), 58.9 (t, C-6), 27.7 (q, OCH₃). Analytical data for second isomer: [α]_D -93.3° (*c* 0.14, MeOH); ¹H NMR (300 MHz, CD₃OD): δ 4.59 (d, 1H, *J* 9.0), 3.75 and 3.72 (each d, each 1H, *J* = 11.7 and 11.7), 3.62 (apparent d, 2H, *J* 4.7), 3.36 (s, 3H, OCH₃), 3.16 (dd, 1H, *J* 9.0 and 4.7, H-2); ¹³C NMR (125 MHz, CD₃OD): δ 100.5 (s, C-5), 86.3 (d, C-1), 73.4, 72.7, 71.5 (each d), 50.8 (t, C-6), 47.8 (q, OCH₃).

NMR data for 1-deoxynojirimycin **1** (salt)

¹H NMR (300 MHz, D₂O): δ 3.78 (dd, 1H, *J* = 3.0 and 12.0), 3.59 (dd, 1H, *J* = 6.0 and 12.0), 3.45 (ddd, 1H, *J* = 5.1, 10.9 and 9.5), 3.28 (apparent t, 1H, *J* = 9.5), 3.21 (apparent t, 1H, *J* = 9.5), 3.10 (dd, 1H, *J* =

5.1 and 12.4), 2.55 (ddd, 1H, $J = 9.5, 6.0$ and 3.0), 2.45 (dd, 1H, $J = 10.9$ and 12.4); ^{13}C NMR (75MHz, D_2O) δ 49.2, 61.3, 62.8, 70.3, 71.4, 79.3.

NMR data for 1-deoxygalactostatin 3

^1H NMR (D_2O , 300 MHz); δ 4.10 (apparent d, 1H, $J = 3.0$), 3.99 (ddd, 1H, $J = 11.5, 10.0, 5.0$), 3.79 (dd, 1H, $J = 11.5, 7.0$), 3.74 (dd, 1H, $J = 11.5, 10.0$), 3.56 (dd, 1H, $J = 1.5$ and 7.0) 3.41 (dd, 1H, $J = 10.0$ and 3.0), 3.30 (dd, 1H, $J = 12.5$ and 5.0), 2.79 (apparent t, 1H, $J = 11.5$).

NMR data for 1,6-dideoxygalactostatin 4

$[\alpha]_{\text{D}} + 34^\circ$ (c 0.1, MeOH); ^1H NMR (D_2O , 300MHz); δ 3.89-4.00 (m, 2H), 3.57 (dd, 1H, J 9.9, 3.1), 3.37 (m, 2H), 2.77 (t, 1H, J 11.7), 1.27 (d, 3H, J 6.7).

NMR Spectra

 ^1H -NMR (CDCl_3) of 9a ^{13}C -NMR (CDCl_3) of 9a

^1H -NMR (300 MHz, 4.5-6.0 ppm, CDCl_3) of 10aJUS-294-1
STANDARD 1H OBSERVE

Pulse Sequence: zgpg30
Solvent: cdcl_3
Temp: 30.0 C / 293.1 K
File: P_Murphy_03_01
INDRA-500 "sunec001"
PULSE SEQUENCE
Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 3.784 sec
Width 4798.2 Hz
S repetitions
OBSERVE M1 299.906279 MHz
DATA PROCESSING
FT size 65536
Total time 8 min, 38 sec

^{13}C -NMR (CDCl_3) of 10a

^1H -NMR (300 MHz, 4.5-6.0 ppm, CDCl_3) of 10a

^1H NMR (300 MHz, 3-6 ppm, CDCl_3) of 13 **^{13}C NMR of 13 (D_2O)**

^1H NMR (300 MHz, D_2O) of 14 **^{13}C NMR of 14 (D_2O)**

^1H NMR (270 MHz, D_2O), Deoxymannojirimycin (free amine)

^1H NMR (270 MHz, D_2O), Deoxymannojirimycin (HCl salt)

^1H NMR (270 MHz, D_2O), Deoxymannojirimycin (crude product unpurified)

^1H NMR (270 MHz, D_2O), 1-Deoxymannojirimycin (HCl salt) from Sigma

^1H NMR (500 MHz, D_2O) of synthesized deoxymannojirimycin (HCl salt) 2 after chromatography and salt formation

¹H NMR (300 MHz, CDCl₃) of Epoxides 11b

^{13}C NMR (300 MHz, CDCl_3) of Epoxides 11b

^1H -NMR (300 MHz, CDCl_3) of 19

[illegible]

^1H -NMR (300 MHz, CDCl_3) of 12 (Isomer 1)

¹H-NMR (300 MHz, CDCl₃) of 13 (Isomer 2)

¹H-NMR (D₂O, 300 MHz) 1-Deoxynojirimycin (crude product after hydrogenation)

¹H-NMR (D₂O, 300 MHz) 1-Deoxynojirimycin 1 (HCl salt, after purification)

COMPARISON

¹H-NMR (D₂O, 300 MHz) 1-Deoxynojirimycin (HCl salt, Sigma, authentic sample)

^1H -NMR (D_2O , 300 MHz) 1-Deoxynojirimycin (amine, after chromatography)

¹³C-NMR (300 MHz, CDCl₃) of 22 (one diastereoisomer)

Pulse 40.0 usg.vvv
Acq. time 1.199 sec
Width 18850.1 Hz
896 repetitions
OBSERVE C13, 75.4089450 MHz
DECOUPLE H1, 299.8897221 MHz
Power 35 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 1.0 Hz
FT size 65536
Total time 1 hr, 13 min, 36 sec

¹H-NMR (300 MHz, CDCl₃) of 23 (one diastereoisomer)

^{13}C -NMR (300 MHz, CDCl_3) of acetylated 21 (one diastereoisomer)

¹H-NMR (D₂O, 300 MHz) 1-Deoxygalactostatin 3 and 1-Deoxyaltronojirimycin 25 (4:1

^1H -NMR (D_2O , 300 MHz) 1-Deoxy-D-galactostatin 3 (authentic sample)

¹H-NMR (D₂O, 300 MHz) 1,6-dideoxy-D-galactonojirimycin 4 (HCl salt)

¹H-NMR (D₂O, 300 MHz) 1-Deoxy-L-fuconojirimycin (enantiomer of 4, HCl salt) authentic sample