A Highly Efficient, Asymmetric Synthesis of Benzothiadiazine-substituted Tetramic Acids: Potent Inhibitors of Hepatitis C Virus RNA-dependent RNA Polymerase

Duke M. Fitch,^{*,†} Karen A. Evans,[‡] Deping Chai,[†] and Kevin J. Duffy[†]

Departments of [†]Medicinal Chemistry and [‡]Discovery Research, GlaxoSmithKline Pharmaceuticals, 1250 South Collegeville Road, P.O. Box 5089, Collegeville, Pennsylvania 19426-0989

General Information. Starting materials were either commercially available or prepared as reported previously in the literature, unless otherwise noted. Solvents and reagents were used without further purification. Reactions were monitored by TLC, performed on silica gel glass plates containing F-254 indicator (Kieselgel 60 F₂₅₄, Merck or Uniplate Silica Gel GF, Analtech). Visualization on TLC was achieved by UV light, potassium permanganate or iodine indicator. Column chromatography was performed with Merck 40-63 mesh silica gel. Reaction temperatures refer to the oil bath or cold bath temperature. ¹H NMR spectra were recorded on a Bruker ARX-300 or Avance-400 instrument. Chemical shifts (δ) are reported in ppm downfield from internal TMS standard or from solvent references. Mass spectra were recorded on a SciEX API 150EX electrospray LC-mass spectrometer.

1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl-acetic acid (4). A solution of ethyl 1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl-acetate (prepared by the method of Kovalenko, S. N.; Chernykh, V. P.; Shkarlat, A. E.; Ukrainets, I. V.; Gridasov, V. I.; Rudnev, S. A. *Chem. Heterocycl. Compd. (Engl. Trans.)* **1998**, 34, 791) (10.2 g, 38.0 mmol) in tetrahydrofuran (50.0 mL)

was treated with 6M aqueous hydrochloric acid (100.0 mL). After stirring 1 wk at ambient temperature, the reaction mixture was diluted with brine, extracted thrice with EtOAc, dried over magnesium sulfate, filtered, and concentrated in vacuo to give the title compound as a white solid (8.30 g, 91%). ¹H NMR (400MHz, D₆-DMSO) δ 13.0 (br s, 1H), 12.9 (s, 1H), 7.80 (dd, *J* = 1, 8 Hz, 1H), 7.68 (dt, *J* = 1, 7 Hz, 1H), 7.45 (dt, *J* = 1, 7 Hz, 1H), 7.31 (d, *J* = 8 Hz, 1H), 3.57 (s, 2H). MS(ES⁺) *m/e* 241 [M+H]⁺.

Method A

Methyl (S)-2-(3-methylbut-1-yl-amino)-3-phenylpropanoate (3a). A solution of L-phenylalanine methyl ester hydrochloride (1.00 g, 4.64 mmol) in anhydrous methanol (10.0 mL) was treated with triethylamine (0.711 mL, 5.10 mmol), isovaleraldehyde (0.547 mL, 5.10 mmol), and sodium cyanoborohydride (0.204 g, 3.25 mmol). After stirring 3 d at ambient temperature, the reaction mixture was concentrated in vacuo, diluted with brine, extracted twice with EtOAc, dried over

magnesium sulfate, filtered, concentrated in vacuo, and purified *via* flash column chromatography (10-30% EtOAc/hexanes) to give the title compound as a clear, colorless oil (1.05 g, 91%). ¹H NMR (300MHz, CDCl₃) δ 7.35-7.18 (m, 5H), 3.74 (d, *J* = 2 Hz, 1H), 3.64 (s, 3H), 3.52 (t, *J* = 7 Hz, 1H), 2.97

(dd, $J = 3, 9$ Hz, 1H), 2.92 (dd, $J = 3, 9$ Hz, 1H), 2.58 (ddd, $J = 7, 7, 11$ Hz, 1H), 2.46 (ddd, $J = 7, 7, 11$ Hz, 1H), 1.60-1.50 (m, 1H), 1.37-1.28 (m, 2H), 0.85 (d, $J = 7$ Hz, 6H). MS(ES⁺) m/e 250 [M+H]⁺.

Method B

Methyl (R)-2-(3-methylbut-1-yl-amino)-3-phenylpropanoate (ent-3a). A slurry of D-phenylalanine (0.223 g, 1.35 mmol) in anhydrous methanol (5.0 mL) was treated with isovaleraldehyde (0.145 mL, 1.35 mmol) and sodium cyanoborohydride (0.085 g, 1.35 mmol). After stirring 2 d at ambient temperature, concentrated sulfuric acid (2.0 mL) was added and the reaction mixture was heated to reflux. After stirring 24 h at reflux, the reaction mixture was concentrated in vacuo, neutralized with solid sodium carbonate, diluted with brine, extracted thrice with methylene chloride, dried over magnesium sulfate, filtered, concentrated in vacuo, and purified *via* flash column chromatography (10-30% EtOAc/hexanes) to give the title compound as a clear, colorless oil (0.246 g, 73%). See Method A for characterization data.

Method C

Methyl (S)-[[2-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-ethanoyl]-(3-methylbut-1-yl)- amino]-3-phenylpropanoate (5a). A solution of **3a** (0.094 g, 0.379 mmol) and **4** (0.100 g, 0.416 mmol) in anhydrous methylene chloride (3.0 mL) and anhydrous dimethylformamide (1.0 mL) was treated with dicyclohexylcarbodiimide (1M solution in methylene chloride, 0.416 mL, 0.416 mmol). After stirring 2 h at ambient temperature, the reaction mixture was filtered and concentrated in vacuo to give **5** as a yellow oil. ¹H NMR (300MHz, D₆-DMSO) δ 12.2 (s, 1H), 7.83 (dd, $J = 1, 8$ Hz, 1H), 7.70 (dt, $J = 1, 8$ Hz, 1H), 7.47 (dt, $J = 1, 7$ Hz, 1H), 7.37 (d, $J = 8$ Hz, 1H), 7.37-7.15 (m, 5H), 4.28 (dd, $J = 5, 10$ Hz, 1H), 3.77-3.64 (m, 1H), 3.59 (s, 3H), 3.42 (s, 2H), 3.17 (dd, $J = 5, 12$ Hz, 1H), 3.09 (dd, $J = 10, 12$ Hz, 1H), 2.72-2.42 (m, 1H), 1.42-1.30 (m, 1H), 1.27-1.17 (m, 2H), 0.76 (d, $J = 7$ Hz, 3H), 0.73 (d, $J = 7$ Hz, 3H). MS(ES⁺) m/e 472 [M+H]⁺.

(S)-5-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-1,5-dihydropyrrol-2-one (6a). A solution of **5a** (0.070 g, 0.148 mmol) in anhydrous 2-methyl-2-propanol (5.0 mL) was treated with potassium *tert*-butoxide (0.042 g, 0.371 mmol). After stirring 45 min at ambient temperature, the reaction mixture was quenched with 1M aqueous hydrochloric acid, diluted with brine, extracted thrice with methylene chloride, dried over magnesium sulfate, filtered, concentrated in vacuo, and purified *via* flash column chromatography (60-100% EtOAc/hexanes) to give **6a** as a white solid (0.039 g, 60%, 2 steps). ¹H NMR (300MHz, D₆-DMSO) δ 12.7 (s, 1H), 7.61 (dd, $J = 1, 8$ Hz, 1H), 7.50 (dt, $J = 1, 8$ Hz, 1H), 7.29 (dt, $J = 1, 8$ Hz, 1H), 7.29-7.10 (m, 6H), 3.87 (t, $J = 5$ Hz, 1H), 3.62 (ddd, $J = 8, 8, 14$ Hz, 1H), 3.12 (dd, $J = 4, 14$ Hz, 1H),

2.87 (dd, $J = 6, 14$ Hz, 1H), 2.72 (ddd, $J = 7, 7, 14$ Hz, 1H), 1.40-1.26 (m, 1H), 1.26-1.15 (m, 2H), 0.79 (d, $J = 7$ Hz, 3H), 0.76 (d, $J = 7$ Hz, 3H). $[\alpha]_D = -120^\circ$ ($c = 1.0$, CH_2Cl_2). MS(ES+) m/e 440 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{23}\text{H}_{25}\text{N}_3\text{SO}_4$: C, 62.85; H, 5.73; N, 9.56. Found: C, 62.87; H, 5.70; N, 9.52. The product was shown to be of greater than 96% ee by chiral HPLC analysis (Chiralpak AD column, 0.1% trifluoroacetic acid in ethanol, 1 mL/min, T_r (S) = 13.8 min, T_r (R) = 25.9 min).

Method D

(S)-5-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-1,5-dihydropyrrol-2-one (6a). As an alternative to the procedure from Method C, **6a** could be prepared according to the following method: A solution of **3a** (0.052 g, 0.208 mmol) and **4** (0.050 g, 0.208 mmol) in anhydrous methylene chloride (3.0 mL) and anhydrous dimethylformamide (0.5 mL) was treated with dicyclohexylcarbodiimide (1M solution in methylene chloride, 0.210 mL, 0.210 mmol). After stirring 30 min at ambient temperature, triethylamine

(0.087 mL, 0.624 mmol) was added. After stirring 90 min at ambient temperature, the reaction mixture was filtered, concentrated in vacuo, and purified *via* flash column chromatography (60-100% EtOAc/hexanes) to give **6a** as a white solid (0.074 g, 81%). *See above for characterization data.* The product was shown to be of greater than 99% ee by chiral HPLC analysis.

3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-1,5-dihydropyrrol-2-one (6b). Following Method A, except substituting glycine methyl ester hydrochloride for L-phenylalanine methyl ester hydrochloride, followed by Method C, **6b** was obtained as a yellow solid. ^1H NMR (400MHz, $\text{D}_6\text{-DMSO}$) δ 12.5 (s, 1H), 7.78 (dd, $J = 1, 8$ Hz, 1H), 7.65 (ddd, $J = 1, 7, 8$ Hz, 1H), 7.40 (ddd, $J = 1, 7, 8$ Hz, 1H), 7.39 (dd, $J = 1, 8$ Hz, 1H), 3.97 (s, 2H), 3.42 (t, $J = 7$ Hz, 2H), 1.63-1.56 (m, 1H), 1.54-1.45 (m, 2H), 0.97 (d, $J = 7$ Hz, 6H). MS(ES $^+$) m/e 350 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{16}\text{H}_{19}\text{N}_3\text{SO}_4$: C, 55.00; H, 5.48; N, 12.03. Found: C, 54.30; H, 5.55; N, 11.87.

(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-methyl-1-(3-methylbutyl)-1,5-dihydropyrrol-2-one (6c). Following Method A, except substituting L-alanine methyl ester hydrochloride for L-phenylalanine methyl ester hydrochloride, followed by Method C, **6c** was obtained as a grey solid. ^1H NMR (400MHz, $\text{D}_6\text{-DMSO}$) δ 12.8 (s, 1H), 7.66 (dd, $J = 1, 8$ Hz, 1H), 7.55 (ddd, $J = 1, 7, 8$ Hz, 1H), 7.25 (ddd, $J = 1, 7, 8$ Hz, 1H), 7.23 (dd, $J = 1, 8$ Hz, 1H), 3.65 (ddd, $J = 6, 10, 14$ Hz, 1H), 3.55 (q, $J = 7$ Hz, 1H), 3.03 (ddd, $J = 4, 10, 14$ Hz, 1H), 1.60-1.54 (m, 1H), 1.43-1.40 (m, 2H), 1.21 (d, $J = 7$ Hz, 3H), 0.97 (d, $J = 7$ Hz, 3H), 0.96 (d, $J = 7$ Hz, 3H). $[\alpha]_{\text{D}}^{20} = +23.8^\circ$ ($c = 2.4$, CH_2Cl_2). MS(ES $^+$) m/e 364 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{17}\text{H}_{21}\text{N}_3\text{SO}_4$: C, 56.18; H, 5.82; N, 11.61. Found: C, 55.88; H, 5.86; N, 12.03.

3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-5,5-dimethyl-1,5-dihydropyrrol-2-one (6d). Following Method B, except substituting 2-aminoisobutyric acid for D-phenylalanine, followed by Method C, **6d** was obtained as a grey solid. ^1H NMR (400MHz, $\text{D}_6\text{-DMSO}$) δ 12.4 (s, 1H), 7.72 (dd, $J = 1, 8$ Hz, 1H), 7.59 (ddd, $J = 1, 7, 8$ Hz, 1H), 7.34 (dd, $J = 7, 8$ Hz, 1H), 7.33 (d, $J = 8$ Hz, 1H), 3.28 (dd, $J = 6, 8$ Hz, 2H), 1.66-1.54 (m, 1H), 1.54-1.39 (m, 2H), 1.29 (s, 6H), 0.93 (d, $J = 7$ Hz, 6H). MS(ES+) m/e 378 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{18}\text{H}_{23}\text{N}_3\text{SO}_4$: C, 57.28; H, 6.14; N, 11.13. Found: C, 57.15; H, 6.28; N, 10.64.

6-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-7-hydroxy-4-(3-methyl-butyl)-4-aza-spiro[2.4]hept-6-en-5-one (6e). Following Method B, except substituting 1-amino-1-cyclopropanecarboxylic acid for D-phenylalanine, followed by Method C, **6e** was obtained as a white solid. ^1H NMR (400MHz, $\text{D}_6\text{-DMSO}$) δ 12.4 (s, 1H), 7.71 (d, $J = 8$ Hz, 1H), 7.58 (dd, $J = 7, 8$ Hz, 1H), 7.33 (dd, $J = 7, 8$ Hz, 1H), 7.32 (d, $J = 8$ Hz, 1H), 3.07 (t, $J = 8$ Hz, 2H), 1.69-1.48 (m, 1H), 1.43 (s, 2H), 1.40-1.28 (m, 2H), 1.26 (s, 2H), 0.88 (d, $J = 7$ Hz, 6H). MS(ES+) m/e 376 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{18}\text{H}_{21}\text{N}_3\text{SO}_4$: C, 57.58; H, 5.64; N, 11.19. Found: C, 57.38; H, 6.04; N, 11.11.

(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-5-phenyl-1,5-dihydropyrrol-2-one (6f). Following Method A, except substituting L-phenylglycine methyl ester hydrochloride for L-phenylalanine methyl ester hydrochloride, followed by Method C, **6f** was obtained as a grey solid. ^1H NMR (400MHz, $\text{D}_6\text{-DMSO}$) δ 12.6 (s, 1H), 7.71 (dd, $J = 1, 8$ Hz, 1H), 7.58 (ddd, $J = 1, 7, 8$ Hz, 1H), 7.45-7.26 (m, 7H), 4.79 (s, 1H), 3.72 (ddd, $J = 6, 10, 14$ Hz, 1H), 2.75 (ddd, $J = 4, 10, 14$ Hz, 1H), 1.54-1.48 (m, 1H), 1.47-1.29 (m, 2H), 0.87 (d, $J = 6$ Hz, 3H), 0.86 (d, $J = 6$ Hz, 3H). $[\alpha]_{\text{D}} = +127.8^\circ$ ($c = 1.0$, CH_2Cl_2). MS(ES+) m/e 426 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{22}\text{H}_{23}\text{N}_3\text{SO}_4$: C, 62.10; H, 5.45; N, 9.88. Found: C, 62.51; H, 5.24; N, 9.68.

(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-5-(1-methylethyl)-1,5-dihydropyrrol-2-one (6g). Following Method A, except substituting L-valine methyl ester hydrochloride for L-phenylalanine methyl ester hydrochloride, followed by Method C, **6g** was obtained as an off-white solid. ^1H NMR (300MHz, $\text{D}_6\text{-DMSO}$) δ 12.4 (s, 1H), 7.70 (dd, $J = 1, 8$ Hz, 1H), 7.58 (dt, $J = 1, 8$ Hz, 1H), 7.32 (t, $J = 7$ Hz, 1H), 7.32 (d, $J = 8$ Hz, 1H), 3.92 (d, $J = 5$ Hz, 1H), 3.78 (ddd, $J = 8, 8, 14$ Hz, 1H), 2.97 (ddd, $J = 5, 9, 14$ Hz, 1H), 2.23 (dq, $J = 5, 7, 7$ Hz, 1H), 1.52-1.43 (m, 1H), 1.43-1.35 (m, 2H), 1.01 (d, $J = 7$ Hz, 3H), 0.90 (d, $J = 6$ Hz, 3H), 0.89 (d, $J = 7$ Hz, 3H), 0.88 (d, $J = 7$ Hz, 3H). $[\alpha]_{\text{D}} = -47.5^\circ$ ($c = 1.0$, CH_2Cl_2). MS(ES+) m/e 392 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{19}\text{H}_{25}\text{N}_3\text{SO}_4$: C, 58.29; H, 6.44; N, 10.73. Found: C, 57.96; H, 6.47; N, 10.68.

(R)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-5-(1-methylethyl)-1,5-dihydropyrrol-2-one (6h).

Following Method B, except substituting D-valine for D-phenylalanine, followed by Method C, **6h** was obtained as an off-white solid. *See 6g for NMR data.* $[\alpha]_D = +45.0^\circ$ ($c = 1.0$, CH_2Cl_2). MS(ES+) m/e 392 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{19}\text{H}_{25}\text{N}_3\text{SO}_4$: C, 58.29; H, 6.44; N, 10.73. Found: C, 57.76; H, 6.48; N, 10.64.

(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-methyl-1-(3-methylbutyl)-5-(1-methylethyl)-1,5-dihydropyrrol-2-one (6i).

Following Method B, except substituting L- α -methyl-valine for D-phenylalanine, followed by Method C, **6i** was obtained as a white solid. ^1H NMR (400MHz, D_6 -DMSO) δ 12.7 (s, 1H), 7.64 (d, $J = 8$ Hz, 1H), 7.52 (t, $J = 8$ Hz, 1H), 7.25 (t, $J = 8$ Hz, 1H), 7.21 (d, $J = 8$ Hz, 1H), 3.65 (ddd, $J = 8, 8, 14$ Hz, 1H), 3.10 (ddd, $J = 5, 9, 14$ Hz, 1H), 2.25-2.18 (m, 1H), 1.52-1.43 (m, 1H), 1.43-1.35 (m, 2H), 1.23 (s, 3H), 0.99 (d, $J = 7$ Hz, 3H), 0.92 (d, $J = 6$ Hz, 3H), 0.91 (d, $J = 7$ Hz, 3H), 0.77 (d, $J = 7$ Hz, 3H). $[\alpha]_D = -10.5^\circ$ ($c = 1.0$, CH_2Cl_2).

MS(ES+) m/e 406 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{20}\text{H}_{27}\text{N}_3\text{SO}_4$: C, 59.24; H, 6.71; N, 10.36. Found: C, 58.45; H, 6.75; N, 10.14.

(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-5-(2-methylpropyl)-1,5-dihydropyrrol-2-one (6j).

Following Method A, except substituting L-leucine methyl ester hydrochloride for L-phenylalanine methyl ester hydrochloride, followed by Method C, **6j** was obtained as an off-white solid. ^1H NMR (300MHz, D_6 -DMSO) δ 12.8 (s, 1H), 7.60 (d, $J = 8$ Hz, 1H), 7.50 (dt, $J = 1, 8$ Hz, 1H), 7.21 (dt, $J = 1, 7$ Hz, 1H), 7.14 (d, $J = 8$ Hz, 1H), 3.73 (ddd, $J = 8, 8, 14$ Hz, 1H), 3.49 (t, $J = 5$ Hz, 1H), 2.84 (ddd, $J = 6, 7, 14$ Hz, 1H), 1.88-1.77 (m, 1H), 1.58-1.40 (m, 3H), 1.38-1.27 (m, 2H), 0.89 (d, $J = 7$ Hz, 3H), 0.88 (d, $J = 7$ Hz, 3H), 0.87 (d, $J = 7$ Hz, 3H), 0.86 (d, $J = 7$ Hz, 3H). $[\alpha]_D = -43.8^\circ$ ($c = 1.0$, CH_2Cl_2).

MS(ES+) m/e 406 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{20}\text{H}_{27}\text{N}_3\text{SO}_4$: C, 59.24; H, 6.71; N, 10.36. Found: C, 58.37; H, 6.78; N, 10.21.

(S)-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-1,5-dihydropyrrol-2-one (6k).

Following Method A, except substituting L-cyclohexylglycine methyl ester hydrochloride for L-phenylalanine methyl ester hydrochloride, followed by Method C, **6k** was obtained as an off-white solid. ^1H NMR (400MHz, D_6 -DMSO) δ 12.3 (s, 1H), 7.72 (dd, $J = 1, 8$ Hz, 1H), 7.60 (ddd, $J = 1, 7, 8$ Hz, 1H), 7.36 (d, $J = 8$ Hz, 1H), 7.35 (ddd, $J = 1, 7, 8$

Hz, 1H), 3.86 (d, $J = 2$ Hz, 1H), 3.74 (ddd, $J = 8, 8, 14$ Hz, 1H), 3.02 (ddd, $J = 5, 9, 14$ Hz, 1H), 1.94-1.88 (m, 1H), 1.79-1.68 (m, 2H), 1.67-1.61 (m, 1H), 1.60-1.49 (m, 2H), 1.52-1.37 (m, 4H), 1.29-0.99 (m, 4H), 0.91 (d, $J = 7$ Hz, 3H), 0.90 (d, $J = 7$ Hz, 3H). $[\alpha]_D = -16.6^\circ$ ($c = 1.0$, CH_2Cl_2). MS(ES+) m/e 432 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{22}\text{H}_{29}\text{N}_3\text{SO}_4$: C, 61.23; H, 6.77; N, 9.74. Found: C, 61.21; H, 6.77; N, 9.76.

(S)-5-tert-butyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-1,5-dihydropyrrol-2-one (6l). Following Method A, except substituting *L*-tert-leucine methyl ester hydrochloride for *L*-phenylalanine methyl ester hydrochloride, followed by Method C, **6l** was obtained as a white solid. ^1H NMR (400MHz, D_6 -DMSO) δ 12.5 (s, 1H), 7.65 (dd, $J = 1, 8$ Hz, 1H), 7.57 (ddd, $J = 1, 7, 8$ Hz, 1H), 7.31 (dd, $J = 1, 8$ Hz, 1H), 7.31 (ddd, $J = 1, 7, 8$ Hz, 1H), 3.81 (ddd, $J = 6, 10, 14$ Hz, 1H), 3.49 (s, 1H), 3.05 (ddd, $J = 4, 10, 14$ Hz, 1H), 1.55-1.39 (m, 2H), 1.39-1.25 (m, 1H), 1.03 (s, 9H), 0.89 (d, $J = 6$ Hz, 3H), 0.88 (d, $J = 6$ Hz, 3H). $[\alpha]_D = -142.1^\circ$ ($c = 1.0$, CH_2Cl_2). MS(ES+) m/e 406 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{20}\text{H}_{27}\text{N}_3\text{SO}_4$: C, 59.24; H, 6.71; N, 10.36. Found: C, 59.45; H, 6.68; N, 10.24.

(S)-5-tert-butyl-1-(3,3-dimethylbutyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one (10a). Following Method A, except substituting *L*-tert-leucine methyl ester hydrochloride for *L*-phenylalanine methyl ester hydrochloride and 3,3-dimethylbutyraldehyde for isovaleraldehyde, followed by Method D, **10a** was obtained as a white solid. ^1H NMR (400MHz, D_6 -DMSO) δ 12.9 (s, 1H), 7.60 (d, $J = 8$ Hz, 1H), 7.48 (ddd, $J = 1, 7, 8$ Hz, 1H), 7.20 (t, $J = 8$ Hz, 1H), 7.15 (d, $J = 8$ Hz, 1H), 3.78 (ddd, $J = 6, 10, 14$ Hz, 1H), 3.13 (s, 1H), 2.94 (ddd, $J = 4, 10, 14$ Hz, 1H), 1.55-1.42 (m, 1H), 1.31-1.18 (m, 1H), 0.98 (s, 9H), 0.88 (s, 9H). MS(ES+) m/e 420 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{21}\text{H}_{29}\text{N}_3\text{SO}_4$: C, 60.12; H, 6.97; N, 10.02. Found: C, 60.08; H, 6.98; N, 10.04.

(S)-5-tert-butyl-1-(3,3-dimethylbutyl)-4-hydroxy-3-(7-hydroxy-1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-1,5-dihydropyrrol-2-one (10b). Following Method A, except substituting *L*-tert-leucine methyl ester hydrochloride for *L*-phenylalanine methyl ester hydrochloride and 3,3-dimethylbutyraldehyde for isovaleraldehyde, followed by Method D, except substituting (7-OH)**4** (see patent application WO03099801 for experimental procedures for the synthesis of the 7-hydroxy variant of **4**) for **4**, **10a** was obtained as an off-white solid. ^1H NMR (400MHz, D_6 -DMSO) δ 12.6 (s, 1H), 9.77 (s, 1H), 7.09 (d, $J = 8$ Hz, 1H), 6.96 (s, 1H), 6.95 (d, $J = 8$ Hz, 1H), 3.76 (ddd, $J = 5, 5, 14$ Hz, 1H), 3.25 (s, 1H), 2.96 (ddd, $J = 4, 5, 14$ Hz, 1H), 1.51 (ddd, $J = 5, 5, 12$ Hz, 1H), 1.20 (ddd, $J = 4, 5, 12$ Hz,

1H), 0.99 (s, 9H), 0.88 (s, 9H). MS(ES+) m/e 436 [M+H]⁺. Anal calcd for C₂₁H₂₉N₃SO₅: C, 57.91; H, 6.71; N, 9.65. Found: C, 57.12; H, 6.77; N, 9.52.

(S)-4-hydroxy-3-(7-hydroxy-1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-methyl-1-(3-methylbutyl)-5-(1-methylethyl)-1,5-dihydropyrrol-2-one (10c). Following Method B, except substituting L-alpha-methyl-valine for D-phenylalanine, followed by Method D, except substituting **(7-OH)4** for **4**, **10c** was obtained as a grey solid. ¹H NMR (400MHz, D₆-DMSO) δ 12.6 (s, 1H), 9.72 (s, 1H), 7.04 (d, *J* = 8 Hz, 1H), 6.95 (s, 1H), 6.93 (d, *J* = 8 Hz, 1H), 3.20 (ddd, *J* = 6, 10, 13 Hz, 1H), 3.05 (ddd, *J* = 6, 8, 13 Hz, 1H), 1.90 (qq, *J* = 7, 7 Hz,

1H), 1.56 (ddqq, *J* = 7, 7, 13, 13 Hz, 1H), 1.49-1.40 (m, 2H), 1.18 (s, 3H), 0.97 (d, *J* = 7 Hz, 3H), 0.91 (d, *J* = 7 Hz, 3H), 0.90 (d, *J* = 7 Hz, 3H), 0.75 (d, *J* = 7 Hz, 3H). MS(ES+) m/e 422 [M+H]⁺. Anal calcd for C₂₀H₂₇N₃SO₅: C, 56.99; H, 6.46; N, 9.97. Found: C, 57.10; H, 6.32; N, 9.88.

tert-butyl 3-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-2-oxo-1,8-diaza-spiro[4.5]dec-3-ene-8-carboxylate (10d). Following Method B, except substituting 1-tert-butyl 4-amino-piperidine-1,4-dicarboxylate for D-phenylalanine, followed by Method D, **10d** was obtained as a light yellow solid. ¹H NMR (400MHz, D₆-DMSO) δ 12.8 (s, 1H), 7.60 (dd, *J* = 1, 8 Hz, 1H), 7.48 (ddd, *J* = 1, 7, 8 Hz, 1H), 7.20 (ddd, *J* = 1, 7, 8 Hz, 1H), 7.14 (d, *J* = 8 Hz, 1H), 3.87-3.75 (m, 2H), 3.70-3.52 (m, 2H), 3.08 (t, *J* = 8 Hz, 2H), 1.75-1.65 (m, 2H), 1.61-1.50 (m, 1H), 1.49-1.36 (m, 2H), 1.42 (s, 9H),

0.90 (d, *J* = 7 Hz, 6H). MS(ES+) m/e 519 [M+H]⁺. Anal calcd for C₂₅H₃₄N₄SO₆: C, 57.90; H, 6.61; N, 10.80. Found: C, 57.72; H, 6.74; N, 10.22.

ethyl 2-(benzyl)amino-2-(1-methyl-1H-pyrrol-2-yl)acetate. A solution of 1-methylpyrrole (0.178 mL, 2.0 mmol), benzylamine (0.218 mL, 2.0 mmol), and ethyl glyoxylate (50% solution in toluene, 0.408 g, 2.0 mmol) in methanol (1.0 mL) was heated to 100 °C under microwave irradiation for 1h followed by concentration in vacuo. Purification *via* flash column chromatography (0-20% EtOAc/hexanes) gave the title compound as a light yellow oil (0.298 g, 55%). ¹H NMR (400MHz, CDCl₃) δ 7.38-7.22 (m, 5H), 6.57 (t, *J* = 2 Hz, 1H), 6.04 (dd, *J* = 2, 3 Hz, 1H), 6.03 (dd, *J* = 2, 3 Hz, 1H), 4.38 (s, 1H), 4.23 (m, 2H), 3.81 (d, *J* = 13 Hz, 1H), 3.71 (d, *J* = 13 Hz, 1H), 3.56 (s, 3H), 2.17 (br s, 1H), 1.27 (t, *J* = 7 Hz, 3H). MS(ES+) m/e 273 [M+H]⁺.

1'-benzyl-3'-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4'-hydroxy-1-methyl-1',5'-dihydro-1H,5'H-2,5'-bipyrrol-2'-one (10e). Following Method D, except substituting ethyl 2-(benzyl)amino-2-(1-methyl-1H-pyrrol-2-yl)acetate for **3a**, **10e** was obtained as a white solid. ¹H NMR (400MHz, D₆-DMSO) δ 12.7 (s, 1H), 7.63 (d, *J* = 8 Hz, 1H),

7.51 (dt, $J = 2, 8$ Hz, 1H), 7.29 (dt, $J = 1, 8$ Hz, 2H), 7.23 (dt, $J = 1, 8$ Hz, 1H), 7.22 (t, $J = 7$ Hz, 1H), 7.14 (d, $J = 8$ Hz, 1H), 7.08 (dd, $J = 1, 8$ Hz, 2H), 6.64 (t, $J = 2$ Hz, 1H), 5.92 (t, $J = 3$ Hz, 1H), 5.87 (dd, $J = 2, 3$ Hz, 1H), 4.90 (d, $J = 15$ Hz, 1H), 4.49 (s, 1H), 3.75 (d, $J = 15$ Hz, 1H), 3.35 (s, 3H). MS(ES+) m/e 449 $[M+H]^+$. Anal calcd for $C_{23}H_{20}N_4SO_4$: C, 61.59; H, 4.49; N, 12.49. Found: C, 61.72; H, 4.32; N, 12.38.

methyl 2-(benzylamino)-2-(pyridin-2-yl)acetate. A solution of methyl 2-bromo-2-(pyridin-2-yl)acetate (prepared by the method of Edwards, O.E.; Chaput, M.; Clarke, F. H.; Singh, T. *Can. J. Chem.* **1954**, *34*, 785) (0.69 g, 3.0 mmol) in tetrahydrofuran (10.0 mL) was treated with benzylamine (0.49 mL, 4.5 mmol) and potassium carbonate (1.08 g, 7.8 mmol). After stirring 16 h at ambient temperature, the reaction mixture was filtered, concentrated in vacuo, and purified *via* flash column chromatography (0-20% EtOAc/hexanes) to give the title compound as a light yellow oil (0.43 g, 56%). MS(ES+) m/e 257 $[M+H]^+$.

1-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(pyridin-2-yl)-1,5-dihydropyrrol-2-one (10f). Following Method D, except substituting methyl 2-(benzylamino)-2-(pyridin-2-yl)acetate for **3a**, **10f** was obtained as a yellow solid. 1H NMR (400MHz, D_6 -DMSO) δ 12.6 (s, 1H), 8.50 (ddd, $J = 1, 2, 4$ Hz, 1H), 7.74 (dt, $J = 2, 8$ Hz, 1H), 7.63 (dd, $J = 1, 8$ Hz, 1H), 7.51 (dt, $J = 2, 8$ Hz, 1H), 7.29 (ddd, $J = 1, 4, 8$ Hz, 1H), 7.28 (t, $J = 8$ Hz, 2H), 7.26 (dd, $J = 1, 8$ Hz, 1H), 7.23 (t, $J = 8$ Hz, 1H), 7.21 (dd, $J = 1, 8$ Hz, 1H), 7.18 (d, $J = 8$ Hz, 1H), 7.11 (d, $J = 1, 8$ Hz, 2H), 5.03 (d, $J = 15$ Hz, 1H), 4.43 (s, 1H), 3.63 (d, $J = 15$ Hz, 1H). MS(ES+) m/e 447 $[M+H]^+$. Anal calcd for $C_{23}H_{18}N_4SO_4$: C, 61.87; H, 4.06; N, 12.55. Found: C, 61.85; H, 4.06; N, 12.51.

1-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(pyridin-3-yl)-1,5-dihydropyrrol-2-one (10g). Following the procedures of **10f**, except substituting methyl 2-bromo-2-(pyridin-3-yl)acetate for methyl 2-bromo-2-(pyridin-2-yl)acetate, **10g** was obtained as a light brown solid. 1H NMR (400MHz, D_6 -DMSO) δ 12.6 (s, 1H), 8.48 (d, $J = 4$ Hz, 1H), 8.36 (s, 1H), 7.63 (d, $J = 8$ Hz, 1H), 7.51 (t, $J = 7$ Hz, 1H), 7.50 (d, $J = 8$ Hz, 1H), 7.34 (dd, $J = 4, 7$ Hz, 1H), 7.28 (t, $J = 7$ Hz, 2H), 7.23 (d, $J = 8$ Hz, 1H), 7.22 (t, $J = 7$ Hz, 1H), 7.16 (d, $J = 8$ Hz, 1H), 7.11 (d, $J = 7$ Hz, 2H), 5.02 (d, $J = 15$ Hz, 1H), 4.39 (s, 1H), 3.75 (d, $J = 15$ Hz, 1H). MS(ES+) m/e 447 $[M+H]^+$. Anal calcd for $C_{23}H_{18}N_4SO_4$: C, 61.87; H, 4.06; N, 12.55. Found: C, 61.05; H, 4.16; N, 12.38.

1-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(pyridin-4-yl)-1,5-dihydropyrrol-2-one (10h). Following the procedures of **10f**, except substituting methyl 2-bromo-2-(pyridin-4-yl)acetate for methyl 2-bromo-2-(pyridin-2-yl)acetate, **10h** was obtained as a light yellow solid. ^1H NMR (400MHz, D_6 -DMSO) δ 12.5 (s, 1H), 8.50 (d, J = 5 Hz, 2H), 7.63 (d, J = 8 Hz, 1H), 7.51 (t, J = 8 Hz, 1H), 7.28 (t, J = 7 Hz, 2H), 7.24 (t, J = 7 Hz, 1H), 7.23 (d, J = 8 Hz, 1H), 7.17 (d, J = 5 Hz, 2H), 7.16 (t, J = 8 Hz, 1H), 7.11 (d, J = 7 Hz, 2H),

5.07 (d, J = 15 Hz, 1H), 4.34 (s, 1H), 3.73 (d, J = 15 Hz, 1H). MS(ES+) m/e 447 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{23}\text{H}_{18}\text{N}_4\text{SO}_4$: C, 61.87; H, 4.06; N, 12.55. Found: C, 61.62; H, 4.10; N, 12.49.

1-benzyl-3-(7-hydroxy-1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(pyridin-2-yl)-1,5-dihydropyrrol-2-one (10i). Following Method D, except substituting methyl 2-(benzyl)amino-2-(pyridin-2-yl)acetate for **3a** and (7-OH)**4** for **4**, **10i** was obtained as a light yellow solid. ^1H NMR (400MHz, D_6 -DMSO) δ 12.6 (s, 1H), 9.75 (s, 1H), 8.50 (ddd, J = 1, 2, 4 Hz, 1H), 7.73 (dt, J = 1, 7 Hz, 1H), 7.64 (s, 1H), 7.41 (s, 1H), 7.30-7.25 (m, 1H), 7.27 (d, J = 7 Hz, 2H), 7.22 (dd, J = 1, 7 Hz, 1H), 7.20-7.11 (m, 1H), 7.08 (d, J = 7 Hz, 2H), 6.96 (dd, J = 1, 7 Hz, 1H), 5.03 (d,

J = 15 Hz, 1H), 4.43 (s, 1H), 3.63 (d, J = 15 Hz, 1H). MS(ES+) m/e 463 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{23}\text{H}_{18}\text{N}_4\text{SO}_5$: C, 59.73; H, 3.92; N, 12.11. Found: C, 59.93; H, 4.11; N, 11.89.

2-(benzyl)amino-2-(pyridin-2-yl)propanenitrile. A solution of 2-acetylpyridine (0.443 mL, 4.0 mmol) in methanol (5.0 mL) was added to a solution of sodium cyanide (0.490 mg, 10.0 mmol) and benzylamine (1.09 mL, 10.0 mmol) in H_2O (10.0 mL). After stirring 2 h at ambient temperature, the reaction mixture was concentrated in vacuo, diluted with brine, extracted twice with methylene chloride, dried over magnesium sulfate, filtered, and concentrated in vacuo. Purification *via* flash column chromatography (0-20% EtOAc/hexanes) gave the title compound as a clear, colorless

oil (0.313 g, 33%). MS(ES+) m/e 238 $[\text{M}+\text{H}]^+$.

methyl N-benzyl-2-(pyridin-2-yl)alaninate. A solution of 2-(benzyl)amino-2-(pyridin-2-yl)propanenitrile (0.313 g, 1.32 mmol) in methanol (5.0 mL) was treated with HCl (2 N solution in Et_2O , 5 mL, 10 mmol) followed by concentrated aqueous HCl (10.0 mL). After stirring 24 h at reflux, the reaction mixture was concentrated in vacuo, neutralized with solid sodium carbonate, diluted with brine, extracted thrice with methylene chloride, dried over magnesium sulfate, filtered, concentrated in vacuo, and purified *via* flash column chromatography (10-30% EtOAc/hexanes)

to give the title compound as a clear, colorless oil (0.107 g, 30%). MS(ES+) m/e 271 $[M+H]^+$.

1-benzyl-3-(7-hydroxy-1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-methyl-5-(pyridin-2-yl)-1,5-dihydropyrrol-2-one (10j). Following Method D, except substituting methyl *N*-benzyl-2-(pyridin-2-yl)alaninate for **3a** and **(7-OH)4** for **4**, **10j** was obtained as a white solid. ^1H NMR (400MHz, D_6 -DMSO) δ 12.6 (s, 1H), 9.74 (s, 1H), 8.49 (dd, J = 1, 4 Hz, 1H), 7.66 (dt, J = 1, 8 Hz, 1H), 7.62 (s, 1H), 7.41 (s, 1H), 7.29-7.09 (m, 8H), 4.73 (d, J = 16 Hz, 1H), 3.78 (d, J = 16 Hz, 1H), 1.43 (s, 3H). MS(ES+) m/e 477

$[M+H]^+$. Anal calcd for $\text{C}_{24}\text{H}_{20}\text{N}_4\text{SO}_5$: C, 60.49; H, 4.23; N, 11.76. Found: C, 59.93; H, 4.30; N, 11.65.

2-{3-[(S)-(5-tert-butyl-1-(3,3-dimethylbutyl)-4-Hydroxy-2-oxo-2,5-dihydro-1H-pyrrol-3-yl)-1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-7-yloxy}-acetamide (11). A solution of **10b** (0.100 g, 0.230 mmol) in anhydrous acetonitrile (10.0 mL) was treated with 2-bromoacetamide (0.024 g, 0.253 mmol) and solid potassium carbonate (0.095 g, 0.689 mmol) and heated to reflux. After stirring 18 h at reflux, the reaction mixture was filtered, concentrated in vacuo, and purified *via* flash column chromatography (0-10% MeOH/EtOAc) to give

the title compound as a white solid (0.103 g, 91%). ^1H NMR (400MHz, D_6 -DMSO) δ 12.8 (s, 1H), 7.62 (s, 1H), 7.40 (s, 1H), 7.19 (d, J = 9 Hz, 1H), 7.15 (dd, J = 3, 9 Hz, 1H), 7.11 (d, J = 3 Hz, 1H), 3.77 (ddd, J = 5, 5, 13 Hz, 1H), 3.18 (s, 1H), 2.95 (ddd, J = 4, 4, 13 Hz, 1H), 1.51 (ddd, J = 4, 4, 11 Hz, 1H), 1.20 (ddd, J = 5, 5, 12 Hz, 1H), 0.99 (s, 9H), 0.88 (s, 9H). MS(ES+) m/e 493 $[M+H]^+$. Anal calcd for $\text{C}_{23}\text{H}_{32}\text{N}_4\text{SO}_6$: C, 56.08; H, 6.55; N, 11.37. Found: C, 56.10; H, 6.55; N, 11.35.

4-amino-5-tert-butyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-1-(3-methylbutyl)-1,5-dihydropyrrol-2-one (13). A solution of pivalaldehyde (0.217 mL, 2.00 mmol) in methanol (1.0 mL) was added to a solution of sodium cyanide (0.245 mg, 5.00 mmol) and isoamyl amine (0.580 mL, 5.00 mmol) in H_2O (5.0 mL). After stirring 30 min. at ambient temperature, the reaction mixture was concentrated in vacuo, diluted with brine, extracted thrice with methylene chloride, dried over magnesium sulfate, filtered, and concentrated in vacuo. A solution of the resultant residue (~0.053 g crude, 0.290 mmol) and **4** (0.070 g, 0.290 mmol) in anhydrous methylene chloride (3.0 mL) and anhydrous dimethylformamide (0.5 mL)

was treated with dicyclohexylcarbodiimide (1M solution in methylene chloride, 0.290 mL, 0.290 mmol). After stirring 30 min at ambient temperature, triethylamine (0.122 mL, 0.870 mmol) was added. After stirring overnight at ambient temperature, the reaction mixture was filtered, concentrated in

vacuo, and purified *via* flash column chromatography (10-70% EtOAc/hexanes) to give **13** as a white solid (0.059 g, 50%, 2 steps). ^1H NMR (400MHz, D_6 -DMSO) δ 12.2 (s, 1H), 7.81 (dd, $J = 1, 8$ Hz, 1H), 7.70 (ddd, $J = 1, 7, 8$ Hz, 1H), 7.46 (t, $J = 8$ Hz, 1H), 7.33 (d, $J = 8$ Hz, 1H), 3.84 (ddd, $J = 6, 10, 14$ Hz, 1H), 3.45 (ddd, $J = 4, 10, 14$ Hz, 1H), 3.31 (s, 1H), 1.72-1.51 (m, 3H), 1.03 (s, 9H), 0.94 (d, $J = 6$ Hz, 3H), 0.93 (d, $J = 6$ Hz, 3H). MS(ES+) m/e 405 $[\text{M}+\text{H}]^+$. Anal calcd for $\text{C}_{20}\text{H}_{28}\text{N}_4\text{SO}_3$: C, 59.38; H, 6.98; N, 13.85. Found: C, 58.94; H, 7.01; N, 13.75.

Solid-Phase Chemistry (Scheme 2) General Information. The platform shaker employed for array production using Mettler-Toledo Bohdan Mini-Blocks was an Innova 2180 Platform Shaker. Liquid transfers during array production were done using a Tecan Genesis RSP100 liquid handler whenever possible. LCMS method: This method provides a sample cycle of 5 min. inject to inject. The system uses a reverse phase HPLC method. The stationary phase is Aquasil (C18) and the column dimensions are 1mm x 40mm. The gradient is linear starting at 4.5% to 90% CH_3CN (0.02% TFA) in 3.2 min with a 0.4 min. hold and 1.4 min. re-equilibration; detection by MS, UV at 214 nm, and an evaporative light-scattering detector (ELSD). Preparative HPLC purifications (reverse phase) were done on a Gilson using Unipoint software, typically with a YMC 20 x 50 combiprep column ODS-A, 5u. The larger size is a 30 x 75 column of the same material. A 5-minute run (25mL/min, 10% ACN/ H_2O , 0.1% TFA to 90% ACN/ H_2O , 0.1% TFA) with UV detection at 254 nm was used unless otherwise specified.

The following compounds were prepared as part of a solid-phase array according to the general procedure:

- a) To the appropriate Fmoc-protected alpha-amino acid on Wang resin (~0.1 g., 150-300 μm bead diameter) was added 20% piperidine/dimethylformamide solution (1 .5 mL). The slurry was orbitally mixed at ambient temperature for 30 min and filtered. The resin was washed with dimethylformamide, methanol, methylene chloride, and methanol (3 x 2-3 mL each). The resin was dried in vacuo.
- b) To the appropriate resin swelled in trimethyl orthoformate (2.0 mL) was added the appropriate aldehyde (20 equiv, 0.5 mL) as a 4 M solution in trimethyl orthoformate. The reaction was orbitally mixed at ambient temperature for 13 h. The reaction was filtered and the resin was washed with dimethylformamide, methylene chloride, and trimethyl orthoformate (3 x 2-3 mL each). The resin was swollen in trimethyl orthoformate (2.0 mL). Sodium cyanoborohydride (10 equiv, 1 mL) was added as a 0.1 M dispersion in trimethyl orthoformate, followed by acetic acid (20.0 μL). The reaction was vigorously mixed for an additional 3 h at ambient temperature and filtered. The resin was washed with dimethylformamide, methanol, 10% triethylamine/methylene chloride, methanol, methylene chloride, methanol, and diethyl ether (3 x 2-3 mL each). The resin was dried in vacuo.

c) To the appropriate resin swelled in dimethylformamide (1.5 mL) was added 1-hydroxy-7-azabenzotriazole (HOAT) (5 equiv, 0.5 mL) as a 0.01 M solution in dimethylformamide, the compound from example 1(b) (5 equiv, 0.5 mL) as a 0.01 M solution in dimethylformamide, and 1,3-diisopropylcarbodiimide (DIC) (5 equiv, 98.5 μ L). The reaction was orbitally mixed at 775 rpm at ambient temperature for 3.5 d. The reaction was filtered and washed with dimethylformamide, methanol, methylene chloride, and methanol (3 x 2-3 mL each).

d) To the appropriate resin suspended in a solution of 2:1 tetrahydrofuran/2-methyl-2-propanol (1.5 mL) was added potassium *tert*-butoxide (2.5 equiv, 0.5 mL) as a 0.5 M solution in 2-methyl-2-propanol. The reaction was orbitally mixed for 10 min at ambient temperature, and then quenched with acetic acid (50.0 μ L). The reaction was orbitally mixed for another 10 min and filtered into appropriate collection vessels. The resin was washed with methylene chloride, methanol, tetrahydrofuran, methanol, methylene chloride, and methanol (3 x 1-2 mL each). The filtrate and washings were combined and concentrated in vacuo.

e) Examples 105-128 required a subsequent deprotection according to the following general procedures:

Examples 105-110. To the compounds from examples 105-110(d) was added trifluoroacetic acid (TFA) (1 mL). The reaction was orbitally shaken (200 rpm) at ambient temperature for 1 h. The solutions were then evaporated in vacuo to afford the crude deprotected product.

Examples 111-128. To the compounds from examples 111-128(d) was added 1:1 methylene chloride/(95% trifluoroacetic acid/water) (2 mL). The reaction was orbitally shaken (200 rpm) at ambient temperature for 1 h. The solutions were then evaporated in vacuo to afford the crude deprotected product.

All products were subjected to purification via reverse-phase HPLC and met with satisfactory LCMS analysis as follows:

Example #	Name	[M+H] ⁺
1	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(2-methanesulfinyl-ethyl)-1-pyridin-4-ylmethyl-1,5-dihydropyrrol-2-one	461
2	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(2-methanesulfinyl-ethyl)-1-[1,2,3]thiadiazol-4-ylmethyl-1,5-dihydropyrrol-2-one	468
3	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(2-methanesulfinyl-ethyl)-1-pentyl-1,5-dihydropyrrol-2-one	440
4	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(2-methanesulfinyl-ethyl)-1-(3-methylbutyl)-1,5-dihydropyrrol-2-one	440
5	(S)-1-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(2-methanesulfinyl-ethyl)-1,5-dihydropyrrol-2-one	460
6	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(2-methanesulfinyl-ethyl)-1-(2-methylpropyl)-1,5-dihydropyrrol-2-one	426
7	(S)-5- <i>tert</i> -butyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(2-methylpropyl)-1,5-dihydropyrrol-2-one	392
8	(S)-1-benzyl-5- <i>tert</i> -butyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	426
9	(S)-5- <i>tert</i> -butyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-pentyl-1,5-dihydropyrrol-2-one	406

10	(S)-5-cyclopropylmethyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(2-methylpropyl)-1,5-dihydropyrrol-2-one	390
11	(S)-1-benzyl-5-cyclopropylmethyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	424
12	(S)-5-cyclopropylmethyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-1,5-dihydropyrrol-2-one	404
13	(S)-5-cyclopropylmethyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-pentyl-1,5-dihydropyrrol-2-one	404
14	(S)-5-cyclopropylmethyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-[1,2,3]thiadiazol-4-ylmethyl-1,5-dihydropyrrol-2-one	432
15	(S)-5-cyclopropylmethyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-pyridin-4-ylmethyl-1,5-dihydropyrrol-2-one	425
16	(S)-1-benzyl-5-tert-butoxymethyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	456
17	(S)-5-tert-butoxymethyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-[1,2,3]thiadiazol-4-ylmethyl-1,5-dihydropyrrol-2-one	464
18	(S)-5-tert-butoxymethyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-pyridin-4-ylmethyl-1,5-dihydropyrrol-2-one	457
19	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(2-methanesulfanyl-ethyl)-1-(2-methylpropyl)-1,5-dihydropyrrol-2-one	410
20	(S)-1-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(2-methanesulfanyl-ethyl)-1,5-dihydropyrrol-2-one	444
21	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(2-methanesulfanyl-ethyl)-1-pentyl-1,5-dihydropyrrol-2-one	424
22	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(1-methylethyl)-1-(2-methylpropyl)-1,5-dihydropyrrol-2-one	378
23	(S)-1-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(1-methylethyl)-1,5-dihydropyrrol-2-one	412
24	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(1-methylethyl)-1-pentyl-1,5-dihydropyrrol-2-one	392
25	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(1-methylethyl)-1-[1,2,3]thiadiazol-4-ylmethyl-1,5-dihydropyrrol-2-one	420
26	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(1-methylethyl)-1-pyridin-4-ylmethyl-1,5-dihydropyrrol-2-one	413
27	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(1H-imidazol-2-ylmethyl)-5-(1-methylethyl)-1,5-dihydropyrrol-2-one	402
28	(S)-5-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(2-methylpropyl)-1,5-dihydropyrrol-2-one	426
29	(S)-1,5-dibenzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	460
30	(S)-5-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-pentyl-1,5-dihydropyrrol-2-one	440

31	(S)-5-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-[1,2,3]thiadiazol-4-ylmethyl-1,5-dihydropyrrol-2-one	468
32	(S)-5-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-pyridin-4-ylmethyl-1,5-dihydropyrrol-2-one	461
33	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(2-methylpropyl)-5-(1-methylpropyl)-1,5-dihydropyrrol-2-one	392
34	(S)-1-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(1-methylpropyl)-1,5-dihydropyrrol-2-one	426
35	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-5-(1-methylpropyl)-1,5-dihydropyrrol-2-one	406
36	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(1-methylpropyl)-1-pentyl-1,5-dihydropyrrol-2-one	406
37	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(1-methylpropyl)-1-[1,2,3]thiadiazol-4-ylmethyl-1,5-dihydropyrrol-2-one	434
38	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(1-methylpropyl)-1-pyridin-4-ylmethyl-1,5-dihydropyrrol-2-one	427
39	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-bis-(2-methylpropyl)-1,5-dihydropyrrol-2-one	392
40	(S)-1-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(2-methylpropyl)-1,5-dihydropyrrol-2-one	426
41	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(2-methylpropyl)-1-pentyl-1,5-dihydropyrrol-2-one	406
42	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(2-methylpropyl)-1-[1,2,3]thiadiazol-4-ylmethyl-1,5-dihydropyrrol-2-one	434
43	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(2-methylpropyl)-1-(pyridin-4-yl)methyl-1,5-dihydropyrrol-2-one	427
44	(S)-5-benzyl-1-(3,3-dimethylbutyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	454
45	(S)-5-benzyl-1-(2,2-dimethylpropyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	440
46	(S)-1-(3,4-dichlorophenyl)methyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1,5-dihydropyrrol-2-one	532
47	(S)-1-(3,4-difluorophenyl)methyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1,5-dihydropyrrol-2-one	500
48	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1-(thien-2-yl)methyl-1,5-dihydropyrrol-2-one	470
49	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1-(2-trifluoromethylphenyl)methyl-1,5-dihydropyrrol-2-one	532
50	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1-(3-trifluoromethylphenyl)methyl-1,5-dihydropyrrol-2-one	532
51	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1-(4-trifluoromethylphenyl)methyl-1,5-dihydropyrrol-2-one	532

52	(S)-1-(3,3-dimethylbutyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1,5-dihydropyrrol-2-one	458
53	(S)-1-(cyclopropylmethyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1,5-dihydropyrrol-2-one	428
54	(S)-1-(2-cyclopropyl)ethyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1,5-dihydropyrrol-2-one	442
55	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1-(thien-3-yl)methyl-1,5-dihydropyrrol-2-one	470
56	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1-(pyridin-2-yl)methyl-1,5-dihydropyrrol-2-one	465
57	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1-(pyridin-3-yl)methyl-1,5-dihydropyrrol-2-one	465
58	(S)-1-(2-chlorophenyl)methyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1,5-dihydropyrrol-2-one	498
59	(S)-1-(3-chlorophenyl)methyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1,5-dihydropyrrol-2-one	498
60	(S)-1-(4-chlorophenyl)methyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1,5-dihydropyrrol-2-one	498
61	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-1-(2-fluorophenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	482
62	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-1-(3-fluorophenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	482
63	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-1-(4-fluorophenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	482
64	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-1-(2-methoxyphenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	494
65	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-1-(3-methoxyphenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	494
66	(S)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-1-(4-methoxyphenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	494
67	(S)-1-(2-bromophenyl)methyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1,5-dihydropyrrol-2-one	542/544
68	(S)-1-(3-bromophenyl)methyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1,5-dihydropyrrol-2-one	542/544
69	(S)-1-(4-bromophenyl)methyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-5-(4-fluorophenyl)-4-hydroxy-1,5-dihydropyrrol-2-one	542/544

70	(S)-5-tert-butyl-1-(3,3-dimethylbutyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	420
71	(S)-5-tert-butyl-1-(cyclopropylmethyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	390
72	(S)-5-tert-butyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(pyridin-2-yl)methyl-1,5-dihydropyrrol-2-one	427
73	(S)-5-tert-butyl-1-(3-chlorophenyl)methyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	460
74	(S)-5-tert-butyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-1-(4-fluorophenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	444
75	(S)-5-tert-butyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(2-methoxyphenyl)methyl-1,5-dihydropyrrol-2-one	456
76	(S)-5-tert-butyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methoxyphenyl)methyl-1,5-dihydropyrrol-2-one	456
77	(S)-5-cyclohexyl-1-(3,4-dichlorophenyl)methyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	520
78	(S)-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-1-(3,4-difluorophenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	488
79	(S)-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(thien-2-yl)methyl-1,5-dihydropyrrol-2-one	458
80	(S)-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-1-(2-fluorophenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	520
81	(S)-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-1-(3-fluorophenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	520
82	(S)-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-1-(4-fluorophenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	520
83	(S)-5-cyclohexyl-1-(3,3-dimethylbutyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	446
84	(S)-5-cyclohexyl-1-(cyclopropylmethyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	416
85	(S)-5-cyclohexyl-1-(2-cyclopropyl)ethyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	430
86	(S)-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(thien-3-yl)methyl-1,5-dihydropyrrol-2-one	458
87	(S)-1-(2-chlorophenyl)methyl-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	486
88	(S)-1-(3-chlorophenyl)methyl-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	486
89	(S)-1-(4-chlorophenyl)methyl-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	486
90	(S)-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-1-(2-fluorophenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	470
91	(S)-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-1-(3-fluorophenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	470
92	(S)-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-1-(4-fluorophenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	470
93	(S)-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(2-methoxyphenyl)methyl-1,5-dihydropyrrol-2-one	482
94	(S)-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methoxyphenyl)methyl-1,5-dihydropyrrol-2-one	482
95	(S)-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(4-methoxyphenyl)methyl-1,5-dihydropyrrol-2-one	482

96	(S)-1-(2-bromophenyl)methyl-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	530/532
97	(S)-1-(3-bromophenyl)methyl-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	530/532
98	(S)-1-(4-bromophenyl)methyl-5-cyclohexyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	530/532
99	(S)-5-tert-butyl-1-(3,4-dichlorophenyl)methyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	494
100	(S)-5-tert-butyl-1-(3,4-difluorophenyl)methyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	462
101	(S)-5-tert-butyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-trifluoromethylphenyl)methyl-1,5-dihydropyrrol-2-one	494
102	(S)-5-tert-butyl-1-(2-cyclopropyl)ethyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	404
103	(S)-5-tert-butyl-1-(4-chlorophenyl)methyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	460
104	(S)-5-tert-butyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-1-(3-fluorophenyl)methyl-4-hydroxy-1,5-dihydropyrrol-2-one	444
105	3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(4-hydroxy-benzyl)-1-(3-methylbutyl)-1,5-dihydropyrrol-2-one	456
106	3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(4-hydroxy-benzyl)-1-pentyl-1,5-dihydropyrrol-2-one	456
107	3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(4-hydroxy-benzyl)-1-(2-methylpropyl)-1,5-dihydropyrrol-2-one	442
108	1-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(4-hydroxy-benzyl)-1,5-dihydropyrrol-2-one	476
109	3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(4-hydroxy-benzyl)-1-pyridin-4-ylmethyl-1,5-dihydropyrrol-2-one	477
110	3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-5-(4-hydroxy-benzyl)-1-[1,2,3]thiadiazol-4-ylmethyl-1,5-dihydropyrrol-2-one	484
111	[4-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-3-hydroxy-1-(3-methylbutyl)-5-oxo-2,5-dihydro-1H-pyrrol-2-yl]-acetic acid	408
112	[4-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-3-hydroxy-1-pentyl-5-oxo-2,5-dihydro-1H-pyrrol-2-yl]-acetic acid	408
113	[4-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-3-hydroxy-1-(2-methylpropyl)-5-oxo-2,5-dihydro-1H-pyrrol-2-yl]-acetic acid	394
114	[1-benzyl-4-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-3-hydroxy-5-oxo-2,5-dihydro-1H-pyrrol-2-yl]-acetic acid	428
115	[4-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-3-hydroxy-1-pyridin-4-ylmethyl-5-oxo-2,5-dihydro-1H-pyrrol-2-yl]-acetic acid	429
116	[4-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-3-hydroxy-1-[1,2,3]thiadiazol-4-ylmethyl-5-oxo-2,5-dihydro-1H-pyrrol-2-yl]-acetic acid	436
117	[4-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-3-hydroxy-1-(3-methylbutyl)-5-oxo-2,5-dihydro-1H-pyrrol-2-yl]-propionic acid	422
118	[4-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-3-hydroxy-1-pentyl-5-oxo-2,5-dihydro-1H-pyrrol-2-yl]-propionic acid	422
119	[4-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-3-hydroxy-1-(2-methylpropyl)-5-oxo-2,5-dihydro-1H-pyrrol-2-yl]-propionic acid	408
120	[1-benzyl-4-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-3-hydroxy-5-oxo-2,5-dihydro-1H-pyrrol-2-yl]-propionic acid	442

121	[4-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-3-hydroxy-1-pyridin-4-ylmethyl-5-oxo-2,5-dihydro-1H-pyrrol-2-yl]-propionic acid	443
122	[4-(1,1-Dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-3-hydroxy-1-[1,2,3]thiadiazol-4-ylmethyl-5-oxo-2,5-dihydro-1H-pyrrol-2-yl]-propionic acid	450
123	5-(4-amino-butyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(3-methylbutyl)-1,5-dihydropyrrol-2-one	421
124	5-(4-amino-butyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-pentyl-1,5-dihydropyrrol-2-one	421
125	5-(4-amino-butyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-(2-methylpropyl)-1,5-dihydropyrrol-2-one	407
126	5-(4-amino-butyl)-1-benzyl-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1,5-dihydropyrrol-2-one	441
127	5-(4-amino-butyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-pyridin-4-ylmethyl-1,5-dihydropyrrol-2-one	442
128	5-(4-amino-butyl)-3-(1,1-dioxo-1,4-dihydro-1-benzo[1,2,4]thiadiazin-3-yl)-4-hydroxy-1-[1,2,3]thiadiazol-4-ylmethyl-1,5-dihydropyrrol-2-one	449