Supplementary Data to Accompany:

A Mild and Efficient One-Step Synthesis of Quinolines

Brian R. McNaughton[§] and Benjamin L. Miller^{‡*}

Spectral characterization for compound **9**:

¹³CNMR (75 MHz, CDCl₃): 158.8, 154.6, 145.3, 144.3, 131.5, 129.7, 122.4, 120.6, 106.8, 60.9, 56.0, 33.1, 29.1, 23.1, 22.9.

^1H NMR (400 MHz, CDCl_3): δ 8.08 (s), 7.31 (s), 3.99 (s), 3.97 (s), 2.99 (t, $J=6.4$), 2.00 (m), 1.93 (m).

FT-IR (thin film, from CDCl_3): 2931, 1613, 1479, 1410, 1374, 1248, 1152, 1040, 1004, 619 cm^{-1}

HRMS: $\text{M}+\text{H}$ $\text{C}_{15}\text{H}_{17}\text{ClNO}_2$ calculated 278.0948; found 278.0947.

The remaining compounds exhibited characteristics identical to those previously reported in the literature:

Compound	Reference
2	Fakhfakh, M. A.; Franck, X.; Fournet, A.; Hocquemiller, R.; Figaderre, B. <i>Tetrahedron Lett.</i> 2001 , 42, 3847-3850
3	Russel, G. A.; Rafaratnam, R.; Wang, L.; Shi, B. Z.; Kim, B. H.; Yao, C. F. <i>J. Am. Chem. Soc.</i> 1993 , 115, 10596-10604.
5	Jones, W. D.; Joseman, J. <i>J. Org. Chem.</i> 2003 , 68, 3563-3568.
6	Jacquigon, P.; Buu-Hoi, N. P.; Dufuir, M. <i>Bull. Soc. Chim. Fr.</i> 1966 , 9, 2765-2767
7	Olson, R. J. <i>Tetrahedron Lett.</i> 1991 , 32, 5235-5238
8	Curran, D. P.; Kuo, S. C. <i>J. Org. Chem.</i> 1984 , 49, 2063-2065
10	Petrow, V. A. <i>J. Chem. Soc.</i> 1942 , 693-695
11	Katritzky, A. R.; Arend, M. <i>J. Org. Chem.</i> 1998 , 63, 9989-9991
12	Nicolaou, K. C.; Safina, B. S.; Funke, C.; Zak, M. Zecri, F. J. <i>Angew. Chem. Int. Ed. Engl.</i> 2002 , 41, 1937-1940.