A new and efficient method for *o*-Quinone methide intermediate generation: application to the biomimetic synthesis of (±)-Alboatrin.

Raphaël Rodriguez, Robert M. Adlington, John. E. Moses, Andrew cowley and
Jack E. Baldwin*

*Chemistry Research Laboratory, University of Oxford, Mansfield Road, Oxford OX1
3TA, United Kingdom*

Supporting informations

General experimental details:

All solvents and reagents were purified by standard techniques reported in Perrin, D.D.; Amarego, W. L. F., *Purification of Laboratory Chemicals*, 3rd edition, Pergamon Press, Oxford, 1988 or used as supplied from commercial sources as appropriate. Solvents were removed under reduced pressure using a Buchi R110 or R114 Rotavapor fitted with a water condenser. Final traces of solvent were removed from samples using an Edwards E2M5 high vacuum pump with pressures below 2mmHg.

All experiments were carried out under inert atmosphere unless otherwise stated.

¹H NMR spectra were recorded at 200, 400 and 500 MHz using, Bruker DPX 200, DQX400 and Bruker AMX500 instruments. For ¹H spectra recorded in CDCl₃, CD₃OD, C₆D₆, chemical shifts are quoted in parts per million (ppm) and are referenced to the residual solvent peak. The following abbreviations are used: s, singlet; d, doublet; t, triplet; m, multiplet; br, broad. Proton assignments and stereochemistry are supported by ¹H-¹H COSY and NOESY where necessary. Data are reported in the following manner: chemical shift (integration, multiplicity, coupling constant if appropriate). Coupling constants (*J*) are reported in Hertz to the nearest 0.5Hz.

¹³C NMR spectra were recorded at 50.2, 100.6 and 125.8 MHz using Bruker DPX200, Bruker DQX400, and Bruker AMX500 instruments. Carbon spectra assignments are supported by DEPT-135 spectra, ¹³C-¹H (HMQC) correlations where necessary. Chemical shifts are quoted in ppm and are referenced to the appropriate residual solvent peak.

Flash column chromatography was carried out using SorbsilTM C60 (40-63 mm, 230-40 mesh) silica gel. Thin layer chromatography was carried out on glass plates pre-coated with Merck silica gel 60 F₂₅₄ which were visualised by quenching of u.v. fluorescence or by staining with 10% w/v ammonium molybdate in 2M sulphuric acid or 1% w/v potassium permanganate in aqueous alkaline solution followed by heat, as appropriate.

Melting points were recorded using a Cambridge Instruments Gallen™ III Kofler Block melting apparatus or a Buchi 510 capillary apparatus and are uncorrected. Infrared spectra were recorded either as a thin film between NaCl plates on a Perkin-Elmer Paragon 1000 Fourier Transform spectrometer with internal referencing. Absorption maxima are reported in wavenumbers (cm^{-1}).

Low resolution mass spectra were recorded on V. G. Micromass ZAB 1F and V. G. Masslab instruments as appropriate with modes of ionisation being indicated as CI, EI, ES or APCI with only molecular ions. High resolution mass spectrometry was measured on a Waters 2790-Micromass LCT electrospray ionisation mass spectrometer and on a VG autospec chemical ionisation mass spectrometer.

Experimental Procedures:

2-methyleneacetoxy-3-methyl-5-acetoxyphenol (**6a**)

To a stirred solution of 2,4-diacetoxy-6-methyl-benzaldehyde (**5**) (3.8 g, 16.1 mmol) in dry THF (90 ml) was slowly added a 2 M THF solution of borane-DMS complex (8.0 ml, 16.1 mmol) at 0 °C under nitrogen. The reaction was stirred for one hour at room temperature and was then quenched at 0 °C with water (2 ml). The mixture was evaporated to dryness under reduced pressure to give a crude which was purified by flash silica gel chromatography (70 : 30 30-40 P.E. : EtOAc). The title compound was obtained as a white solid which was crystallised in ether to afford (**6a**) as a white crystal (3.2 g, 84 %). R_F 0.3 (70 : 30 30-40 P.E. : EtOAc).

Mp = 95-96 °C; $\nu_{\text{max}}/\text{cm}^{-1}$ (NaCl) 3413, 1779, 1735, 1708, 1599, 1370, 1209, 1133; δ_{H} (400 MHz, CDCl_3) 2.10 (3H, s), 2.27 (3H, s), 2.38 (3H, s), 5.12 (2H, s), 6.53 (1H, s), 6.54 (1H, s) 8.23 (1H, s); δ_{C} (100 MHz, CDCl_3) 19.6, 21.0, 21.2, 59.6, 109.3, 115.8, 118.8, 141.2, 152.1, 157.3, 169.5, 174.3; HRMS (ES-) Calculated for $\text{C}_{12}\text{H}_{13}\text{O}_5$ ($[\text{M} - \text{H}]$): 237.0763, Found: 237.0763.

(±)-Acetylalboatrin (**7**)

(±)-Acetylalboatrin (**7**)

(±)-Acetyl-*epi*-alboatrin

Spiroacetal (**8**)

In an sealed tube was stirred 2-methyleneacetoxy-3-methyl-5-acetoxyphenol (**6a**) (200 mg, 0.84 mmol) with (±)-4,5-dihydro-2,4-dimethylfuran (**3**) (82 mg, 0.84 mmol) in benzene (1 ml) at 80 °C, under argon for 36 hours. After evaporation of benzene under reduced pressure, the colourless oil obtained was purified by flash silica gel

chromatography (98 : 2 30-40 P.E. : EtOAc) to give a viscous colourless oil (57 mg, 25 %) of spiroacetal (**8**) as an inseparable mixture [3:2] of two diastereoisomers [R_F 0.5 (90 : 10 30-40 P.E. : EtOAc)], and a white solid (158 mg, 68 %) of ($\pm$)-**acetylalboatrin (7)** and ($\pm$)-**acetyl-*epi*-alboatrin** as a mixture [12.6:1]. R_F 0.3 (90 : 10 30-40 P.E. : EtOAc).

Data for ($\pm$)-(7): $\nu_{\max}/\text{cm}^{-1}$ (NaCl) 3054, 2987, 2305, 1760, 1603, 1421, 1262, 1215, 896, 752; δ_H (200 MHz, CDCl_3) 1.04 (3H, d, J 6.0 Hz), 1.50 (3H, s), 1.93 (1H, ddd, J 11.0, 4.5, 3.0 Hz), 2.12-2.17 (1H, m), 2.22 (3H, s), 2.26 (3H, s), 2.71-2.73 (2H, m), 3.52 (1H, t, J 8.0 Hz), 4.17 (1H, t, J 8.0 Hz), 6.42 (1H, d, J 2.0 Hz), 6.49 (1H, d, J 2.0 Hz); δ_C (100 MHz, CDCl_3) 15.9, 19.5, 21.2, 22.0, 22.8, 35.4, 47.9, 74.1, 107.1, 108.3, 114.9, 115.3, 138.1, 149.5, 153.6, 169.7; HRMS (ES+) Calculated for $\text{C}_{16}\text{H}_{21}\text{O}_4$ ($[\text{M} + \text{H}]^+$): 277.1440, Found: 277.1431.

($\pm$)-Alboatrin (**1**)

To a stirred solution of ($\pm$)-acetylalboatrin (**7**) and ($\pm$)-acetyl-*epi*-alboatrin [12.6:1] (480 mg, 1.74 mmol) in 9 ml of DCM/MeOH/ H_2O (12/7/1) was quickly added potassium carbonate (721 mg, 5.21 mmol) at room temperature under nitrogen. The mixture was stirred for 6 hours and was then extracted with ethyl acetate (3 x 50 ml). The combined organic layers were washed with a saturated solution of ammonium chloride (3 x 50 ml), dried over magnesium sulphate and concentrated under reduced pressure. The crude obtained was purified by flash silica gel chromatography (98 : 2 30-40 P.E. : EtOAc) to give a white solid. The solid was crystallised in ether to afford ($\pm$)-**Alboatrin (1)** (368 mg, 90 %) as white crystals. R_F 0.1 (90 : 10 30-40 P.E. : EtOAc).

$\text{Mp} = 148\text{-}149\text{ }^\circ\text{C}$; $\nu_{\max}/\text{cm}^{-1}$ (NaCl) 3400, 2958, 1620, 1599, 1494, 1462, 1335, 1204, 1148, 1117, 986, 845; δ_H (200 MHz, CDCl_3) 1.04 (3H, d, J 6.5 Hz), 1.51 (3H, s), 1.93 (1H, ddd, J 11.0, 4.5, 3.5 Hz), 2.01-2.14 (1H, m), 2.19 (3H, s), 2.68 (2H, br s), 3.51 (1H, t, J 8.5 Hz), 4.17 (1H, t, J 8.5 Hz), 4.80 (1H, br s), 6.26 (1H, s), 6.29 (1H, s); δ_C (100 MHz, CDCl_3) 16.0, 19.4, 21.5, 23.2, 35.5, 48.4, 74.0, 101.9, 107.5, 109.6, 110.0, 138.2, 153.7, 155.1; HRMS (ES-) Calculated for $\text{C}_{14}\text{H}_{17}\text{O}_3$ ($[\text{M} - \text{H}]^-$): 233.1178, Found: 233.1180.

Compound (9)

In a sealed tube was stirred 2-methyleneacetoxy-3-methyl-5-acetoxyphenol (**6a**) (203 mg, 0.85 mmol) in 4,5-dihydro-2-methylfuran (1.0 ml, 10.9 mmol) at 100 °C, under argon for 12 hours. After evaporation of excess of reagent under reduced pressure, the yellow oil obtained was purified by flash silica gel chromatography (95 : 5 30-40 P.E. : EtOAc) to give a white solid (175 mg, 78 %). Mp = 115-117 °C; R_F 0.2 (90 : 10 30-40 P.E. : EtOAc); $\nu_{\max}/\text{cm}^{-1}$ (NaCl) 3056, 2982, 1757, 1597, 1482, 1368, 1265, 1213, 1109, 736; δ_H (400 MHz, CDCl_3) 1.51 (3H, s), 1.75-1.85 (1H, m), 2.01-2.09 (1H, m), 2.21 (3H, s), 2.26 (3H, s), 2.43-2.49 (1H, m), 2.78 (1H, s), 2.79 (1H, s), 3.95 (1H, dd, J 16.5, 8.5 Hz), 4.03 (1H, td, J 8.5, 3.0 Hz), 6.43 (1H, br s), 6.50 (1H, br s); δ_C (100 MHz, CDCl_3) 19.4, 21.2, 22.4, 23.5, 29.0, 40.5, 66.7, 106.3, 108.2, 115.0, 115.3, 138.1, 149.5, 153.8, 169.7; HRMS (ES⁺) Calculated for $\text{C}_{15}\text{H}_{19}\text{O}_4$ ($[\text{M} + \text{H}]^+$): 263.1283, Found: 263.1289.

Compound (10)

In a sealed tube was stirred 2-methyleneacetoxy-3-methyl-5-acetoxyphenol (**6a**) (203 mg, 0.85 mmol) in 3,4-dihydro-2H-pyran (1.0 ml, 10.9 mmol) at 100 °C, under argon for 12 hours. After evaporation of excess of reagent under reduced pressure, the yellow oil obtained was purified by flash silica gel chromatography (95 : 5 30-40 P.E. : EtOAc) to give a colourless oil (161 mg, 72 %). R_F 0.3 (90 : 10 30-40 P.E. : EtOAc); $\nu_{\max}/\text{cm}^{-1}$ (NaCl) 3055, 2934, 1759, 1598, 1422, 1265, 1217, 1093, 897, 739; δ_H (400 MHz, CDCl_3) 1.62-1.70 (5H, m), 2.18 (3H, s), 2.26 (3H, s), 2.50 (1H, dd, J 16.5, 4.5 Hz), 2.72 (1H, dd, J 16.5, 6.5 Hz), 3.69-3.73 (1H, m), 3.98-4.04 (1H, m), 5.28 (1H, d, J 2.5 Hz), 6.49 (1H, s), 6.50 (1H, s); δ_C (100 MHz, CDCl_3) 19.4, 21.2, 23.6, 24.4, 26.5, 31.6, 62.7, 96.3, 107.8, 115.6, 116.4, 138.3, 149.4, 153.3, 169.7; HRMS (CI⁺) Calculated for $\text{C}_{15}\text{H}_{19}\text{O}_4$ ($[\text{M} + \text{H}]^+$): 263.1283, Found: 263.1288.

Compound (11)

In a sealed tube was stirred 2-methyleneacetoxy-3-methyl-5-acetoxyphenol (**6a**) (203 mg, 0.85 mmol) in (1R)-(+)- α -pinene (1.0 ml, 6.3 mmol) at 140 °C, under argon for 12 hours. After evaporation of excess of pinene under reduced pressure, the yellow oil obtained was purified by flash silica gel chromatography (99 : 1 30-40 P.E. : EtOAc) to give a viscous colourless oil (81 mg, 30 %). $[\alpha]_D^{25} = -2.9$ ($c = 1$, CHCl₃); R_F 0.6 (98 : 2 30-40 P.E : EtOAc); $\nu_{\max}/\text{cm}^{-1}$ (NaCl) 2922, 1758, 1559, 1483, 1369, 1216, 1127, 1015, 739; δ_H (500 MHz, CDCl₃) 1.04 (3H, s), 1.16 (1H, d, J 10.5 Hz), 1.28 (3H, s), 1.31 (3H, s), 1.32-1.35 (1H, m), 1.81-1.84 (1H, m), 2.04-2.15 (2H, m), 2.17 (1H, t, J 5.5 Hz), 2.24 (3H, s), 2.25 (3H, s), 2.43 (1H, dd, J 15.0, 5.0 Hz), 2.47-2.53 (1H, m), 2.73 (1H, dd, J 15.0, 6.0 Hz), 6.40 (1H, d, J 2.5 Hz), 6.49 (1H, d, J 2.5 Hz); δ_C (125 MHz, CDCl₃) 19.2, 21.2, 23.2, 26.8, 27.5, 28.0, 29.3, 34.2, 34.5, 40.2, 40.4, 54.9, 84.0, 108.3, 114.9, 121.3, 136.8, 149.3, 157.3, 169.7; HRMS (ES+) Calculated for C₂₀H₂₇O₃ ($[M + H]^+$): 315.1960, Found: 315.1963.

Compound (12)

In a sealed tube was stirred 2-methyleneacetoxy-3-methyl-5-acetoxyphenol (**6a**) (203 mg, 0.85 mmol) in 1-methylcyclohexene (1.0 ml, 8.5 mmol) at 140 °C, under argon for 12 hours. After evaporation of excess of methylcyclohexene under reduced pressure, the yellow oil was purified by flash silica gel chromatography (99 : 1 30-40 P.E. : EtOAc) to give a white solid (141 mg, 60 %). Mp = 100-101 °C; R_F 0.6 (98 : 2 30-40 P.E : EtOAc); $\nu_{\max}/\text{cm}^{-1}$ (NaCl) 2933, 1758, 1596, 1370, 1265, 1218, 1125, 737; δ_H (400 MHz, CDCl₃) 1.17 (3H, s), 1.24-1.31 (2H, m), 1.36-1.44 (2H, m), 1.48-1.52 (1H, m), 1.57-1.75 (3H, m), 1.93 (1H, bd, J 13.5 Hz), 2.18 (3H, s), 2.24 (1H, d, J 17.0 Hz), 2.25 (3H, s), 2.81 (1H, dd, J 17.0, 6.5 Hz), 6.40 (1H, d, J 2.0 Hz), 6.44 (1H, d, J 2.0 Hz); δ_C (100 MHz, CDCl₃) 19.4, 21.2, 21.7, 25.4, 25.6, 27.1, 28.8, 36.8, 38.3, 74.7, 108.0, 114.3, 116.4, 138.4, 149.1, 153.8, 169.8; HRMS (CI+) Calculated for C₁₇H₂₃O₃ ($[M + H]^+$): 275.1647, Found: 275.1644.