A New and Efficient Synthesis of 4-Functionalized Benzo[*b*]furans from 2,3-Dihalophenols

Roberto Sanz,^{,†} M^a Pilar Castroviejo,[†] Yolanda Fernández,[†] and Francisco J. Fañanás[‡]*

Departamento de Química, Área de Química Orgánica, Facultad de Ciencias, Universidad de Burgos,

Pza. Missael Bañuelos s/n, 09001-Burgos, Spain, and Instituto Universitario de Química

Organometálica “Enrique Moles”, Universidad de Oviedo, C/ Julián Clavería, 8, 33006-Oviedo, Spain

rsd@ubu.es

Supporting Information

Index

General Considerations	S-2
Typical Procedures and Spectroscopic Data	S-2
Spectra	S-19
References	S-59

General Considerations. All reactions were carried out under a nitrogen atmosphere in oven-dried glassware with magnetic stirring. Temperatures are reported as bath temperatures. THF were continuously refluxed and freshly distilled from sodium under nitrogen. TMEDA were refluxed over potassium under nitrogen using benzophenone as indicator and distilled at reduced pressure. Solvents used in extraction and purification were distilled prior to use. TLC was performed on alumina-backed plates coated with silica gel 60 with F₂₅₄ indicator (Merck) and compounds were visualized by UV light (254 nm) or iodine. Flash column chromatography was carried out on silica gel 60, 230-400 mesh (Merck). Melting points were obtained with open capillary tubes and are uncorrected. ¹H NMR spectra were recorded at 400 MHz or at 300 MHz. Chemical shifts are reported in ppm from tetramethylsilane with the residual solvent resonance as the internal standard (CHCl₃: δ 7.26). Data are reported as follows: chemical shift, multiplicity (s: singlet, d: doublet, dd: double doublet, dt: double triplet, t: triplet, q: quartet, br: broad, m: multiplet), coupling constants (*J* in Hz) and integration. ¹³C NMR spectra were recorded at 100.6 MHz or at 75.4 MHz using broadband proton decoupling. Chemical shifts are reported in ppm with the solvent resonance as internal standard (CDCl₃: δ 76.95). Low-resolution electron impact mass spectra (EI-LRMS) were obtained at 70 eV on mass spectrometers and only the molecular ions and/or base peaks in MS are given. High-resolution mass spectrometry (HRMS) was carried out on a mass spectrometer. Infrared spectra were recorded with a FT-IR spectrophotometer. Elemental analyses were performed on a microanalyzer. All commercially available reagents were used without purification unless otherwise indicated and were purchased from standard chemical suppliers.

Typical Procedure for the Synthesis of *O*-3-Halophenyl *N,N*-Diethylcarbamates **1.** Carbamates **1** were prepared by either these methods. Method A:¹ To a well stirred suspension of NaH (0.96 g, 40 mmol) in THF (20 mL), a solution of 3-halophenol (20 mmol) in THF (5 mL) was dropwise added at room temperature. After stirring the reaction mixture for 2 h, *N,N*-diethylcarbamoyl chloride (5.42 g, 40 mmol) in THF (7 mL) was added. Stirring was continued for another 8 h. Usual aqueous work up afforded the crude carbamate which was purified by column chromatography (eluent: hexane/EtOAc) on silica gel to afford compounds **1**. Method B: A solution of 3-halophenol (20 mmol) in dry THF (20

mL) was treated with *n*-BuLi (8 mL of a 2.5M solution in hexane, 20 mmol) at -78°C . After 10 min at low temperature the mixture was allowed to reach room temperature and stirred for 30 min. Then, a solution of *N,N*-diethylcarbamoyl chloride (2.71 g, 20 mmol) in THF (20 mL) was added and the resulting mixture was heated at reflux for 36 h. After cooling to room temperature, H_2O was added and the resulting carbamate was extracted with EtOAc (3 x 20 mL). The combined organic layers were dried (anhydrous Na_2SO_4), and evaporated under reduced pressure. The crude was purified by column chromatography (eluent: hexane/EtOAc) on silica gel to afford compounds **1**.

3-Fluorophenyl *N,N*-diethylcarbamate (1a**):**² 3-Fluorophenol (2.24 g, 20 mmol) was used according to Method A or Method B. Purification by column chromatography (eluent: hexane/EtOAc, 10/1) on silica gel gave 3.54 g (84%), according to Method A, or 3.76 g (89%), according to Method B, of **1a** as a colorless oil. R_f 0.22 (hexane/EtOAc, 10/1); ^1H RMN (400 MHz, CDCl_3) δ 7.34-7.26 (m, 1H), 6.95-6.87 (m, 3H), 3.43 (q, $J = 7.2$ Hz, 2H), 3.38 (q, $J = 7.2$ Hz, 2H), 1.25 (t, $J = 7.2$ Hz, 3H), 1.20 (t, $J = 7.2$ Hz, 3H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 162.8 (d, $J = 247.5$ Hz), 153.6, 152.5 (d, $J = 10.7$ Hz), 129.9 (d, $J = 9.2$ Hz), 117.5 (d, $J = 3.8$ Hz), 112.0 (d, $J = 21.4$ Hz), 109.8 (d, $J = 23.7$ Hz), 42.3, 41.9, 14.2, 13.3; EI-LRMS m/z 211 (M^+ , 30), 72 (100); IR (neat) 3073, 2976, 2933, 1720, 1417, 1258, 1155, 970 cm^{-1} . HRMS calcd. for $\text{C}_{11}\text{H}_{14}\text{FNO}_2$: 211.1009. Found: 211.1002.

3-Chlorophenyl *N,N*-diethylcarbamate (1b**):**³ 3-Chlorophenol (2.57 g, 20 mmol) was used according to Method A or Method B. Purification by column chromatography (eluent: hexane/EtOAc, 10/1) on silica gel gave 3.87 g (85%), according to Method A, or 3.96 g (87%), according to Method B, of **1b** as a colorless oil. R_f 0.25 (hexane/EtOAc, 10/1); ^1H RMN (400 MHz, CDCl_3) δ 7.31-7.25 (m, 1H), 7.21-7.16 (m, 2H), 7.08-7.03 (m, 1H), 3.49-3.35 (m, 4H), 1.30-1.17 (m, 6H); ^{13}C -RMN (100.6 MHz, CDCl_3) δ 153.6, 152.0, 134.3, 129.9, 125.2, 122.4, 120.2, 42.3, 41.9, 14.2, 13.3; EI-LRMS m/z 229 ($\text{M}^+ + 2$, 3), 227 (M^+ , 8), 100 (100); IR (neat) 3075, 2977, 2935, 1720, 1412, 1258, 1150, 965 cm^{-1} . HRMS calcd. for $\text{C}_{11}\text{H}_{14}\text{ClNO}_2$: 227.0713. Found: 227.0721.

3-Bromophenyl *N,N*-diethylcarbamate (1c**):** 3-Bromophenol (3.46 g, 20 mmol) was used according to Method A or Method B. Purification by column chromatography (eluent: hexane/EtOAc, 12/1) on

silica gel gave 4.95 g (91%), according to Method A, or 4.84 g (89%), according to Method B, of **1c** as a colorless oil. R_f 0.28 (hexane/EtOAc, 10/1); ^1H RMN (400 MHz, CDCl_3) δ 7.35-7.31 (m, 2H), 7.24-7.20 (m, 1H), 7.12-7.08 (m, 1H), 3.45-3.36 (m, 4H), 1.29-1.18 (m, 6H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 153.7, 152.1, 130.3, 128.3, 125.3, 122.2, 120.7, 42.4, 42.0, 14.3, 13.4; EI-LRMS m/z 273 ($\text{M}^+ + 2$, 4), 271 (M^+ , 4), 100 (100); IR (neat) 3072, 2974, 2934, 1720, 1413, 1261, 1149 cm^{-1} . HRMS calcd. for $\text{C}_{11}\text{H}_{14}\text{BrNO}_2$: 271.0288. Found: 271.0281.

3-Iodophenyl *N,N*-diethylcarbamate (1d): 3-Iodophenol (4.40 g, 20 mmol) was used according to Method A. Purification by column chromatography (eluent: hexane/EtOAc, 15/1) on silica gel gave **1d** (6.06 g, 95%) as a colorless oil. R_f 0.17 (hexane/EtOAc, 15/1); ^1H RMN (400 MHz, CDCl_3) δ 7.53 (t, J = 1.5 Hz, 1H), 7.52-7.49 (m, 1H), 7.14-7.04 (m, 2H), 3.45-3.33 (m, 4H), 1.29-1.15 (m, 6H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 153.7, 151.9, 134.2, 131.0, 130.5, 121.5, 93.4, 42.4, 42.0, 14.3, 13.4; EI-LRMS m/z 319 (M^+ , 12), 100 (100); IR (neat) 3064, 2974, 2933, 1720, 1411, 1261, 1152, 961 cm^{-1} . HRMS calcd. for $\text{C}_{11}\text{H}_{14}\text{INO}_2$: 319.0069. Found: 319.0080.

Typical Procedure for the Synthesis of 3-Chloro- and 3-Fluoro-2-halophenyl *N,N*-Diethylcarbamates 3a-e. To a well stirred solution of *s*-BuLi (9.23 mL of a 1.3M solution in cyclohexane/hexane, 12 mmol), TMEDA (1.81 mL, 12 mmol) in dry THF (40 mL) kept at -78°C under N_2 atmosphere, the corresponding carbamate **1a,b** (10 mmol) was added and the mixture was stirred at -78°C for 2 h. Then, the corresponding electrophile (12 mmol) was added and stirring continued at low temperature for further 30 min. The reaction mixture was then allowed to warm to room temperature and after quenching with H_2O , or aqueous solution of $\text{Na}_2\text{S}_2\text{O}_3$ in the case of using iodine as electrophile, it was extracted with EtOAc (3 x 25 mL). The combined organic layers were dried (Na_2SO_4), and evaporated under reduced pressure. The crude was purified by column chromatography (eluent: hexane/EtOAc) on silica gel to afford compounds **3a-e**.

3-Fluoro-2-iodophenyl *N,N*-diethylcarbamate (3a): The treatment of **1a** (2.11 g, 10 mmol) according to the typical procedure, using iodine (3.05 g, 12 mmol) as electrophile and purification by column chromatography (eluent: hexane/EtOAc, 10/1) on silica gel, gave **3a** (2.96 g, 88%) as a

colorless oil. R_f 0.17 (hexane/EtOAc, 10/1); ^1H RMN (400 MHz, CDCl_3) δ 7.35-7.28 (m, 1H), 7.03-7.00 (m, 1H), 6.89-6.95 (m, 1H), 3.53 (q, $J = 7.2$ Hz, 2H), 3.40 (q, $J = 7.2$ Hz, 2H), 1.33 (t, $J = 7.2$ Hz, 3H), 1.23 (t, $J = 7.2$ Hz, 3H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 162.5 (d, $J = 245.3$ Hz), 153.3, 153.2, 129.9 (d, $J = 9.2$ Hz), 119.0 (d, $J = 3.8$ Hz), 112.4 (d, $J = 24.4$ Hz), 80.1 (d, $J = 26$ Hz), 42.4, 42.2, 14.4, 13.3; EI-LRMS m/z 337 (M^+ , 23), 100 (100); IR (neat) 3076, 2975, 2935, 1727, 1460, 1414, 1260, 1225, 1153, 1045, 994 cm^{-1} . HRMS calcd. for $\text{C}_{11}\text{H}_{13}\text{FINO}_2$: 336.9975. Found: 336.9962.

2-Bromo-3-fluorophenyl *N,N*-diethylcarbamate (3b): The treatment of **1a** (2.11 g, 10 mmol) according to the typical procedure, using 1,2-dibromo-1,1,2,2-tetrachloroethane (3.91 g, 12 mmol) as electrophile and purification by column chromatography (eluent: hexane/EtOAc, 10/1) on silica gel, gave **3b** (2.44 g, 84%) as a colorless oil. R_f 0.22 (hexane/EtOAc, 10/1); ^1H RMN (300 MHz, CDCl_3) δ 7.32-7.22 (m, 1H), 7.06-6.94 (m, 2H), 3.49 (q, $J = 7.2$ Hz, 2H), 3.39 (q, $J = 7.2$ Hz, 2H), 1.30 (t, $J = 7.2$ Hz, 3H), 1.21 (t, $J = 7.2$ Hz, 3H); ^{13}C RMN (75.4 MHz, CDCl_3) δ 159.9 (d, $J = 247.8$ Hz), 152.8, 150.4 (d, $J = 2.9$ Hz), 128.3 (d, $J = 9.3$ Hz), 119.6 (d, $J = 3.5$ Hz), 113.2 (d, $J = 22.5$ Hz), 105.0 (d, $J = 22.2$ Hz), 42.6, 42.2, 14.3, 13.3; EI-LRMS m/z 289 (M^+ , 0.3), 100 (100); IR (neat) 3083, 2977, 2935, 1725, 1468, 1419, 1264, 1233, 1151, 1049, 998 cm^{-1} . HRMS calcd. for $\text{C}_{11}\text{H}_{13}\text{BrFNO}_2$: 289.0114. Found: 289.0123.

2-Chloro-3-fluorophenyl *N,N*-diethylcarbamate (3c): The treatment of **1a** (2.11 g, 10 mmol) according to the typical procedure, using hexachloroethane (2.84 g, 12 mmol) as electrophile and purification by column chromatography (eluent: hexane/EtOAc, 10/1) on silica gel, gave **3c** (2.04 g, 83%) as a colorless oil. R_f 0.19 (hexane/EtOAc, 10/1); ^1H RMN (300 MHz, CDCl_3) δ 7.28-7.18 (m, 1H), 7.09-6.99 (m, 2H), 3.49 (q, $J = 7.2$ Hz, 2H), 3.40 (q, $J = 7.2$ Hz, 2H), 1.31 (t, $J = 7.2$ Hz, 3H), 1.22 (t, $J = 7.2$ Hz, 3H); ^{13}C RMN (75.4 MHz, CDCl_3) δ 158.9 (d, $J = 249.5$ Hz), 152.8, 149.2 (d, $J = 2.3$ Hz), 127.2 (d, $J = 9.2$ Hz), 119.6 (d, $J = 3.5$ Hz), 115.6 (d, $J = 19.2$ Hz), 113.3 (d, $J = 21.2$ Hz), 42.6, 42.2, 14.2, 13.4; EI-LRMS m/z 210 ($\text{M}^+ - \text{Cl}$, 2), 100 (100); IR (neat) 3086, 2977, 2936, 1727, 1467, 1418, 1263, 1236, 1154, 1049, 999 cm^{-1} . HRMS calcd. for $\text{C}_{11}\text{H}_{13}\text{ClFNO}_2$: 245.0619. Found: 245.0625.

3-Chloro-2-iodophenyl *N,N*-diethylcarbamate (3d): The treatment of **1b** (2.28 g, 10 mmol) according to the typical procedure, using iodine (3.05 g, 12 mmol) as electrophile and purification by column chromatography (eluent: hexane/EtOAc, 10/1) on silica gel, gave **3d** (3.04 g, 86%) as a white solid. Mp 69-71 °C; ¹H RMN (400 MHz, CDCl₃) δ 7.33-7.25 (m, 2H), 7.06 (dd, *J* = 7.3, 2.2 Hz, 1H), 3.54 (q, *J* = 7.2 Hz, 2H), 3.41 (q, *J* = 7.2 Hz, 2H), 1.33 (t, *J* = 7.2 Hz, 3H), 1.23 (t, *J* = 7.2 Hz, 3H); ¹³C RMN (100.6 MHz, CDCl₃) δ 153.4, 152.7, 139.5, 129.5, 126.1, 121.3, 96.8, 42.4, 42.1, 14.4, 13.3; EI-LRMS *m/z* 353 (M⁺, 0.1), 100 (100); IR (KBr) 3075, 2975, 2934, 1707, 1415, 1237, 1153, 1041, 964 cm⁻¹. Anal. Calcd. for C₁₁H₁₃ClINO₂: C, 37.37; H, 3.71; N, 3.96. Found: C, 37.49; H, 3.85; N, 3.89.

2-Bromo-3-chlorophenyl *N,N*-diethylcarbamate (3e): The treatment of **1b** (2.28 g, 10 mmol) according to the typical procedure, using 1,2-dibromoethane (3.91 g, 12 mmol) as electrophile and purification by column chromatography (eluent: hexane/EtOAc, 10/1) on silica gel, gave **3e** (1.96 g, 64%) as a white solid. Mp 56-58 °C; ¹H RMN (300 MHz, CDCl₃) δ 7.31 (dd, *J* = 7.9, 1.6 Hz, 1H), 7.24 (t, *J* = 7.9 Hz, 1H), 7.12 (dd, *J* = 7.9, 1.6 Hz, 1H), 3.49 (q, *J* = 7.2 Hz, 2H), 3.38 (q, *J* = 7.2 Hz, 2H), 1.30 (t, *J* = 7.2 Hz, 3H), 1.21 (t, *J* = 7.2 Hz, 3H); ¹³C RMN (75.4 MHz, CDCl₃) δ 152.8, 150.4, 135.6, 128.2, 127.1, 122.4, 118.1, 42.5, 42.2, 14.3, 13.4; EI-LRMS *m/z* 226 (M⁺-Br, 11), 100 (100); IR (KBr) 3082, 2984, 2937, 1722, 1419, 1245, 1158, 963 cm⁻¹. Anal. Calcd. for C₁₁H₁₃BrClINO₂: C, 43.09; H, 4.27; N, 4.57. Found: C, 43.01; H, 4.29; N, 4.59.

Typical Procedure for the Synthesis of 3-Bromo- and 3-Iodo-2-halophenyl *N,N*-diethylcarbamates 3f-k. *n*-BuLi (4.4 mL of a 2.5M sol. in hexane, 11 mmol) was added to a solution of *i*-Pr₂NH (1.54 mL, 11 mmol) in THF (30 mL) at 0 °C. After 30 min at 0 °C the LDA solution was cooled at -78 °C and the corresponding carbamate **1c,d** (10 mmol) was added. The resulting solution was stirred for 30 min at -78 °C and then, the corresponding electrophile (12 mmol) was added. After 30 min at low temperature the reaction mixture was allowed to warm to room temperature, H₂O was added and THF evaporated under reduced pressure. The aqueous phase was extracted with EtOAc (3 x 25 mL) and the combined organic layers were washed with 1M HCl, dried over anhydrous Na₂SO₄, and

evaporated under reduced pressure. The crude was purified by column chromatography (eluent: hexane/EtOAc) on silica gel to afford compounds **3f-k**.

3-Bromo-2-iodophenyl *N,N*-diethylcarbamate (3f): The treatment of **1c** (2.72 g, 10 mmol) according to the typical procedure, using iodine (3.05 g, 12 mmol) as electrophile and purification by column chromatography (eluent: hexane/EtOAc, 10/1) on silica gel, gave **3f** (3.34 g, 84%) as a white solid. Mp 69-71 °C; ¹H RMN (400 MHz, CDCl₃) δ 7.47 (dd, *J* = 8.1, 1.5 Hz, 1H), 7.22 (t, *J* = 8.1 Hz, 1H), 7.08 (dd, *J* = 8.1, 1.5 Hz, 1H), 3.52 (q, *J* = 7.1 Hz, 2H), 3.38 (q, *J* = 7.1 Hz, 2H), 1.31 (t, *J* = 7.1 Hz, 3H), 1.22 (t, *J* = 7.1 Hz, 3H); ¹³C RMN (100.6 MHz, CDCl₃) δ 153.4, 152.8, 130.6, 130.0, 129.7, 121.8, 99.9, 42.5, 42.2, 14.5, 13.4; EI-LRMS *m/z* 399 (M⁺, 0.1), 100 (100); IR (KBr) 3074, 2976, 2934, 1777, 1413, 1237, 1151, 1049, 960 cm⁻¹. Anal. Calcd. for C₁₁H₁₃BrINO₂: C, 33.19; H, 3.29; N, 3.52. Found: C, 33.25; H, 3.25; N, 3.48.

2,3-Dibromophenyl *N,N*-diethylcarbamate (3g): The treatment of **1c** (2.72 g, 10 mmol) according to the typical procedure, using 1,2-dibromoethane (3.91 g, 12 mmol) as electrophile and purification by column chromatography (eluent: hexane/EtOAc, 10/1) on silica gel, gave **3g** (2.49 g, 71%) as a white solid. Mp 50-52 °C; ¹H RMN (300 MHz, CDCl₃) δ 7.49-7.45 (m, 1H), 7.22-7.12 (m, 2H), 3.49 (q, *J* = 7.2 Hz, 2H), 3.39 (q, *J* = 7.2 Hz, 2H), 1.30 (t, *J* = 7.2 Hz, 3H), 1.21 (t, *J* = 7.2 Hz, 3H); ¹³C RMN (75.4 MHz, CDCl₃) δ 152.8, 150.3, 130.4, 128.7, 125.8, 123.0, 120.3, 42.5, 42.2, 14.3, 13.4; EI-LRMS *m/z* 353 (M⁺+4, 0.1), 351 (M⁺+2, 0.2), 349 (M⁺, 0.1), 100 (100); IR (KBr) 3076, 2989, 2978, 2935, 1715, 1417, 1242, 1149 cm⁻¹. HRMS calcd. for C₁₁H₁₃Br₂NO₂: 348.9313. Found: 348.9319.

3-Bromo-2-chlorophenyl *N,N*-diethylcarbamate (3h): The treatment of **1c** (2.72 g, 10 mmol) according to the typical procedure, using hexachloroethane (2.84 g, 12 mmol) as electrophile and purification by column chromatography (eluent: hexane/EtOAc, 10/1) on silica gel, gave **3h** (2.82 g, 92%) as a white solid. Mp 39-41 °C; ¹H RMN (400 MHz, CDCl₃) δ 7.48 (dd, *J* = 8.0, 1.7 Hz, 1H), 7.20 (dd, *J* = 8.0, 1.7 Hz, 1H), 7.14 (t, *J* = 8.0 Hz, 1H), 3.49 (q, *J* = 7.1 Hz, 2H), 3.40 (q, *J* = 7.1 Hz, 2H), 1.30 (t, *J* = 7.1 Hz, 3H), 1.22 (t, *J* = 7.1 Hz, 3H); ¹³C RMN (100.6 MHz, CDCl₃) δ 152.9, 148.9, 130.5, 128.6, 127.8, 123.4, 123.2, 42.6, 42.2, 14.3, 13.4; EI-LRMS *m/z* 270 (M⁺-Cl, 4), 100 (100); IR (KBr)

3078, 2978, 2934, 1716, 1415, 1243, 1156, 960 cm^{-1} . Anal. Calcd. for $\text{C}_{11}\text{H}_{13}\text{BrClINO}_2$: C, 43.09; H, 4.27; N, 4.57. Found: C, 43.21; H, 4.24; N, 4.45.

2,3-Diiodophenyl *N,N*-diethylcarbamate (3i): The treatment of **1d** (3.19 g, 10 mmol) according to the typical procedure, using iodine (3.05 g, 12 mmol) as electrophile and purification by column chromatography (eluent: hexane/EtOAc, 10/1) on silica gel, gave **3i** (3.29 g, 74%) as a white solid. Mp 75-77 $^{\circ}\text{C}$; ^1H RMN (400 MHz, CDCl_3) δ 7.71 (dd, $J = 7.1, 2.2$ Hz, 1H), 7.13-7.05 (m, 2H), 3.51 (q, $J = 7.2$ Hz, 2H), 3.39 (q, $J = 7.2$ Hz, 2H), 1.30 (t, $J = 7.2$ Hz, 3H), 1.22 (t, $J = 7.2$ Hz, 3H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 152.8, 152.5, 136.3, 130.4, 122.4, 108.7, 106.1, 42.4, 42.2, 14.5, 13.4; EI-LRMS m/z 445 (M^+ , 12), 100 (100); IR (KBr): 3062, 2973, 2932, 1718, 1415, 1233, 1215, 1152, 958, 763 cm^{-1} . Anal. calcd. for $\text{C}_{11}\text{H}_{13}\text{I}_2\text{NO}_2$: C, 29.69; H, 2.94; N, 3.15. Found: C, 29.64; H, 2.96; N, 3.17.

2-Bromo-3-iodophenyl *N,N*-diethylcarbamate (3j): The treatment of **1d** (3.19 g, 10 mmol) according to the typical procedure, using 1,2-dibromoethane (3.91 g, 12 mmol) as electrophile and purification by column chromatography (eluent: hexane/EtOAc, 10/1) on silica gel, gave **3j** (2.39 g, 60%) as a white solid. Mp 52-54 $^{\circ}\text{C}$; ^1H RMN (300 MHz, CDCl_3) δ 7.70 (dd, $J = 8.0, 1.6$ Hz, 1H), 7.18 (dd, $J = 8.0, 1.6$ Hz, 1H), 7.03 (t, $J = 8.0$ Hz, 1H), 3.48 (q, $J = 7.2$ Hz, 2H), 3.38 (q, $J = 7.2$ Hz, 2H), 1.29 (t, $J = 7.2$ Hz, 3H), 1.20 (t, $J = 7.2$ Hz, 3H); ^{13}C RMN (75.4 MHz, CDCl_3) δ 152.9, 149.3, 137.0, 129.3, 124.8, 123.9, 102.1, 42.5, 42.2, 14.4, 13.4; EI-LRMS m/z 399 ($\text{M}^+ + 2$, 0.1), 397 (M^+ , 0.1), 100 (100); IR (KBr) 2973, 2934, 1706, 1405, 1263, 1238, 1151, 956, 760 cm^{-1} . HRMS calcd. for $\text{C}_{11}\text{H}_{13}\text{BrINO}_2$: 396.9174. Found: 396.9178.

2-Chloro-3-iodophenyl *N,N*-diethylcarbamate (3k): The treatment of **1d** (3.19 g, 10 mmol) according to the typical procedure, using hexachloroethane (2.84 g, 12 mmol) as electrophile and purification by column chromatography (eluent: hexane/EtOAc, 10/1) on silica gel, gave **3k** (2.48 g, 70%) as a white solid. Mp 49-51 $^{\circ}\text{C}$; ^1H RMN (300 MHz, CDCl_3) δ 7.70 (dd, $J = 8.1, 1.6$ Hz, 1H), 7.20 (dd, $J = 8.1, 1.6$ Hz, 1H), 6.96 (t, $J = 8.1$ Hz, 1H), 3.47 (q, $J = 7.2$ Hz, 2H), 3.38 (q, $J = 7.2$ Hz, 2H), 1.28 (t, $J = 7.2$ Hz, 3H), 1.20 (t, $J = 7.2$ Hz, 3H); ^{13}C RMN (75.4 MHz, CDCl_3) δ 152.9, 147.8, 136.9, 132.2, 128.5, 124.2, 97.0, 42.6, 42.2, 14.3, 13.4; EI-LRMS m/z 353 (M^+ , 0.4), 100 (100); IR (KBr)

3069, 2976, 2931, 1715, 1410, 1240, 1149, 956, 760 cm^{-1} . Anal. Calcd. for $\text{C}_{11}\text{H}_{13}\text{ClINO}_2$: C, 37.37; H, 3.71; N, 3.96. Found: C, 37.31; H, 3.66; N, 3.92.

Typical Procedure for the Deprotection of 2,3-Dihalophenyl *N,N*-Diethylcarbamates **3.** To a solution of the corresponding carbamate **3** (5 mmol) in EtOH (50 mL) a large excess of NaOH (2 g, 0.05 mol) was added. The mixture was refluxed for 5-8 h (completion of the hydrolysis was monitored by GC-MS). After cooling to room temperature most of the EtOH was evaporated under reduced pressure, the residue was diluted with Et_2O and the excess of NaOH was neutralized at 0 $^{\circ}\text{C}$ using a 1M solution of HCl. The aqueous solution was extracted with Et_2O (3 x 20 mL) and the combined organic phase was washed with brine, dried (anhydrous Na_2SO_4), and evaporated under reduced pressure. The crude was purified by column chromatography (eluent: hexane/EtOAc) on silica gel to afford compounds **4**.

3-Fluoro-2-iodophenol (4a): 3-Fluoro-2-iodophenyl *N,N*-diethylcarbamate **3a** (1.68 g, 5 mmol) was used according to the typical procedure. Purification by column chromatography (eluent: hexane/EtOAc, 5/1) on silica gel, gave **4a** (0.78 g, 66%) as a brown liquid. R_f 0.20 (hexane/EtOAc, 5/1); ^1H RMN (400 MHz, CDCl_3) δ 7.23-7.16 (m, 1H), 6.82-6.78 (m, 1H), 6.68-6.62 (m, 1H), 5.80 (br s, 1H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 162.2 (d, $J = 244.3$ Hz), 156.8 (d, $J = 4.2$ Hz), 130.6 (d, $J = 9.9$ Hz), 110.7 (d, $J = 3.1$ Hz), 107.7 (d, $J = 23.8$ Hz), 74.2 (d, $J = 27.8$ Hz); EI-LRMS m/z 238 (100) (M^+ , 100); IR (neat) 3478, 2978, 2927, 1584, 1454, 1188, 992, 771 cm^{-1} . HRMS calcd. for $\text{C}_6\text{H}_4\text{FIO}$: 237.9291. Found: 237.9298.

2-Bromo-3-fluorophenol (4b):⁴ 2-Bromo-3-fluorophenyl *N,N*-diethylcarbamate **3b** (1.45 g, 5 mmol) was used according to the typical procedure. Purification by column chromatography (eluent: hexane/EtOAc, 5/1) on silica gel, gave **4b** (0.71 g, 75%) as a yellow liquid. R_f 0.31 (hexane/EtOAc, 5/1); ^1H RMN (300 MHz, CDCl_3) δ 7.21-7.11 (m, 1H), 6.81 (dt, $J = 8.2, 1.4$ Hz, 1H), 6.70 (dt, $J = 8.2, 1.4$ Hz, 1H), 6.12 (br s, 1H); ^{13}C RMN (75.4 MHz, CDCl_3) δ 159.6 (d, $J = 276.7$ Hz), 154.2 (d, $J = 2.9$ Hz), 129.0 (d, $J = 9.4$ Hz), 111.6 (d, $J = 3.2$ Hz), 108.1 (d, $J = 22.1$ Hz), 98.2 (d, $J = 23.2$ Hz); EI-LRMS m/z 192 ($\text{M}^+ + 2$, 99), 190 (M^+ , 100); IR (neat) 3412, 2963, 2933, 1599, 1464, 1193, 1002, 771 cm^{-1} . HRMS calcd. for $\text{C}_6\text{H}_4\text{BrFO}$: 188.9430. Found: 188.9425.

2-Chloro-3-fluorophenol (4c): 2-Chloro-3-fluorophenyl *N,N*-diethylcarbamate **3c** (1.23 g, 5 mmol) was used according to the typical procedure. Purification by column chromatography (eluent: hexane/EtOAc, 5/1) on silica gel, gave **4c** (0.33 g, 45%) as a colorless liquid. R_f 0.25(hexane/EtOAc, 5/1); ^1H RMN (300 MHz, CDCl_3) δ 7.17-7.07 (m, 1H), 6.83 (dd, $J = 8.5, 1.4$ Hz, 1H), 6.73 (dt, $J = 8.5, 1.4$ Hz, 1H), 5.61 (br s, 1H); ^{13}C RMN (75.4 MHz, CDCl_3) δ 158.6 (d, $J = 248.2$ Hz), 153.1 (d, $J = 1.5$ Hz), 128.0 (d, $J = 9.5$ Hz), 111.7 (d, $J = 3.2$ Hz), 108.3 (d, $J = 20.9$ Hz), 108.1 (d, $J = 19.5$ Hz); EI-LRMS m/z 148 ($\text{M}^+ + 2$, 32), 146 (M^+ , 100); IR (neat) 3412, 2970, 2937, 1467, 1195, 1004, 772 cm^{-1} . HRMS calcd. for $\text{C}_6\text{H}_4\text{ClFO}$: 145.9935. Found: 145.9938.

3-Chloro-2-iodophenol (4d): 3-Chloro-2-iodophenyl *N,N*-diethylcarbamate **3d** (1.77 g, 5 mmol) was used according to the typical procedure. Purification by column chromatography (eluent: hexane/EtOAc, 5/1) on silica gel, gave **4d** (1.12 g, 88%) as a brownish solid. Mp 54-56 $^\circ\text{C}$ (lit.⁵ mp 56 $^\circ\text{C}$); ^1H RMN (400 MHz, CDCl_3) δ 7.17 (t, $J = 8.2$ Hz, 1H), 7.03 (dd, $J = 8.2, 1.5$ Hz, 1H), 6.88 (dd, $J = 8.2, 1.5$ Hz, 1H), 5.55 (br s, 1H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 156.6, 136.6, 130.3, 121.6, 111.9, 91.3; EI-LRMS m/z 256 ($\text{M}^+ + 2$, 31), 254 (M^+ , 100); IR (KBr) 3275, 1568, 1435, 1272, 896, 767 cm^{-1} . Anal. Calcd. for $\text{C}_6\text{H}_4\text{ClIO}$: C, 28.32; H, 1.58. Found: C, 28.30; H, 1.55.

2-Bromo-3-chlorophenol (4e): 2-Bromo-3-chlorophenyl *N,N*-diethylcarbamate **3e** (1.53 g, 5 mmol) was used according to the typical procedure. Purification by column chromatography (eluent: hexane/EtOAc, 5/1) on silica gel, gave **4e** (0.85 g, 82%) as a white solid. Mp 53-55 $^\circ\text{C}$ (lit.⁵ mp 55.5 $^\circ\text{C}$); ^1H RMN (300 MHz, CDCl_3) δ 7.15 (t, $J = 8.0$ Hz, 1H), 7.03 (dd, $J = 8.0, 1.4$ Hz, 1H), 6.92 (dd, $J = 8.0, 1.4$ Hz, 1H), 5.80 (br s, 1H); ^{13}C RMN (75.4 MHz, CDCl_3) δ 153.9, 134.6, 129.0, 122.2, 114.2, 111.3; EI-LRMS m/z 210 ($\text{M}^+ + 4$, 24), 208 ($\text{M}^+ + 2$, 100), 206 (M^+ , 77); IR (KBr) 3393, 3086, 1452, 1426, 1159, 1030, 774 cm^{-1} . Anal. Calcd. for $\text{C}_6\text{H}_4\text{BrClO}$: C, 34.74; H, 1.94. Found: C, 34.79; H, 1.91.

3-Bromo-2-iodophenol (4f): 3-Bromo-2-iodophenyl *N,N*-diethylcarbamate **3f** (1.99 g, 5 mmol) was used according to the typical procedure. Purification by column chromatography (eluent: hexane/EtOAc, 5/1) on silica gel, gave **4f** (1.33 g, 89%) as a brown solid. Mp 83-85 $^\circ\text{C}$ (lit.⁶ mp 85 $^\circ\text{C}$); ^1H RMN (400 MHz, CDCl_3) δ 7.20 (dd, $J = 8.1, 1.5$ Hz, 1H), 7.12 (t, $J = 8.1$ Hz, 1H), 6.92 (dd, $J = 8.1,$

1.5 Hz, 1H), 5.54 (br s, 1H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 156.6, 130.8, 129.6, 125.0, 113.3, 94.4; EI-LRMS m/z 300 ($\text{M}^+ + 2$, 98), 298 (M^+ , 100); IR (KBr) 3383, 1432, 1272, 872, 764 cm^{-1} . Anal. Calcd. for $\text{C}_6\text{H}_4\text{BrIO}$: C, 24.11; H, 1.35. Found: C, 24.07; H, 1.37.

2,3-Dibromophenol (4g): 2,3-Dibromophenyl *N,N*-diethylcarbamate **3g** (1.75 g, 5 mmol) was used according to the typical procedure. Purification by column chromatography (eluent: hexane/EtOAc, 5/1) on silica gel, gave **4g** (1.18 g, 94%) as a white solid. Mp 64-66 °C (lit.⁷ mp 68-69 °C); ^1H RMN (300 MHz, CDCl_3) δ 7.20 (dd, J = 8.0, 1.7 Hz, 1H), 7.1 (t, J = 8.0 Hz, 1H), 6.97 (dd, J = 8.0, 1.7 Hz, 1H), 5.68 (br s, 1H); ^{13}C RMN (75.4 MHz, CDCl_3) δ 153.8, 129.5, 125.5, 124.6, 114.8, 113.6; EI-LRMS m/z 254 ($\text{M}^+ + 4$, 49), 252 ($\text{M}^+ + 2$, 100), 250 (M^+ , 49); IR (KBr) 3383, 3075, 2978, 2935, 1447, 1421, 1283, 1200, 1159, 1026, 872, 769 cm^{-1} . Anal. Calcd. for $\text{C}_6\text{H}_4\text{Br}_2\text{O}$: C, 28.61; H, 1.60. Found: C, 28.59; H, 1.62.

3-Bromo-2-chlorophenol (4h): 3-Bromo-2-chlorophenyl *N,N*-diethylcarbamate **3h** (1.53 g, 5 mmol) was used according to the typical procedure. Purification by column chromatography (eluent: hexane/EtOAc, 5/1) on silica gel, gave **4h** (0.85 g, 82%) as a white solid. Mp 65-67 °C (lit.⁶ mp 60 °C); ^1H RMN (300 MHz, CDCl_3) δ 7.20 (dd, J = 8.0, 1.6 Hz, 1H), 7.06 (t, J = 8.0 Hz, 1H), 6.98 (dd, J = 8.0, 1.6 Hz, 1H), 5.65 (br s, 1H); ^{13}C RMN (75.4 MHz, CDCl_3) δ 152.6, 128.7, 125.5, 122.3, 121.0, 115.1; EI-LRMS m/z 210 ($\text{M}^+ + 4$, 26), 208 ($\text{M}^+ + 2$, 100), 206 (M^+ , 80); IR (KBr) 3388, 1570, 1442, 1283, 882, 764 cm^{-1} ; Anal. Calcd. for $\text{C}_6\text{H}_4\text{BrClO}$: C, 34.74; H, 1.94. Found: C, 34.69; H, 1.91.

2,3-Diiodophenol (4i): 2,3-Diiodophenyl *N,N*-diethylcarbamate **3i** (2.22 g, 5 mmol) was used according to the typical procedure. Purification by column chromatography (eluent: hexane/EtOAc, 5/1) on silica gel, gave **4i** (1.42 g, 82%) as a white solid. Mp 102-104 °C; ^1H RMN (400 MHz, CDCl_3) δ 7.44 (dd, J = 7.8, 1.5 Hz, 1H), 7.00 (t, J = 7.8 Hz, 1H), 6.94 (dd, J = 7.8, 1.5 Hz, 1H), 5.47 (br s, 1H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 156.0, 131.8, 131.4, 114.0, 107.4, 100.7; EI-LRMS m/z 346 (M^+ , 100); IR (KBr) 3387, 1574, 1424, 1268, 856, 761 cm^{-1} . Anal. Calcd. for $\text{C}_6\text{H}_4\text{I}_2\text{O}$: C, 20.83; H, 1.17. Found: C, 20.88; H, 1.13.

2-Bromo-3-iodophenol (4j): 2-Bromo-3-iodophenyl *N,N*-diethylcarbamate **3j** (1.99 g, 5 mmol) was used according to the typical procedure. Purification by column chromatography (eluent: hexane/EtOAc, 5/1) on silica gel, gave **4j** (1.04 g, 70%) as a white solid. Mp 69-71 °C; ¹H RMN (300 MHz, CDCl₃) δ 7.44-7.39 (m, 1H), 7.02-6.92 (m, 2H), 5.32 (br s, 1H); ¹³C RMN (75.4 MHz, CDCl₃) δ 152.9, 132.1, 130.3, 118.2, 115.6, 100.5; EI-LRMS *m/z* 300 (M⁺+2, 98), 298 (M⁺, 100); IR (KBr) 3476, 1565, 1447, 1421, 1277, 1205, 1164, 1016, 861, 779 cm⁻¹. Anal. Calcd. for C₆H₄BrIO: C, 24.11; H, 1.35. Found: C, 24.07; H, 1.38.

2-Chloro-3-iodophenol (4k): 2-Chloro-3-iodophenyl *N,N*-diethylcarbamate **3k** (1.77 g, 5 mmol) was used according to the typical procedure. Purification by column chromatography (eluent: hexane/EtOAc, 5/1) on silica gel, gave **4k** (1.08 g, 85%) as a white solid. Mp 72-74 °C; ¹H RMN (300 MHz, CDCl₃) δ 7.41 (dd, *J* = 8.0, 1.6 Hz, 1H), 7.00 (dd, *J* = 8.0, 1.6 Hz, 1H), 6.91 (t, *J* = 8.0 Hz, 1H), 5.61 (br s, 1H); ¹³C RMN (75.4 MHz, CDCl₃) δ 151.7, 131.9, 129.6, 124.7, 116.1, 97.5; EI-LRMS *m/z* 256 (M⁺+2, 32), 254 (M⁺, 100); IR (KBr) 3512, 1565, 1452, 1283, 1211, 1164, 1036, 861, 769 cm⁻¹. Anal. Calcd. for C₆H₄ClIO: C, 28.32; H, 1.58. Found: C, 28.35; H, 1.53.

Typical Procedure for the Synthesis of 4-Halobenzo[*b*]furans 5 and Benzo[*b*]furan 6a. To a stirred solution of the corresponding phenol **4a,d,f,i** (1 mmol) and piperidine (0.085 g, 1 mmol) in DMF (1 mL), the corresponding 1-alkyne (1.2 mmol or 2.4 mmol for **4i**), Pd(OAc)₂(PPh₃)₂ (0.02 mmol or 0.04 mmol for **4i**) and CuI (0.04 mmol or 0.08 mmol for **4i**) were added under N₂ atmosphere. The mixture was heated at 60 °C for 5 h and then, it was cooled to room temperature and diluted with H₂O (15 mL). The aqueous solution was extracted with Et₂O (3 x 20 mL) and the combined organic phase was dried (anhydrous Na₂SO₄), and evaporated under reduced pressure. The crude was purified by column chromatography (eluent: hexane) on silica gel to afford compounds **5**.

4-Fluoro-2-phenylbenzo[*b*]furan (5a): The reaction of **4a** (0.238 g, 1 mmol), piperidine (0.085 g, 1 mmol), phenylacetylene (0.123 g, 1.2 mmol), Pd(OAc)₂(PPh₃)₂ (0.015 g, 0.02 mmol) and CuI (0.008 g, 0.04 mmol) in DMF (1mL) afforded **5a** (0.168 g, 79%) as a white solid. Mp 55-57 °C; ¹H RMN (400 MHz, CDCl₃) δ 7.88-7.83 (m, 2H), 7.48-7.42 (m, 2H), 7.39-7.31 (m, 2H), 7.24-7.17 (m, 1H), 7.06 (s,

1H), 6.96-6.90 (m, 1H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 156.8 (d, $J = 9.9$ Hz), 156.1, 156.0 (d, $J = 250.2$ Hz), 130.0, 129.0, 128.7, 125.1, 124.7 (d, $J = 7.3$ Hz), 118.6 (d, $J = 22.2$ Hz), 108.6 (d, $J = 18.7$ Hz), 107.5 (d, $J = 3.8$ Hz), 97.4 (d, $J = 1.6$ Hz); EI-LRMS m/z 212 (M^+ , 100); IR (KBr) 3122, 3093, 3035, 1482, 1260, 1235, 1050, 759, 687 cm^{-1} . Anal. Calcd. for $\text{C}_{14}\text{H}_9\text{FO}$: C, 79.23; H, 4.27. Found: C, 79.30; H, 4.23.

2-Butyl-4-fluorobenzo[b]furan (5b): The reaction of **4a** (0.238 g, 1 mmol), piperidine (0.085 g, 1 mmol), 1-hexyne (0.099 g, 1.2 mmol), $\text{Pd}(\text{OAc})_2(\text{PPh}_3)_2$ (0.015 g, 0.02 mmol) and CuI (0.008 g, 0.04 mmol) in DMF (1mL) afforded **5b** (0.115 g, 60%) as a colorless liquid. R_f 0.57(hexane); ^1H RMN (400 MHz, CDCl_3) δ 7.22 (d, $J = 8.2$ Hz, 1H), 7.17-7.10 (m, 1H), 6.92-6.85 (m, 1H), 6.47 (s, 1H), 2.77 (t, $J = 7.6$ Hz, 2H), 1.79-1.70 (m, 2H), 1.49-1.38 (m, 2H), 0.97 (t, $J = 7.5$ Hz, 3H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 159.9, 156.7 (d, $J = 10$ Hz), 155.5 (d, $J = 248.5$ Hz), 123.5 (d, $J = 7.4$ Hz), 118.0 (d, $J = 22.2$ Hz), 108.1 (d, $J = 19.1$ Hz), 107.2 (d, $J = 3.8$ Hz), 98.1 (d, $J = 1.5$ Hz), 29.8, 28.2, 22.4, 13.9; EI-LRMS m/z 192 (M^+ , 26), 149 (100); IR (neat) 2958, 2931, 2871, 1592, 1492, 1437, 1025, 769 cm^{-1} . HRMS calcd. for $\text{C}_{12}\text{H}_{13}\text{FO}$: 192.0950. Found: 192.0959.

4-Chloro-2-phenylbenzo[b]furan (5c): The reaction of **4d** (0.254 g, 1 mmol), piperidine (0.085 g, 1 mmol), phenylacetylene (0.123 g, 1.2 mmol), $\text{Pd}(\text{OAc})_2(\text{PPh}_3)_2$ (0.015 g, 0.02 mmol) and CuI (0.008 g, 0.04 mmol) in DMF (1mL) afforded **5c** (0.139 g, 61%) as a white solid. Mp 66-68 $^\circ\text{C}$; ^1H RMN (400 MHz, CDCl_3) δ 7.91-7.87 (m, 2H), 7.51-7.38 (m, 4H), 7.28-7.19 (m, 2H), 7.12 (s, 1H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 156.6, 155.1, 129.9, 129.1, 128.9, 128.8, 125.8, 125.1, 124.8, 122.9, 109.8, 99.8; EI-LRMS m/z 230 ($\text{M}^+ + 2$, 31), 228 (M^+ , 100); IR (KBr) 1473, 1426, 944, 754 cm^{-1} . Anal. Calcd. for $\text{C}_{14}\text{H}_9\text{ClO}$: C, 73.53; H, 3.97. Found: C, 73.61; H, 3.94.

2-Butyl-4-chlorobenzo[b]furan (5d): The reaction of **4d** (0.254 g, 1 mmol), piperidine (0.085 g, 1 mmol), 1-hexyne (0.099 g, 1.2 mmol), $\text{Pd}(\text{OAc})_2(\text{PPh}_3)_2$ (0.015 g, 0.02 mmol) and CuI (0.008 g, 0.04 mmol) in DMF (1mL) afforded **5d** (0.152 g, 73%) as a colorless liquid. R_f 0.55 (hexane); ^1H RMN (400 MHz, CDCl_3) δ 7.33 (dt, $J = 7.9, 1.0$ Hz, 1H), 7.14 (t, $J = 7.9$ Hz, 1H), 7.20 (dd, $J = 7.9, 1.0$ Hz, 1H), 6.50 (s, 1H), 2.79 (t, $J = 7.5$ Hz, 2H), 1.80-1.71 (m, 2H), 1.50-1.40 (m, 2H), 0.99 (t, $J = 7.5$ Hz, 3H);

^{13}C RMN (100.6 MHz, CDCl_3) δ 160.7, 154.9, 128.5, 125.2, 123.7, 122.4, 109.4, 100.6, 29.8, 28.2, 22.4, 13.9; EI-LRMS m/z 210 ($\text{M}^+ + 2$, 8), 208 (M^+ , 24), 165 (100); IR (neat) 2957, 2927, 2870, 1583, 1476, 1425, 1260, 768 cm^{-1} . HRMS calcd. for $\text{C}_{12}\text{H}_{13}\text{ClO}$: 208.0655. Found: 208.0662.

4-Bromo-2-phenylbenzo[*b*]furan (5e): The reaction of **4f** (0.299 g, 1 mmol), piperidine (0.085 g, 1 mmol), phenylacetylene (0.123 g, 1.2 mmol), $\text{Pd}(\text{OAc})_2(\text{PPh}_3)_2$ (0.015 g, 0.02 mmol) and CuI (0.008 g, 0.04 mmol) in DMF (1mL) afforded **5e** (0.197 g, 72%) as a white solid. Mp 66-68 $^\circ\text{C}$; ^1H RMN (400 MHz, CDCl_3) δ 7.91-7.87 (m, 2H), 7.58-7.45 (m, 3H), 7.44-7.38 (m, 2H), 7.16 (t, $J = 8.1$ Hz, 1H), 7.07 (s, 1H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 156.5, 154.5, 130.8, 129.9, 129.1, 128.9, 126.0, 125.1, 113.9, 110.3, 102.3, 101.4; EI-LRMS m/z 274 ($\text{M}^+ + 2$, 96), 272 (M^+ , 100); IR (KBr) 3083, 1423, 1268, 1248, 1160, 1022, 908, 751, 685 cm^{-1} . HRMS calcd. for $\text{C}_{14}\text{H}_9\text{BrO}$: 271.9837. Found: 271.9878. Anal. Calcd. for $\text{C}_{14}\text{H}_9\text{BrO}$: C, 61.57; H, 3.32. Found: C, 61.49; H, 3.28.

4-Bromo-2-butylbenzo[*b*]furan (5f): The reaction of **4f** (0.299 g, 1 mmol), piperidine (0.085 g, 1 mmol), 1-hexyne (0.099 g, 1.2 mmol), $\text{Pd}(\text{OAc})_2(\text{PPh}_3)_2$ (0.015 g, 0.02 mmol) and CuI (0.008 g, 0.04 mmol) in DMF (1mL) afforded **5f** (0.157 g, 62%) as a colorless liquid. R_f 0.64 (hexane); ^1H RMN (400 MHz, CDCl_3) δ 7.37-7.32 (m, 2H), 7.07 (t, $J = 8.0$ Hz, 1H), 6.43 (s, 1H), 2.78 (t, $J = 7.6$ Hz, 2H), 1.79-1.70 (m, 2H), 1.49-1.38 (m, 2H), 0.97 (t, $J = 7.5$ Hz, 3H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 160.6, 154.4, 130.5, 125.4, 124.1, 113.2, 109.9, 102.1, 29.7, 28.2, 22.4, 13.9; EI-LRMS m/z 254 ($\text{M}^+ + 2$, 21), 252 (M^+ , 24), 211 (100). HRMS calcd. for $\text{C}_{12}\text{H}_{13}\text{BrO}$: 252.0150. Found: 252.0145.

2-Phenyl-4-(phenylethynyl)benzo[*b*]furan (6a): The reaction of **4i** (0.346 g, 1 mmol), piperidine (0.170 g, 2 mmol), phenylacetylene (0.246 g, 2.4 mmol), $\text{Pd}(\text{OAc})_2(\text{PPh}_3)_2$ (0.030 g, 0.04 mmol) and CuI (0.016 g, 0.08 mmol) in DMF (2mL) afforded **6** (0.180 g, 61%) as a white solid. Mp 78-80 $^\circ\text{C}$; ^1H RMN (400 MHz, CDCl_3) δ 7.93-7.89 (m, 2H), 7.65-7.60 (m, 2H), 7.52-7.33 (m, 8H), 7.27-7.23 (m, 2H); ^{13}C RMN (100.6 MHz, CDCl_3) δ 156.8, 154.6, 131.9, 131.5, 130.4, 129.1, 128.7, 128.6, 126.6, 125.3, 124.3, 123.5, 115.8, 111.7, 101.3, 92.8, 87.4; EI-LRMS m/z 294 (M^+ , 100); IR (KBr) 3078, 3054, 3031, 1487, 1260, 920, 787 cm^{-1} . Anal. Calcd. for $\text{C}_{22}\text{H}_{14}\text{O}$: C, 89.77; H, 4.79. Found: C, 89.80; H, 4.74.

Typical Procedure for the Synthesis of 4-Alkylbenzo[*b*]furans 6b,c.⁸ After flushing with nitrogen, Et₂O (5 mL), NiCl₂(dppp) (2.71 mg, 1 mol%) and benzofuran **5c** (0.114 g, 0.5 mmol) were placed in a flask. Then a solution of alkylmagnesium halide (~1.2 mL of a 0.5M solution in Et₂O) was added at room temperature. The resulting solution was heated under reflux for an hour. After cooling to room temperature the reaction mixture was poured into 1M HCl (10 mL). The aqueous solution was extracted with hexane (3 x 10 mL) and the combined organic phase was dried (anhydrous Na₂SO₄), and evaporated under reduced pressure. The crude was purified by column chromatography (eluent: hexane) on silica gel to afford compounds **6b,c**.

4-Heptyl-2-phenylbenzo[*b*]furan (6b): The reaction of **5c** (0.114 g, 0.5 mmol), NiCl₂(dppp) (2.71 mg, 0.0055 mmol) and heptyl magnesium bromide (1.2 mL, 0.6 mmol) in Et₂O (5mL) afforded **6b** (0.133 g, 91%) as a colorless liquid. *R_f* 0.39 (hexane); ¹H RMN (300 MHz, CDCl₃) δ 7.91-7.85 (m, 2H), 7.49-7.41 (m, 2H), 7.39-7.31 (m, 2H), 7.21 (t, *J* = 7.8 Hz, 1H), 7.06 (s, 1H), 7.03 (d, *J* = 7.1 Hz, 1H), 2.87 (t, *J* = 7.7 Hz, 2H), 1.79-1.67 (m, 2H), 1.45-1.23 (m, 8H), 0.89 (t, *J* = 7.1 Hz, 3H); ¹³C RMN (75.4 MHz, CDCl₃) δ 155.4, 155.0, 136.1, 130.7, 128.9, 128.6, 128.5, 125.0, 124.3, 122.4, 108.8, 100.1, 33.5, 32.0, 30.9, 29.7, 29.4, 22.8, 14.3; EI-LRMS *m/z* 292 (M⁺, 61), 207 (100); IR (neat) 3428, 2953, 2930, 2848, 1424, 1252, 778, 758, 684 cm⁻¹. HRMS calcd. for C₂₁H₂₄O: 292.1827. Found: 292.1826.

4-Butyl-2-phenylbenzo[*b*]furan (6c): The reaction of **5c** (0.114 g, 0.5 mmol), NiCl₂(dppp) (2.71 mg, 0.0055 mmol) and butyl magnesium bromide (1.2 mL, 0.6 mmol) in Et₂O (5mL) afforded **6c** (0.100 g, 80%) as a colorless liquid. *R_f* 0.30 (hexane); ¹H RMN (400 MHz, CDCl₃) δ 7.90 (d, *J* = 8.0 Hz, 2H), 7.46 (t, *J* = 7.8 Hz, 2H), 7.40-7.33 (m, 2H), 7.25-7.19 (m, 1H), 7.08 (s, 1H), 7.05 (d, *J* = 7.4 Hz, 1H), 2.89 (t, *J* = 7.7 Hz, 2H), 1.78-1.69 (m, 2H), 1.49-1.39 (m, 2H), 0.98 (t, *J* = 7.4 Hz, 3H); ¹³C RMN (75.4 MHz, CDCl₃) δ 155.4, 155.0, 136.1, 130.7, 128.9, 128.6, 128.5, 125.0, 124.3, 122.4, 108.8, 100.1, 33.2, 33.0, 22.8, 14.2; EI-LRMS *m/z* 250 (M⁺, 44), 207 (100); IR (neat) 2958, 2924, 2857, 1487, 1429, 1250, 758, 691 cm⁻¹. HRMS calcd. for C₁₈H₁₈O: 250.1358. Found: 250.1349.

Typical Procedure for the Synthesis of 4-Functionalized Benzo[*b*]furans 6d,e. *t*-Butyllithium (0.66 mL of a 1.5M solution in pentane, 1 mmol) was added at -78 °C to a stirred solution of benzofuran

5e (0.137 g, 0.5 mmol) in Et₂O (5 mL). After 30 min at this temperature, the corresponding electrophile (0.55 mmol) was added and the reaction mixture was stirred from –78 °C to room temperature overnight. The mixture was quenched with H₂O, the aqueous solution was extracted with Et₂O (3 x 20 mL) and the combined organic phase was dried (anhydrous Na₂SO₄), and evaporated under reduced pressure. The crude was purified by column chromatography (eluent: hexane) on silica gel to afford compounds **6d,e**.

4-(Diphenylhydroxymethyl)-2-phenylbenzo[*b*]furan (6d): The treatment of **5e** (0.137 g, 0.5 mmol) according to the typical procedure, using benzophenone (0.100 g, 0.55 mmol) as electrophile gave **6d** (0.137 g, 73%) as a white solid. Mp 159-161 °C; ¹H RMN (300 MHz, CDCl₃) δ 7.79-7.73 (m, 2H), 7.50 (d, *J* = 8.3 Hz, 1H), 7.44-7.28 (m, 13H), 7.17 (t, *J* = 7.8 Hz, 1H), 6.81 (d, *J* = 7.4 Hz, 1H), 6.74 (s, 1H), 3.12 (br s, 1H); ¹³C RMN (75.4 MHz, CDCl₃) δ 155.6, 155.4, 146.2, 139.9, 130.4, 128.8, 128.6, 128.2, 128.1, 127.9, 127.5, 125.1, 123.5, 123.2, 111.1, 102.7, 82.5; EI-LRMS *m/z* 376 (M⁺, 100); IR (KBr) 3420, 3054, 3027, 2925, 2848, 1491, 1442, 1420, 1031, 759, 705, 688 cm⁻¹. HRMS calcd. for C₂₇H₂₀O₂: 376.1463. Found: 376.1469.

2-Phenyl-4-tributylstannylbenzo[*b*]furan (6e): The treatment of **5e** (0.137 g, 0.5 mmol) according to the typical procedure, using tributyltin chloride (0.179 g, 0.55 mmol) as electrophile gave **6e** (0.188 g, 78%) as a colorless liquid. *R_f* 0.41 (hexane); ¹H RMN (300 MHz, CDCl₃) δ 7.90-7.84 (m, 2H), 7.50-7.42 (m, 3H), 7.38-7.21 (m, 3H), 6.93 (s, 1H), 1.65-1.52 (m, 6H), 1.40-1.31 (m, 6H), 1.21-1.17 (m, 6H), 0.89 (t, *J* = 7.3 Hz, 9H); ¹³C RMN (75.4 MHz, CDCl₃) δ 155.8, 153.9, 136.1, 135.1, 131.3, 130.8, 128.9, 128.6, 125.0, 124.0, 111.0, 103.4, 29.3, 27.5, 13.8, 10.1; EI-LRMS *m/z* 484 (M⁺, 2), 313 (100); IR (neat) 2953, 2921, 2848, 1461, 1400, 1244, 757, 687 cm⁻¹. HRMS calcd. for C₂₆H₃₆OSn: 484.1788. Found: 484.1774.

Typical Procedure for the Synthesis of 4-Arylbenzo[*b*]furans 6f,g.^{8,9} In a Schlenk tube benzofuran **5e** (0.109 g, 0.4 mmol), the corresponding arylboronic acid (0.6 mmol) and [Pd(PPh₃)₄] (9.2 mg, 2 mol%) were introduced under nitrogen. Then, DME (5 mL) was added followed by the addition of Na₂CO₃ (0.064 g, 0.6 mmol) in H₂O (2 mL). The reaction mixture was vigorously stirred and heated at

80 °C. The progress of the reaction was monitored by GC-MS and after 24 h another load of catalyst was added. After completion of the reaction (48 h), the mixture was diluted with CH₂Cl₂ (10 mL) and the organic layer was washed with brine (2 x 10 mL). The combined organic extracts were dried (anhydrous Na₂SO₄) and evaporated under reduced pressure. The residue was purified by column chromatography (eluent: hexane) on silica gel to afford compounds **6f,g**.

2,4-Diphenylbenzo[*b*]furan (6f): The reaction of **5e** (0.109 g, 0.4 mmol), phenylboronic acid (0.073 g, 0.6 mmol) and [Pd(PPh₃)₄] (2 x 9.2 mg, 4 mol%) in DME (5 mL) followed by the addition of Na₂CO₃ (0.064 g, 0.6 mmol) in H₂O (2 mL) afforded **6f** (0.078 g, 72%) as a yellow oil. *R_f* 0.30 (hexane/EtOAc, 100/1); ¹H RMN (300 MHz, CDCl₃) δ 7.96-7.91 (m, 2H), 7.77-7.71 (m, 2H), 7.62-7.36 (m, 9H), 7.28 (s, 1H); ¹³C RMN (75.4 MHz, CDCl₃) δ 156.3, 155.3, 140.3, 135.1, 130.5, 128.9, 128.8, 128.7, 128.6, 127.7, 127.5, 125.0, 124.6, 122.7, 110.2, 100.9; EI-LRMS *m/z* 270 (M⁺, 100); IR (neat) 3053, 3030, 1473, 1415, 1253, 791, 759, 697 cm⁻¹. HRMS calcd. for C₂₀H₁₄O: 270.1045. Found: 270.1034.

4-(3-Nitrophenyl)-2-phenylbenzo[*b*]furan (6g): The reaction of **5e** (0.109 g, 0.4 mmol), 3-nitrophenylboronic acid (0.100 g, 0.6 mmol) and [Pd(PPh₃)₄] (2 x 9.2 mg, 4 mol%) in DME (5 mL) followed by the addition of Na₂CO₃ (0.064 g, 0.6 mmol) in H₂O (2 mL) afforded **6g** (0.082 g, 65%) as a brown solid. Mp 115-117 °C; ¹H RMN (400 MHz, CDCl₃) δ 8.54(t, *J* = 2.0 Hz, 1H), 8.27 (ddd, *J* = 8.3, 2.4, 0.9 Hz, 1H), 8.00 (dt, *J* = 7.7, 1.3 Hz, 1H), 7.91-7.87 (m, 2H), 7.69 (t, *J* = 7.9 Hz, 1H), 7.60 (dt, *J* = 7.9, 0.9 Hz, 1H), 7.50-7.33 (m, 5H), 7.16 (s, 1H); ¹³C RMN (100.6 MHz, CDCl₃) δ 157.2, 156.4, 148.8, 142.0, 134.5, 132.3, 130.1, 129.9, 129.1, 129.0, 127.7, 125.2, 124.8, 123.3, 122.9, 122.3, 111.5, 100.0; EI-LRMS *m/z* 315 (M⁺, 100); IR (KBr) 3032, 2919, 2853, 1532, 1342, 1255, 753, 738, 682 cm⁻¹. HRMS calcd. for C₂₀H₁₃NO₂: 315.0895. Found: 315.0891.

Synthesis of 4-(1-Hexynyl)-2-phenylbenzo[*b*]furan (6h):⁹ In a Schlenk tube benzofuran **5e** (0.095 g, 0.35 mmol), 1-hexyne (0.043 g, 0.52 mmol), PdCl₂(PPh₃)₂ (7.7 mg, 5 mol%), CuI (6.75 mg, 10 mol%) and PPh₃ (4.6 mg, 5 mol%) were introduced under nitrogen. Then, anhydrous Et₃N (4 mL) and DMF (2 mL) were added. The reaction mixture was vigorously stirred and heated at 70 °C for 96 h. The progress of the reaction was monitored by GC-MS and after 48 h another load of alkyne and catalysts was added.

After completion of the reaction, the mixture was poured into water and extracted with CH_2Cl_2 (2 x 10 mL). The combined organic layers were washed with brine (2 x 10 mL), dried over anhydrous Na_2SO_4 and evaporated under reduced pressure. The residue was purified by column chromatography (eluent: hexane) on silica gel to afford **6h** (0.071 g, 74%) as a colorless liquid. R_f 0.40 (hexane/EtOAc, 100/1); ^1H RMN (300 MHz, CDCl_3) δ 7.93-7.87 (m, 2H), 7.51-7.42 (m, 3H), 7.41-7.33 (m, 1H), 7.30 (dd, J = 7.6, 1.0 Hz, 1H), 7.21 (d, J = 8.3 Hz, 1H), 7.17 (s, 1H), 2.55 (t, J = 7.0 Hz, 2H), 1.75-1.50 (m, 4H), 1.01 (t, J = 7.3 Hz, 3H); ^{13}C RMN (75.4 MHz, CDCl_3) δ 156.3, 154.5, 131.4, 130.4, 128.9, 128.8, 126.4, 125.1, 124.1, 116.7, 110.8, 101.2, 94.0, 78.5, 31.1, 22.2, 19.5, 13.8; EI-LRMS m/z 274 (M^+ , 100); IR (neat) 2959, 2924, 2870, 1422, 1261, 1157, 784, 761, 684 cm^{-1} . HRMS calcd. for $\text{C}_{20}\text{H}_{18}\text{O}$: 274.1358. Found: 274.1369.

1a

RMN ^1H spectrum

RMN ^{13}C spectrum

1b

RMN ^1H spectrum

RMN ^{13}C spectrum

1c

RMN ^1H spectrum

RMN ^{13}C spectrum

1d

RMN ^1H spectrum

RMN ^{13}C spectrum

3a

RMN ^1H spectrum

RMN ^{13}C spectrum

3b

RMN ^1H spectrum

RMN ^{13}C spectrum

3c

RMN ^1H spectrum

RMN ^{13}C spectrum

3d

RMN ^1H spectrum

RMN ^{13}C spectrum

3e

RMN ^1H spectrum

RMN ^{13}C spectrum

3f

RMN ^1H spectrum

RMN ^{13}C spectrum

3g

RMN ^1H spectrum

RMN ^{13}C spectrum

3h

RMN ^1H spectrum

RMN ^{13}C spectrum

3i

RMN ^1H spectrum

RMN ^{13}C spectrum

3j

RMN ^1H spectrum

RMN ^{13}C spectrum

3k

RMN ^1H spectrum

RMN ^{13}C spectrum

163.133
161.222
156.711
156.735
130.853
130.572
110.710
110.711
107.513
107.516
77.008
77.160
76.979
76.812
76.571
76.401
76.101

PPM

4b

RMN ^1H spectrum

RMN ^{13}C spectrum

4c

RMN ^1H spectrum

RMN ^{13}C spectrum

4d

RMN ^1H spectrum

RMN ^{13}C spectrum

4e

RMN ^1H spectrum

RMN ^{13}C spectrum

4f

RMN ^1H spectrum

RMN ^{13}C spectrum

4g

RMN ^1H spectrum

RMN ^{13}C spectrum

4h

RMN ^1H spectrum

RMN ^{13}C spectrum

4i

RMN ^1H spectrum

RMN ^{13}C spectrum

4j

RMN ^1H spectrum

RMN ^{13}C spectrum

4k

RMN ^1H spectrum

RMN ^{13}C spectrum

5a

RMN ^1H spectrum

RMN ^{13}C spectrum

5b

RMN ^1H spectrum

RMN ^{13}C spectrum

5c

RMN ^1H spectrum

RMN ^{13}C spectrum

5d

RMN ^1H spectrum

RMN ^{13}C spectrum

5e

RMN ^1H spectrum

RMN ^{13}C spectrum

5f

RMN ^1H spectrum

RMN ^{13}C spectrum

6a

RMN ^1H spectrum

RMN ^{13}C spectrum

6b

RMN ^1H spectrum

RMN ^{13}C spectrum

6c

RMN ^1H spectrum

RMN ^{13}C spectrum

6d

RMN ^1H spectrum

RMN ^{13}C spectrum

6e

RMN ^1H spectrum

RMN ^{13}C spectrum

6f

RMN ^1H spectrum

RMN ^{13}C spectrum

6g

RMN ^1H spectrum

RMN ^{13}C spectrum

6h

RMN ^1H spectrum

RMN ^{13}C spectrum

REFERENCES.

- (1) Mukherjee, C.; Kamila, S.; De, A. *Synlett* **2003**, 1474-1478.
- (2) Chauder, B. A.; Kalinin, A. V.; Snieckus, V. *Synthesis* **2001**, 140-144.
- (3) Kalinin, A. V.; da Silva, A. J. M.; Lopes, C. C.; Lopes, R. S. C.; Snieckus, V. *Tetrahedron Lett.* **1998**, 39, 4995-4998.
- (4) Hodgson, H. H.; Nixon, J. *J. Chem. Soc.* **1929**, 1632-1639.
- (5) Hodgson, H. H.; Kershaw, A. *J. Chem. Soc.* **1930**, 1419-1425.
- (6) Hodgson, H. H.; Nixon, J. *J. Chem. Soc.* **1934**, 137-138.
- (7) Henley, R. V.; Turner, E. E. *J. Chem. Soc.* **1930**, 928-934.
- (8) Brandsma, L.; Vasilevsky, S. F.; Verkruijsse, H. D. In *Application of Transition Metal Catalysts in Organic Synthesis*; Springer-Verlag, Berlin 1998; Ch. 11.
- (9) Witulski, B.; Azcon J. R.; Alayrac, C.; Arnautu, A.; Collot, V.; Rault, S. *Synthesis* **2005**, 771-780.