Supporting Information for

A Novel Asymmetric Vinylogous Tin-Pummerer Rearrangement

José Luis García Ruano,^{a*} José Aleman,^a and Albert Padwa^{b*}
joseluis.garcia.ruano@acm.es

a Departamento de Química Orgánica, Universidad Autónoma de Madrid, Cantoblanco,
28049-Madrid, Spain

b Department of Chemistry, Emory University, Atlanta, Georgia 30322
chemap@emory.edu

Experimental section.

General Methods.

¹H NMR spectra were acquired at 300 or 400 MHz and ¹³C NMR were acquired at 75 or 100 MHz (unless otherwise indicated). Chemical shifts (δ) are reported in ppm relative to CDCl₃ (7.26 and 77.0 ppm). Mass spectra (MS) were determined at ionizing voltage of 70 eV. All reactions were carried out in anhydrous solvents and under an argon atmosphere. THF and Et₂O were distilled from sodium-benzophenone under argon while CH₂Cl₂ was distilled from P₂O₅. Flash column chromatography was performed using silica gel Merk-60 (230-400 mesh). *n*-BuLi (2.5 M solution in hexanes) were purchased from Aldrich.

(S)-1-Methyl-2-(*p*-tolylsulfinyl)benzene (11).¹ White solid; mp: 76-78 °C; [α]_D²⁰ = -91.0 (*c* 2, acetone); ¹H NMR (300 MHz) δ 7.94 (dd, *J* = 9.3, 1.6 Hz, 1H), 7.40-7.30 (m, 4H), 7.20 (d, *J* = 8.5 Hz, 2H), 7.12 (d, *J* = 7.6 Hz, 1H), 2.32 (s, 3H), and 2.28 (s, 3H).

(S)-1-Ethyl-2-(*p*-tolylsulfinyl)benzene (12).¹ White solid; mp: 42-43 °C; [α]_D²⁰ -106.4 (*c* 1, CHCl₃); ¹H NMR (300 MHz) δ 7.94 (m, 1H), 7.52-7.38 (m, 4H), 7.24 (m, 3H), 2.75 (m, 2H), 2.35 (s, 3H), and 1.14 (t, *J* = 7.5 Hz, 3H).

(S₈)-Tributyl-[2-(*p*-tolylsulfinyl)benzyl]stannane (7). A solution of *n*-BuLi (0.60 mmol, 2.3 M in hexane) was added to ⁱPr₂NH (0.89 mmol) in THF (3 mL) at 0 °C. After stirring for 30 min, the mixture was cooled to -78 °C. A solution of sulfoxide **11** (0.50 mmol) in THF (2 mL) was added. After 1 h stirring, *n*-Bu₃SnCl (1mmol) was added at -78 °C. When the reaction was complete (15 min), the mixture was hydrolyzed (saturated

¹ Sulfoxides **11** and **12** were prepared following described procedures; see a) García Ruano, J. L.; Aranda, M.; Carreño, M. C.; Toledo, M. A. *Angew. Chem. Int. Ed. Engl.* **2000**, 39, 2376. b) García Ruano, J. L.; Alemán J.; Soriano, J. F. *Org. Lett.*, **2003**, 5, 677.

NH₄Cl), extracted (3x10 mL Et₂O), washed (2x10 mL NaCl sat.), dried (MgSO₄) and the solvent was evaporated. The residue was purified by flash-column chromatography: *n*-hexane-AcOEt 15:1; yield: 92%; colorless oil; $[\alpha]_D^{20} = -269.3$ (*c* 1.0, CHCl₃); IR (NaCl): 2923, 1588, 1492, 1464, 1084, and 1034 cm⁻¹; ¹H NMR (400 MHz) δ 7.84 (dd, *J* = 6.8, 1.2 Hz, 1H), 7.46 (d, *J* = 8.4 Hz, 2H), 7.30-7.10 (m, 4H), 6.95 (d, *J* = 6.8 Hz, 1H), 2.35 (s, 3H), 2.29 (d, *J* = 12.0 Hz, 1H), 2.20 (d, *J* = 12.0 Hz, 1H), 1.44-1.30 (m, 6H), 1.29-1.10 (m, 12H), and 0.84 (m, 9H); ¹³C NMR (100 MHz) δ 142.4, 141.5, 141.4, 138.9, 130.7, 129.9, 128.6, 126.2, 124.6, 124.0, 28.8, 27.7, 27.2, 13.6, 13.5, and 10.1; Anal. Calcd for C₂₆H₄₀OSSn: C, 60.13; H, 7.76; Found: C, 59.97; H, 7.85.

(*S,S,S*)-Tributyl-{1-[2-(*p*-tolylsulfinyl)phenyl]ethyl}stannane (10). A solution of *n*-BuLi (0.60 mmol, 2.3 M in hexane) was added to ^{*i*}Pr₂NH (0.89 mmol) in THF (3 mL) at 0 °C. After stirring for 30 min, the mixture was cooled to -78 °C. A solution of sulfoxide **12** (0.50 mmol) in THF (2 mL) was then added. After stirring for 1 h, Bu₃SnCl (1.50 mmol) was added at -78 °C. When the reaction was complete (15 min), the mixture was hydrolyzed (saturated NH₄Cl), extracted (3x10 mL Et₂O), washed (2x10 mL NaCl sat.), dried (MgSO₄) and the solvent was evaporated. The residue was purified by flash-column chromatography: *n*-hexane-AcOEt 25:1; yield: 70% (82% combined isolated yield with minor diastereoisomer); colorless oil; IR (NaCl): 2956, 2925, 1641, and 1464 cm⁻¹; $[\alpha]_D^{20} = -258.8$ (*c* 1.8, CHCl₃); ¹H NMR (300 MHz) δ 7.94 (dd, *J* = 7.8, 1.5 Hz, 1H), 7.45-7.41 (m, 2H), 7.36 (dt, *J* = 7.4, 1.4 Hz, 1H), 7.24-7.20 (m, 3H), 7.10 (dd, *J* = 7.8, 1.2 Hz, 1H), 2.70 (q, *J* = 7.5 Hz, 1H), 2.34 (s, 3H), 1.45-1.19 (m, 18H), 0.92 (d, *J* = 7.5 Hz, 3H), and 0.85 (t, *J* = 7.3 Hz, 9H); ¹³C NMR (75 MHz) δ 146.9, 142.3, 141.6, 139.1, 130.9, 130.0, 126.2, 125.7, 124.4, 123.9, 29.0, 27.3, 23.1, 21.4, 16.9, 13.7, and 9.4; MS

(EI): m/z 534 (M^+ , 5), 477 (30), 235 (63), 225 (46), 179 (100), 177 (87), and 91 (48); HMRS (EI): $C_{27}H_{42}OSSn$: 534.1978. Found: 534.1963; Anal. Calcd for $C_{27}H_{42}OSSn$: C, 60.80; H, 7.94; S 6.01; Found: C, 60.24; H, 7.44; S, 5.79.

Protocol for Pummerer reaction (Table 1 and 2).

To a stirring solution of the appropriate sulfoxide (0.20 mmol) in 2 mL of CH_2Cl_2 at rt (except entries 3 and 5 of Table 2 which were done in refluxing CH_2Cl_2), 0.30 mmol of the corresponding electrophile was added. The reaction was followed by TLC and when completed, the solvent was removed under reduced pressure and the resulting crude mixture was chromatographed (solvent is indicated in each case).

Protocol for Pummerer reaction (Solvent free conditions).

To the appropriate sulfoxide (0.20 mmol) at rt was added acetyl chloride or trifluoroacetic anhydride (0.25 mmol). The reaction was followed by TLC, and when completed, the resulting crude mixture was chromatographed (solvent is indicated in each case).

(2-*p*-Tolylsulfanyl)benzyl acetate (9). Electrophile: acetyl chloride; Chromatography: *n*-hexane-AcOEt: 50:1, yield: 73%; colorless oil; IR (NaCl): 2853, 1742, 1491, 1463, and 1225 cm^{-1} ; 1H NMR (400 MHz) δ 7.44–7.38 (m, 1H), 7.30–7.22 (m, 3H), 7.17 (d, $J = 8.0$ Hz, 2H), 7.09 (d, $J = 8.0$ Hz, 2H), 5.25 (s, 2H), 2.31 (s, 3H), and 2.05 (s, 3H); ^{13}C NMR (100 MHz) δ 170.7, 137.2, 136.3, 135.8, 132.4, 131.8, 131.2, 131.0, 129.4, 129.0, 127.4, 64.6, 21.0, and 20.8; Anal. Calcd for $C_{16}H_{16}O_2S$: C, 70.56; H, 5.67; Found: C, 70.90; H, 5.75.

(2-*p*-Tolylsulfanyl)benzyl acrylate (13). Electrophile: acryloyl chloride; Chromatography: *n*-hexane-AcOEt 50:1, yield: 72%; colorless oil; IR (NaCl): 2920,

1724, 1634, 1491, 1405, and 1177 cm^{-1} ; ^1H NMR (400 MHz) δ 7.43–7.39 (m, 2H), 7.30–7.22 (m, 3H), 7.17 (d, J = 8.4 Hz, 2H), 7.09 (d, J = 8.4 Hz, 2H), 6.42 (dd, J = 17.6, 1.6 Hz, 1H), 6.13 (dd, J = 17.6, 10.4 Hz, 1H), 5.80 (dd, J = 10.4, 1.6 Hz, 1H), 5.34 (s, 2H), and 2.31 (s, 3H); ^{13}C NMR (100 MHz) δ 165.9, 137.2, 136.3, 135.7, 132.4, 131.5, 131.2, 131.1, 130.0, 129.3, 129.0, 128.2, 127.4, 64.6, and 21.0; Anal. Calcd for $\text{C}_{17}\text{H}_{16}\text{O}_2\text{S}$; C, 69.80; H, 5.67; Found: C, 69.49; H, 5.70

(2-*p*-Tolylsulfanyl)benzyl (2-Phenyl)acetate (14). Electrophile: (2-phenyl)acetyl chloride; Chromatography: *n*-hexane-AcOEt 60:1, yield: 78%; colorless oil: IR (NaCl): 2955, 1737, 1492, 1454, 1250, and 1145 cm^{-1} ; ^1H NMR (400 MHz) δ 7.35–7.20 (m, 9H), 7.16 (d, J = 8.0 Hz, 2H), 7.09 (d, J = 8.0 Hz, 2H), 5.27 (s, 2H), 3.62 (s, 2H), and 2.31 (s, 3H); ^{13}C NMR (100 MHz) δ 171.2, 137.2, 136.3, 135.5, 133.9, 132.5, 131.6, 131.1, 130.0, 129.3, 129.2, 128.9, 128.5, 127.4, 127.1, 64.9, 41.2, and 21.1; Anal. Calcd for $\text{C}_{22}\text{H}_{20}\text{O}_2\text{S}$; C, 75.83; H, 5.79; Found: C, 75.69; H, 5.91.

(2-*p*-Tolylsulfanyl)benzyl Benzoate (15). Electrophile: benzoyl chloride; Chromatography: *n*-hexane-AcOEt 100:1; yield: 67%; colorless oil: IR (NaCl): 1717, 1603, 1491, 1450, 1269 cm^{-1} ; ^1H NMR (400 MHz) δ 8.03 (d, J = 8.4 Hz, 1H), 7.56–7.50 (m, 2H), 7.41 (t, J = 8.0 Hz, 2H), 7.31–7.21 (m, 4H), 7.20 (d, J = 8.0 Hz, 2H), 7.08 (d, J = 8.0 Hz, 2H), 5.51 (s, 2H), and 2.29 (s, 3H); ^{13}C NMR (100 MHz) δ 166.3, 137.2, 136.5, 135.7, 132.9, 132.6, 131.7, 131.1, 130.0 (2C), 129.7 (2C), 129.5, 129.0, 128.3, 127.5, 65.0, and 21.0; Anal. Calcd for $\text{C}_{21}\text{H}_{18}\text{O}_2\text{S}$; C, 75.42; H, 5.42; Found: C, 75.18; H, 5.40.

(2-*p*-Tolylsulfanyl)benzyl Trifluoroacetate (16). Electrophile: trifluoroacetyl anhydride; Chromatography: *n*-hexane-AcOEt 50:1, yield: 76%; colorless oil; IR (NaCl): 2925, 1786, 1593, 1492, and 1148 cm^{-1} ; ^1H NMR (300 MHz) δ 7.43 (m, 1H),

7.36–7.28 (m, 3H), 7.19 (d, $J = 8.2$ Hz, 2H), 7.12 (d, $J = 8.2$ Hz, 2H), 5.52 (s, 2H), and 2.33 (s, 3H); ^{13}C NMR (75 MHz) δ 157.5 (q, $J_{\text{C-F}} = 41.8$ Hz), 137.6, 136.3, 133.8, 133.8, 131.6, 131.3, 130.2, 129.9, 120.7, 120.6, and 115.0 (q, $J_{\text{C-F}} = 284.2$ Hz), 67.7, 21.0; Anal. Calcd for $\text{C}_{16}\text{H}_{13}\text{F}_3\text{O}_2\text{S}$; C, 58.89; H, 4.02; S, 9.83; Found: C, 59.01; H, 4.15; S, 10.01.

(2-*p*-Tolylsulfanyl)benzyl chloride (17). Electrophile: thionyl chloride; Chromatography: *n*-hexane-AcOEt 80:1, yield: 85%; colorless oil; IR (NaCl): 2917, 1582, 1490, 1441, and 1405 cm^{-1} ; ^1H NMR (400 MHz) δ 7.46 (d, $J = 6.8$ Hz, 1H), 7.29–7.19 (m, 4H), 7.12 (d, $J = 8.0$ Hz, 2H), 4.79 (s, 3H), and 2.33 (s, 3H); ^{13}C NMR (100 MHz) δ 137.5, 137.4, 136.2, 132.2, 131.6, 131.2, 130.2, 130.1, 129.3, 127.4, 44.5, and 21.1; Anal. Calcd for $\text{C}_{14}\text{H}_{13}\text{ClS}$; C, 67.59; H, 5.27; Found: 67.48; H, 5.33.

(2-*p*-Tolylsulfanyl)benzyl bromide (18). Electrophile: thionyl bromide; Chromatography: *n*-hexane-AcOEt 70:1, yield: 81%; colorless oil; IR (NaCl): 3018, 2918, 1587, 1491, 1463, and 1218 cm^{-1} ; ^1H NMR (400 MHz) δ 7.42 (dd, $J = 7.6, 1.6$ Hz, 1H), 7.25 (dd, $J = 7.6, 1.6$ Hz, 2H), 7.21–7.10 (m, 5H), 4.70 (s, 3H), and 2.33 (s, 3H); ^{13}C NMR (100 MHz) δ 137.9, 137.7, 137.0, 132.2 (2C), 131.2, 130.9, 130.4, 129.6, 127.5, 32.1, and 21.3; Anal. Calcd for $\text{C}_{14}\text{H}_{13}\text{BrS}$; C, 57.35; H, 4.47; Found: C, 57.69; H, 4.56.

(1*R*)-1-{2-[*p*-Tolylsulfanyl]phenyl}ethyl acetate (23). Electrophile: acetyl chloride; Chromatography: *n*-hexane-AcOEt 50:1; yield: 73%; colorless oil; $[\alpha]_{\text{D}}^{20} = +44.0$ (c 1.1, CHCl_3); IR (NaCl): 2926, 1733, 1492, 1438, 1370, and 1237 cm^{-1} ; ^1H NMR (300 MHz) δ 7.45 (d, $J = 7.8$ Hz, 1H), 7.30–7.05 (m, 7H), 6.37 (q, $J = 6.6$ Hz, 1H), 2.30 (s, 3H), 2.04 (s, 3H), and 1.48 (s, 3H); ^{13}C NMR (75 MHz) δ 169.9, 142.5, 137.0, 133.8, 132.6, 131.9, 131.3, 129.9, 128.2, 127.6, 125.8, 69.8, 21.5, 21.2, and 21.0; MS (EI): m/z 286 (M^+ , 87), 227 (100), 225 (31), 211 (48), 151 (100), and 91 (31); HMRS (EI): $\text{C}_{17}\text{H}_{18}\text{O}_2\text{S}$: 286.1027.

Found 286.1016; Anal. Calcd for C₁₇H₁₈O₂S: C, 71.30; H, 6.34; Found: C, 71.45; H, 6.66.

(1R)-1-[2-(*p*-Tolyl sulfanyl)phenyl]ethyl acrylate (24). Electrophile: acryloyl chloride; Chromatography: *n*-hexane-AcOEt 50:1; yield: 80%; colorless oil; $[\alpha]_D^{20} = +30.2$ (*c* 0.55, CHCl₃); IR (NaCl): 2855, 1725, 1637, 1620, 1492, 1469, and 1404 cm⁻¹; ¹H NMR (300 MHz) δ 7.47 (d, *J* = 7.6 Hz, 1H), 7.29-7.08 (m, 7H), 6.47 (q, *J* = 6.5 Hz, 1H), 6.43 (dd, *J* = 17.4, 1.6 Hz, 1H), 6.16 (dd, *J* = 17.4, 10.4 Hz, 1H), 5.83 (dd, *J* = 10.4, 1.6 Hz, 1H), 2.32 (s, 3H), and 1.54 (d, *J* = 6.5 Hz, 3H); ¹³C NMR (75 MHz) δ 165.2, 142.6, 137.1, 133.9, 132.6, 131.9, 131.4, 130.7, 130.0, 128.7, 128.3, 127.6, 125.8, 70.1, 29.7, and 21.7; MS (EI) *m/z* 298 (M⁺, 56), 227 (100), 135 (17), and 91 (18); Anal. Calcd for C₁₈H₁₈O₂S: C, 72.45; H, 6.08; S 10.75; Found: C, 72.31; H, 6.07; S, 10.80.

(1R)-1-[2-(*p*-Tolyl sulfanyl)phenyl]ethyl (2-phenyl)acetate (25). Electrophile: (2-phenyl)acetyl chloride; Chromatography: *n*-hexane-AcOEt 60:1; yield: 70%; colorless oil; $[\alpha]_D^{20} = -34.0$ (*c* 0.89, CHCl₃); IR(NaCl): 2980, 2927, 1718, 1493, 1454, and 1177 cm⁻¹; ¹H NMR (300 MHz) δ 7.40-7.10 (m, 13H), 6.42 (q, *J* = 6.9 Hz, 1H), 3.63 (s, 2H), 2.32 (s, 3H), and 1.52 (d, *J* = 6.9 Hz, 3H); ¹³C NMR (75 MHz) δ 165.1, 142.5, 137.1, 133.8, 132.5, 131.9, 131.4, 130.7, 130.7, 129.9 (2C), 128.6, 128.2, 127.6, 125.8, 70.1, 29.6, 21.7, and 21.0; MS (EI) *m/z* 348 (M⁺, 73), 211 (67), 151 (99), 135 (66), 105 (100), and 77 (87); HMRS (EI): C₂₂H₂₀O₂S: 348.1184. Found 348.1185.

(1R)-1-[2-(*p*-Tolyl sulfanyl)phenyl]ethyl benzoate (26). Electrophile: benzoyl chloride; Chromatography: *n*-hexane-AcOEt 60:1; yield: 75%; colorless oil; $[\alpha]_D^{20} = -13.6$ (*c* 0.70, CHCl₃); IR (NaCl): 2980, 1731, 1493, and 1193 cm⁻¹; ¹H NMR (400 MHz) δ 8.08 (dd, *J* = 8.4, 1.2 Hz, 2H), 7.54 (t, *J* = 9.2 Hz, 2H), 7.43 (t, *J* = 7.6 Hz, 2H), 7.30-7.15 (m, 7H),

7.08 (d, $J = 8.0$ Hz, 2H), 6.61 (q, $J = 6.4$ Hz, 1H), 2.29 (s, 3H), and 1.61 (d, $J = 6.4$ Hz, 3H); ^{13}C NMR (100 MHz) δ 165.4, 142.7, 137.1, 133.9, 132.8, 132.8, 132.6, 131.9, 131.4, 130.4, 130.0, 129.6, 128.3, 127.6, 125.8, 70.6, 21.8, and 21.0; MS (EI) m/z 362 (M^+ , 61), 225 (51), 151 (99), 135 (40), and 91 (100); HMRS (EI): $\text{C}_{23}\text{H}_{22}\text{O}_2\text{S}$: 362.1340. Found: 362.1345.

(1*R*)-1-[2-(Tolyl sulfanyl) phenyl] ethyl trifluoroacetate (27). Electrophile: trifluoroacetyl anhydride; Chromatography: *n*-hexane-AcOEt 50:1; yield: 63%; colorless oil; IR (NaCl): 2927, 1785, 1492, 1220, and 1159 cm^{-1} ; ^1H NMR (300 MHz) δ 7.52 (dd, 1H, $J = 7.7, 1.5$ Hz), 7.42-7.20 (m, 5H), 7.16 (d, $J = 8.0$ Hz, 1H), 6.60 (q, $J = 6.5$ Hz, 1H), 2.37 (s, 3H), and 1.65 (d, $J = 6.5$ Hz, 3H); ^{13}C NMR (75 MHz) δ 156.4 (q, $J_{\text{C-F}} = 41.9$ Hz), 140.3, 137.4, 134.0, 133.1, 131.5, 131.2, 130.1, 129.1, 128.1, 125.7, 114.5 (q, $J_{\text{C-F}} = 284.3$ Hz), 74.4, 21.3, and 21.0; Anal. Calcd for $\text{C}_{17}\text{H}_{15}\text{O}_2\text{S}$: C, 59.99; H, 4.44; S, 9.42; Found C, 60.00; H, 4.63; S 9.31.

1-[2-(*p*-Tolyl-sulfanyl)-phenyl]-1-ethylchloride (28). Electrophile: thionyl chloride; Chromatography: *n*-hexane-AcOEt 80:1; yield: 70%; colorless oil; IR (NaCl): 1491, 1467, 1437, and 1043 cm^{-1} ; ^1H NMR (400 MHz) δ 7.66 (d, $J = 8.0$ Hz, 1H), 7.32 (td, $J = 7.2, 1.2$ Hz, 1H), 7.27-7.14 (m, 4H), 7.10 (d, $J = 8.8$ Hz, 2H), 5.78 (q, $J = 6.8$ Hz, 1H), 2.31 (s, 3H), and 1.77 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (100 MHz) δ 143.5, 137.1, 133.6, 132.8, 130.8, 130.1 (2C), 128.7, 128.1, 127.2, 55.3, 26.2, and 21.1; MS (EI) m/z 262 (M^+ , 9), 226 (61), 211 (46), 170 (49), and 135 (60); Anal. Calcd for $\text{C}_{15}\text{H}_{15}\text{ClS}$: C, 68.55; H, 5.75; Found C, 68.78; H, 5.41; *ee* = 2 HPLC: Column Daicel Chiralpak AD; Hexane/*i*-PrOH 100:0; flow rate 0.1mL/min; t_R = 35.0 major isomer, t_R = 44.5 minor isomer)

1-(2-(*p*-Tolyl-sulfanyl)-phenyl)-1-ethylbromide (29). Electrophile: thionyl bromide; Chromatography: *n*-hexane-AcOEt 100:1; yield: 71%; colorless oil; IR (NaCl): 2920, 1587, 1490, 1435, and 809 cm⁻¹; ¹H RMN (400 MHz) 7.67 (dd, *J* = 7.6, 0.8 Hz, 1H), 7.30 (td, *J* = 6.8, 1.6 Hz, 1H), 7.25-7.14 (m, 4H), 7.11 (d, *J* = 8.0 Hz, 2H), 5.88 (q, *J* = 6.8 Hz, 1H), 2.33 (s, 3H), and 2.01 (d, *J* = 6.8 Hz, 3H); ¹³C NMR (400 MHz) δ 143.5, 137.3, 134.1, 132.7, 131.7, 131.2, 130.1, 128.8, 128.0, 127.5, 46.0, 26.4, and 21.1; Anal. Calcd for C₁₅H₁₅BrS: C, 58.64; H, 4.92; S, 10.44; Found C, 58.40; H, 5.05; S 10.42; *ee* = 1 HPLC: Column Daicel Chiralpak AD; Hexane/*i*-PrOH 100:0 flow rate 0.1mL/min; *t_R* = 44.0 major isomer, *t_R* = 54.4 minor isomer).

***Protocol for Mosher esters.*²**

To a solution of the corresponding ester (0.010 mmol) in 2 mL THF under argon atmosphere was added slowly 0.015 mmol of LiAlH₄. When the reaction was complete, it was quenched with 1 mL of sat. NH₄Cl and the mixture was extracted with Et₂O (2x2 mL), washed (2x2 NaCl sat.), dried with MgSO₄ anhydrous, the solvent was evaporated and the mixture was chromatographed using 2:1 Hexane/AcOEt.

(1*R*)-1-[2-(*p*-Tolylsulfanyl)phenyl]ethan-1-ol (34). Chromatography: *n*-hexane-AcOEt 2:1; yield: 82%; colorless oil; [α]_D²⁰ = -60 (c=0.76, CHCl₃); IR (NaCl): 3385 (bs), 2970, 2924, 2855, 1590, 1491, 1090, and 1016 cm⁻¹; ¹H NMR (300 MHz) δ 7.58 (dd, *J* = 7.3 Hz, 0.8 Hz, 1H), 7.29 (dt, *J* = 6.5, 2.4 Hz, 1H), 7.21-7.07 (m, 6H), 5.39 (dq, *J* = 6.5, 3.6 Hz, 1H), 2.30 (s, 3H), 2.03 (d, *J* = 3.6 Hz, 1H), and 1.45 (d, *J* = 6.5 Hz, 3H); ¹³C NMR (75 MHz) δ 146.1, 137.1, 132.8, 132.5, 132.0, 130.8, 130.1, 128.0, 127.8, 125.8, 67.3, 24.0, and 21.0; MS (FAB) *m/z* 245 (M+1, 21), 154 (100), 136 (71), and 107 (28).

² Ohtani, I.; Kusumi, T.; Kashman, Y.; Kakisawa, H. *J. Am. Chem. Soc.* **1991**, *113*, 4092.

MTPA esters of 34

To solution of compound **8** (0.05 mmol) in CH₂Cl₂ (1 mL) was added Et₃N (14 μL, 0.1 mmol), DMAP (0.01 mmol) and (*S*)-MTPA-Cl or (*R*)-MTPA-Cl (0.1 mmol) under an argon atmosphere. When the reaction was complete, the mixture was hydrolyzed (HCl 5%, 1 mL), extracted (3x1 mL, CH₂Cl₂), washed (2x2 mL NaHCO₃ sat), dried (Na₂SO₄) and the solvent was evaporated.

(1*R*)-1-[2-(*p*-Tolylsulfanyl)phenyl]ethan-1-ol O-(*R*)-MTPA ester (35a). ¹H NMR (400 MHz) δ 7.47 (d, *J* = 7.2 Hz, 1H), 7.41-7.12 (m, 7H), 7.10 (d, *J* = 8.0 Hz, 2H), 6.63 (q, *J* = 6.8 Hz, 1H), 3.55 (s, 3H), 2.31 (s, 3H), and 1.53 (d, *J* = 6.8 Hz, 3H).

(1*R*)-1-[2-(*p*-Tolylsulfanyl)phenyl]ethan-1-ol O-(*S*)-MTPA ester (35b). ¹H NMR (400 MHz) δ 7.48 (d, *J* = 8.0 Hz, 1H), 7.42-7.16 (m, 7H), 7.10 (d, *J* = 8.0 Hz, 2H), 6.68 (q, *J* = 6.8 Hz, 1H), 3.50 (s, 3H), 2.32 (s, 3H), and 1.47 (d, *J* = 6.8 Hz, 3H).