Supporting information for:

**A Novel Class of P-O Monophosphite Ligands Derived from
D-Mannitol: Broad Applications in Highly Enantioselective
Rh-Catalyzed Hydrogenations**

Hanmin Huang, Zhuo Zheng^{*}, Huili Luo, Changmin Bai, Xinquan Hu, and Huilin Chen

Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian, 116023, P. R. China

CONTENTS

(A) GC Analytical Data and graphic of Products (S1-S21)

- (1) GC analytical data and graphic of hydrogenation products of dehydroamino acid esters (S2-S7).**
- (2) GC analytical data and graphic of hydrogenation products of enamides (S8-S16).**
- (3) GC analytical data and graphic of hydrogenation products of β -(acylamino)acrylates (S17-S20).**
- (4) GC analytical data and graphic of hydrogenation products of dimethyl itaconate (S20-S21).**

(B) NMR of ligands 3a-k (S22-S54)

(A) Determination of enantiomeric excesses of hydrogenation products

Gas chromatograms to determine enantiomeric excesses for α -amino acid

esters

97.4% ee (S)

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		13.850	49212.7	924453.3	49.9843
2	2		15.292	49317.8	925032.5	50.0157
Total				98530.5	1849485.8	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		13.798	429.6	5782.6	1.3096
2	2		15.173	29040.5	435776.9	98.6904
Total				29470.1	441559.5	100.0000

The ee's were determined by GC analysis using a Chrompack chiral fused silica 25m x 0.25mm Chirasil-L-Val column. Oven temperature is 160 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.²

96.6% ee (S)

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		13.383	31004.8	414951.4	50.0374
2	2		14.883	29685.8	414330.9	49.9626
Total				60690.6	829282.3	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		13.215	1615.7	23539.5	1.6975
2	2		14.873	55865.8	1363169.1	98.3025
Total				57481.5	1386708.6	100.0000

The ee's were determined by GC analysis using a Chrompack chiral fused silica 25m x 0.25mm chirasil-L-Val column. Oven temperature is 160 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.²

97.1% ee (S)

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		15.542	45915.9	978946.3	49.9581
2	2		17.517	44190.3	980589.6	50.0419
Total				90106.2	1959535.9	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		15.333	767.1	10465.5	1.4205
2	2		17.450	37460.1	726267.9	98.5795
Total				38227.3	736733.4	100.0000

The ee's were determined by GC analysis using a Chrompack chiral fused silica 25m x 0.25mm chirasil-L-Val column. Oven temperature is 160 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.²

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		18.942	37876.9	1091534.7	49.9581
2	2		21.092	37863.0	1093367.5	50.0419
Total				75739.9	2184902.2	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		18.717	456.4	7363.9	1.0011
2	2		20.967	30081.1	728182.3	98.9989
Total				30537.5	735546.2	100.0000

The ee's were determined by GC analysis using a Chrompack chiral fused silica 25m x 0.25mm chirasil-L-Val column. Oven temperature is 160 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.²

96.8% ee (S)

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		6.625	127435.5	1216486.3	50.0248
2	2		7.292	134030.2	1215279.1	49.9752
Total				261465.7	2431765.4	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		6.617	2883.3	18551.9	1.6037
2	2		7.358	133246.6	1138232.9	98.3963
Total				136130.0	1156784.7	100.0000

The ee's were determined by GC analysis using a Chrompack chiral fused silica 25m x 0.25mm chirasil-L-Val column. Oven temperature is 160 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.²

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		8.148	25847.6	299481.0	49.9823
2	2		8.882	23963.6	299693.4	50.0177
Total				49811.2	599174.4	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		8.083	24639.1	283613.5	98.8594
2	2		8.825	321.0	3272.2	1.1406
Total				24960.1	286885.8	100.0000

The ee's were determined by GC analysis using a Suplco Chiral select 1000 column, 30m x 0.25mm. Oven temperature is 110 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.²

Gas chromatograms to determine enantiomeric excesses for hydrogenation

products of enamides

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		9.708	121147.8	1531439.8	50.0049
2	2		10.417	125660.2	1531136.8	49.9951
Total				246808.0	3062576.6	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		9.700	171136.1	2216830.3	99.9668
2	2		10.310	93.8	735.2	0.0332
Total				171230.0	2217565.5	100.0000

The ee's were determined by GC analysis using a Supelco Chiral select 1000 column, 30m x 0.25mm. Oven temperature is 150 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.³

99.8% (S)

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		16.750	51782.8	1088866.1	49.8791
2	2		18.325	50357.4	1094144.6	50.1209
Total				102140.2	2183010.7	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		16.625	31308.8	612241.8	99.9229
2	2		18.125	38.0	472.5	0.0771
Total				31346.8	612714.3	100.0000

The ee's were determined by GC analysis using a Supelco Chiral select 1000 column, 30m x 0.25mm. Oven temperature is 130 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.³

96.0% ee (S)

No.	PeakNo	ID.Name	R.Time	PeakHeight	PeakArea	Conc
1	1		21.192	18702.0	404951.2	50.0749
2	2		22.333	17778.9	403739.6	49.9251
Total				36480.9	808690.8	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		21.158	14736.4	324997.0	97.9945
2	2		22.275	331.2	6651.3	2.0055
Total				15067.6	331648.3	100.0000

The ee's were determined by GC analysis using a Supelco Chiral select 1000 column, 30m x 0.25mm. Oven temperature is 150 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.^{1c}

99.9% ee (S)

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		12.908	73039.0	1162483.5	50.0218
2	2		13.808	74755.0	1161469.0	49.9782
Total				147793.9	2323952.5	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		12.900	79022.4	1241855.7	99.9464
2	2		13.692	76.2	666.1	0.0536
Total				79098.7	1242521.8	100.0000

The ee's were determined by GC analysis using a Supelco Chiral select 1000 column, 30m x 0.25mm. Oven temperature is 140 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.³

99.4% ee (S)

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		45.408	26139.1	1595895.3	50.0633
2	2		47.850	28605.1	1591859.5	49.9367
Total				54744.1	3187754.7	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		45.058	27676.5	1600975.0	99.7256
2	2		46.918	99.6	4404.5	0.2744
Total				27776.1	1605379.5	100.0000

The ee's were determined by GC analysis using a Supelco Chiral select 1000 column, 30m x 0.25mm. Oven temperature is 140 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.³

99.2% ee (S)

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		20.692	37405.7	1109405.7	49.9869
2	2		22.292	37970.5	1109988.8	50.0131
Total				75376.2	2219394.5	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		20.917	61900.0	2064421.4	99.6108
2	2		22.033	355.9	8065.1	0.3892
Total				62255.9	2072486.5	100.0000

The ee's were determined by GC analysis using a Supelco Chiral select 1000 column, 30m x 0.25mm. Oven temperature is 130 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.³

99.5% ee (S)

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		22.477	34670.7	1557429.8	49.9596
2	2		24.025	47787.9	1559950.7	50.0404
Total				82458.6	3117380.5	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		22.392	238.1	6062.4	0.2554
2	2		24.008	55470.0	2367358.8	99.7446
Total				55708.1	2373421.2	100.0000

The ee's were determined by GC analysis using a Chrompack chiral fused silica 25m x 0.25mm chirasil-L-Val column. Oven temperature is 170 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.³

99.7% ee (S)

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		24.242	34615.9	999145.1	50.0058
2	2		25.475	37626.3	998913.5	49.9942
Total				72242.3	1998058.5	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		24.808	105928.2	4243092.4	99.8492
2	2		25.508	363.8	6406.9	0.1508
Total				106292.0	4249499.3	100.0000

The ee's were determined by GC analysis using a Supleco Chiral select 1000 column, 30m x 0.25mm. Oven temperature is 150 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.³

99.9% ee (S)

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		43.150	15099.1	702170.3	50.0181
2	2		45.333	15299.0	701661.5	49.9819
Total				30398.2	1403831.8	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		43.258	22639.2	1067953.0	99.9712
2	2		45.147	12.5	307.8	0.0288
Total				22651.6	1068260.8	100.0000

The ee's were determined by GC analysis using a Supelco Chiral select 1000 column, 30m x 0.25mm. Oven temperature is 150 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.³

Gas chromatograms to determine enantiomeric excesses for hydrogenation

products of β -(acylamino)acrylates

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		9.517	45580.0	741115.5	49.9963
2	2		10.425	43138.3	741224.2	50.0037
Total				88718.3	1482339.7	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		9.433	37557.8	618948.0	99.9182
2	2		10.355	55.6	506.9	0.0818
Total				37613.4	619454.9	100.0000

The ee's were determined by GC analysis using a Supelco Chiral select 1000 column, 30m x 0.25mm. Oven temperature is 110 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.^{1d}

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		6.375	79704.9	719528.9	50.0069
2	2		7.292	72345.8	719330.7	49.9931
Total				152050.8	1438859.6	100.0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		6.317	65159.8	613897.2	99.3672
2	2		7.250	400.5	3909.8	0.6328
Total				65560.3	617807.0	100.0000

The ee's were determined by GC analysis using a Supelco Chiral select 1000 column, 30m x 0.25mm. Oven temperature is 130 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.^{1d}

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		4. 692	34963. 0	233310. 9	49. 9584
2	2		5. 342	30921. 4	233699. 7	50. 0416
Total				65884. 4	467010. 6	100. 0000

No.	PeakNo	ID. Name	R. Time	PeakHeight	PeakArea	Conc
1	1		4. 658	62379. 9	439628. 8	99. 5825
2	2		5. 308	259. 5	1843. 2	0. 4175
Total				62639. 3	441472. 0	100. 0000

The ee's were determined by GC analysis using a Supelco Chiral select 1000 column, 30m x 0.25mm. Oven temperature is 130 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.^{1d}

The ee's were determined by GC analysis using a Supelco γ -DEX 225 column, 30m x 0.25mm. Oven temperature is 125 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.^{1d}

Gas chromatograms to determine enantiomeric excesses for hydrogenation products of dimethyl itaconate

The ee's were determined by GC analysis using a Supelco γ -DEX 225 column, 30m x 0.25mm. Oven temperature is 80 °C. Absolute configuration was determined by comparison of the sign of the optical rotation with reported value.⁴

References

- (1) (a) Herbst, R. M.; Shemin, D. *Organic Syntheses, Collect. Vol. II*, Wiley, New York, N. Y., **1943**, p 1; (b) Burk, M. J.; Casy, G.; Johnson, N. B. *J. Org. Chem.* **1998**, *63*, 6084; (c) Zhang, Z.; Zhu, G.; Jiang, Q.; Xiao, D.; Zhang, X. *J. Org. Chem.* **1999**, *64*, 1774; (d) Zhu, G.; Chen, Z.; Zhang, X. *J. Org. Chem.* **1999**, *64*, 6907.
- (2) Burk, M. J.; Feaster, J. E.; Nugent, W. A.; Harlow, R. L. *J. Am. Chem. Soc.* **1993**, *115*, 10125.
- (3) (a) Burk, M. J.; Wang, Y. M.; Lee, J. R. *J. Am. Chem. Soc.* **1996**, *118*, 5142; (b) Zhu, G.; Casalnuovo, A. L.; Zhang, X. *J. Org. Chem.* **1998**, *63*, 8100.
- (4) Berens, U.; Burk, M. J.; Gerlach, A.; Hems, W. *Angew. Chem. Int. Ed. Engl.* **2000**, *39*, 1981.

(B) NMR for 3a-k

3a
³¹P NMR (DMSO-d₆)

4

F2 - Acquisition Parameters

Date_ 500000
Time_ 14.12
INSTRUM DRX400
PROBHD BBI 5mm Z
PULPROG zg
TD 16384
SOLVENT Aceton
NS 16
DS 0
SWH 6009.615 Hz
FIDRES 0.366798 Hz
AQ 1.3631988 sec
RG 32
DW 83.200 usec
DE 14.29 usec
TE 300.0 K
D1 2.00000000 sec
P1 8.00 usec
DE 14.29 usec
SF01 400.1336012 MHz
NUC1 1H
P11 7.50 dB

F2 - Processing parameters
SI 16384
SF 400.1308803 MHz
WDW GM
SSB 0
LB -0.50 Hz
GB 0.2
PC 1.20

1D NMR plot parameters
CX 20.00 cm
F1P 9.000 ppm
F1 3601.18 Hz
F2P -0.200 ppm
F2 -80.03 Hz
PPMCM 0.46000 ppm/cm
HZCM 184.06021 Hz/cm

F2 - Acquisition Parameters
Date_ 50000

Date_	500000
Time_	14.34
INSTRUM	DRX400
PROBHD	BBI 5mm Z
PULPROG	zgdc
TD	16384
SOLVENT	Aceton
NS	45366

F2 - Processing parameters

SI	16384	
SF	100.6130378	MHz
WDW	EM	
SSB	0	
LB	6.00	Hz
GB	0	
PC	1.30	

1D NMR plot parameters

CX	20.00	cm
F1P	200.000	ppm
F1	20122.61	Hz
F2P	0.000	ppm
F2	0.00	Hz
PPMCM	10.00000	ppm/cm

```

F2 - Acquisition Parameters
Date_      500000
Time_      16.00
INSTRUM    DRX400
PROBHD     BBI 5mm Z
PULPROG    zgpg30
TD         16384
SOLVENT    DMSO
NS         400
DS         0
SWH         43103.449 Hz
FIDRES     2.630826 Hz
AQ         0.1901044 sec
RG         8192
DW         11.600 usec
DE         51.00 usec
TE         300.0 K
d11        0.0300000 sec
PL12       25.60 dB
CPDPRG2    waltz16
PCPD2      80.00 usec
SF02       400.1320000 MHz
NUC2       31P
PL2        7.50 dB
D1         1.00000000 sec
P1         20.00 usec
DE         51.00 usec
SF01       161.9899872 MHz
NUC1       31P
PL1        0.00 dB

F2 - Processing parameters
SI         16384
SF         161.9755552 MHz
WDW        EM
SSB        0
LB         6.00 Hz
GB         0
PC         2.00

1D NMR plot parameters
CX         20.00 cm
F1P        220.000 ppm
F1         35634.63 Hz
F2P        -20.000 ppm
F2         -3239.51 Hz
P1P1P1P1  17.000000 ppm/cm

```

¹H NMR (DMSO-d₆)

3e

³¹P NMR (DMSO-d₆)

154.862

ppm

^{13}P NMR (DMSO- d_6)