Supporting Information

A Novel Generation of Indolo-2,3-quinodimethanes

Norikazu Kuroda, Yukie Takahashi, Kazuo Yoshinaga, Chisato Mukai*

Division of Pharmaceutical Sciences, Graduate School of Natural Science and Technology,
Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan

Table of Contents

Characterization data for compounds 9a,b	S2
Characterization data for compounds 9c-f	S3
Characterization data for compounds 9g, 11a,c	S4
Characterization data for compounds 11d-f	S5
Characterization data for compounds 11g, 12f, 14a	S6
Characterization data for compounds 14d,e, 15	S7
Characterization data for compounds 16	S8
¹ H and ¹³ C NMR spectra for compound 9b	S9
¹ H and ¹³ C NMR spectra for compound 9d	S10
¹ H and ¹³ C NMR spectra for compound 9e	S11
¹ H and ¹³ C NMR spectra for compound 9g	S12
¹ H and ¹³ C NMR spectra for compound 12f	S13
¹ H and ¹³ C NMR spectra for compound 15	S14

General procedure for the preparation of allenylaniline derivatives **9a-g and **15**.**

To a solution of *N*-(*tert*-butoxycarbonyl)-2-[1-(*tert*-butyldimethylsilyl)-2,3-butadien-2-yl]aniline derivatives⁵ (0.25 mmol), prepared from the Stille coupling reaction between *N*-(*tert*-butoxycarbonyl)-2-iodoaniline derivatives and 1-(*tert*-butyldimethylsiloxy)-2-tributylstannyl)-2,3-butadiene, in THF (2.5 mL) were added AcOH (0.17 mmol) and TBAF (0.75 mmol) at 0 °C. The reaction mixture was stirred for 3 h, quenched by addition of water, and extracted with AcOEt. The extract was washed with water and brine, dried, and concentrated to dryness. The residue was chromatographed with hexane-AcOEt (4:1) to give *N*-(*tert*-butoxycarbonyl)-2-(1-hydroxy-2,3-butadien-2-yl)aniline derivatives (93%-quant.). To a solution of the alcohol derivatives (1.02 mmol) in pyridine (3.0 mL) was added Ac₂O (5.29 mmol) at 0°C. The reaction mixture was stirred for 1 h, quenched by addition of saturated aqueous NH₄Cl, and extracted with AcOEt. The extract was washed with water and brine, dried, and concentrated to dryness. The residue was chromatographed with hexane-AcOEt (4:1) to give **9b-g** and **15** (74-93%). The mesylate **9a** (73%) was prepared from the alcohol compound by treated with MsCl and Et₃N.

***N*-(*tert*-Butoxycarbonyl)-2-(1-methanesulfonyloxy-2,3-butadien-2-yl)aniline (**9a**).**

Compound **9a** was a colorless oil: IR 3429, 1973(sh), 1946, 1728 cm⁻¹; ¹H NMR δ 7.93 (1H, d, *J* = 8.3 Hz), 7.35-7.20 (2H, m), 7.10 (1H, dt, *J* = 1.3, 7.3 Hz), 6.87 (1H, brs), 5.17 (2H, t, *J* = 2.0 Hz), 5.01 (2H, t, *J* = 2.0 Hz), 2.88 (3H, s), 1.52 (9H, s); ¹³C NMR δ 209.0, 153.0, 136.2, 129.0, 128.1, 123.7, 121.8, 96.8, 80.6, 78.3, 70.4, 38.0, 28.3; MS *m/z* 238 (M⁺-101, 15). Anal. Calcd for C₁₆H₂₁NO₅S: C, 56.62; H, 6.24; N, 4.13. Found: C, 56.33; H, 6.26; N, 4.27.

2-(1-Acetoxy-2,3-butadien-2-yl)-*N*-(*tert*-butoxycarbonyl)aniline (9b**).**

Compound **9b** was a colorless oil: IR 3427, 1948, 1728 cm⁻¹; ¹H NMR δ 8.02 (1H, d, *J* = 8.1 Hz), 7.33-7.12 (3H, m), 7.07-6.99 (1H, m), 5.07 (2H, t, *J* = 2.6 Hz), 4.80 (2H, t, *J* = 2.6 Hz), 2.08 (3H, s), 1.52 (9H, s); ¹³C NMR δ 207.6, 170.5, 152.9, 136.2, 128.7, 123.3, 123.0, 120.4, 98.1, 80.4, 77.9, 64.5, 28.3, 20.8; MS *m/z* 303 (M⁺, 1.0). HRMS calcd. for C₁₇H₂₁NO₄ 303.1471, found 303.1470.

2-(1-Acetoxy-2,3-butadien-2-yl)-N-(tert-butoxycarbonyl)-3-methoxyaniline (9c).

Compound **9c** was a colorless oil: IR 3423, 1954, 1728 cm^{-1} ; ^1H NMR δ 7.70 (1H, d, $J = 8.4$ Hz), 7.24 (1H, t, $J = 8.2$ Hz), 7.02 (1H, brs), 6.59 (1H, d, $J = 8.2$ Hz), 4.98 (2H, t, $J = 2.7$ Hz), 4.75 (2H, t, $J = 2.7$ Hz), 3.81 (3H, s), 2.04 (3H, s), 1.52 (9H, s); ^{13}C NMR δ 207.0, 170.7, 157.0, 152.9, 137.7, 129.3, 112.0, 111.8, 105.1, 94.6, 80.3, 76.9, 64.3, 55.7, 28.3, 20.8; MS m/z 333 (M^+ , 1.1). Anal. Calcd for $\text{C}_{18}\text{H}_{23}\text{NO}_5$: C, 64.85; H, 6.95; N, 4.20. Found: C, 64.54; H, 7.01; N, 4.15.

2-(1-Acetoxy-2,3-butadien-2-yl)-N-(tert-butoxycarbonyl)-4-methoxyaniline (9d).

Compound **9d** was a colorless oil: IR 3429, 3354, 1950, 1724 cm^{-1} ; ^1H NMR δ 7.78 (1H, d, $J = 8.6$ Hz), 6.85-6.77 (3H, m), 5.07 (2H, t, $J = 2.6$ Hz), 4.81 (2H, t, $J = 2.6$ Hz), 3.78 (3H, s), 2.08 (3H, s), 1.51 (9H, s); ^{13}C NMR δ 207.9, 170.5, 155.7, 153.4, 129.2, 123.3, 113.8, 113.6, 98.2, 80.1, 77.9, 64.4, 55.5, 28.3, 20.9; MS m/z 333 (M^+ , 20). HRMS calcd. for $\text{C}_{18}\text{H}_{23}\text{NO}_5$ 333.1576, found 333.1571.

2-(1-Acetoxy-2,3-butadien-2-yl)-N-(tert-butoxycarbonyl)-4,5-methylenedioxyaniline (9e).

Compound **9e** was a colorless oil: IR 3427, 1950, 1724 cm^{-1} ; ^1H NMR δ 7.58-7.42 (1H, m), 6.93 (1H, brs), 6.68 (1H, s), 5.93 (2H, s), 5.04 (2H, t, $J = 2.6$ Hz), 4.74 (2H, t, $J = 2.6$ Hz), 2.08 (3H, s), 1.51 (9H, s); ^{13}C NMR δ 207.9, 170.6, 153.1, 147.6, 143.6, 130.6, 116.7, 107.8, 103.3, 101.3, 98.1, 80.3, 77.7, 64.6, 28.3, 20.9; MS m/z 347 (M^+ , 11). HRMS calcd. for $\text{C}_{18}\text{H}_{21}\text{NO}_6$ 347.1369, found 347.1372.

2-(1-Acetoxy-2,3-butadien-2-yl)-N-(tert-butoxycarbonyl)-4-(ethoxycarbonyl)aniline (9f).

Compound **9f** was a colorless solid: mp 104-106 $^{\circ}\text{C}$ (from Hexane- CH_2Cl_2); IR 3420, 1932, 1732, 1717(sh) cm^{-1} ; ^1H NMR δ 8.22 (1H, d, $J = 8.7$ Hz), 7.97-7.88 (2H, m), 7.49 (1H, brs), 5.10 (2H, t, $J = 2.7$ Hz), 4.79 (2H, t, $J = 2.7$ Hz), 4.35 (2H, q, $J = 7.1$ Hz), 2.10 (3H, s), 1.54 (9H, s), 1.38 (3H, t, $J = 7.1$ Hz); ^{13}C NMR δ 207.4, 170.5, 166.0, 152.4, 140.6, 130.4, 130.3, 124.4, 122.4, 118.5, 97.6, 81.1, 78.3,

64.3, 60.8, 28.3, 20.8, 14.7; MS m/z 375 (M^+ , 1.2). Anal. Calcd for $C_{20}H_{25}NO_6$: C, 63.99; H, 6.71; N, 3.73. Found: C, 63.96; H, 6.72; N, 3.80.

2-(1-Aceoxy-2,3-butadien-2-yl)-N-(tert-butoxycarbonyl)-4-chloroaniline (9g).

Compound **9g** was a colorless oil: IR 3425, 1948, 1726 cm^{-1} ; 1H NMR δ 8.00 (1H, d, J = 8.9 Hz), 7.25-7.18 (3H, m), 5.10 (2H, t, J = 2.6 Hz), 4.76 (2H, t, J = 2.6 Hz), 2.09 (3H, s), 1.52 (9H, s); ^{13}C NMR δ 207.6, 170.5, 152.7, 134.9, 128.5, 128.3, 127.9, 124.9, 121.5, 97.4, 80.7, 78.4, 64.1, 28.2, 20.8; MS m/z 337 (M^+ , 5.5). HRMS calcd. for $C_{17}H_{20}ClNO_4$ 337.1081, found 337.1078.

(2*R,3*R**)-2,3-Bis(methoxycarbonyl)-9-(tert-butoxycarbonyl)-1,2,3,4-tetrahydrocarbazole (11a).**

Compound **11a** was a colorless solid: mp 119-121 °C (from Hexane- CH_2Cl_2); IR 1732 cm^{-1} ; 1H NMR δ 8.09 (1H, d, J = 8.3 Hz), 7.37 (1H, d, J = 7.6 Hz), 7.29-7.20 (2H, m), 3.76 (3H, s), 3.75 (3H, s), 3.61-3.53 (1H, m), 3.23-3.09 (4H, m), 2.86-2.76 (1H, m), 1.67 (9H, s); ^{13}C NMR δ 174.6, 174.3, 150.1, 135.7, 132.4, 128.4, 123.7, 122.4, 117.4, 115.3, 114.1, 83.6, 51.9, 51.9, 42.2, 41.0, 28.0, 27.9, 13.6; MS m/z 387 (M^+ , 5.1). Anal. Calcd for $C_{21}H_{25}NO_6$: C, 65.10; H, 6.50; N, 3.62. Found: C, 64.81; H, 6.53; N, 3.64.

(2*R,3*R**)-2,3-Bis(methoxycarbonyl)-9-(tert-butoxycarbonyl)-5-methoxy-1,2,3,4-tetrahydrocarbazole (11c).**

Compound **11c** was a colorless solid: mp 138-152 °C (from Hexane-AcOEt); IR 1728 cm^{-1} ; 1H NMR δ 7.69 (1H, d, J = 8.2 Hz), 7.14 (1H, t, J = 8.2 Hz), 6.62 (1H, d, J = 8.2 Hz), 3.86 (3H, s), 3.76 (3H, s), 3.74 (3H, s), 3.58-3.40 (2H, m), 3.12-2.87 (4H, m), 1.65 (9H, s); ^{13}C NMR δ 175.2, 174.7, 153.6, 150.4, 137.3, 130.6, 124.5, 118.3, 114.3, 108.6, 103.4, 83.8, 55.2, 52.1, 52.0, 42.2, 41.7, 28.2, 26.2; MS m/z 417 (M^+ , 10). Anal. Calcd for $C_{22}H_{27}NO_7$: C, 63.30; H, 6.52; N, 3.36. Found: C, 63.11; H, 6.56; N, 3.26.

(2*R,3*R**)-2,3-Bis(methoxycarbonyl)-9-(*tert*-butoxycarbonyl)-6-methoxy-1,2,3,4-tetrahydrocarbazole (11d).**

Compound **11d** was a colorless solid: mp 128-138 °C (from Hexane-AcOEt); IR 1732 cm^{-1} ; ^1H NMR δ 7.96 (1H, d, J = 8.9 Hz), 6.88-6.82 (2H, m), 3.84 (3H, s), 3.76 (3H, s), 3.75 (3H, s), 3.59-3.51 (1H, m), 3.15-3.04 (4H, m), 2.80-2.72 (1H, m), 1.66 (9H, s); ^{13}C NMR δ 174.8, 174.5, 155.9, 150.2, 133.3, 130.5, 129.4, 116.3, 114.1, 112.1, 100.5, 83.7, 55.6, 52.1, 52.1, 42.4, 41.2, 28.2, 28.1, 23.8; MS m/z 417 (M^+ , 14). Anal. Calcd for $\text{C}_{22}\text{H}_{27}\text{NO}_7$: C, 63.30; H, 6.52; N, 3.36. Found: C, 63.08; H, 6.54; N, 3.44.

(2*R,3*R**)-2,3-Bis(methoxycarbonyl)-9-(*tert*-butoxycarbonyl)-6,7-methylenedioxy-1,2,3,4-tetrahydrocarbazole (11e).**

Compound **11e** was a colorless solid: mp 168-170 °C (from Hexane- AcOEt); IR 1733 cm^{-1} ; ^1H NMR δ 7.64 (1H, s), 6.75 (1H, s), 5.96 (2H, s), 3.76 (3H, s), 3.75 (3H, s), 3.53-3.48 (1H, m), 3.16-2.98 (4H, m), 2.76-2.67 (1H, m), 1.65 (9H, s); ^{13}C NMR δ 174.9, 174.6, 150.2, 145.7, 144.2, 131.2, 130.5, 122.7, 114.3, 101.0, 97.7, 96.8, 83.9, 52.1, 52.1, 42.4, 41.2, 28.2, 23.8; MS m/z 431 (M^+ , 19). Anal. Calcd for $\text{C}_{22}\text{H}_{25}\text{NO}_8$: C, 61.25; H, 5.84; N, 3.25. Found: C, 61.16; H, 5.90; N, 3.22.

(2*R,3*R**)-2,3-Bis(methoxycarbonyl)-9-(*tert*-butoxycarbonyl)-6-ethoxycarbonyl-1,2,3,4-tetrahydrocarbazole (11f).**

Compound **11f** was a colorless solid: mp 126-131 °C (from Hexane- AcOEt); IR 1732 cm^{-1} ; ^1H NMR δ 8.14-8.08 (2H, m), 8.00-7.92 (1H, m), 4.40 (2H, q, J = 7.1 Hz), 3.76 (3H, s), 3.76 (3H, s), 3.60-3.52 (1H, m), 3.27-3.13 (4H, m), 2.91-2.82 (1H, m), 1.68 (9H, s), 1.42 (3H, t, J = 7.1 Hz); ^{13}C NMR δ 174.6, 174.3, 166.9, 150.0, 138.7, 134.0, 128.4, 125.3, 124.9, 119.7, 115.1, 114.7, 84.5, 60.8, 52.2, 52.2, 42.2, 41.0, 28.2, 27.9, 23.5, 14.4; MS m/z 459 (M^+ , 5.0). Anal. Calcd for $\text{C}_{24}\text{H}_{29}\text{NO}_8$: C, 62.73; H, 6.36; N, 3.05. Found: C, 62.49; H, 6.43; N, 2.98.

(2*R,3*R**)-2,3-Bis(methoxycarbonyl)-9-(*tert*-butoxycarbonyl)-6-chloro-1,2,3,4-tetrahydrocarbazole (11g).**

Compound **11g** was a colorless solid: mp 136-138 °C (from Hexane- AcOEt); IR 1732 cm^{-1} ; ^1H NMR δ 8.00 (1H, d, $J = 8.9$ Hz), 7.33-7.18 (2H, m), 3.76 (3H, s), 3.75 (3H, s), 3.56-3.49 (1H, m), 3.20-3.03 (4H, m), 2.81-2.73 (1H, m), 1.66 (9H, s); ^{13}C NMR δ 174.6, 174.3, 150.0, 134.3, 134.1, 129.9, 128.3, 124.0, 117.4, 116.5, 113.7, 84.3, 52.2, 52.2, 42.2, 41.0, 28.2, 27.9, 23.4; MS m/z 421 (M^+ , 24). Anal. Calcd for $\text{C}_{21}\text{H}_{24}\text{ClNO}_6$: C, 59.79; H, 5.73; N, 3.32. Found: C, 59.98; H, 5.78; N, 3.30.

Dimer 12f was a colorless oil: IR 1728, 1710(sh) cm^{-1} ; ^1H NMR δ 8.20 (1H, s), 8.16 (1H, d, $J = 8.6$ Hz), 8.01 (1H, dd, $J = 1.5, 8.9$ Hz), 7.95 (1H, dd, $J = 1.8, 8.5$ Hz), 7.88 (1H, d,

$J = 8.5$ Hz), 7.66 (1H, d, $J = 1.2$ Hz), 7.51 (1H, s), 4.66 (1H, d, $J = 1.8$ Hz), 4.42 (2H, q, $J = 7.1$ Hz), 4.27-4.20 (2H, m), 3.41 (2H, dd, $J = 18.6, 42.4$ Hz), 3.02-2.83 (2H, m), 2.02-1.91 (2H, m), 1.66 (9H, s), 1.65 (9H, s), 1.43 (3H, t, $J = 7.0$ Hz), 1.22 (3H, t, $J = 7.0$ Hz); ^{13}C NMR δ 167.1, 166.2, 153.2, 151.4, 150.2, 144.9, 139.0, 136.4, 135.2, 130.3, 128.7, 125.4, 125.3, 124.9, 124.2, 120.1, 116.7, 115.1, 115.0, 94.3, 84.5, 83.4, 60.8, 60.7, 46.6, 37.9, 34.8, 31.6, 28.3, 28.3, 22.6, 17.6, 14.4, 14.1, 0.0; MS m/z 630 (M^+ , 1.1). FABHRMS calcd. for $\text{C}_{36}\text{H}_{42}\text{N}_2\text{O}_8$ 630.2941, found 630.2922.

(2*R,3*S**)-2,3-Bis(methoxycarbonyl)-9-(*tert*-butoxycarbonyl)-1,2,3,4-tetrahydrocarbazole (14a).**

Compound **14a** was a colorless solid: mp 115-118 °C (from Hexane- CH_2Cl_2); IR 1728 cm^{-1} ; ^1H NMR δ 8.11 (1H, d, $J = 7.9$ Hz), 7.42 (1H, d, $J = 7.3$ Hz), 7.28-7.20 (2H, m), 3.71 (3H, s), 3.70 (3H, s), 3.62-3.55 (1H, m), 3.41-3.29 (3H, m), 3.24-3.17 (1H, m), 3.03-2.96 (1H, m), 1.68 (9H, s); ^{13}C NMR δ 173.3, 173.2, 150.4, 136.2, 133.2, 129.7, 129.0, 123.9, 122.6, 117.7, 115.5, 114.5, 83.7, 52.2, 52.1, 52.1, 41.1, 40.0, 28.3, 26.5, 21.9; MS m/z 387 (M^+ , 19). Anal. Calcd for $\text{C}_{21}\text{H}_{25}\text{NO}_6$: C, 65.10 H, 6.50; N, 3.62. Found: C, 64.85; H, 6.72; N, 3.61.

2,3-Bis(methoxycarbonyl)-9-(*tert*-butoxycarbonyl)-1,4-dihydrocarbazole (14d).

Compound **14d** was a colorless solid: mp 133-137 °C (from Hexane-CH₂Cl₂); IR 1724, 1664 cm⁻¹; ¹H NMR δ 8.13 (1H, d, *J* = 7.3 Hz), 7.45-7.37 (1H, m), 7.35-7.22 (2H, m), 4.05 (2H, t, *J* = 7.9 Hz), 3.86 (3H, s), 3.85 (3H, s), 3.75-3.67 (2H, m), 1.69 (9H, s); ¹³C NMR δ 173.3, 167.1, 150.2, 132.2, 130.2, 127.7, 126.1, 124.6, 123.3, 120.0, 119.4, 116.0, 84.5, 52.4, 52.0, 38.1, 28.3, 28.2, 22.9; MS *m/z* 385 (M⁺, 27). Anal. Calcd for C₂₁H₂₃NO₆: C, 65.44 H, 6.02; N, 3.63. Found: C, 65.34; H, 6.08; N, 3.63.

2-(Methoxycarbonyl)-9-(*tert*-butoxycarbonyl)-1,4-dihydrocarbazole (R¹=H, R²=CO₂Me) and 3-(Methoxycarbonyl)-9-(*tert*-butoxycarbonyl)-1,4-dihydrocarbazole (R¹=CO₂Me, R²=H) (14e) .

Compound **14e** was a colorless solid: mp 116-123 °C (from Hexane-CH₂Cl₂); IR 1724 cm⁻¹; ¹H NMR δ 8.19 (1H, d, *J* = 8.5 Hz), 7.40 (1H, d, *J* = 7.3 Hz), 7.32-7.22 (3H, m), 3.93 (2H, dt, *J* = 1.8, 7.9 Hz), 3.84 (20/100 x 3H, brs), 3.83 (80/100 x 3H, brs), 3.63-3.54 (2H, m), 1.72 (9H, brs); ¹³C NMR δ 167.4, 167.3, 150.8, 150.4, 139.0, 138.0, 135.8, 135.7, 129.6, 129.0, 128.5, 128.4, 128.4, 127.7, 123.3, 122.7, 120.4, 119.3, 118.1, 117.7, 115.6, 115.5, 113.2, 111.8, 84.5, 83.8, 52.2, 52.2, 28.3, 28.3, 28.3, 28.3, 27.1, 24.1; MS *m/z* 327 (M⁺, 26). Anal. Calcd for C₁₉H₂₁NO₄: C, 69.71 H, 6.47; N, 4.28. Found: C, 69.58; H, 6.50; N, 4.40.

2-(4-Acetoxy-1,2-pentadien-3-yl)-*N*-(*tert*-butoxycarbonyl)aniline (15).

Compound **15** was a colorless oil: IR 3427, 3325, 1954, 1724 cm⁻¹; ¹H NMR δ 8.08 (1H, d, *J* = 8.3 Hz), 7.45 (1H, brs), 7.29-7.25 (1H, m), 7.15 (1H, dd, *J* = 1.5, 7.8, Hz), 7.00 (1H, dd, *J* = 1.1, 7.5, Hz), 5.45-5.39 (1H, m), 5.01 (2H, d, *J* = 2.4 Hz), 2.07 (3H, s), 1.53 (9H, s), 1.37 (3H, d, *J* = 6.3 Hz); ¹³C NMR δ 205.8, 170.4, 153.0, 136.6, 129.6, 128.7, 123.3, 122.6, 119.6, 103.8, 80.2, 78.2, 70.3, 28.3, 21.1, 29.1; MS *m/z* 317 (M⁺, 2.6). HRMS Calcd for C₁₈H₂₃NO₄ 317.1627, found 317.1632.

Compound **16** was a colorless solid: mp 117-121 °C (from Hexane-CH₂Cl₂); IR 1732 cm⁻¹; ¹H NMR δ 8.15-8.07 (1H, m), 7.52 (60/100H, d, *J* = 7.3 Hz), 7.43 (40/100H, d, *J* = 6.7 Hz), 7.29-7.19 (2H, m), 3.78-3.73 (6H, m), 3.65-3.06 (4H, m), 2.99-2.92 (40/100H, m), 2.78-2.73 (60/100H, m), 1.17 (40/100 x 9H, s), 1.16 (60/100 x 9H, s), 1.47 (60/100H x 3H, d, *J* = 7.3 Hz), 1.16 (40/100 x 3H, d, *J* = 6.7 Hz); ¹³C NMR δ 175.4, 175.0, 174.1, 173.8, 150.4, 150.3, 136.3, 136.1, 132.4, 131.9, 128.2, 128.0, 123.9, 123.6, 122.6, 122.4, 120.0, 119.2, 119.1, 117.7, 115.7, 115.6, 83.9, 83.9, 52.1, 52.1, 50.8, 45.9, 43.1, 37.9, 31.5, 29.2, 28.2, 28.2, 28.2, 28.1, 19.6, 19.6, 16.8; MS *m/z* 402 (M⁺+1, 8.4). Anal. Calcd for C₂₂H₂₇NO₆: C, 65.82 H, 6.78; N, 3.49. Found: C, 65.75; H, 6.91; N, 3.46.