Supporting Information

A Novel Route to Dipeptides via Non-condensation of Amino Acids: 2-aminoperfluoropropane as a Synthone for Trifluoroalane Dipeptides

Yong Guo, Fujiwara Kana and Kenji Uneyama*

Department of Applied Chemistry, Faculty of Engineering, Okayama University,

Okayama, 700-8653 JAPAN

Experiment Section

¹H NMR spectra were recorded at 300, 500 or 600 MHz, ¹⁹F NMR spectra were recorded at 282 MHz using CDCl₃. The chemical shifts are reported in δ (ppm) related to the CHCl₃ (7.26 ppm) and C₆F₆ (0 ppm). Coupling constants (J) are reported in hertz (Hz). All air and/or moisture sensitive reactions were carried out under argon atmosphere with dry, freshly distilled solvents. Dried THF was distilled from sodium and benzophenone. DMF was dried over calcium hydride. Diastereomeric excess values were measured on Chiralel OD-H column by HPLC or were determined by ¹⁹F NMR if two diastereomers were of obviously different chemical shifts.

Synthesis of *N*-(*p*-methoxyphenyl) bis(trifluoromethyl)imine (9)

In a 3000 mL flask, a solution of *p*-anisidine (123 g, 1.0 mol) in 1500 mL dichloromethylene was cooled at about -40 °C, and then was bubbled into hexafluoroacetone gas which was generated by slowly dropping hexafluoroacetone trihydrate (265 g, 1.2 mol) to concentrated sulfuric acid (133 mL) at 90 °C over a period of 8 hours. A moisture-sensitive pale violet precipitate which was adduct of hexafluoroacetone with *p*-anisidine was gradually formed. Then, triethylamine (415 mL, 3.0 mmol) was added to the solution and the precipitate was dissolved. After that the flask was equipped with a condenser and phosphorous oxychloride (153 g, 1 mmol) was added dropwise at such a rate that gentle reflux was maintained. The solution turned to yellow and some precipitate formed. After removal of the solvent and the tertiary amine,

the residue was subjected to distillation under reduced pressure to give the crude product.

Distillation again afforded a yellow liquid **1** (158 g, 0.58 mmol, 58%). Bp: 78-81 °C / 28 torr or 105-110 °C / 50 torr; IR (neat) 1600, 1506, 1250, 1164 cm⁻¹; ¹H NMR (CDCl₃) δ 3.84 (s, 3H), 6.94 (d, J = 9, 2H), 7.00 (d, J = 9, 2H); ¹⁹F NMR (CDCl₃) δ 91.8 (s), 98.8 (s). Anal. Calcd for C₁₀H₇F₆NO: C, 44.29; H, 2.60; N, 5.17. Found: C, 44.08; H, 2.60; N, 5.45.

***N*-trimethylsilyl-*N*-*p*-methoxyphenyl-1-trifluoromethyl-2,2-difluoroethenamine (**7**)**

Under argon atmosphere, in a 1000 mL flask, were added Mg (4.86 g, 200 mmol), **9** (27.1 g, 100 mmol) and 400 mL anhydrous THF. The suspension was cooled at 0 °C and stirred. Chlorotrimethylsilane (50 mL, 400 mmol) was dropped into the suspension. The starting material **9** was completely consumed within 30 minutes determined by TLC monitoring. After decantation of Mg followed by evaporation of the solvent and chlorotrimethylsilane, subsequent addition of hexane to the residue to precipitate inorganic salt and filtration, the filtrate was evaporated and distilled under reduced pressure to afford product **7** (30.9 g, 95 mmol, 95%). Bp: 75 °C/1 torr; IR (neat) 2964, 1734, 1512, 1238 cm⁻¹; ¹H NMR δ 0.22 (s, 9H), 3.78 (s, 3H), 6.80 (d, J = 9, 2H), 6.98 (d, J = 9, 2H); ¹⁹F NMR δ 76.3 (m, 1F), 78.3 (m, 1F), 97.4 (dd, J = 23.4, 11.6, 3F). Anal. Calcd for C₁₃H₁₆F₅NOSi: C, 47.99; H, 4.96; N, 4.31. Found: C, 47.60; H, 4.71; N, 4.56.

***N*-*p*-methoxyphenyl-3,3,3-trifluoroalanylglycine benzyl ester (11a)**

Under argon atmosphere in a 30 mL flask, were added **7** (325 mg, 1 mmol), DMF (1.4 mL), **10a** (505 mg, 1.5 mmol) and triethylamine (506 mg, 5 mmol) subsequently. The mixture was stirred at room temperature for 1.5 h. Then, into the mixture cooled at to 0 °C, was added 1.5 mL water and then hydrochloric acid was dropped into until the pH of the solution was 3. After stirring for 1 h, the mixture was poured into 10 mL water and extracted with ether (10 mL × 3). The combined organic phase was washed with saturated sodium hydrogen carbonate aqueous solution (10 mL) and saturated brine (5 mL × 3), and dried over anhydrous magnesium sulfate. After removal of the solvent, the residue was subjected to column chromatography on silica gel (Hexane/EtOAc = 3/1) to afford a solid **11a** (362 mg, 0.91 mmol, 91%) which could be recrystallized from hexane and ethyl ether. Mp 97-99 °C; IR (KBr) 3336, 1740, 1658, 1512, 1240 cm⁻¹; ¹H NMR (CDCl₃) δ 3.75 (s, 3H), 4.10-4.18 (m, 2H), 4.28-4.32 (m, 1H), 4.36 (d, J = 6.0, 1H), 5.18 (s, 2H), 6.67-6.70 (m, 2H), 6.78-6.81 (m, 2H), 7.02 (br, 1H), 7.32-7.36 (m, 5H); ¹⁹F NMR (CDCl₃) δ 89.0 (d, J = 7.1). Anal. Calcd for C₁₉H₁₉F₃N₂O₄: C, 57.57; H, 4.83; N, 7.07. Found: C, 57.32; H, 4.68; N, 6.95.

***N*-*p*-methoxyphenyl-3,3,3-trifluoroalanylalanine methyl ester (11b)**

Under argon atmosphere in a 30 mL flask, were added **7** (325mg, 1mmol), DMF (1.4 mL), **10b** (208 mg, 1.5 mmol) and triethylamine (0.69 mL, 5 mmol) subsequently. The mixture was stirred at room temperature for 1.5 h. Then, into the mixture cooled at 0 °C, was added 1.4 mL water and then hydrochloric acid was dropped into until the pH of the solution was 3. After stirring for 1 h, the mixture was poured into 10 mL water and extracted with ethyl ether (10 mL × 3). The combined organic phase was washed with saturated sodium hydrogen carbonate aqueous solution (10 mL) and saturated brine (5 mL × 3), and dried over anhydrous magnesium sulfate. After removal of the solvent, the residue was subjected to column chromatography on silica gel (Hexane/EtOAc = 3/1) to afford a solid **11b** (287 mg, 0.86 mmol, 86%). Mp 73-75 °C; IR (KBr) 3316, 1740, 1662, 1522, 1248 cm⁻¹; ¹H NMR (CDCl₃) δ 1.43 (t, J = 6.9, 3H), 3.74 (s, 3H), 3.76 (m, 3H), 4.23-4.29 (m, 1H), 4.38 (br, 1H), 4.60-4.66 (m, 1H), 6.67-6.72 (m, 2H), 6.79-6.82 (m, 2H), 6.94 (d, J = 6.0, 0.44H), 7.04 (d, J = 6.0, 0.56H); ¹⁹F NMR (CDCl₃) δ 89.0 (d, J = 6.8). Anal. Calcd for C₁₄H₁₇F₃N₂O₄: C, 50.30; H, 5.13; N, 8.38. Found: C, 50.30; H, 5.23; N, 8.14.

***N*-*p*-methoxyphenyl-3,3,3-trifluoroalanylphenylalanine ethyl ester (**11c**)**

Under argon atmosphere in a 100 mL flask, were added **7** (1.63 g, 5 mmol), DMF (7.1 mL), **10c** (1.72 g, 7.5 mmol) and triethylamine (3.5 mL, 25 mmol) subsequently. The mixture was stirred at room temperature for 1.5 h. Then, into the mixture cooled at 0 °C, was added 7.1 mL water and then hydrochloric acid was dropped into until the pH of the solution was 3. After stirring for 3 h, the mixture was poured into 100 mL water and extracted with ethyl ether (50 mL × 3). The combined organic phase was washed with saturated sodium hydrogen carbonate aqueous solution (30 mL) and saturated brine (10 mL × 3), and dried over anhydrous magnesium sulfate. After removal of the solvent, the residue was subjected to column chromatography on silica gel (Hexane/EtOAc = 3/1) to afford a solid **11c** (1.96 g, 4.63 mmol, 93%). Mp 104-106 °C; IR (neat) (KBr) 3348, 1734, 1664, 1520, 1248 cm⁻¹; ¹H NMR (CDCl₃) δ 1.20-1.27 (m, 3H), 3.10-3.19 (m, 2H), 3.76 (s, 3H), 4.14-4.22 (m, 3H), 4.25-4.37 (m, 1H), 4.84-4.93 (m, 1H), 6.61-6.68 (m, 2H), 6.77-6.81 (m, 2H), 6.85-6.89 (m, 1H), 6.98-7.00 (m, 2H) 7.18-7.23 (m, 3H); ¹⁹F NMR (CDCl₃) δ 89.5 (d, J = 7.1, 0.62 × 3F), 89.6 (d, J = 7.1, 0.38 × 3F). Anal. Calcd for C₂₁H₂₃F₃N₂O₄: C, 59.43; H, 5.46; N, 6.60. Found: C, 59.42; H, 5.55; N, 6.56.

***N*-*p*-methoxyphenyl-3,3,3-trifluoroalanylleucine ethyl ester (**11d**)**

Under argon atmosphere in a 30 mL flask, were added **7** (325mg, 1mmol), DMF (1.4 mL), **10b** (294 mg, 1.5 mmol) and triethylamine (0.69 mL, 5 mmol) subsequently. The mixture was stirred at room temperature for 1.5 h. Then, into the mixture cooled at 0 °C, was added 1.4 mL water and then hydrochloric acid was dropped into until the pH of the solution was 1. After stirring for 1 h, the mixture was poured into 10 mL water and extracted with ethyl ether (10 mL × 3). The combined organic phase was washed with saturated sodium hydrogen carbonate aqueous solution (10 mL) and saturated brine (5 mL × 3), and dried over anhydrous magnesium sulfate. After removal of the solvent, the residue was subjected to column chromatography on silica gel (Hexane/EtOAc = 3/1) to afford a liquid **11d** (331 mg, 0.85 mmol, 85%). IR 3348, 1740, 1668, 1516, 1238 cm⁻¹; ¹H NMR (CDCl₃) δ 0.87-0.89 (m, 6H), 1.21-1.27 (m, 3H), 1.53-1.68 (m, 3H), 3.75 (s, 3H), 4.13-4.20 (m, 2H), 4.24-4.33 (m, 1H), 4.35-4.47(m, 1H), 4.62-4.68 (m, 1H), 6.66-6.71 (m, 2H), 6.78-6.81 (m, 2H), 6.95 (d, J = 8.4, 1H); ¹⁹F NMR (CDCl₃) _ 89.0 (d, J = 6.8 Hz, 0.72 × 3F), 89.1 (d, J = 7.1, 0.28 × 3F) (slightly overlapped). Anal. Calcd for C₁₈H₂₅F₃N₂O₄: C, 55.38; H, 6.45; N, 7.18. Found: C. 55.32; H, 6.42; N, 7.21.

***N*-*p*-methoxyphenyl-3,3,3-trifluoroalanylserine methyl ester (**11e**)**

Under argon atmosphere in a 30 mL flask, were added **7** (325mg, 1mmol), DMF (1.4 mL), **10e** (234 mg, 1.5 mmol) and triethylamine (0.69 mL, 5 mmol) subsequently. The mixture was stirred at room temperature for 1.5 h. Then, into the mixture cooled at 0 °C, was added 1.4 mL water and then hydrochloric acid was dropped into until the pH of the solution was 3. After stirring for 1 h, the mixture was poured into 10 mL water and extracted with ethyl ether (10 mL $\times$ 3). The combined organic phase was washed with saturated sodium hydrogen carbonate aqueous solution (10 mL) and saturated brine (5 mL $\times$ 3), and dried over anhydrous magnesium sulfate. After removal of the solvent, the residue was subjected to column chromatography on silica gel (Hexane/EtOAc = 2/1) to afford a solid **11e** (280 mg, 0.80 mmol, 80%). Mp 89-95 °C; IR (KBr) 3548, 3300, 1736, 1664, 1518, 1238 cm^{-1} ; ^1H NMR (CDCl_3) δ 3.75-3.79 (m, 6H), 3.89-3.94 (m, 1H), 4.00-4.08 (m, 1H), 4.30-4.43 (m, 2H), 4.68-4.72 (m, 1H), 6.70-6.73 (m, 2H), 6.80-6.83 (m, 2H), 7.28-7.34 (m, 1H); ^{19}F NMR (CDCl_3) δ -89.0 (two groups of overlapped doublet, J = 7.1). Anal. Calcd for $\text{C}_{14}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_5$: C, 48.00; H, 4.89; N, 8.00. Found: C, 48.20; H, 4.86; N, 7.96.

***N*-*p*-methoxyphenyl-3,3,3-trifluoroalanyltyrosine methyl ester (**11f**)**

Under argon atmosphere in a 30 mL flask, were added **7** (325mg, 1mmol), DMF (1.4 mL), **10f** (346mg, 1.5 mmol) and triethylamine (0.69 mL, 5 mmol) subsequently. The mixture was stirred at room temperature for 1.5 h. Then, into the mixture cooled at 0 °C, was added 1.4 mL water and then hydrochloric acid was dropped into until the pH of the solution was 3. After stirring for 1 h, the mixture was poured into 10 mL water and extracted with ethyl ether (10 mL $\times$ 3). The combined organic phase was washed with saturated sodium hydrogen carbonate aqueous solution (10 mL) and saturated brine (5 mL $\times$ 3), and dried over anhydrous magnesium sulfate. After removal of the solvent, the residue was subjected to column chromatography on silica gel (Hexane/EtOAc = 2/1) to afford a solid **11f** (290 mg, 0.68 mmol, 68%). Mp 166-168 °C; IR (KBr) 3324, 1736, 1676, 1520, 1238 cm^{-1} ; ^1H NMR (CDCl_3) δ 3.01-3.11 (m, 2H), 3.70-3.72 (3H), 3.75 (s, 3H), 4.18-4.38 (m, 2H), 4.81-4.92 (m, 1H), 6.60-6.68 (m, 4H), 6.78-6.83 (m, 4H), 6.89 (br, 1H); ^{19}F NMR (CDCl_3) δ 89.0 (d, $J = 7.1$, $0.59 \times 3\text{F}$), 89.1 (d, $J = 7.1$, $0.41 \times 3\text{F}$). Anal. Calcd for $\text{C}_{20}\text{H}_{21}\text{F}_3\text{N}_2\text{O}_5$: C, 56.34; H, 4.96; N, 6.57. Found: C, 55.93; H, 4.82; N, 6.50.

***N*-*p*-methoxyphenyl-3,3,3-trifluoroalanylmethionine methyl ester (**11g**)**

Under argon atmosphere in a 30 mL flask, were added **7** (325mg, 1mmol), DMF (1.4 mL), **10g** (330 mg, 1.5 mmol) and triethylamine (0.69 mL, 5 mmol) subsequently. The mixture was stirred at room temperature for 1.5 h. Then, into the mixture cooled at 0 °C, was added 1.4 mL water and then hydrochloric acid was dropped into until the pH of the solution was 1. After stirring for 1 h, the mixture was poured into 10 mL water and extracted with ethyl ether (10 mL × 3). The combined organic phase was washed with saturated sodium hydrogen carbonate aqueous solution (10 mL) and saturated brine (5 mL × 3), and dried over anhydrous magnesium sulfate. After removal of the solvent, the residue was subjected to column chromatography on silica gel (Hexane/EtOAc = 3/1) to afford a solid **11g** (311 mg, 0.79 mmol, 79%). Mp 74-79 °C; IR (KBr) 3336, 1738, 1654, 1520, 1244 cm⁻¹; ¹H NMR (CDCl₃) δ 1.98-2.48 (m, 7H), 3.73-3.76 (m, 6H), 4.25-4.33 (m, 2H), 4.73-4.79 (m, 1H), 6.68-6.72 (m, 2H), 6.79-6.82 (m, 2H), 7.10-7.25 (m, 1H); ¹⁹F NMR (CDCl₃) - 89.0 (two groups of overlapped doublet, J=7.1). Anal. Calcd for C₁₆H₂₁F₃N₂O₄S: C, 48.72; H, 5.37; N, 7.10. Found: C. 48.38; H, 5.43; N, 6.90

***N*-*p*-methoxyphenyl-3,3,3-trifluoroalanyltryptophan methyl ester (**11h**)**

Under argon atmosphere in a 30 mL flask, were added **7** (325mg, 1mmol), DMF (1.4 mL), **10h** (382 mg, 1.5 mmol) and triethylamine (0.69 mL, 5 mmol) subsequently. The mixture was stirred at room temperature for 1.5 h. Then, into the mixture cooled at 0°C , was added 1.4 mL water and then hydrochloric acid was dropped into until the pH of the solution was 3. After stirring for 1 h, the mixture was poured into 10 mL water and extracted with ethyl ether (10 mL $\times$ 3). The combined organic phase was washed with saturated sodium hydrogen carbonate aqueous solution (10 mL) and saturated brine (5 mL $\times$ 3), and dried over anhydrous magnesium sulfate. After removal of the solvent, the residue was subjected to column chromatography on silica gel (Hexane/EtOAc = 3/1) to afford a liquid **11h** (354 mg, 0.79 mmol, 79%). Mp $177\text{--}180^\circ\text{C}$; IR (KBr) 3448, 1740, 1658, 1516, 1246 cm^{-1} ; ^1H NMR (CDCl_3) δ 3.30-3.43 (m, 2H), 3.68-3.76 (m, 6H), 4.15-4.24 (m, 2H), 4.93-4.99 (m, 1H), 6.55-6.64 (m, 2H), 6.74-6.81 (m, 2H), 6.84-6.91 (m, 1H), 6.95-7.00 (m, 1H), 7.14-7.20 (m, 1H), 7.33-7.42 (m, 1H), 8.00 (br, 1H). ^{19}F NMR (CDCl_3) of the recrystallized product δ 89.0 (d, $J = 7.1$, $0.65 \times 3\text{F}$), 89.2 (d, $J = 7.1$, $0.35 \times 3\text{F}$). Anal. Calcd for $\text{C}_{22}\text{H}_{22}\text{F}_3\text{N}_3\text{O}_4$: C, 58.79; H, 4.93; N, 9.35. Found: C, 58.09; H, 4.97; N, 9.15.

***N*-*p*-methoxyphenyl-3,3,3-trifluoroalanylproline methyl ester (11i)**

Under argon atmosphere in a 30 mL flask, were added **7** (325mg, 1mmol), DMF (1.4 mL), **10i** (249 mg, 1.5 mmol) and triethylamine (0.69 mL, 5 mmol) subsequently. The mixture was stirred at room temperature for 1.5 h. Then, into the mixture cooled at 0 °C, was added 1.4 mL water and then hydrochloric acid was dropped into until the pH of the solution was 3. After stirring for 1 h, the mixture was poured into 10 mL water and extracted with ethyl ether (10 mL × 3). The combined organic phase was washed with saturated sodium hydrogen carbonate aqueous solution (10 mL) and saturated brine (5 mL × 3), and dried over anhydrous magnesium sulfate. After removal of the solvent, the residue was subjected to column chromatography on silica gel (Hexane/EtOAc = 2/1) to afford a liquid **11i** (248 mg, 0.69 mmol, 69%). IR (Film) 3340, 2960, 1746, 1664, 1516, 1244 cm⁻¹; ¹H NMR (CDCl₃) 1.94-2.14 (m, 4H), 2.19-2.32 (m, 1H), 3.68-3.83 (m, 8H), 4.52-4.65 (m, 3H), 6.66-6.84 (m, 4H); ¹⁹F NMR (CDCl₃) δ 88.7 (d, J = 4.5, 0.7 × 3), 88.8 (d, J = 4.8, 0.3 × 3F), (partly overlapped). Anal. Calcd for C₂₁H₂₀F₃N₃O₄: Calcd for C₁₆H₁₉F₃N₂O₄: C, 53.33; H, 5.31; N, 7.77. Found: C, 53.22; H, 5.32; N, 7.65.

3-phenyl-2-[1,3,3,3-tetrafluoro-2-(4-methoxyphenylamino)propylideneamino]-propionic acid ethyl ester (13c)

Under argon atmosphere in a 50 mL flask, were added **7** (1.74 g, 5.35 mmol), DMF (15 mL), **10c** (1.73 g, 7.5 mmol) and triethylamine (3.5 mL, 25 mmol) subsequently. The mixture was stirred at room temperature for 1.5 hours. Then, 150 mL water was added into the mixture and the mixture was extracted with ethyl ether (100 mL $\times$ 3). The combined organic phase was washed with saturated brine (30 mL $\times$ 3), and dried over anhydrous magnesium sulfate. After removal of the solvent, the residue was subjected to column chromatography on silica gel (Hexane/EtOAc = 5/1) to afford a yellow liquid **13c** (1.78 g, 4.2 mmol, 79%).

Competitive reaction of enamine **7** with benzyl amine and benzyl alcohol

Under argon atmosphere in a 30 mL flask, were added **7** (100 mg, 0.31 mmol), DMF (1 mL), benzyl amine (50 mg, 0.47 mmol) and benzyl alcohol (50 mg, 0.47 mmol) and triethylamine (0.21 mL, 1.6 mmol) subsequently. The mixture was stirred at room temperature for 2 h. Then, into the mixture cooled at 0 °C, was added 1 mL water and then perchloric acid was dropped into until the pH of the solution was 1. After stirring for 5 h, the mixture was poured into 10 mL water and extracted with ethyl ether (10 mL $\times$

3). The combined organic phase was washed with saturated sodium hydrogen carbonate aqueous solution (10 mL) and saturated brine (5 mL x 3), and dried over anhydrous magnesium sulfate. After removal of the solvent, the residue was subjected to column chromatography on silica gel (Hexane/EtOAc = 4/1) to afford a white solid **14** (102 mg, 0.30 mmol, 98%) without **15**. **14**: Mp 110-112 °C; IR (neat) cm^{-1} 3320, 1660, 1522; ^1H NMR (CDCl_3) δ 3.76 (s, 3H), 4.26 (q, $J = 7.5$ Hz, 1H), 4.39 (br, 1H), 4.50-4.58 (m, 2H), 6.66-6.78 (m, 3H), 6.78-6.82 (m, 2H), 7.18-7.21 (m, 2H), 7.26-7.35 (m, 3H); ^{19}F NMR (CDCl_3) δ 89.0 (d, $J = 7.1$ Hz). Anal. Calcd for $\text{C}_{17}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_2$: C, 60.35; H, 5.06; N, 8.28. Found: C, 60.65; H, 5.45; N, 8.58.

3,3,3-trifluoroalanylphenylalanine ethyl ester (**16ca** and **16cb**)

To a solution of **11c** (1.58 g, 3.73 mmol) in 37 mL acetonitrile at 0 °C, was dropped an aqueous solution of CAN (6.13 g, 11.2 mmol) in 13 mL water. The reaction took place at 0 °C for 0.5 h and at room temperature for another 2.5 h. Then, the reaction solution was neutralized with NaHCO_3 aqueous solution, reduced by $\text{Na}_2\text{S}_2\text{O}_3$ (630 mg, 4 mmol) and stirred for 30 minutes. After filtration, the filtrate was extracted by ethyl acetate, washed with brine and dried over MgSO_4 . After evaporation of the solvent, the residue was subjected column chromatography over silica gel (hexane/AcOEt = 2.5/1 to 2/1) to separate the either diastereomer of the product **16ca** (601 mg, 1.88 mmol, 50%) and **16cb** (354 mg, 1.11 mmol, 30%). **16ca**: Mp 89-91 °C; IR (KBr) cm^{-1} 3344,

1736, 1666; ^1H NMR (CDCl_3) δ : 1.24-1.29 (t, $J = 7.1\text{ Hz}$, 3H), 3.10-3.24 (m, 2H), 3.86 (q, $J = 7.8\text{ Hz}$, 1H), 4.20 (q, $J = 7.2\text{ Hz}$, 2H), 4.86 (dt, $J = 7.8, 5.9\text{ Hz}$, 1H), 7.08-7.11 (m, 2H), 7.21-7.34 (m, 5H); ^{19}F NMR (CDCl_3) δ 87.1 (d, $J = 9.3\text{ Hz}$). Anal. Calcd for $\text{C}_{14}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_3$: C, 52.83; H, 5.38; N, 8.80. Found: C, 52.90; H, 5.51; N, 8.89. **16cb**: Mp 86-88 $^\circ\text{C}$; IR (KBr) 3332, 1734, 1660 cm^{-1} ; ^1H NMR (CDCl_3) δ 1.24-1.28 (m, 3H), 3.11-3.19 (m, 2H), 3.82 (q, $J = 7.6\text{ Hz}$, 1H), 4.17-4.21 (m, 2H), 4.85-4.89 (m, 1H), 7.10-7.12 (m, 3H), 7.23-7.30 (m, 3H); ^{19}F NMR (CDCl_3) δ 87.0 (d, $J = 7.1\text{ Hz}$). Anal. Calcd for $\text{C}_{14}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_3$: C, 52.83; H, 5.38; N, 8.80. Found: C, 52.91; H, 5.32; N, 9.05.

3,3,3-trifluoroalanylphenylalanine ethyl ester hydrochloric acid salt (**17ca**)

To a solution of **16ca** (200 mg, 0.63 mmol) in ethyl acetate was added a solution of hydrochloric acid in methanol (1M, 1 mL). After evaporation of all solvent and recrystallization in methanol and ethyl acetate, a white crystalline compound (200 mg, 0.56 mmol, 90%) was obtained. Mp 150-157 $^\circ\text{C}$; IR (KBr) cm^{-1} 3268, 2996, 1706, 1578, 1504; ^1H NMR ($d\text{-DMSO}$) δ 1.09 (t, $J = 7.1$, 3H), 3.04 (d, $J = 7$, 2H), 4.04 (q, $J = 7.2$, 2H), 4.57 (q, $J = 7.1$, 1H), 4.90 (q, $J = 7.5$, 1H), 7.17-7.40 (m, 5H), 9.42 (d, $J = 6.8$, 1H); ^{19}F NMR ($d\text{-DMSO}$, trifluoroacetic acid (86.0 ppm) as internal standard) δ 91.8 (d, $J = 9.3$). Calcd for $\text{C}_{14}\text{H}_{18}\text{ClF}_3\text{N}_2\text{O}_3$: C, 47.40; H, 5.11; N, 7.90. Found: C, 47.31; H, 5.29; N, 8.16.

HPLC analysis of the ratio of diastereomers of trifluoroalanine dipeptides **11**:

The HPLC analysis of the ratio of diastereomers of trifluoroalanine dipeptides **11** was performed by Shimazu LC-10AT with a chiral column of OD-H (Daicel Chem. Co Ltd.).

The elution solvent was a mixture of hexane:i-PrOH=9:1 which was passed on the flow rate of 0.5 ml/min.

X-ray crystallographic analysis of **17ca**

ORTEP drawing of X-ray structure of **17ca**

Cl F data of **17ca**

data_General

_audit_creation_date 'Mon Jun 6 14:34:12 2005'

_audit_creation_method 'by teXsan'

_audit_update_record ?

#-----

PROCESSING SUMMARY (IUCr Office Use Only)

_journal_date_rcd_electronic ?

_journal_date_from_coeditor ?

_journal_date_accepted ?


```

_journal_coeditor_code      ?
#-----
# SUBMISSION DETAILS
_publ_contact_author_name   ' ENTER NAME'
_publ_contact_author_address
;
    ENTER ADDRESS
;
_publ_contact_author_email   ' ENTER EMAIL ADDRESS '
_publ_contact_author_fax     ' ENTER FAX NUMBER '
_publ_contact_author_phone   ' ENTER PHONE NUMBER '
_publ_contact_letter
;
    ENTER TEXT OF LETTER
;
_publ_requested_journal      ' ENTER JOURNAL NAME HERE'
_publ_requested_category     ' CHOOSE FI FM FO CI CM CO or AD'

_publ_requested_coeditor_name      ?
#-----
# TITLE AND AUTHOR LIST

_publ_section_title
;
    ENTER SECTION TITLE
;

_publ_section_title_footnote
;
    ENTER FOOTNOTE TO TITLE OF PAPER
;
loop_
_publ_author_name
_publ_author_footnote
_publ_author_address

' FIRST AUTHORS NAME '
;
    FIRST AUTHORS FOOTNOTES
;
;

```

FIRST AUTHORS ADDRESS

;

_publ_section_synopsis

;

ENTER SYNOPSIS

;

#-----

TEXT

_publ_section_abstract

;

ENTER ABSTRACT

;

_publ_section_comment

;

ENTER TEXT

;

_publ_section_acknowledgements

;

ENTER ACKNOWLEDGEMENTS

;

_publ_section_references

;

ENTER OTHER REFERENCES

Molecular Structure Corporation, Rigaku Corporation. (2000). teXsan.

Single Crystal Structure Analysis Software. Version 1.11.

MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.

S18

Rigaku, 3-9-12 Akishima, Tokyo, Japan.;

_publ_section_figure_captions;

ENTER FIGURE CAPTIONS;

_publ_section_exptl_prep;

ENTER COMPOUND PREPARATION DETAILS;

_publ_section_exptl_refinement

;

ENTER SPECIAL DETAILS OF THE REFINEMENT;

```
#-----
data_katagiri96
#-----
# CHEMICAL DATA
_chemical_formula_sum      'C14 H18 Cl F3 N2 O3 '
_chemical_formula_moiety    '?'
_chemical_formula_weight    354.76
_chemical_melting_point     ?
_Chemical_absolute_configuration  rm
#-----
# CRYSTAL DATA
_symmetry_cell_setting      orthorhombic
_symmetry_space_group_name_H-M  'P 21 21 21 '
_symmetry_Int_Tables_number  19
loop_
_symmetry_equiv_pos_as_xyz
x,y,z
1/2-x,-y,1/2+z
1/2+x,1/2-y,-z
-x,1/2+y,1/2-z
_cell_length_a              5.1724(5)
_cell_length_b              16.492(3)
_cell_length_c              19.460(2)
_cell_angle_alpha           90
                             S19
_cell_angle_beta            90
_cell_angle_gamma           90
_cell_volume                1659.9(4)
_cell_formula_units_Z       4
_cell_measurement_reflns_used  12256
_cell_measurement_theta_min  1.6
_cell_measurement_theta_max  27.5
_cell_measurement_temperature  120.2
#-----
_exptl_crystal_description   'needle'
_exptl_crystal_colour        'colorless'
_exptl_crystal_size_max     0.100
_exptl_crystal_size_mid     0.100
_exptl_crystal_size_min     0.100
_exptl_crystal_size_rad     ?
```

```

_exptl_crystal_density_diffn    1.419
_exptl_crystal_density_meas    ?
_exptl_crystal_density_method  'not measured'
_exptl_absorpt_coefficient_mu   0.275
_exptl_absorpt_correction_type  integration
_exptl_absorpt_process_details  '(Higashi, 1999)'
_exptl_absorpt_correction_T_min 0.974
_exptl_absorpt_correction_T_max 0.991

```

#-----

EXPERIMENTAL DATA

```

_diffn_radiation_type          'Mo Kαa'
_diffn_radiation_wavelength    0.7107
_diffn_measurement_device_type  'Rigaku RAXIS-IV Imaging Plate'
_diffn_measurement_method      'w
_diffn_detector_area_resol_mean 10.00
_diffn_reflns_number           9790
_diffn_reflns_av_R_equivalents 0.050
_diffn_reflns_theta_max        27.48
_diffn_measured_fraction_theta_max 0.9816
_diffn_reflns_theta_full       27.48
                                S20
_diffn_measured_fraction_theta_full 0.9816
_diffn_reflns_limit_h_min      -6
_diffn_reflns_limit_h_max      6
_diffn_reflns_limit_k_min      -21
_diffn_reflns_limit_k_max      21
_diffn_reflns_limit_l_min      -24
_diffn_reflns_limit_l_max      25

```

#-----

REFINEMENT DATA

```
_refine_special_details
```

;

Refinement using reflections with $F^2 > 0.0 \sigma(F^2)$. The weighted R-factor

(wR), goodness of fit (S) and R-factor (gt) are based on F, with F set to zero for negative F. The threshold expression of $F^2 > 2.0 \sigma(F^2)$ is used only for calculating R-factor (gt).

;

```

_reflns_number_total           2193
_reflns_number_gt               1606
_reflns_threshold_expression     $F^2 > 2.0 \sigma(F^2)$ 

```

```

_refine_ls_structure_factor_coef    F
_refine_ls_R_factor_gt              0.0787
_refine_ls_wR_factor_ref            0.1217
_refine_ls_hydrogen_treatment      refall
_refine_ls_number_reflns            2113
_refine_ls_number_parameters         281
_refine_ls_goodness_of_fit_ref       1.112
_refine_ls_weighting_scheme          calc
_refine_ls_weighting_details
  'w = 1/[ $\sigma^2(F_o) + 0.00608|F_o|^2$ ]'
_refine_ls_shift/su_max              0.0766
_refine_diff_density_max             0.53
_refine_diff_density_min            -0.41
_refine_ls_extinction_method          none
_refine_ls_extinction_coef            ?

                                         S21
_refine_ls_abs_structure_details      rm
_refine_ls_abs_structure_Flack        0.2(2)
loop_
  _atom_type_symbol
  _atom_type_description
  _atom_type_scatter_dispersion_real
  _atom_type_scatter_dispersion_imag
  _atom_type_scatter_source
  'C' 'C' 0.003 0.002
;International Tables for Crystallography
(1992, Vol. C, Tables 4.2.6.8 and 6.1.1.1)
;
  'H' 'H' 0.000 0.000
;International Tables for Crystallography
(1992, Vol. C, Table 6.1.1.2)
;
  'O' 'O' 0.011 0.006
;International Tables for Crystallography
(1992, Vol. C, Tables 4.2.6.8 and 6.1.1.1)
;
  'N' 'N' 0.006 0.003
;International Tables for Crystallography
(1992, Vol. C, Tables 4.2.6.8 and 6.1.1.1)
;
  'Cl' 'Cl' 0.148 0.159

```

;International Tables for Crystallography
(1992, Vol. C, Tables 4.2.6.8 and 6.1.1.1)

;

'F' 'F' 0.017 0.010

;International Tables for Crystallography
(1992, Vol. C, Tables 4.2.6.8 and 6.1.1.1)

;

#-----

ATOMIC COORDINATES AND DISPLACEMENT PARAMETERS

loop_

S22

_atom_site_label

_atom_site_type_symbol

_atom_site_fract_x

_atom_site_fract_y

_atom_site_fract_z

_atom_site_U_iso_or_equiv

_atom_site_adp_type

_atom_site_occupancy

_atom_site_calc_flag

_atom_site_refinement_flags

_atom_site_disorder_assembly

_atom_site_disorder_group

Cl(1)	Cl	0.4067(3)	-0.11604(9)	0.47536(9)	0.0226(3)	Uani	1.00	d...
F(4)	F	0.6595(10)	-0.3217(3)	0.6939(3)	0.044(1)	Uani	1.00	d...
F(5)	F	0.2413(9)	-0.3259(3)	0.6911(3)	0.039(1)	Uani	1.00	d...
F(6)	F	0.439(1)	-0.2280(3)	0.7437(2)	0.044(1)	Uani	1.00	d...
O(3)	O	0.8700(8)	-0.2007(3)	0.5834(3)	0.022(1)	Uani	1.00	d...
O(5)	O	1.236(1)	-0.0433(4)	0.7206(3)	0.037(1)	Uani	1.00	d...
O(6)	O	0.8422(9)	-0.0166(4)	0.7584(3)	0.032(1)	Uani	1.00	d...
N(1)	N	0.419(1)	-0.2800(4)	0.5623(3)	0.021(1)	Uani	1.00	d...
N(2)	N	0.671(1)	-0.1028(4)	0.6458(3)	0.021(1)	Uani	1.00	d...
C(1)	C	0.682(1)	-0.1748(4)	0.6156(3)	0.017(1)	Uani	1.00	d...
C(2)	C	0.878(1)	-0.0436(4)	0.6402(4)	0.023(1)	Uani	1.00	d...
C(4)	C	1.008(1)	-0.0338(4)	0.7094(4)	0.021(1)	Uani	1.00	d...
C(5)	C	0.446(1)	-0.2752(5)	0.6886(4)	0.030(1)	Uani	1.00	d...
C(6)	C	0.771(1)	0.0367(4)	0.6113(4)	0.023(2)	Uani	1.00	d...
C(7)	C	0.975(1)	0.0988(4)	0.5960(4)	0.023(1)	Uani	1.00	d...
C(8)	C	0.445(1)	-0.2276(4)	0.6231(4)	0.021(1)	Uani	1.00	d...
C(9)	C	0.927(2)	-0.0140(6)	0.8292(4)	0.034(2)	Uani	1.00	d...
C(10)	C	0.697(2)	-0.0244(5)	0.8746(4)	0.034(2)	Uani	1.00	d...

C(11)	C	0.987(2)	0.1691(5)	0.6367(5)	0.035(2)	Uani 1.00 d ...
C(12)	C	1.141(2)	0.0895(5)	0.5420(4)	0.031(2)	Uani 1.00 d ...
C(13)	C	1.324(2)	0.1484(6)	0.5263(6)	0.042(2)	Uani 1.00 d ...
C(14)	C	1.334(2)	0.2185(6)	0.5662(6)	0.045(2)	Uani 1.00 d ...
C(15)	C	1.174(2)	0.2279(6)	0.6219(6)	0.045(2)	Uani 1.00 d ...

S23

H(1)	H	0.88(1)	0.171(4)	0.684(3)	0.000(9)	Uiso 1.00 calc ...
H(2)	H	1.11(4)	0.08(1)	0.51(1)	0.155(2)	Uiso 1.00 calc ...
H(3)	H	1.43(2)	0.142(5)	0.487(5)	0.032(7)	Uiso 1.00 calc ...
H(4)	H	1.49(3)	0.257(8)	0.564(6)	0.071(5)	Uiso 1.00 calc ...
H(5)	H	1.19(1)	0.262(4)	0.643(3)	-0.008(9)	Uiso 1.00 calc ...
H(6)	H	0.70(1)	0.023(4)	0.565(4)	0.004(9)	Uiso 1.00 calc ...
H(7)	H	0.65(1)	0.051(4)	0.642(4)	0.008(9)	Uiso 1.00 calc ...
H(8)	H	1.01(1)	-0.047(4)	0.835(4)	0.005(9)	Uiso 1.00 calc ...
H(9)	H	1.03(1)	0.038(4)	0.852(3)	0.002(9)	Uiso 1.00 calc ...
H(10)	H	0.57(1)	0.012(4)	0.864(3)	-0.008(9)	Uiso 1.00 calc ...
H(11)	H	0.61(1)	-0.078(5)	0.855(4)	0.011(9)	Uiso 1.00 calc ...
H(12)	H	0.72(1)	-0.018(3)	0.916(3)	-0.007(9)	Uiso 1.00 calc ...
H(13)	H	0.99(1)	-0.062(4)	0.608(3)	0.002(9)	Uiso 1.00 calc ...
H(14)	H	0.362(9)	-0.209(3)	0.622(2)	-0.034(9)	Uiso 1.00 calc ...
H(15)	H	0.56(1)	-0.097(3)	0.668(3)	-0.011(9)	Uiso 1.00 calc ...
H(16)	H	0.32(2)	-0.121(6)	0.534(5)	0.047(6)	Uiso 1.00 calc ...
H(17)	H	0.52(1)	-0.304(3)	0.559(2)	-0.028(8)	Uiso 1.00 calc ...
H(18)	H	0.27(3)	-0.294(7)	0.558(7)	0.063(5)	Uiso 1.00 calc ...

loop_

_atom_site_aniso_label

_atom_site_aniso_U_11

_atom_site_aniso_U_22

_atom_site_aniso_U_33

_atom_site_aniso_U_12

_atom_site_aniso_U_13

_atom_site_aniso_U_23

Cl(1)	0.0219(6)	0.0172(7)	0.0287(8)	0.0004(6)	-0.0008(6)	0.0002(7)
F(4)	0.040(2)	0.046(3)	0.045(3)	0.007(2)	-0.007(2)	0.016(3)
F(5)	0.038(2)	0.041(3)	0.037(3)	-0.024(2)	0.002(2)	0.009(2)
F(6)	0.067(4)	0.043(3)	0.021(2)	-0.020(3)	0.004(2)	-0.006(2)
O(3)	0.014(2)	0.020(2)	0.031(3)	-0.002(2)	-0.002(2)	-0.006(2)
O(5)	0.026(2)	0.056(4)	0.031(3)	0.004(3)	0.001(2)	-0.008(3)
O(6)	0.023(2)	0.043(3)	0.030(2)	0.006(2)	0.001(2)	-0.013(3)
N(1)	0.008(2)	0.023(3)	0.031(3)	0.001(2)	0.001(2)	0.001(2)

S24

N(2)	0.020(2)	0.023(3)	0.021(3)	-0.007(2)	0.006(2)	-0.006(2)
C(1)	0.015(2)	0.018(2)	0.018(3)	-0.002(2)	-0.002(2)	0.004(2)
C(2)	0.029(3)	0.012(2)	0.028(3)	-0.005(2)	0.003(3)	-0.005(2)
C(4)	0.025(2)	0.013(3)	0.027(3)	-0.002(2)	0.005(2)	0.008(3)
C(5)	0.031(3)	0.031(3)	0.028(3)	-0.012(2)	0.003(3)	-0.005(2)
C(6)	0.024(3)	0.018(3)	0.028(4)	0.001(2)	0.009(3)	0.000(3)
C(7)	0.024(3)	0.024(3)	0.021(3)	-0.007(2)	-0.008(2)	0.006(2)
C(8)	0.012(3)	0.016(3)	0.036(3)	0.004(2)	0.002(3)	-0.001(2)
C(9)	0.032(3)	0.041(4)	0.029(3)	0.003(4)	0.000(3)	-0.007(4)
C(10)	0.041(4)	0.040(4)	0.022(4)	0.005(4)	0.005(3)	-0.005(4)
C(11)	0.044(4)	0.024(3)	0.038(4)	0.001(3)	0.007(4)	-0.004(3)
C(12)	0.035(4)	0.033(4)	0.023(3)	0.004(3)	-0.001(3)	0.001(3)
C(13)	0.029(3)	0.042(4)	0.054(5)	0.010(3)	0.010(4)	0.020(3)
C(14)	0.035(4)	0.035(4)	0.065(5)	-0.009(4)	-0.005(4)	0.025(3)
C(15)	0.054(5)	0.026(4)	0.054(5)	-0.015(4)	-0.009(4)	0.001(4)

#-----

_computing_data_collection	'PROCESS-AUTO'
_computing_cell_refinement	'PROCESS-AUTO'
_computing_data_reduction	'teXsan Ver. 1.11'
_computing_structure_solution	SIR97
_computing_structure_refinement	'teXsan Ver. 1.10'
_computing_publication_material	'teXsan Ver. 1.11'
_computing_molecular_graphics	?

#-----

_geom_special_details

?

;

loop_

_geom_bond_atom_site_label_1

_geom_bond_atom_site_label_2

_geom_bond_distance

_geom_bond_site_symmetry_1

_geom_bond_site_symmetry_2

_geom_bond_publ_flag

F(4) C(5) 1.350(10) .. yes

S25

F(5) C(5) 1.348(8) .. yes

F(6) C(5) 1.327(9) .. yes

O(3) C(1) 1.232(8) .. yes

O(5) C(4) 1.208(10) .. yes

O(6)	C(4)	1.314(9)	.. yes
O(6)	C(9)	1.45(1)	.. yes
N(1)	C(8)	1.47(1)	.. yes
N(2)	C(1)	1.326(9)	.. yes
N(2)	C(2)	1.454(9)	.. yes
C(1)	C(8)	1.513(9)	.. yes
C(2)	C(4)	1.51(1)	.. yes
C(2)	C(6)	1.54(1)	.. yes
C(5)	C(8)	1.50(1)	.. yes
C(6)	C(7)	1.498(9)	.. yes
C(7)	C(11)	1.41(1)	.. yes
C(7)	C(12)	1.37(1)	.. yes
C(9)	C(10)	1.49(1)	.. yes
C(11)	C(15)	1.40(1)	.. yes
C(12)	C(13)	1.39(1)	.. yes
C(13)	C(14)	1.39(2)	.. yes
C(14)	C(15)	1.37(2)	.. yes

#-----

loop_

_geom_angle_atom_site_label_1

_geom_angle_atom_site_label_2

_geom_angle_atom_site_label_3

_geom_angle

_geom_angle_site_symmetry_1

_geom_angle_site_symmetry_2

_geom_angle_site_symmetry_3

_geom_angle_publ_flag

C(4)	O(6)	C(9)	120.1(6)	... yes
------	------	------	----------	---------

C(1)	N(2)	C(2)	122.4(6)	... yes
------	------	------	----------	---------

O(3)	C(1)	N(2)	124.7(6)	... yes
------	------	------	----------	---------

O(3)	C(1)	C(8)	119.4(6)	... yes
------	------	------	----------	---------

S26

N(2)	C(1)	C(8)	115.9(6)	... yes
------	------	------	----------	---------

N(2)	C(2)	C(4)	109.4(6)	... yes
------	------	------	----------	---------

N(2)	C(2)	C(6)	109.9(6)	... yes
------	------	------	----------	---------

C(4)	C(2)	C(6)	113.1(6)	... yes
------	------	------	----------	---------

O(5)	C(4)	O(6)	122.2(8)	... yes
------	------	------	----------	---------

O(5)	C(4)	C(2)	125.4(7)	... yes
------	------	------	----------	---------

O(6)	C(4)	C(2)	112.3(6)	... yes
------	------	------	----------	---------

F(4)	C(5)	F(5)	106.7(6)	... yes
------	------	------	----------	---------

F(4)	C(5)	F(6)	107.1(7)	... yes
------	------	------	----------	---------

F(4)	C(5)	C(8)	111.5(7)	... yes
F(5)	C(5)	F(6)	108.3(7)	... yes
F(5)	C(5)	C(8)	110.7(6)	... yes
F(6)	C(5)	C(8)	112.4(7)	... yes
C(2)	C(6)	C(7)	114.1(6)	... yes
C(6)	C(7)	C(11)	118.9(7)	... yes
C(6)	C(7)	C(12)	121.2(7)	... yes
C(11)	C(7)	C(12)	119.9(7)	... yes
N(1)	C(8)	C(1)	109.5(6)	... yes
N(1)	C(8)	C(5)	112.2(6)	... yes
C(1)	C(8)	C(5)	112.4(6)	... yes
O(6)	C(9)	C(10)	108.7(7)	... yes
C(7)	C(11)	C(15)	119.1(9)	... yes
C(7)	C(12)	C(13)	121.3(8)	... yes
C(12)	C(13)	C(14)	118.9(9)	... yes
C(13)	C(14)	C(15)	120.6(8)	... yes
C(11)	C(15)	C(14)	120.2(10)	... yes

#-----

loop_

_geom_torsion_atom_site_label_1

_geom_torsion_atom_site_label_2

_geom_torsion_atom_site_label_3

_geom_torsion_atom_site_label_4

_geom_torsion

_geom_torsion_site_symmetry_1

_geom_torsion_site_symmetry_2

S27

_geom_torsion_site_symmetry_3

_geom_torsion_site_symmetry_4

_geom_torsion_publ_flag

F(4)	C(5)	C(8)	N(1)	67.0(8)	 yes
F(4)	C(5)	C(8)	C(1)	-56.8(9)	 yes
F(5)	C(5)	C(8)	N(1)	-51.6(9)	 yes
F(5)	C(5)	C(8)	C(1)	-175.4(6)	 yes
F(6)	C(5)	C(8)	N(1)	-172.7(7)	 yes
F(6)	C(5)	C(8)	C(1)	63.4(9)	 yes
O(3)	C(1)	N(2)	C(2)	4(1)	 yes
O(3)	C(1)	C(8)	N(1)	-30.9(9)	 yes
O(3)	C(1)	C(8)	C(5)	94.4(8)	 yes
O(5)	C(4)	O(6)	C(9)	-4(1)	 yes
O(5)	C(4)	C(2)	N(2)	123.6(8)	 yes

O(5)	C(4)	C(2)	C(6)	-113.6(9)	yes
O(6)	C(4)	C(2)	N(2)	-53.7(8)	yes
O(6)	C(4)	C(2)	C(6)	69.1(8)	yes
N(1)	C(8)	C(1)	N(2)	149.8(7)	yes
N(2)	C(1)	C(8)	C(5)	-84.9(8)	yes
N(2)	C(2)	C(6)	C(7)	-173.4(6)	yes
C(1)	N(2)	C(2)	C(4)	-111.6(7)	yes
C(1)	N(2)	C(2)	C(6)	123.7(7)	yes
C(2)	N(2)	C(1)	C(8)	-176.8(7)	yes
C(2)	C(4)	O(6)	C(9)	173.1(7)	yes
C(2)	C(6)	C(7)	C(11)	-111.3(8)	yes
C(2)	C(6)	C(7)	C(12)	71.2(9)	yes
C(4)	O(6)	C(9)	C(10)	-158.9(7)	yes
C(4)	C(2)	C(6)	C(7)	64.0(8)	yes
C(6)	C(7)	C(11)	C(15)	-178.9(8)	yes
C(6)	C(7)	C(12)	C(13)	177.7(8)	yes
C(7)	C(11)	C(15)	C(14)	3(1)	yes
C(7)	C(12)	C(13)	C(14)	0(1)	yes
C(11)	C(7)	C(12)	C(13)	0(1)	yes
C(11)	C(15)	C(14)	C(13)	-3(1)	yes
C(12)	C(7)	C(11)	C(15)	-1(1)	yes

S28

C(12)	C(13)	C(14)	C(15)	2(1)	yes
C(12)	C(13)	C(14)	C(15)	2(1)	yes

#-----

loop_

_geom_contact_atom_site_label_1

_geom_contact_atom_site_label_2

_geom_contact_distance

_geom_contact_site_symmetry_1

_geom_contact_site_symmetry_2

_geom_contact_publ_flag

Cl(1)	N(1)	3.136(7)	. 3_446 ?
-------	------	----------	-----------

Cl(1)	N(1)	3.191(7)	.. ?
-------	------	----------	------

Cl(1)	C(1)	3.227(7)	.. ?
-------	------	----------	------

Cl(1)	O(3)	3.238(5)	. 3_446 ?
-------	------	----------	-----------

Cl(1)	N(1)	3.240(6)	. 3_546 ?
-------	------	----------	-----------

Cl(1)	C(8)	3.420(9)	.. ?
-------	------	----------	------

Cl(1)	O(3)	3.480(5)	.. ?
-------	------	----------	------

Cl(1)	N(2)	3.594(7)	.. ?
-------	------	----------	------

F(4)	F(5)	3.011(7)	. 1_655 ?
------	------	----------	-----------

F(5)	C(9)	3.25(1)	. 4_646 ?
F(5)	O(6)	3.323(8)	. 4_646 ?
F(5)	O(3)	3.513(7)	. 1_455 ?
F(5)	C(10)	3.53(1)	. 4_646 ?
F(5)	C(11)	3.55(1)	. 4_646 ?
F(6)	O(5)	3.252(9)	. 1_455 ?
F(6)	C(15)	3.37(1)	. 4_746 ?
O(3)	N(1)	2.865(9)	. 3_546 ?
O(3)	C(8)	3.103(9)	. 1_655 ?
O(3)	N(1)	3.154(9)	. 1_655 ?
O(5)	N(2)	2.855(9)	. 1_655 ?
O(5)	O(6)	3.252(8)	. 1_655 ?
C(6)	C(13)	3.39(1)	. 1_455 ?

#-----

loop_

_geom_hbond_atom_site_label_D

S29

_geom_hbond_atom_site_label_H

_geom_hbond_atom_site_label_A

_geom_hbond_site_symmetry_D

_geom_hbond_site_symmetry_H

_geom_hbond_site_symmetry_A

_geom_hbond_site_distance_DH

_geom_hbond_site_distance_HA

_geom_hbond_site_distance_DA

_geom_hbond_angle_DHA

_geom_hbond_publ_flag

N(1)	H(17)	F(4)	...	0.66(5)	2.73(4)	2.928(8)	100(4)	no
N(1)	H(17)	O(3)	...	0.66(5)	2.52(4)	2.706(7)	98(4)	no
N(2)	H(15)	F(6)	...	0.71(5)	2.69(5)	3.055(7)	114(4)	no
N(2)	H(15)	O(6)	...	0.71(5)	2.64(5)	2.759(8)	92(4)	no
F(4)	O(3)	3.130(7)	...	-0.2830	0.6586	0.7268	92(4)	no
F(4)	N(1)	2.928(8)	...	-0.3075	0.6490	0.5774	92(4)	no
F(5)	N(1)	2.773(8)	...	-0.3094	0.6446	0.3054	92(4)	no
F(6)	N(2)	3.055(7)	...	-0.1870	0.7117	0.5147	92(4)	no
O(3)	F(4)	3.130(7)	...	-0.2393	0.6187	0.8028	92(4)	no
O(3)	N(1)	2.706(7)	...	-0.2300	0.5756	0.7034	92(4)	no
N(1)	F(5)	2.773(8)	...	-0.2966	0.6087	0.3550	92(4)	no
N(1)	H(17)	Cl(1)	..	3_566	0.66(5)	2.48(5)	3.136(6)	169(4) no
N(1)	H(18)	O(3)	..	1_455	0.8(1)	2.6(1)	3.154(7)	123(>>) no
N(2)	H(15)	O(5)	..	1_455	0.71(5)	2.17(5)	2.855(8)	162(5) no

#-----

S30