A One-Step Synthesis of 2,3-Dihydro-4*H*-pyran-4-ones from 3-Ethoxy- α,β -Unsaturated Lactones

Jeffrey D. Winkler* and Kyungsoo Oh

Department of Chemistry, University of Pennsylvania, Philadelphia, PA 19104

winkler@sas.upenn.edu

Table of Contents:

Experimental Section	S02-S10
¹ H NMR Spectrum of 2a	S11
¹³ C NMR Spectrum of 2a	S12
¹ H NMR Spectrum of 2b	S13
¹³ C NMR Spectrum of 2b	S14
¹ H NMR Spectrum of 2c	S15
¹³ C NMR Spectrum of 2c	S16
¹ H NMR Spectrum of 2d	S17
¹³ C NMR Spectrum of 2d	S18
¹ H NMR Spectrum of 2e <i>threo</i>	S19
¹³ C NMR Spectrum of 2e <i>threo</i>	S20
¹ H NMR Spectrum of 2e <i>erythro</i>	S21
¹³ C NMR Spectrum of 2e <i>erythro</i>	S22
¹ H NMR Spectrum of 2f	S23
¹³ C NMR Spectrum of 2f	S24
¹ H NMR Spectrum of 3a	S25
¹³ C NMR Spectrum of 3a	S26
¹ H NMR Spectrum of 3b	S27
¹³ C NMR Spectrum of 3b	S28
¹ H NMR Spectrum of 3c	S29
¹³ C NMR Spectrum of 3c	S30
¹ H NMR Spectrum of 3d	S31
¹³ C NMR Spectrum of 3d	S32
¹ H NMR Spectrum of 3e	S33
¹³ C NMR Spectrum of 3e	S34
¹ H NMR Spectrum of 3f	S35
¹³ C NMR Spectrum of 3f	S36
¹ H NMR Spectrum of 3g	S37
¹³ C NMR Spectrum of 3g	S38

Supporting Information

Preparation of Brassard's diene¹

To a stirred solution of ethyl acetoacetate (52 g, 0.40 mol) and triethyl orthoformate (59.2 g, 0.40 mol) at room temperature was added concentrated sulfuric acid (12 drops). The reaction mixture was stirred for 24 hours at ambient temperature before quenched with quinoline (20 drops). The reaction mixture was poured into a mixture of water and diethyl ether (300 ml/300 ml) and extracted with diethyl ether (50 ml×3). The combined organic solution was washed with saturated brine and then dried over anhydrous magnesium sulfate and concentrated in vacuo. The residue was purified by flash chromatography on silica gel (eluent 100 % hexanes) to give the vinyl ether as a clear liquid (35.0 g, 55%).

3-Ethoxy-but-2-enoic acid ethyl ester

¹H NMR (500MHz, CDCl₃): 4.98 (1H, s), 4.11 (2H, q, 7.0Hz), 3.80 (2H, q, 7.0Hz), 2.28 (3H, s), 1.32 (3H, t, 7.0Hz), 1.24 (3H, t, 7.0Hz);

¹³C NMR (125MHz, CDCl₃): 172.2, 168.0, 91.0, 63.6, 59.2, 19.1, 14.4, 14.1;

(1,3-Diethoxy-buta-1,3-dienyloxy)-trimethyl-silane (1)

To a stirred solution of lithium diisopropylamide (0.25 mol) in dry tetrahydrofuran (100 ml) at -78°C, was added the vinyl ether (36.0 g, 0.23 mol) in dry tetrahydrofuran (50 ml) dropwise. The resulting mixture was stirred for 30 minutes at same temperature and trimethylsilyl chloride (37.3 ml, 0.25 mol) was added. The mixture was allowed to warm up to ambient temperature slowly then dry pentane was added to precipitate the lithium salt. After centrifuging the resulting solution, the yellowish solution was collected and the solvent was removed to give the Brassard's diene 1 as a yellowish liquid (50.0 g, 94 %).

IR (neat) 2978, 2900, 1658, 1626;

¹ a) Savard, J.; Brassard, P. *Tetrahedron Lett.* **1979**, 20, 4911-4914. b) Kaweck, R. *Tetrahedron* **2001**, 57, 8385-8390. c) Kaweck, R. *Synthesis* **2001**, 828-830. d) Fan, Q.; Lin, L.; Liu, J.; Huang, Y.; Feng, X.; Zhang, G. *Org. Lett.* **2004**, 6, 2185-2188.

^1H NMR (500MHz, CDCl_3): 4.31 (1H, s), 3.99 (1H, s), 3.94 (1H, s), 3.79 (2H, q, 7.0Hz), 3.74 (2H, q, 7.0Hz), 1.28 (6H, m), 0.25 (9H, s);

^{13}C NMR (125MHz, CDCl_3): 158.1, 157.6, 79.0, 76.0, 63.5, 62.0, 14.6, 14.3, 0.4;

General procedure for Hetero Diels-Alder reaction

To a stirred solution of heptaldehyde (0.55 ml, 3.95 mmol) in dry dichloromethane (15 ml) at -78°C , was added diethyl aluminium chloride-1 M in hexane (4.34 ml, 4.35 mmol) dropwise. After stirring for 10 minutes at -78°C , Brassard's diene (2.0 g, 8.68 mmol) was added slowly. The reaction mixture was stirred for one hour at -78°C and slowly warmed up to ambient temperature and stirred for 18 hours. The reaction mixture was re-cooled to 0°C and carefully quenched by addition of water. The mixture was partitioned between diethyl ether and water and extracted with diethyl ether (20 ml $\times$ 3). The combined organic solution was washed with saturated brine and then dried over anhydrous magnesium sulfate and concentrated in vacuo. The residue was purified by flash chromatography on silica gel (eluent 90/10 % hexanes/Ethyl Acetate) to give the pyran-2-one **2a** as a yellowish gum (0.60g, 67%).

4-Ethoxy-6-hexyl-5,6-dihydro-pyran-2-one (**2a**)

IR (neat) 2930, 2860, 1711, 1624, 1463;

^1H NMR (500MHz, CDCl_3): 5.10 (1H, d, 1.3Hz), 4.36 (1H, m), 3.94 (2H, m), 2.45 (1H, dd, 17.0Hz, 12.0Hz), 2.30 (1H, 17.0Hz, 3.8Hz), 1.80 (1H, m), 1.63 (1H, m), 1.50 (1H, m), 1.37 (3H, t, 7.0Hz), 1.28 (7H, m), 0.88 (3H, t, 7.0Hz);

^{13}C NMR (125MHz, CDCl_3): 172.1, 167.9, 90.4, 75.9, 64.7, 34.6, 33.2, 31.6, 28.9, 24.7, 22.5, 14.0, 13.9;

HRMS calcd for $\text{C}_{13}\text{H}_{23}\text{O}_3$ 227.1647, found 227.1651 (MH^+)

e.e. determination conditions: Chiralcel AS, Hexane:IPA= 90:10, flow=0.5 ml/min, t=46.2 min, 66.0 min.

4-Ethoxy-6-phenyl-5,6-dihydro-pyran-2-one (**2b**)

IR (neat) 2984, 2935, 1710, 1622, 1452;

¹H NMR (500MHz, CDCl₃): 7.41-7.26 (5H, m), 5.41 (1H, dd, 12.1 Hz, 3.8Hz), 5.20 (1H, s), 3.98 (2H, m), 2.80 (1H, dd, 17.1Hz, 12.1 Hz), 2.58 (1H, dd, 17.1Hz, 3.9Hz), 1.38 (3H, t, 7.0Hz);

¹³C NMR (125MHz, CDCl₃): 171.6, 167.0, 138.2, 128.6, 128.5, 125.9, 90.7, 77.0, 64.8, 35.1, 13.9;

HRMS calcd for C₁₃H₁₅O₃ 219.1021, found 219.1014 (MH⁺)

e.e. determination conditions: Chiralcel AD, Hexane:IPA= 90:10, flow=1.0 ml/min, t=25.2 min, 29.4 min. (90 % e.e. using Feng's conditions, 5.3% e.e. using (+)-Eu(hfc)₃ at -20 to 0 °C)²

6-Cyclohexyl-4-ethoxy-5,6-dihydro-pyran-2-one (**2c**)

IR (neat) 2927, 2854, 1709, 1623, 1448;

¹H NMR (500MHz, CDCl₃): 5.05 (1H, s), 4.08 (1H, m), 3.89 (2H, m), 2.47 (1H, dd, 16.9Hz, 12.6Hz), 2.21 (1H, dd, 16.9Hz, 3.6Hz), 1.92 (1H, d, 12.2Hz), 1.73 (2H, d, 12.8Hz), 1.61 (3H, m), 1.33 (3H, t, 7.0Hz), 1.23-1.01 (5H, m);

¹³C NMR (125MHz, CDCl₃): 172.2, 167.6, 90.3, 79.7, 64.5, 41.3, 30.4, 28.2, 27.9, 26.1, 25.7, 25.6, 13.8;

HRMS calcd for C₁₃H₂₁O₃ 225.1491, found 225.1500 (MH⁺)

² Fan, Q.; Lin, L.; Liu, J.; Huang, Y.; Feng, X.; Zhang, G. *Org. Lett.* **2004**, *6*, 2185-2188.

6-*tert*-Butyl-4-ethoxy-5,6-dihydro-pyran-2-one (**2d**)

IR (neat) 2962, 1710, 1624, 1475;

^1H NMR (500MHz, CDCl_3): 5.04 (1H, dd, 7.1Hz, 1.7Hz), 3.97 (1H, m), 3.91 (2H, m), 2.47 (1H, m), 2.18 (1H, m), 1.33 (3H, q, 7.0Hz), 0.95 (9H, m);

^{13}C NMR (125MHz, CDCl_3): 172.4, 167.7, 90.2, 83.0, 64.6, 33.5, 28.5, 25.3, 13.8;

HRMS calcd for $\text{C}_{11}\text{H}_{19}\text{O}_3$ 199.1334, found 199.1335 (MH^+)

6-(2-Benzyloxy-octyl)-4-ethoxy-5,6-dihydro-pyran-2-one (**2e**)

IR (neat) 2928, 2859, 1709, 1623, 1456;

^1H NMR (500MHz, CDCl_3): 2:1 mixture of threo/erythro threo: 7.53-7.25 (5H, m), 5.09 (1H, s), 4.61 (1H, d, 11.1Hz), 4.58 (1H, m), 4.47 (1H, d, 11.1Hz), 3.93 (2H, m), 3.84 (1H, m), 2.44 (1H, dd, 17.0Hz, 12.0Hz), 2.30 (1H, dd, 17.0Hz, 3.9Hz), 1.90 (1H, m), 1.70 (1H, m), 1.62 (1H, m), 1.42 (1H, m), 1.36 (3H, t, 7.0Hz), 1.33-1.25 (8H, m), 0.88 (3H, t, 6.8Hz); erythro: 7.35-7.25 (5H, m), 5.08 (1H, s), 4.56 (1H, m), 4.54 (1H, d, 11.6Hz), 4.45 (1H, d, 11.6Hz), 3.93 (2H, m), 3.62 (1H, m), 2.45 (1H, m), 2.26 (1H, m), 2.10 (1H, m), 1.80 (1H, m), 1.60 (2H, m), 1.36 (3H, t, 7.0Hz), 1.35-1.24 (8H, m), 0.87 (3H, t, 6.8Hz);

^{13}C NMR (125MHz, CDCl_3): threo: 172.0, 167.4, 138.6, 128.3, 127.8, 127.5, 90.4, 75.0, 72.8, 71.8, 64.6, 40.4, 34.2, 33.7, 31.7, 29.4, 24.7, 22.5, 14.0, 13.9; erythro: 172.0, 167.4, 128.3, 127.8, 127.7, 127.5, 90.4, 74.9, 73.2, 70.6, 64.6, 38.6, 33.4, 33.2, 31.7, 29.3, 24.9, 22.5, 14.0, 13.9;

HRMS calcd for $\text{C}_{22}\text{H}_{33}\text{O}_4$ 361.2379, found 361.2387 (MH^+)

4-Ethoxy-1-oxa-spiro[5.5]undec-3-en-2-one (**2f**)

IR (neat) 2935, 2861, 1705, 1628, 1447;

^1H NMR (500MHz, CDCl_3): 5.07 (1H, s), 3.92 (2H, q, 7.0Hz), 2.41 (2H, s), 1.94 (2H, m), 1.75 (2H, m), 1.54-1.45 (6H, m), 1.37 (3H, t, 7.0Hz);

^{13}C NMR (125MHz, CDCl_3): 170.3, 168.8, 89.9, 79.0, 64.4, 38.0, 36.1, 25.5, 21.5, 13.9;

HRMS calcd for $\text{C}_{12}\text{H}_{19}\text{O}_3$ 211.1334, found 211.1334 (MH^+)

General procedure for the preparation of pyran-3-ones

To a stirred solution of lithium diisopropylamide (2.65 mmol) in dry tetrahydrofuran (6 ml) at -78°C , was added ethyl isobutyrate (0.35 ml, 2.65 mmol) dropwise. The resulting mixture was stirred for one hour at -78°C then the pyran-2-one (0.30 g, 1.33 mmol) in dry tetrahydrofuran (10 ml) was added via a cannula. The resulting solution was stirred for two hours then rapidly warmed up to -20 - 30°C then water was added. The mixture was partitioned between diethyl ether and water and extracted with diethyl ether (10 ml $\times$ 3). The combined organic solution was washed with saturated brine and then dried over anhydrous magnesium sulfate and concentrated in vacuo. The residue was purified by flash chromatography on silica gel (eluent 70/30 % hexanes/diethyl ether) to give the pyran-3-one **3g** as a brownish liquid (0.36g, 92%).

2-(6-Hexyl-4-oxo-5,6-dihydro-4H-pyran-2-yl)-2-methyl-propionic acid ethyl ester (**3g**)

IR (neat) 2931, 2862, 1738, 1672, 1605, 1464;

^1H NMR (500MHz, CDCl_3): 5.45 (1H, s), 4.31 (1H, m), 4.13 (2H, m), 2.43 (1H, dd, 16.9Hz, 12.4Hz), 2.38 (1H, dd, 16.9Hz, 4.5Hz), 1.75 (1H, m), 1.65 (1H, m), 1.40 (3H, s), 1.37 (3H, s), 1.50-1.25 (8H, m), 1.24 (3H, t, 7.1Hz), 0.88 (3H, t, 6.8Hz);

^{13}C NMR (125MHz, CDCl_3): 193.6, 178.3, 173.5, 102.3, 79.8, 61.2, 47.9, 41.2, 34.2, 31.6, 28.9, 24.8, 23.2, 23.0, 22.5, 14.1, 14.0;

HRMS calcd for $\text{C}_{17}\text{H}_{29}\text{O}_4$ 297.2066, found 297.2056 (MH^+)

6-Butyl-2-hexyl-2,3-dihydro-pyran-4-one (3a)

IR (neat) 2930, 2864, 1671, 1606, 1461, 1400;

^1H NMR (500MHz, CDCl_3): 5.30 (1H, s), 4.32 (1H, m), 2.39 (1H, dd, 16.5Hz, 12.0Hz), 2.35 (1H, dd, 16.5Hz, 5.0Hz), 2.22 (2H, m), 1.80 (1H, m), 1.65 (1H, m), 1.54-1.30 (12H, m), 0.91 (3H, t, 7.3Hz), 0.89 (3H, t, 6.9Hz);

^{13}C NMR (125MHz, CDCl_3): 193.2, 177.9, 103.9, 79.2, 41.0, 34.4, 34.4, 31.6, 28.9, 28.4, 24.8, 22.5, 22.1, 14.0, 13.7;

HRMS calcd for $\text{C}_{15}\text{H}_{27}\text{O}_2$ 239.2011, found 239.2009 (MH^+)

6-Allyl-2-hexyl-2,3-dihydro-pyran-4-one (3b)

IR (neat) 2927, 2857, 1671, 1606, 1394;

^1H NMR (500MHz, CDCl_3): 5.80 (1H, m), 5.21 (1H, s), 5.18 (1H, m), 5.15 (1H, m), 4.34 (1H, m), 2.98 (2H, d, 6.8), 2.41 (1H, dd, 16.9Hz, 11.8Hz), 2.36 (1H, dd, 16.9Hz, 3.8Hz), 1.80 (1H, m), 1.67 (2H, m), 1.50-1.29 (7H, m), 0.88 (3H, t, 6.7Hz);

^{13}C NMR (125MHz, CDCl_3): 193.1, 175.4, 131.2, 118.7, 104.1, 79.4, 41.0, 38.9, 34.3, 31.6, 28.9, 24.7, 22.4, 13.9;

HRMS calcd for $\text{C}_{14}\text{H}_{23}\text{O}_2$ 223.1698, found 223.1687 (MH^+)

2-Hexyl-6-vinyl-2,3-dihydro-pyran-4-one (3c)

IR (neat) 2928, 2860, 1666, 1633, 1566, 1461, 1411;

^1H NMR (500MHz, CDCl_3): 6.20 (1H, dd, 17.2Hz, 10.6Hz), 6.04 (1H, dd, 17.2Hz, 1.3Hz), 5.53 (1H, dd, 10.6Hz, 0.9Hz), 5.39 (1H, s), 4.39 (1H, m), 2.48 (1H, dd, 16.8Hz, 12.0Hz), 2.43 (1H, ddd, 16.8Hz, 4.6Hz, 0.8Hz), 1.86 (1H, m), 1.68 (1H, m), 1.52 (1H, m), 1.45 (1H, m), 1.37-1.25 (6H, m), 0.89 (3H, t, 6.8Hz);

^{13}C NMR (125MHz, CDCl_3): 193.8, 167.7, 131.2, 122.7, 105.8, 79.1, 41.6, 34.5, 31.6, 29.0, 24.9, 22.5, 14.0;

HRMS calcd for $\text{C}_{13}\text{H}_{21}\text{O}_2$ 209.1541, found 209.1536 (MH^+)

2-Hexyl-6-phenyl-2,3-dihydro-pyran-4-one (3d)

IR (neat) 3064, 2928, 2859, 1663, 1599, 1571, 1491, 1451;

^1H NMR (500MHz, CDCl_3): 7.73 (2H, m), 7.48-7.41 (3H, m), 6.00 (1H, s), 4.53 (1H, m), 2.56 (1H, dd, 16.7Hz, 13.0Hz), 2.50 (1H, dd, 16.7Hz, 3.3Hz), 1.96 (1H, m), 1.76 (1H, m), 1.61 (1H, m), 1.51 (1H, m), 1.41-1.25 (6H, m), 0.91 (3H, t, 6.9Hz);

^{13}C NMR (125MHz, CDCl_3): 193.5, 170.2, 132.8, 131.5, 128.6, 126.4, 101.9, 79.6, 41.4, 34.5, 31.6, 29.0, 25.0, 22.5, 14.0;

HRMS calcd for $\text{C}_{17}\text{H}_{23}\text{O}_2$ 259.1698, found 259.1705 (MH^+)

2-Hexyl-6-isopropenyl-2,3-dihydro-pyran-4-one (3e)

IR (neat) 2929, 2860, 1667, 1573, 1459;

^1H NMR (500MHz, CDCl_3): 5.85 (1H, s), 5.56 (1H, s), 5.33 (1H, s), 4.36 (1H, m), 2.46 (1H, dd, 16.8Hz, 12.1Hz), 2.41 (1H, dd, 16.8Hz, 4.6Hz), 1.91 (3H, s), 1.85 (1H, m), 1.66 (1H, m), 1.51 (1H, m), 1.44 (1H, m), 1.37-1.30 (6H, m), 0.89 (3H, t, 6.8Hz);

^{13}C NMR (125MHz, CDCl_3): 194.2, 169.7, 137.0, 120.3, 103.0, 79.0, 41.4, 34.5, 31.6, 28.9, 25.0, 22.5, 18.7, 14.0;

HRMS calcd for $\text{C}_{14}\text{H}_{23}\text{O}_2$ 223.1698, found 223.1693 (MH^+)

2-Hexyl-6-(2-methyl-propenyl)-2,3-dihydro-pyran-4-one (3f)

IR (neat) 2928, 2860, 1641, 1564, 1454, 1403;

^1H NMR (500MHz, CDCl_3): 5.66 (1H, m), 5.29 (1H, s), 4.37 (1H, m), 2.44 (1H, dd, 16.7Hz, 13.2Hz), 2.04 (3H, d, 0.8Hz), 1.89 (1H, d, 1.0Hz), 1.84 (1H, m), 1.67 (1H, m), 1.50 (1H, m), 1.42-1.24 (7H, m), 0.88 (3H, t, 6.8Hz);

^{13}C NMR (125MHz, CDCl_3): 193.5, 170.7, 148.7, 119.5, 105.5, 79.2, 41.2, 34.6, 31.6, 28.9, 28.5, 25.0, 22.5, 21.0, 14.0;

HRMS calcd for $\text{C}_{15}\text{H}_{25}\text{O}_2$ 237.1855, found 237.1858 (MH^+)

Asymmetric aldol reaction in the presence of the Jacobsen's catalyst

To a stirred solution of pivaldehyde (36 μl , 0.33 mmol) in dry acetone (4ml) and 4Å powered molecular sieves was added the Jacobsen's catalyst³ (25 mg, 25 mol %) in one portion. The resulting mixture was stirred for 1 hour at ambient temperature then furan (100 mg, 0.42 mmol) was added dropwise. The resulting mixture was stirred at ambient temperature for 7 days and the solvent was removed under reduced pressure. The resulting yellowish oil was purified by flash chromatography on silica gel (eluent 93/7 %

³ Dossetter, A. G.; Jamison, T. F.; Jacobsen, E. N. *Angew. Chem. Int. Ed.* **1999**, 38, 2398-2400.

hexanes/diethyl ether) to give the furan-3-one⁴ as a brownish liquid (0.018g, 20%, 97 % e.e.).

IR (neat) 3406, 2957, 2867, 1686, 1593, 1466;

¹H NMR (500MHz, CDCl₃): 5.47 (1H, s), 4.56 (1H, s), 3.78 (1H, d, 7.2Hz), 2.53 (2H, t, 7.5Hz), 1.73 (1H, d, 7.8 Hz), 1.66 (2H, m), 1.37 (2H, m), 1.30 (4H, m), 1.04 (9H,s), 0.89 (3H, t, 6.8Hz);

¹³C NMR (125MHz, CDCl₃): 204.9, 196.0, 104.3, 85.6, 78.3, 35.4, 31.7, 31.0, 28.9, 26.6, 26.1, 22.6, 14.1;

HRMS calcd for C₁₅H₂₆O₃Na 277.1780, found 277.1777 (MNa⁺)

e.e. determination conditions: Chiralcel OD-H, Hexane:IPA= 97:3, flow=0.7 ml/min, t=14.3 min, 16.5 min.

⁴ Winkler, J. D.; Oh, K.; Asselin, S. M. *Org. Lett.* **2005**, 7, 387.

CCCCCCCC1C(=O)OC(=C)C1OCC
 4-Ethoxy-6-hexyl-5,6-dihydro-2H-pyran-2-one

5.0
4.5
4.0
3.5
3.0
2.5
2.0
1.5
1.0
ppm

4-Ethoxy-6-hexyl-5,6-dihydro-pyran-2-one (**2a**)

¹³C NMR Spectrum of 2a

4-Ethoxy-6-hexyl-5,6-dihydro-pyran-2-one (**2a**)

¹H NMR Spectrum of 2b

¹³C NMR Spectrum of 2b

4-Ethoxy-6-phenyl-5,6-dihydro-pyran-2-one (**2b**)

¹H NMR Spectrum of 2c

6-Cyclohexyl-4-ethoxy-5,6-dihydro-pyran-2-one (2c)

¹³C NMR Spectrum of 2c

6-Cyclohexyl-4-ethoxy-5,6-dihydro-pyran-2-one (**2c**)

¹H NMR Spectrum of 2d

6-*tert*-Butyl-4-ethoxy-5,6-dihydro-pyran-2-one (2d)

^{13}C NMR Spectrum of 2d

6-*tert*-Butyl-4-ethoxy-5,6-dihydro-pyran-2-one (2d)

¹H NMR Spectrum of 2e *threo*

^{13}C NMR Spectrum of **2e** *threo*

¹H NMR Spectrum of 2e *erythro*

^{13}C NMR Spectrum of 2e *erythro*

6-(2-Benzyloxy-octyl)-4-ethoxy-5,6-dihydro-pyran-2-one (2e)

¹H NMR Spectrum of 2f

4-Ethoxy-1-oxa-spiro[5.5]undec-3-en-2-one (2f)

¹³C NMR Spectrum of 2f

4-Ethoxy-1-oxa-spiro[5.5]undec-3-en-2-one (**2f**)

¹H NMR Spectrum of 3a

6-Butyl-2-hexyl-2,3-dihydro-pyran-4-one (**3a**)

^{13}C NMR Spectrum of 3a

6-Butyl-2-hexyl-2,3-dihydro-pyran-4-one (**3a**)

¹H NMR Spectrum of 3b

6-Allyl-2-hexyl-2,3-dihydro-pyran-4-one (3b)

^{13}C NMR Spectrum of 3b

6-Allyl-2-hexyl-2,3-dihydro-pyran-4-one (**3b**)

¹H NMR Spectrum of 3c

2-Hexyl-6-vinyl-2,3-dihydro-pyran-4-one (3c)

¹³C NMR Spectrum of 3c

¹H NMR Spectrum of 3d

2-Hexyl-6-phenyl-2,3-dihydro-pyran-4-one (**3d**)

¹³C NMR Spectrum of 3d

2-Hexyl-6-phenyl-2,3-dihydro-pyran-4-one (**3d**)

¹H NMR Spectrum of 3e

2-Hexyl-6-isopropenyl-2,3-dihydro-pyran-4-one (**3e**)

^{13}C NMR Spectrum of 3e

2-Hexyl-6-isopropenyl-2,3-dihydro-pyran-4-one (**3e**)

¹H NMR Spectrum of 3f

2-Hexyl-6-(2-methyl-propenyl)-2,3-dihydro-pyran-4-one (**3f**)

^{13}C NMR Spectrum of 3f

2-Hexyl-6-(2-methyl-propenyl)-2,3-dihydro-pyran-4-one (**3f**)

¹H NMR Spectrum of 3g

2-(6-Hexyl-4-oxo-5,6-dihydro-4*H*-pyran-2-yl)
-2-methyl-propionic acid ethyl ester (**3g**)

¹³C NMR Spectrum of 3g

2-(6-Hexyl-4-oxo-5,6-dihydro-4*H*-pyran-2-yl)
-2-methyl-propionic acid ethyl ester (**3g**)