A Pentagonal Pyramidal Zr Imido Complex for the Hydroamination of Unactivated Alkenes

Robert K. Thomson, Jason A. Bexrud, and Laurel L. Schafer*

*Department of Chemistry, University of British Columbia, 6174 University Blvd.,
Vancouver, BC, CANADA, V6T 1Z1*

Supporting Information

Experimental

Unless otherwise specified, all reactions and procedures were carried out using standard Schlenk techniques or in a glovebox (Mbraun LabMaster) under an atmosphere of dry, oxygen free nitrogen. Diethyl ether, THF, hexanes, pentane, and toluene were purchased in anhydrous form from Aldrich, sparged with nitrogen, and passed through columns containing activated alumina and Ridox catalyst. C_6D_6 , D_8 -toluene, and $CDCl_3$ were purchased from Cambridge Isotopes and were degassed via successive freeze-pump-thaw cycles, and dried over molecular sieves prior to use. $Zr(NMe_2)_4$, $Ti(NMe_2)_4$, and $Ti(NEt_2)_4$ were purchased from Strem and used as received. Benzoyl chloride and 2,6-diisopropylaniline were purchased from Aldrich and used as received. 2,6-Dimethylaniline was purchased from Aldrich, distilled from molecular sieves under nitrogen, and degassed by freeze-pump-thaw cycles prior to use. Triphenylphosphine oxide was purchased from Aldrich and sublimed prior to use. All aminoalkene hydroamination substrates were synthesized according to literature procedures, and distilled from CaH_2 or molecular sieves, degassed via freeze-pump-thaw cycles, and stored over molecular sieves prior to use.¹ All other chemicals were purchased from Aldrich and used as received. 1H , $^{13}C\{^1H\}$, and $^{31}P\{^1H\}$ NMR spectra were recorded on Bruker AVANCE 300 or 400 instruments operating at 300.0 and 400.0 MHz for 1H , respectively. Elemental analyses and electron impact mass spectrometry (EIMS) were performed at the Department of Chemistry analysis facilities at the University of British Columbia. GCMS data were recorded on an Agilent Technologies 5793 Mass Selective instrument equipped with an electron impact ionization source, and quadrupolar detector.

Synthesis of $[\text{DIPP}(\text{NO})^{\text{Ph}}]_2\text{Ti}(\text{NEt}_2)_2$, **1**

This complex was synthesized as previously reported.²

Synthesis of $[\text{DIPP}(\text{NO})^{\text{Ph}}]_2\text{Zr}(\text{NMe}_2)_2$, **2**

In a 100 mL round bottomed Schlenk flask 2.00 g (7.48 mmol, 1 equiv.) of $\text{Zr}(\text{NMe}_2)_4$ was dissolved in approximately 25 mL of Et_2O . In a separate 250 mL round bottomed Schlenk flask, 4.21 g (15.0 mmol, 2 equiv.) of $[\text{DIPP}(\text{NO})^{\text{Ph}}]\text{H}$ {2,6-diisopropylphenyl(phenyl)amide} was slurried in approximately 75 mL of Et_2O . The ligand slurry was stirred and cooled to -78°C using a dry-ice/isopropanol bath. The ethereal $\text{Zr}(\text{NMe}_2)_4$ solution was transferred via cannula into the ligand slurry. The reaction mixture was allowed to warm to room temperature overnight, during which time the solution became clear and turned a pale yellow color. Excess Et_2O was removed *in vacuo* to afford a pale yellow fluffy solid. The crude product was dissolved in hexanes and filtered through Celite to remove insoluble impurities. The pale yellow filtrate was concentrated to dryness, resulting in a pale yellow solid isolated in 92% yield (5.08 g). Generally, no further purification was required; however, recrystallization from concentrated hexanes or pentane solutions resulted in analytically pure product in moderate yields. ^1H NMR (C_6D_6 , 300 MHz, 25°C): δ 0.91 (d, $J = 6.8$ Hz, $^i\text{Pr}-\text{CH}(\text{CH}_3)_2$, 12H), 1.21 (d, $J = 6.8$ Hz, $^i\text{Pr}-\text{CH}(\text{CH}_3)_2$, 12H), 3.10 (s, $-\text{N}(\text{CH}_3)_2$, 12H), 3.49 (spt, $^i\text{Pr}-\text{CH}(\text{CH}_3)_2$, 4H), 6.89 (m, Ar-*H*, 8H), 7.18 (m, Ar-*H*, 4H), 7.74 (d, Ar-*H*, 4H). ^{13}C NMR (C_6D_6 , 100 MHz, 25°C): δ 24.80, 28.74, 42.41, 124.50, 124.74, 126.30, 128.30, 130.83, 132.10, 133.14, 142.49, 177.65. Anal. Calcd. for $\text{C}_{42}\text{H}_{56}\text{N}_4\text{O}_2\text{Zr}$: C, 68.16; H, 7.63; N, 7.57. Found: C, 68.22; H, 7.69; N, 7.23.

Synthesis of $[\text{DIPP}(\text{NO})^{\text{Ph}}]_2\text{Ti}=\text{N}(\text{2,6-Me}_2\text{C}_6\text{H}_3)(\text{TPPO})$, **3**

In a 250 mL round bottomed Schlenk flask 2.00 g (2.87 mmol, 1 equiv.) of the dimethyl amido variant of **1** and 0.80 g (2.87 mmol, 1 equiv.) of triphenylphosphine oxide (TPPO) were combined, and the mixture was cooled to -78°C . To this mixture of solids was added approximately 100 mL of Et_2O . The resulting slurry was stirred rapidly and the temperature was maintained at -78°C . In a separate 25 mL Schlenk flask 0.40 mL (2.87 mmol, 1 equiv.) of 2,6-dimethylaniline was dissolved in 15 mL Et_2O . The 2,6-

dimethylaniline solution was then transferred via cannula into the reaction mixture. The dark red-orange solution was then allowed to warm to room temperature and stir overnight. During this time, all remaining TPPO was observed to dissolve, and the solution turned a dark red-brown color. Excess Et₂O was removed *in vacuo* to yield a red-brown solid. The crude product was washed with hexanes, filtered over a fritted disk, and the insoluble product was collected, and dried under vacuum. Yield = 2.30 g (80 %). ¹H NMR (C₆D₆, 300 MHz, 25°C): δ 0.81 (d, *J* = 6.5 Hz, ⁱPr-CH(CH₃)₂, 12H), 0.98 (d, *J* = 6.4 Hz, ⁱPr-CH(CH₃)₂, 6H), 1.04 (d, *J* = 6.5 Hz, ⁱPr-CH(CH₃)₂, 6H), 2.25 (s, Ar-(CH₃)₂, 6H), 3.65 (br m, ⁱPr-CH(CH₃)₂, 2H), 4.14 (br m, ⁱPr-CH(CH₃)₂, 2H), 6.62-7.22 (m, Ar-*H*, 24H), 7.74-8.00 (m, Ar-*H*, 10H). ¹³C NMR (C₆D₆, 75 MHz, 25°C): δ 19.29, 24.70, 28.23, 119.71, 124.11, 124.29, 125.62, 127.00, 127.57, 128.69, 128.87, 130.74, 132.87, 133.43, 133.57, 134.50, 143.13. ³¹P{¹H} NMR (C₆D₆, 121 MHz, 25°C): δ 41.31. EIMS (*m/z*): 887 (M⁺-NAr), 727 (M⁺-TPPO), 277 (TPPO⁺). Anal. Calcd. for C₆₄H₆₈N₃O₃PTi: C, 76.40; H, 6.81; N, 4.18. Found: C, 76.80; H, 7.04; N, 4.50.

Synthesis of [^{DIPP}(NO)^{Ph}]₂Zr=N(2,6-Me₂C₆H₃)(Ph₃P=O), **4**

In a 500 mL round bottomed Schlenk flask 5.00 g (6.76 mmol, 1 equiv.) of **2** and 1.88 g (6.76 mmol, 1 equiv.) of triphenylphosphine oxide (TPPO) were combined, and the flask was cooled to -78°C using a dry-ice/isopropanol bath before adding about 100 mL Et₂O via cannula. In a separate 100 mL Schlenk flask 0.82 g (6.76 mmol, 1 equiv.) of 2,6-dimethylaniline was dissolved in approximately 20 mL of Et₂O. To the slurry of **2** and TPPO (TPPO has low solubility in Et₂O) was added the solution of 2,6-dimethylaniline via cannula. The reaction mixture was allowed to warm to room temperature while stirring overnight. The TPPO completely dissolved and the solution changed color from pale yellow to a bright golden yellow. Excess Et₂O was removed under vacuum, and a golden yellow solid was isolated. The crude product was agitated in hexanes solution with vigorous stirring. The golden yellow insoluble product was isolated by filtration (5.75 g, 82% yield). ¹H NMR (C₆D₆, 300 MHz, 25°C): δ 0.90 (d, *J* = 6.8 Hz, ⁱPr-CH(CH₃)₂, 12H), 1.02 (br d, *J* = 4.8 Hz, ⁱPr-CH(CH₃)₂, 6H), 1.15 (br d, *J* = 6.4 Hz, ⁱPr-CH(CH₃)₂, 6H), 2.47 (s, Ar-(CH₃)₂, 6H), 3.57 (br spt, ⁱPr-CH(CH₃)₂, 2H), 3.94 (br spt, ⁱPr-CH(CH₃)₂, 2H), 6.78-7.18 (m, Ar-*H*, 24H), 7.61-7.67 (m, Ar-*H*, 10H). ¹³C

NMR (C_6D_6 , 75 MHz, 25°C): δ 19.89, 24.07, 28.73, 116.39, 123.81, 124.12, 125.44, 127.05, 127.77, 128.73, 128.91, 129.28, 129.82, 130.22, 131.01, 131.93, 132.86, 133.59. $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6 , 121 MHz, 25°C): δ 42.72. EIMS (m/z): 1047 (M^+), 930 ($\text{M}^+ - \text{NAr}$), 277 (TPPO). Anal. Calcd. for $\text{C}_{64}\text{H}_{68}\text{N}_3\text{O}_3\text{PZr}$: C, 73.25; H, 6.53; N, 4.00. Found: C, 73.65; H, 6.93; N, 4.32.

Solution phase behavior of 4

Complex 4 displays C_1 symmetry in solution, with three different isopropyl signals in a 12:6:6 ratio, indicating two separate environments for the N-aryl substituents. One of these aryl units undergoes free rotation about the N- C_{ipso} bond (as shown in Figure 1), and the other exhibits hindered rotation, giving rise to the three observed peaks. This low symmetry is also apparent in the two signals observed for the methine *isopropyl* protons at δ 3.57 and 4.00, with a very broad signal observed for the protons on the ring experiencing hindered rotation. The aryl methyl resonances on the imido moiety are equivalent on the NMR timescale and appear as a singlet at δ 2.47. The diagnostic imido C_{ipso} signal appears at δ 142 in the ^{13}C NMR spectrum. Coordination of TPPO was observed as a downfield shift to δ 42.7 in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum. It was observed that TPPO can be readily displaced upon addition of an excess of another strong donor such as THF or pyridine, and under catalytic hydroamination reaction conditions, free TPPO can be seen by ^{31}P NMR spectroscopy. Kinetic investigations imply that TPPO dissociates during the early stages of catalytic reactivity, but then acts as a competitive donor for the vacant coordination site as substrate concentration decreases over time (see Figures 18 – 21).

Figure 1 – ^1H NMR spectrum of **4**

Figure 2 – $^{13}\text{C}\{^1\text{H}\}$ APT NMR spectrum of **4**

Figure 3 – $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **4**

Representative example of hydroamination procedure (NMR yield)

All catalytic runs with NMR yields were run via the same general procedure, changing only the catalyst employed, the catalyst loading (5 or 10 percent) and the substrate utilized.

Example: *Catalytic hydroamination of 2,2-diphenyl-pent-4-enylamine to generate 2-methyl-4,4-diphenylpyrrolidine.*

150 mg (0.632 mmol) of 2,2-diphenyl-pent-4-enylamine was dissolved in approximately 0.2 mL of D_8 -toluene and loaded into a J-Young sealable NMR tube. Then 33.2 mg (0.032 mmol, 5 mol%) of $[\text{DIPP}(\text{NO})^{\text{Ph}}]_2\text{Zr}=\text{NAr}(\text{TPPO})$ (**4**) was dissolved in approximately 0.5 mL D_8 -toluene, and 35.4 mg (0.2107 mmol, 33 mol%) of 1,3,5-trimethoxybenzene (internal standard) was dissolved in about 0.3 mL D_8 -toluene. The catalyst and internal standard solutions were then transferred to the sealable NMR tube, which was then sealed and shaken vigorously before taking a $t = 0$ ^1H NMR spectrum. The tube was then placed in an oil bath at 110°C , and the reaction progress monitored periodically by ^1H NMR spectroscopy. Yields were determined by comparison of integration values for product peaks to the integration of the internal standard peaks. In most cases the methyl resonance for the internal standard was utilized for integration comparison. In cases where peak overlap with substrate or product precluded the use of

the methyl resonance of the internal standard, the aryl resonance of the internal standard was utilized. In all cases, the total volume of solvent was roughly 1 mL. NMR spectra were all collected with 32 scans and calibrated to residual solvent.

General procedure for aminoalkene hydroamination experiments in which the product was isolated.

All reactions were prepared in an N₂-filled glovebox. A small Schlenk tube equipped with a magnetic stir bar would be charged with the catalyst (0.05 mmol) and the amino olefin (1.0mmol) dissolved in toluene (~ 2 ml). The Schlenk tube would then be sealed, heated to the appropriate temperature, and stirred for the stated duration of time. Following this the solution would be concentrated under reduced pressure and directly subjected to column chromatography (CH₂Cl₂ / 5%MeOH, SiO₂) to afford the purified pyrrolidine or piperidine product.

Figure 4 - Proposed catalytic mechanism for aminoalkene cyclohydroamination:
Reaction proceeds via an imido intermediate, followed by 2 + 2 cycloaddition

Figure 5 – *Alternative catalytic mechanism for aminoalkene cyclohydroamination: Reaction proceeds via alkene insertion into M-N σ -bond*

Representative example of kinetic analysis procedure

Kinetic studies were all performed for the conversion of 2,2-diphenyl-pent-4-enylamine to 2-methyl-4,4-diphenylpyrrolidine. The only variations made were the concentration of the catalyst, or the type of catalyst used. For most kinetic runs, the experiment was repeated to ensure consistency of data (see Figures 8 & 9).

Example: *Catalytic hydroamination of 2,2-diphenyl-pent-4-enylamine to generate 2-methyl-4,4-diphenylpyrrolidine using 10 mol % of $[\text{DIPP}(\text{NO})^{\text{Ph}}]_2\text{Zr}=\text{NAr}(\text{TPPO})$ (**4**) as a catalyst.*

Here 24.6 mg of 1,3,5-trimethoxybenzene and 45.5 mg of $[\text{DIPP}(\text{NO})^{\text{Ph}}]_2\text{Zr}=\text{NAr}(\text{TPPO})$ were dissolved in approximately 0.4 mL D_8 -toluene and added to a 1 mL volumetric flask. Then 110.8 mg of 2,2-diphenyl-pent-4-enylamine was weighed into a glass vial and transferred quantitatively into the 1 mL volumetric flask with approximately 0.3 mL D_8 -toluene. The volume was then topped up to 1 mL, the solution was shaken and transferred quickly to a J-Young sealable NMR tube. The NMR tube was placed in the magnetic field of a cryoprobe equipped NMR spectrometer which had been preheated and equilibrated at 110°C. The NMR tube was allowed to thermally

equilibrate in the magnetic field for 5 minutes prior to collecting the $t = 0$ ^1H NMR spectrum. In general, ^1H NMR spectra were collected every 10 minutes for the first 3 hours, and every hour thereafter. In most cases, the initial data (first 2 to 3 hours) were sufficient for kinetic analysis. Comparison of the integration of product and substrate peaks with internal standard peaks was utilized to determine the percentage of substrate and product at any given time. These values were in turn multiplied by the initial concentration of the substrate and used to plot concentration vs. time.

Kinetic Plots

Figure 6 – Plot of [substrate] vs. time for 10% catalyst loading (4)

Figure 7 – Log plot of [substrate] vs. time for 10% catalyst loading (4)

Figure 8 – Plot of [Substrate] vs Time for 2 different runs at 10% catalyst loading with 4 as catalyst

Figure 9 – Log plot of 2 different runs at 10% catalyst loading with **4** as catalyst

Figure 10 – Plot of [substrate] vs. time for 7.5% catalyst loading (**4**)

Figure 11 – Log plot of [substrate] vs. time for 7.5% catalyst loading (4)

Figure 12 – Plot of [substrate] vs. time for 5.0% catalyst loading (4)

Figure 13 – Log plot of [substrate] vs. time for 5.0% catalyst loading (4)

Figure 14 – Plot of [substrate] vs. time for 2.5% catalyst loading (4)

Figure 15 – Log plot of [substrate] vs. time for 2.5% catalyst loading (4)

Figure 16 – Normalized substrate kinetics plots (4 as catalyst)

Figure 17 – Normalized substrate kinetics log plots (4 as catalyst)

Figure 18 – Comparison of kinetic plots for Complexes 2 and 4 (10% cat.)

Figure 19 – Comparison of kinetic log plots for Complexes **2** and **4** (10% cat.)

Figure 20 – Comparison of kinetic plots for Complexes **2** and **4** (5% cat.)

Figure 21 – Comparison of kinetic log plots for Complexes **2** and **4** (5% cat.)

Figure 22 – Order of catalyst for cyclization with complex **4**

Crystallography

Measurements on a crystal of **4** were made using a Rigaku AFC7 diffractometer fitted with an ADSC CCD area detector collecting phi and omega scans to fill the Ewald sphere. Single crystals were coated with oil and mounted under a stream of cold N₂ onto a glass fiber. Data was collected using D*TREK Area Detector Software,³ processed using the CrystalClear⁴ program, and refined using the WinGX software package⁵ and all programs included within. Full data available in CIF format at Cambridge Crystallographic Database.

Figure 23 - Solid state molecular structure of **4** (simplified structure)

Figure 24 - *Solid state molecular structure of 4 (full structure)*

Figure 25 – Space filling model of **4** viewed from front (amidate ligands = green, imido = purple, TPPO = grey)

Figure 26 – Space filling model of **4** viewed from top (amidate ligands = green, imido = purple, TPPO = grey)

Figure 27 – *Space filling model of 4 viewed from bottom (amidate ligands = green, imido = purple, TPPO = grey)*

Table 1 - *Crystallographic data^a*

	4
--	----------

empirical formula	C ₆₄ H ₆₈ N ₃ O ₃ Zr
fw	1049.44
crystal color, habit	yellow prism
crystal size (mm)	0.25x0.20x0.15
crystal system	monoclinic
space group	P21/n
a (Å)	14.0824(4)
b (Å)	20.0887(6)
c (Å)	22.2465(5)
α (deg)	90
β (deg)	99.8050(10)
γ (deg)	90
V (Å ³)	6201.5(3)
Z	4
D _{calcd} (g/cm ³)	1.405
μ (Mo Kα) (cm ⁻¹)	0.71073
no. of reflections measured	
total	13318
unique	10153
no. of observ. reflns. (I>2.0σ(I))	10153
R; R _w	0.0574; 0.1252

^aData were recorded at 173K using MoKα (λ = 0.71069Å) radiation on a Rigaku/ADSC CCD diffractometer.

Table 2 – Bond distances for 4

Bond	Distance (Å)
Zr(1)-N(3)	1.8531(18)
Zr(1)-O(1)	2.2034(15)
Zr(1)-O(3)	2.2086(14)
Zr(1)-O(2)	2.2391(15)
Zr(1)-N(2)	2.3114(18)
Zr(1)-N(1)	2.3124(18)
P(1)-O(3)	1.5122(15)
P(1)-C(51)	1.799(2)
P(1)-C(45)	1.800(2)
P(1)-C(39)	1.801(2)
O(2)-C(2)	1.290(3)
O(1)-C(1)	1.294(3)
N(1)-C(1)	1.310(3)
N(1)-C(9)	1.424(3)
N(3)-C(57)	1.374(3)

N(2)-C(2)	1.315(3)
N(2)-C(27)	1.425(3)
C(9)-C(10)	1.404(3)
C(9)-C(14)	1.406(3)
C(14)-C(13)	1.395(3)
C(14)-C(15)	1.522(4)
C(10)-C(11)	1.393(3)
C(10)-C(18)	1.525(3)
C(45)-C(46)	1.384(3)
C(45)-C(50)	1.394(3)
C(57)-C(62)	1.410(3)
C(57)-C(58)	1.418(3)
C(3)-C(8)	1.379(3)
C(3)-C(4)	1.388(3)
C(3)-C(1)	1.494(3)
C(2)-C(21)	1.500(3)
C(28)-C(29)	1.399(4)
C(28)-C(27)	1.400(4)
C(28)-C(33)	1.517(4)
C(27)-C(32)	1.409(4)
C(21)-C(22)	1.383(4)
C(21)-C(26)	1.384(3)
C(26)-C(25)	1.384(4)
C(15)-C(16)	1.525(4)
C(15)-C(17)	1.531(4)
C(62)-C(61)	1.391(4)
C(62)-C(64)	1.502(4)
C(52)-C(53)	1.388(3)
C(52)-C(51)	1.394(3)
C(48)-C(49)	1.382(4)
C(48)-C(47)	1.386(4)
C(32)-C(31)	1.397(4)
C(32)-C(36)	1.518(4)
C(39)-C(40)	1.390(3)
C(39)-C(44)	1.399(3)
C(51)-C(56)	1.397(3)
C(4)-C(5)	1.388(3)
C(49)-C(50)	1.382(3)
C(18)-C(19)	1.525(4)
C(18)-C(20)	1.530(4)
C(13)-C(12)	1.382(4)
C(56)-C(55)	1.387(3)
C(5)-C(6)	1.368(4)
C(33)-C(35)	1.526(4)
C(33)-C(34)	1.528(4)

C(47)-C(46)	1.386(3)
C(11)-C(12)	1.379(4)
C(36)-C(38)	1.510(5)
C(36)-C(37)	1.529(4)
C(43)-C(42)	1.376(4)
C(43)-C(44)	1.388(4)
C(58)-C(59)	1.388(4)
C(58)-C(63)	1.499(4)
C(54)-C(53)	1.380(4)
C(54)-C(55)	1.381(4)
C(59)-C(60)	1.385(4)
C(22)-C(23)	1.387(4)
C(40)-C(41)	1.383(4)
C(42)-C(41)	1.371(4)
C(25)-C(24)	1.366(5)
C(29)-C(30)	1.379(4)
C(6)-C(7)	1.375(4)
C(61)-C(60)	1.381(5)
C(8)-C(7)	1.388(4)
C(30)-C(31)	1.369(5)
C(24)-C(23)	1.374(4)

Table 3 – Bond angles for 4

Atoms	Angle (°)
N(3)-Zr(1)-O(1)	105.85(7)
N(3)-Zr(1)-O(3)	100.07(7)
O(1)-Zr(1)-O(3)	78.14(6)
N(3)-Zr(1)-O(2)	96.89(7)
O(1)-Zr(1)-O(2)	134.33(6)
O(3)-Zr(1)-O(2)	136.32(5)
N(3)-Zr(1)-N(2)	116.92(8)
O(1)-Zr(1)-N(2)	134.20(7)
O(3)-Zr(1)-N(2)	79.01(6)
O(2)-Zr(1)-N(2)	57.45(6)
N(3)-Zr(1)-N(1)	103.37(7)
O(1)-Zr(1)-N(1)	57.93(6)
O(3)-Zr(1)-N(1)	134.26(6)
O(2)-Zr(1)-N(1)	78.72(6)
N(2)-Zr(1)-N(1)	121.90(7)
O(3)-P(1)-C(51)	110.75(10)
O(3)-P(1)-C(45)	110.99(10)
C(51)-P(1)-C(45)	107.42(10)
O(3)-P(1)-C(39)	113.62(10)
C(51)-P(1)-C(39)	106.78(10)

C(45)-P(1)-C(39)	106.97(11)
C(2)-O(2)-Zr(1)	96.14(13)
P(1)-O(3)-Zr(1)	147.87(9)
C(1)-O(1)-Zr(1)	95.91(13)
C(1)-N(1)-C(9)	125.46(18)
C(1)-N(1)-Zr(1)	90.51(13)
C(9)-N(1)-Zr(1)	141.55(14)
C(57)-N(3)-Zr(1)	173.54(16)
C(2)-N(2)-C(27)	126.86(19)
C(2)-N(2)-Zr(1)	92.10(13)
C(27)-N(2)-Zr(1)	140.13(15)
C(10)-C(9)-C(14)	121.0(2)
C(10)-C(9)-N(1)	121.1(2)
C(14)-C(9)-N(1)	117.8(2)
C(13)-C(14)-C(9)	118.1(2)
C(13)-C(14)-C(15)	121.0(2)
C(9)-C(14)-C(15)	120.9(2)
C(11)-C(10)-C(9)	118.2(2)
C(11)-C(10)-C(18)	119.7(2)
C(9)-C(10)-C(18)	122.1(2)
C(46)-C(45)-C(50)	119.3(2)
C(46)-C(45)-P(1)	118.80(17)
C(50)-C(45)-P(1)	121.81(17)
N(3)-C(57)-C(62)	121.3(2)
N(3)-C(57)-C(58)	119.7(2)
C(62)-C(57)-C(58)	119.0(2)
C(8)-C(3)-C(4)	118.6(2)
C(8)-C(3)-C(1)	116.7(2)
C(4)-C(3)-C(1)	124.7(2)
O(2)-C(2)-N(2)	114.30(19)
O(2)-C(2)-C(21)	117.1(2)
N(2)-C(2)-C(21)	128.6(2)
C(29)-C(28)-C(27)	118.3(3)
C(29)-C(28)-C(33)	120.5(3)
C(27)-C(28)-C(33)	121.1(2)
C(28)-C(27)-C(32)	121.3(2)
C(28)-C(27)-N(2)	119.8(2)
C(32)-C(27)-N(2)	118.8(2)
C(22)-C(21)-C(26)	118.3(2)
C(22)-C(21)-C(2)	117.0(2)
C(26)-C(21)-C(2)	124.6(2)
C(25)-C(26)-C(21)	120.2(3)
O(1)-C(1)-N(1)	114.46(19)
O(1)-C(1)-C(3)	116.8(2)
N(1)-C(1)-C(3)	128.7(2)

C(14)-C(15)-C(16)	113.5(2)
C(14)-C(15)-C(17)	110.3(2)
C(16)-C(15)-C(17)	110.0(2)
C(61)-C(62)-C(57)	119.2(2)
C(61)-C(62)-C(64)	121.0(2)
C(57)-C(62)-C(64)	119.9(2)
C(53)-C(52)-C(51)	120.3(2)
C(49)-C(48)-C(47)	120.2(2)
C(31)-C(32)-C(27)	117.6(3)
C(31)-C(32)-C(36)	121.0(3)
C(27)-C(32)-C(36)	121.4(2)
C(40)-C(39)-C(44)	119.6(2)
C(40)-C(39)-P(1)	121.35(19)
C(44)-C(39)-P(1)	119.00(18)
C(52)-C(51)-C(56)	119.6(2)
C(52)-C(51)-P(1)	121.32(18)
C(56)-C(51)-P(1)	118.97(17)
C(3)-C(4)-C(5)	120.4(2)
C(50)-C(49)-C(48)	120.0(2)
C(10)-C(18)-C(19)	109.8(2)
C(10)-C(18)-C(20)	112.8(2)
C(19)-C(18)-C(20)	111.8(3)
C(12)-C(13)-C(14)	121.4(2)
C(55)-C(56)-C(51)	119.5(2)
C(6)-C(5)-C(4)	120.4(2)
C(28)-C(33)-C(35)	111.0(3)
C(28)-C(33)-C(34)	113.3(2)
C(35)-C(33)-C(34)	110.3(3)
C(49)-C(50)-C(45)	120.2(2)
C(48)-C(47)-C(46)	119.7(2)
C(12)-C(11)-C(10)	121.5(2)
C(45)-C(46)-C(47)	120.5(2)
C(38)-C(36)-C(32)	110.5(3)
C(38)-C(36)-C(37)	110.1(3)
C(32)-C(36)-C(37)	114.3(3)
C(42)-C(43)-C(44)	120.4(3)
C(59)-C(58)-C(57)	119.4(2)
C(59)-C(58)-C(63)	121.2(2)
C(57)-C(58)-C(63)	119.4(2)
C(53)-C(54)-C(55)	120.5(2)
C(60)-C(59)-C(58)	121.6(3)
C(21)-C(22)-C(23)	120.9(3)
C(43)-C(44)-C(39)	119.4(3)
C(41)-C(40)-C(39)	119.8(3)
C(41)-C(42)-C(43)	120.3(3)

C(24)-C(25)-C(26)	121.1(3)
C(30)-C(29)-C(28)	120.8(3)
C(11)-C(12)-C(13)	119.5(2)
C(54)-C(53)-C(52)	119.7(2)
C(54)-C(55)-C(56)	120.4(3)
C(5)-C(6)-C(7)	119.7(2)
C(42)-C(41)-C(40)	120.6(3)
C(60)-C(61)-C(62)	122.0(3)
C(61)-C(60)-C(59)	118.7(2)
C(3)-C(8)-C(7)	120.7(3)
C(31)-C(30)-C(29)	120.3(3)
C(25)-C(24)-C(23)	119.3(3)
C(30)-C(31)-C(32)	121.5(3)
C(24)-C(23)-C(22)	120.1(3)
C(6)-C(7)-C(8)	120.2(3)

References

- (1) (a) Lauterwasser, F.; Hayes, P. G.; Brase, S.; Piers, W. E.; Schafer, L. L. *Organometallics* **2004**, 23, 2234. (b) Bexrud, J. A.; Beard, J. D.; Leitch, D. C.; Schafer, L. L. *Org. Lett.* **2005**, 7, 1959.
- (2) Zhang, Z.; Schafer, L. L. *Org. Lett.* **2003**, 5, 4733.
- (3) D*TREK: Version 4.13 Molecular Structure Corporation (1996-1998).
- (4) CrystalClear Version 1.3.5b20 Molecular Structure Corporation (2002).
- (5) Farrugia, L.J. *J. Appl. Crystallogr.* **1999**, 32, 83.