A Photochemical Entry to Depsides: Synthesis of Gustastatin

Supporting Information

Jorge García-Fortanet, John R. DeBergh, and Jef K. De Brabander*

*Department of Biochemistry, The University of Texas Southwestern Medical Center at Dallas,
5323 Harry Hines Boulevard, Dallas, TX 75390-9038*

E-mail: jdebra@biochem.swmed.edu

I. General Techniques: Unless otherwise noted, commercially available materials were used without further purification. All solvents were of HPLC or ACS grade. Solvents used for moisture sensitive operations were distilled from drying reagents under a nitrogen atmosphere: Et₂O and THF from sodium benzophenone ketyl; benzene and toluene from sodium; CH₂Cl₂ from CaH₂. Reactions were performed under an atmosphere of nitrogen with magnetic stirring unless noted otherwise. All photochemical reactions were performed with a Rayonett RPR-100 reactor fitted with a test tube carousel and 300 nm bulbs. Flash chromatography (FC) was performed using *E Merck* silicagel 60 (240–400 mesh) according to the protocol of Still, Kahn, and Mitra (*J. Org. Chem.* **1978**, *43*, 2923). Thin layer chromatography was performed using pre-coated plates purchased from *E. Merck* (silicagel 60, PF254, 0.25 mm) that were visualized using a KMnO₄ or Ce (IV) stain.

Nuclear magnetic resonance (NMR) spectra were recorded on a *Varian Inova-400* or *Mercury-300* spectrometer at operating frequencies of 400/300 MHz (¹H NMR) or 100 / 75 MHz (¹³C NMR). Chemical shifts (δ) are given in ppm relative to residual solvent (usually chloroform, δ 7.26 for ¹H NMR or δ 77.23 for proton decoupled ¹³C NMR), and coupling constants (*J*) in Hz. Multiplicity is tabulated as s for singlet, d for doublet, t for triplet, q for quadruplet, and m for multiplet, whereby the prefix app is applied in cases where the true multiplicity is unresolved, and *br* when the signal in question is broadened.

Infrared spectra were recorded on a *Perkin-Elmer I 1000* series FTIR with wavenumbers expressed in cm⁻¹ using samples prepared as thin films between salt plates. UV-VIS spectra were taken in anhydrous diethyl ether and recorded with a Shimadzu UV-1601 spectrometer. Low-resolution mass spectra (LRMS) were recorded on a Shimadzu 2010-LCMS, or at the NIH regional mass spectrometry facility at the University of Washington, St. Louis, MO. High-resolution mass spectra (HRMS) were recorded at the NIH regional mass spectrometry facility at the University of Washington, St. Louis, MO.

II. Experimental Procedures.

5,7-Dihydroxy-2,2-diphenyl-benzo[1,3]dioxin-4-one (i). Prepared according to the procedure described in: Dushin, R. G.; Danishefsky S. J. *J. Am. Chem. Soc.*, **1992**, *114*, 655–659. To a round bottom flask containing 2,4,6-trihydroxybenzoic acid monohydrate (15 g, 71.56 mmol, 1 eq) and benzophenone (39.12 g, 214.68 mmol, 3 eq) was added at 0°C and under N₂ trifluoroacetic acid (118 mL) and trifluoroacetic anhydride (92 mL). The reaction was warmed at rt and then stirred for 24 h. The slightly yellow homogeneous mixture was concentrated and the residue was dissolved in toluene (40 mL) and concentrated again. This operation was repeated 3 times. The crude residue was purified by FC in a rapid fashion to minimize ketal hydrolysis (silicagel, 5% EtOAc/Hexanes) to yield 7.6 g (32%) of **i**: IR (film) 3260, 1667, 1640, 1597, 1480, 1269, 1161, 1094, 695 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 10.20 (1H, s), 7.53–7.55 (4H, m), 7.33–7.36 (6H, m), 6.27 (1H, s), 6.16 (1H, d, *J* = 2.0 Hz), 6.00 (1H, d, *J* = 2.4 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 165.7, 165.1, 163.1, 157.6, 139.0, 129.4, 128.6, 126.3, 107.3, 98.1, 96.2, 94.5; LRMS (ESI) Calcd for C₂₀H₁₃O₅ [M–H]⁻: 333.08; Found: 333.10. HRMS (ESI, added NaO₂CCF₃) Calcd for C₂₀H₁₄O₅Na [MNa]⁺: 357.0739. Found: 357.0727.

7-Benzyloxy-5-hydroxy-2,2-diphenyl-benzo[1,3]dioxin-4-one (ii). Prepared according to the procedure described in: Dushin, R. G.; Danishefsky S. J. *J. Am. Chem. Soc.*, **1992**, *114*, 655–659. To a solution of diol **i** (2.0 g, 5.99 mmol, 1 eq) and benzyl alcohol (651 μL, 6.29 mmol, 1.05 eq) in THF (36 mL) at 0°C was added triphenylphosphine (1.66 g, 6.29 mmol, 1.05 eq), and DIAD (1.22 mL, 6.29 mmol, 1.05 eq) and the mixture was warmed to rt over 2 h. The solution was diluted with EtOAc, washed trice with water and brine, dried over MgSO₄, concentrated, and purified by FC (silicagel, 10% EtOAc/Hexanes) to yield 1.9 g (75%) of **ii**: IR (film) 3209, 3066, 3035, 1695, 1644, 1583, 1503, 1353, 1268, 1103, 1026, 910, 828, 736, 699 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 10.25 (1H, s), 7.54–7.56 (4H, dd, *J* = 2.0 / 8.0 Hz), 7.29–7.36 (11H, m), 6.26 (1H, d, *J* = 2 Hz), 6.12 (1H, d, *J* = 2.4 Hz), 4.99 (2H, s); ¹³C NMR (100 MHz, CDCl₃) δ 166.7, 165.1, 163.1, 157.1, 139.1, 135.4, 129.3, 128.6, 128.5, 128.3, 127.5, 126.3, 107.2, 96.7, 95.8, 94.8, 70.4; LRMS (ESI) Calcd for C₂₇H₁₉O₅ [M–H]⁻: 423.12; Found: 423.25. HRMS (ESI, added NaO₂CCF₃) Calcd for C₂₇H₂₀O₅Na [MNa]⁺: 447.1208. Found: 447.1230.

Trifluoro-methanesulfonic acid 7-benzyloxy-4-oxo-2,2-diphenyl-4H-benzo[1,3]dioxin-5-yl ester (3). To a solution of phenol **ii** (1.9 g, 4.48 mmol, 1 eq) and pyridine (1.81 mL, 22.40 mmol, 5 eq) in CH₂Cl₂ (33 mL) was added trifluoromethanesulfonic anhydride (980 μL, 5.83 mmol, 1.3 eq) dropwise at 0°C. After stirring for 1.5 h at rt and under N₂, the reaction was transferred into a separation funnel containing sat. NaHCO₃, the organic phase was separated and the aqueous phase was washed with CH₂Cl₂ (3 × 50 mL). The combined organic phases were washed with water and brine and dried over MgSO₄. After concentration and purification by FC (silicagel, 20% EtOAc/Hexanes), 4.2 g (94%) of **3** was obtained: IR (film) 3066, 2360, 1746, 1627, 1575, 1431, 1272, 1210, 1153, 1058, 1029, 816, 698 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.52–7.55 (4H, m), 7.32–7.42 (11H, m), 6.73 (1H, d, *J* = 2.4 Hz), 6.54 (1H, d, *J* = 2.4 Hz), 5.10 (2H, s); ¹³C NMR (100 MHz, CDCl₃) δ 164.6, 159.1, 156.9, 150.0, 138.7, 134.5, 129.4, 128.8, 128.7, 128.6, 127.7, 126.4, 107.2, 106.3, 102.4, 71.2; LRMS (ESI) Calcd for C₂₈H₁₉F₃NaO₇S [M+Na]⁺: 579.07;

Found: 579.00. HRMS (ESI, added NaO₂CCF₃) Calcd for C₂₈H₁₉O₇F₃SNa [MNa]⁺: 579.0701. Found: 579.0698.

Potassium *trans*-1-hep-1-enyl Trifluoroborate (7). Prepared according to the procedure described in: Molander, G.A.; Bernardi, C. R. *J. Org. Chem.* **2002**, 67, 8424–8429. To a solution of 1-heptyne (1.34 mL, 10 mmol, 1 eq) in CH₂Cl₂ (5 mL) at 0°C and under N₂ was added Br₂BH·SMe₂ (1M in CH₂Cl₂, 10 mmol, 10 mL, 1 eq). After stirring at rt for 4 h, the light yellow solution was added via syringe to a mixture of diethyl ether/water (10 mL of Et₂O / 4 mL of H₂O) at 0°C. The mixture was stirred at rt for 15 min. The aqueous layer was separated, and the organic phase was washed with cold water and then brine. The organic phase was subsequently dried over MgSO₄ and concentrated to provide 1.05 g (74%) of *trans*-1-hep-1-enyl boronic acid as a white solid.

To a solution of the above prepared *trans*-1-hep-1-enyl boronic acid (1.05 g, 7.41 mmol, 1 eq) in Et₂O (15 mL) was added KHF₂ (1.62 g, 20.75 mmol, 2.8 eq) followed by H₂O (6.7 mL) over 30 min. After stirring at rt for 3 h, the solution was concentrated, and the crude material was dissolved in acetone, filtered, and concentrated. The resulting white solid was purified by dissolving in hot acetone and precipitating with Et₂O to afford 1.26 g (83%) of **7** as a white solid: IR (film) 2959, 2926, 1647, 1111, 933 cm⁻¹; ¹H NMR (400 MHz, DMSO-*d*₆) δ 5.49 (1H, dt, *J* = 6.1 / 17.1 Hz), 5.20–5.27 (1H, dm, *J* = 17.6 Hz), 1.88 (2H, q, *J* = 6.0 Hz), 1.21–1.32 (6H, m), 0.87 (3H, t, *J* = 7.2 Hz); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 133.7, 133.65, 35.4, 31.1, 28.9, 22.1, 14.0.

7-Benzyloxy-5-hept-1-enyl-2,2-diphenyl-benzo[1,3]dioxin-4-one (4).

Prepared according to the procedure described in: Molander, G.A.; Bernardi, C. R. *J. Org. Chem.* **2002**, 67, 8424–8429. A solution of trifluoroborate **7** (102 mg, 0.5 mmol, 1 eq), PdCl₂(dppf)·CH₂Cl₂ (8.1 mg, 0.001 mmol, 2 mol%), triflate **3** (278 mg, 0.5 mmol, 1 eq) in ⁱPrOH/H₂O (2:1, 5 mL, degassed) was added at rt and under N₂ ^tBuNH₂ (158 μL, 1.5 mmol, 3 eq). After refluxing overnight, the reaction mixture was cooled to rt and diluted with water (10 mL), followed by extraction of diethyl ether. The organic layers were combined and washed with brine, dried over MgSO₄, and concentrated. After purification by FC (silicagel, 10% EtOAc/Hexanes), 201 mg (80%) of pure **4** was obtained: IR (film) 3065, 3034, 2927, 2857, 1731, 1605, 1570, 1452, 1452, 1266, 1208, 1160, 970, 697 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.52–7.55 (4H, dd, *J* = 1.2 / 7.6 Hz), 7.25–7.39 (11H, m), 6.74 (1H, d, *J* = 2.4 Hz), 6.57 (1H, d, *J* = 2.0 Hz), 6.11 (1H, dt, *J* = 6.8 / 16.0 Hz), 5.07 (2H, s), 2.20 (2H, q, *J* = 7.2 Hz), 1.44 (2H, app pent, *J* = 7.2 Hz), 1.26–1.32 (4H, m), 0.88 (3H, t, *J* = 6.8 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 163.8, 160.0, 158.9, 144.5, 139.8, 135.7, 135.6, 129.0, 128.7, 128.4, 128.3, 127.9, 127.5, 126.4, 108.9, 105.6, 105.2, 101.1, 70.3, 33.0, 31.4, 28.7, 22.4, 14.0; LRMS (ESI) Calcd for C₃₄H₃₂NaO₄ [MNa]⁺: 527.22; Found: 527.20. HRMS (ESI, added NaO₂CCF₃) Calcd for C₃₄H₃₂O₄Na [MNa]⁺: 527.2198. Found: 527.2212.

7-Benzyloxy-5-(3-pentyl-oxiranyl)-2,2-diphenyl-benzo[1,3]dioxin-4-one (5).

To a solution of alkene **4** (198 mg, 0.393 mmol, 1 eq) in CH₂Cl₂/pH 7 buffer (1:1, 8 mL) was added at 0°C *m*-CPBA (268 mg, ~1.12 mmol, ~2.85 eq). After stirring at rt for 3 h, the organic phase was separated and the aqueous phase was washed with Et₂O (2 × 50 mL). The combined organic phases were washed with

sat. $\text{Na}_2\text{S}_2\text{O}_3$, and brine, dried over MgSO_4 , and concentrated. Purification of the residue by FC (silicagel, 10% EtOAc/Hexanes) afforded 162 mg (79%) of **5**: IR (film) 3065, 2956, 2930, 1731, 1613, 1580, 1453, 1442, 1270, 1160, 698 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.54–7.58 (4H, m), 7.28–7.38 (11H, m), 6.72 (1H, d, $J = 2.4$ Hz), 6.61 (1H, d, $J = 2.4$ Hz), 5.06 (2H, s), 4.32 (1H, d, $J = 1.6$ Hz), 2.64 (1H, td, $J = 2.0 / 5.6$ Hz), 1.64–1.81 (2H, m), 1.54 (2H, m), 1.34 (4H, m), 0.90 (3H, t, $J = 7.2$ Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 164.9, 160.0, 158.8, 145.4, 139.8, 139.5, 135.5, 129.1, 129.1, 128.6, 128.5, 128.3, 127.5, 126.5, 126.2, 106.8, 106.2, 106.0, 102.0, 70.3, 63.4, 57.0, 32.2, 31.5, 25.3, 22.5, 14.0; LRMS (ESI) Calcd for $\text{C}_{34}\text{H}_{32}\text{NaO}_5$ $[\text{MNa}]^+$: 543.21; Found: 543.15. HRMS (ESI, added NaO_2CCF_3) Calcd for $\text{C}_{34}\text{H}_{32}\text{O}_5\text{Na}$ $[\text{MNa}]^+$: 543.2147. Found: 543.2153.

7-Benzyloxy-5-(2-oxo-heptyl)-2,2-diphenyl-benzo[1,3]dioxin-4-one (6). To a solution of epoxide **5** (440 mg, 0.85 mmol, 1 eq), and $\text{Pd}(\text{OAc})_2$ (20 mg, 0.09 mmol, 10% mol) in $t\text{BuOH}$, (degassed, 20 mL) was added at rt and under N_2 , PBu_3 (64 μL , 0.255 mmol, 30 mol%). The reaction mixture was stirred at reflux for 4 h. (*TLC monitoring*). When the starting material was consumed, the solvent was removed under reduced pressure and the residue purified by FC

(silicagel, 15% EtOAc/Hexanes) to afford 370 mg (84%) of pure **6**: IR (film) 3064, 2955, 2930, 1729, 1614, 1578, 1441, 1273, 1162, 1074, 734, 700 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.52–7.55 (4H, dd, $J = 2.0 / 8.0$ Hz), 7.28–7.38 (11H, m), 6.62 (1H, d, $J = 2.4$ Hz), 6.41 (1H, d, $J = 2.0$ Hz), 5.04 (2H, s), 3.94 (2H, s), 2.46 (2H, t, $J = 7.4$ Hz), 1.56 (2H, app pent, $J = 7.6$ Hz), 1.21–1.32 (4H, m), 0.88 (3H, t, $J = 7.2$ Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 206.7, 164.0, 160.4, 159.2, 141.2, 139.6, 135.4, 129.0, 128.6, 128.4, 128.3, 127.5, 126.4, 115.0, 106.6, 106.0, 101.6, 70.3, 48.4, 42.4, 31.3, 23.1, 22.4, 13.9; LRMS (ESI) Calcd for $\text{C}_{34}\text{H}_{32}\text{NaO}_5$ $[\text{MNa}]^+$: 543.21; Found: 543.21. HRMS (ESI, added NaO_2CCF_3) Calcd for $\text{C}_{34}\text{H}_{32}\text{O}_5\text{Na}$ $[\text{MNa}]^+$: 543.2147. Found: 543.2159.

7-Hydroxy-5-(2-oxo-heptyl)-2,2-diphenyl-benzo[1,3]dioxin-4-one (iii). To a suspension of catalyst (10% Pd on carbon, 20 mg, 0.02 mmol, 10 mol%) in EtOAc (1 mL) was added at rt and under H_2 ketone **6** (100 mg, 0.192 mmol, 1 eq) in EtOAc (2 mL). After stirring under H_2 (balloon pressure) at rt for 5 h (*TLC monitoring*), the reaction mixture was filtered through a pad of celite and washed with EtOAc. The filtrate was concentrated and purified by FC (silicagel, 50% EtOAc/Hexanes) to afford 74 mg (90%) of compound **iii**:

IR (film) 3282, 2956, 2931, 1726, 1615, 1451, 1274, 1208, 1164, 1074, 911, 733, 701 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 8.30 (1H, br), 7.49–7.51 (4H, dd, $J = 1.6 / 8.0$ Hz), 7.21–7.29 (6H, m), 6.36 (1H, d, $J = 2.4$ Hz), 6.08 (1H, d, $J = 2.0$ Hz), 3.85 (2H, s), 2.57 (2H, t, $J = 7.4$ Hz), 1.58 (2H, app pent, $J = 7.2$ Hz), 1.22–1.33 (4H, m), 0.87 (3H, t, $J = 7.0$ Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 211.0, 163.4, 161.3, 159.4, 140.2, 139.6, 129.0, 128.4, 126.3, 116.2, 106.0, 105.2, 103.3, 48.5, 42.7, 31.2, 23.1, 22.4, 13.9; LRMS (ESI) Calcd for $\text{C}_{27}\text{H}_{26}\text{NaO}_5$ $[\text{MNa}]^+$: 453.17; Found: 453.10. HRMS (ESI, added NaO_2CCF_3) Calcd for $\text{C}_{27}\text{H}_{26}\text{O}_5\text{Na}$ $[\text{MNa}]^+$: 453.1678. Found: 453.1659.

4-Benzyloxy-2-hydroxy-6-(2-oxo-heptyl)-benzoic acid methyl ester (9). To an oven-dried borosilicate test tube sealed with a rubber septum was added ketone **6** (100 mg, 0.193 mmol, 1eq) in freshly distilled THF (0.1M, 1.9 mL) followed by addition of MeOH (24 μL , 0.577 mmol, 3 eq). This reaction

solution was then placed in a Rayonett reactor and photolyzed for 5 h at 300 nm. The solution was then concentrated and purified by FC (silicagel, 5% EtOAc/Hexanes) to afford 63 mg (89%) of pure ester **9**: IR (film) 2954, 2930, 1719, 1658, 1619, 1579, 1438, 1331, 1258, 1166, 1029, 737, 699 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 11.60 (1H, s), 7.32–7.41 (5H, m), 6.49 (1H, d, J = 2.4 Hz), 6.32 (1H, d, J = 2.4 Hz), 5.05 (2H, s), 3.89 (2H, s), 3.82 (3H, s), 2.41 (2H, t, J = 7.4 Hz), 1.58 (2H, app pent, J = 7.2 Hz), 1.24–1.34 (4H, m), 0.89 (3H, t, J = 7.0 Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 207.4, 170.8, 165.6, 163.2, 138.9, 135.9, 128.6, 128.2, 127.5, 113.4, 105.2, 100.9, 70.0, 51.8, 51.2, 41.9, 31.4, 23.3, 22.4, 13.9; LRMS (ESI) Calcd for $\text{C}_{22}\text{H}_{26}\text{NaO}_5$ $[\text{MNa}]^+$: 393.17; Found: 393.10. HRMS (ESI, added NaO_2CCF_3) Calcd for $\text{C}_{22}\text{H}_{26}\text{O}_5\text{Na}$ $[\text{MNa}]^+$: 393.1678. Found: 393.1677.

6-Benzyloxy-8-hydroxy-3-methoxy-3-pentyl-isochroman-1-one (8). ^1H NMR (300 MHz, CDCl_3) δ 11.20 (1H, d, J = 0.6 Hz), 7.37–7.40 (5H, m), 6.43 (1H, d, J = 2.4 Hz), 6.34 (1H, d, J = 2.1 Hz), 6.06 (2H, s), 3.34 (3H, s), 3.10 (1H, app d, J = 16.2 Hz), 3.04 (1H, app d, J = 16.5 Hz), 2.00 (1H, m), 1.88 (1H, m), 1.35–1.45 (6H, m), 0.92 (3H, t, J = 6.5 Hz); ^{13}C NMR (75 MHz, CDCl_3) δ 168.6, 165.0, 164.3, 144.2, 139.5, 135.8, 128.7, 128.6, 128.3, 127.5, 127.2, 126.9, 107.6, 107.0, 100.2, 70.2, 49.8, 36.2, 35.0, 31.7, 23.1, 22.5, 13.9.

4-Benzyloxy-2-methoxy-6-(2-oxo-heptyl)-benzoic acid methyl ester (10). A solution of phenol **9** (200 mg, 0.541 mmol, 1 eq), Me_3OBF_4 (400 mg, 2.70 mmol, 5 eq), and proton-sponge (694 mg, 3.25 mmol, 6 eq) in CH_2Cl_2 (12 mL) was stirred at rt and under N_2 for 48 h, after which the reaction was diluted with EtOAc (20 mL), filtered through a pad of celite and washed with water (2 $\times$ 50 mL) and aqueous CuSO_4 (1M, 50 mL). The organic phase was dried over MgSO_4 , concentrated and purified by FC (silicagel, 20% EtOAc/Hexanes) to afford 169 mg (81%) of pure **10**: IR (film) 2952, 1720, 1604, 1456, 1430, 1275, 1161, 1098, 1039, 739, 699 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.32–7.42 (5H, m), 6.48 (1H, d, J = 2.4 Hz), 6.39 (1H, d, J = 2.0 Hz), 5.06 (2H, s), 3.84 (3H, s), 3.79 (3H, s), 3.66 (2H, s), 2.40 (2H, t, J = 7.4 Hz), 1.54 (2H, app pent, J = 7.4 Hz), 1.20–1.31 (4H, m), 0.87 (3H, t, J = 7.0 Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 207.5, 168.0, 160.8, 158.9, 136.2, 135.7, 128.6, 128.2, 127.5, 116.3, 108.2, 98.5, 70.1, 56.0, 52.0, 48.2, 41.8, 31.3, 23.3, 22.4, 13.9; LRMS (ESI) Calcd for $\text{C}_{23}\text{H}_{28}\text{NaO}_5$ $[\text{MNa}]^+$: 407.18; Found: 407.10. HRMS (ESI, added NaO_2CCF_3) Calcd for $\text{C}_{23}\text{H}_{28}\text{O}_5\text{K}$ $[\text{MK}]^+$: 423.1574. Found: 423.1584.

4-Hydroxy-2-methoxy-6-(2-oxo-heptyl)-benzoic acid methyl ester (11). To a suspension of catalyst (10% Pd on carbon, 20 mg, 0.02 mmol) in EtOAc (0.5 mL) was added at rt and under H_2 ketone **10** (23.7 mg, 0.0617 mmol, 1 eq) in EtOAc (1 mL). After stirring under H_2 (balloon pressure) at rt for 3 h (*TLC monitoring*), the reaction mixture was filtered through a pad of celite and washed with EtOAc. The filtrate was concentrated and purified by FC (silicagel, 50% EtOAc/Hexanes) to afford 14.6 mg (80%) of compound pure phenol **11**: IR (film) 3365, 2955, 1713, 1609, 1593, 1469, 1434, 1277, 1194, 1165, 1099, 836 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.02 (1H, br), 6.28 (1H, d, J = 2.0 Hz), 6.22 (1H, d, J = 2.0 Hz), 3.83 (3H, s), 3.72 (3H, s), 3.65 (2H, s), 2.47 (2H, t, J = 7.4 Hz), 1.56 (2H, app pent, J = 7.2 Hz), 1.20–1.32 (4H, m), 0.87 (3H, t, J = 7.0 Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 209.6, 168.3, 159.3, 158.9, 135.5, 114.8, 110.2, 98.4, 55.8, 52.1, 48.0, 42.1, 31.2, 23.3, 22.4, 13.9; LRMS (ESI) Calcd for $\text{C}_{16}\text{H}_{22}\text{NaO}_5$ $[\text{MNa}]^+$:

317.14; Found: 317.05. HRMS (ESI, added NaO_2CCF_3) Calcd for $\text{C}_{16}\text{H}_{22}\text{O}_5\text{Na}$ $[\text{MNa}]^+$: 317.1365. Found: 317.1354.

6-Benzyloxy-8-hydroxy-3-pentyl-isochromen-1-one (12). ^1H NMR (400 MHz, CDCl_3) δ 11.12 (1H, s), 7.35–7.42 (5H, m), 6.53 (1H, d, $J = 2.0$ Hz), 6.38 (1H, d, $J = 2.0$ Hz), 6.16 (1H, s), 5.12 (2H, s), 2.48 (2H, t, $J = 7.6$ Hz), 1.68 (2H, app pent, $J = 7.5$ Hz), 1.30–1.36 (4H, m), 0.91 (3H, t, $J = 7.0$ Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 166.5, 165.8, 163.6, 158.1, 139.5, 135.8, 130.1, 128.7, 128.4, 127.5, 103.9, 101.8, 101.0, 70.3, 33.3, 31.1, 26.4, 22.3, 13.9.

6-Benzyloxy-8-methoxy-3-pentyl-isochromen-1-one (13). IR (film) 2929, 2858, 1730, 1664, 1598, 1571, 1456, 1365, 1208, 1167, 997, 692 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.35–7.45 (5H, m), 6.50 (1H, d, $J = 2.4$ Hz), 6.39 (1H, d, $J = 2.4$ Hz), 6.07 (1H, s), 5.13 (2H, s), 3.94 (3H, s), 2.44 (2H, t, $J = 7.6$ Hz), 1.67 (2H, app pent, $J = 7.5$ Hz), 1.25–1.35 (4H, m), 0.90 (3H, t, $J = 6.6$ Hz); ^{13}C NMR (75 MHz, CDCl_3) δ 164.4, 163.2, 159.6, 159.2, 142.4, 135.7, 128.7, 128.4, 127.5, 103.1, 102.9, 100.3, 98.7, 70.3, 56.2, 33.2, 31.1, 26.4, 22.3, 13.9.

7-Methoxy-5-(2-oxo-heptyl)-2,2-diphenyl-benzo[1,3]dioxin-4-one (14). A mixture of phenol **iii** (74 mg, 0.172 mmol, 1 eq), Me_3OBF_4 (127 mg, 0.86 mmol, 5 eq), proton-sponge (221 mg, 1.03 mmol, 6 eq) and 4 Å ms (328 mg) in CH_2Cl_2 (9 mL) was stirred at rt and under N_2 for 16 h, after which the reaction was diluted with EtOAc (20 mL), filtered through a pad of celite and washed with water ($2 \times 50\text{mL}$) and aqueous CuSO_4 (1M, 50 mL). The organic phase was dried over MgSO_4 , concentrated and purified by FC (silicagel, 25% EtOAc/Hexanes) to afford 73 mg (95%) of pure **14**: IR (film) 2930, 2360, 2341, 1727, 1614, 1578, 1451, 1273, 1204, 1158, 1075, 701 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.54–7.56 (4H, dd, $J = 1.4 / 8.0$ Hz), 7.30–7.36 (6H, m), 6.56 (1H, d, $J = 2.4$ Hz), 6.33 (1H, d, $J = 2.4$ Hz), 3.96 (2H, s), 3.82 (3H, s), 2.47 (2H, t, $J = 7.4$ Hz), 1.57 (2H, app pent, $J = 7.2$ Hz), 1.21–1.33 (4H, m), 0.88 (3H, t, $J = 7.2$ Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 206.8, 165.0, 160.5, 159.3, 141.2, 139.7, 129.1, 128.4, 126.4, 114.4, 106.4, 106.0, 100.8, 55.6, 48.4, 42.4, 31.3, 23.2, 22.4, 13.9; LRMS (ESI) Calcd for $\text{C}_{28}\text{H}_{28}\text{NaO}_5$ $[\text{MNa}]^+$: 467.18; Found: 467.15. HRMS (ESI, added NaO_2CCF_3) Calcd for $\text{C}_{28}\text{H}_{28}\text{O}_5\text{Na}$ $[\text{MNa}]^+$: 467.1834. Found: 467.1823.

Gustastatin (1). A solution of phenol **11** (50 mg, 0.170 mmol, 1 eq) and benzodioxinone **14** (151 mg, 0.340 mmol, 2 eq) in benzene (5 mL) was concentrated on the rotavapor (3 times). The residue was dissolved in freshly distilled CH_2Cl_2 (1.7 mL) and transferred to an oven-dried borosilicate test tube sealed with a rubber septum under nitrogen inlet. The nitrogen inlet was removed and the reaction vessel was sealed with parafilm. This solution was then photolyzed in a Rayonett reactor for 2 h at 300 nm. The solution was concentrated and purified by FC (silicagel, 15→30% EtOAc/Hexanes) to afford 42.1 mg (28%) of recovered benzodioxinone **14** and 66.7 mg (71%) of pure gustastatin (**1**) as an amorphous solid: mp 134–138°C; UV (CHCl_3) 246.0, 272.5, 303.5; IR (film) 2950, 2932, 1715, 1657, 1597, 1461, 1428, 1332, 1256, 1160, 114.6, 1074, 1042, 951, 822, 740 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 11.27 (1H, s), 6.59 (1H, d, $J = 1.6$ Hz), 6.56 (1H, d, $J = 1.2$ Hz), 6.46 (1H, d, $J =$

2.4 Hz), 6.29 (1H, d, $J = 2.4$ Hz), 4.06 (2H, s), 3.87 (3H, s), 3.84 (3H, s), 3.82 (3H, s), 3.70 (2H, s), 2.44 (2H, t, $J = 7.6$ Hz), 2.41 (2H, t, $J = 7.6$ Hz), 1.51–1.59 (4H, m), 1.26 (4H, m), 1.22 (4H, m), 0.88 (3H, t, $J = 7.0$ Hz), 0.83 (3H, t, $J = 7.0$ Hz); ^{13}C NMR (75 MHz, CDCl_3) δ 207.4, 206.5, 169.1, 167.4, 166.5, 164.9, 158.3, 151.2, 138.9, 135.4, 121.7, 116.1, 113.3, 104.4, 104.2, 100.0, 56.2, 55.5, 52.3, 51.2, 47.5, 42.5, 42.1, 31.3, 31.2, 23.4, 23.3, 22.4, 13.9, 13.8; LRMS (ESI) Calcd for $\text{C}_{31}\text{H}_{40}\text{O}_9\text{Na}$ $[\text{M}+\text{Na}]^+$: 579.26; Found: 579.20; Calcd for $\text{C}_{31}\text{H}_{39}\text{O}_9$ $[\text{M}-\text{H}]^-$: 555.26; Found: 555.35. HRMS (ESI, added NaO_2CCF_3) Calcd for $\text{C}_{31}\text{H}_{40}\text{O}_9\text{Na}$ $[\text{MNa}]^+$: 579.2570. Found: 579.2587. Elemental Anal. Calcd for $\text{C}_{31}\text{H}_{40}\text{O}_9$: C, 66.89; H, 7.24; O, 25.87. Found: C, 67.07; H, 7.07; O, 25.19.

III. Data comparison of natural and synthetic gustastatin.

Table S1: ^1H NMR Data for natural (500 MHz, CDCl_3) and synthetic (400 MHz, CDCl_3) gustastatin (**1**).

Position	δ in ppm (multiplicity, J in Hz)	
	Natural 1	Synthetic 1
OH2	11.28 (s)	11.27 (s)
H3'	6.59 (d, 1.5)	6.59 (d, 1.6)
H5'	6.57 (d, 1.5)	6.56 (d, 1.2)
H3	6.46 (d, 2.5)	6.46 (d, 2.4)
H5	6.29 (d, 2.5)	6.29 (d, 2.4)
H1''	4.07 (s)	4.06 (s)
CO_2CH_3	3.88 (s)	3.87 (s)
4- OCH_3	3.85 (s)	3.84 (s)
2'- OCH_3	3.83 (s)	3.82 (s)
H1'''	3.70 (s)	3.70 (s)
H3''	2.44 (t, 7.5)	2.44 (t, 7.6)
H3'''	2.41 (t, 7.5)	2.41 (t, 7.6)
H4''	1.56 (pent, 7.5)	
H4'''	1.53 (pent, 7.5)	1.51-1.59 (m)
H6'', H6'''	1.27 (m)	1.26 (m)
H5'', H5'''	1.21 (m)	1.22 (m)
H7'' and H7'''	not reported	0.88 (t, 7.0) / 0.83 (t, 7.0)

Table S2: ^{13}C NMR Data for natural (125 MHz, CDCl_3) and synthetic (75 MHz, CDCl_3) gustastatin (**1**).

Position	δ in ppm	
	Natural 1	Synthetic 1
C2''	207.4	207.4
C2'''	206.4	206.5
C7	169.1	169.1
C7'	167.5	167.4
C2	166.5	166.5
C4	164.9	164.9
C2'	158.4	158.3
C4'	151.3	151.2
C6	138.9	138.9
C6'	135.4	135.4
C1'	121.8	121.7
C5'	116.2	116.1
C5	113.4	113.3
C3'	104.5	104.4
C1	104.3	104.2

Position	δ in ppm	
	Natural 1	Synthetic 1
C3	100.1	100.0
2'-OCH ₃	56.3	56.2
4-OCH ₃	55.5	55.5
CO ₂ CH ₃	52.3	52.3
C1''	51.2	51.2
C1'''	47.5	47.5
C3''	42.5	42.5
C3'''	42.2	42.1
C5''	31.3	31.3
C5'''	31.3	31.2
C4''	23.4	23.4
C4'''	23.3	23.3
C6''	22.4	22.4
C6'''	22.4	22.4
C7''	13.9	13.9
C7'''	13.8	13.8

IV. Proton and Carbon NMR Spectra.

??