Supporting Information

A Positional Scanning Approach to the Discovery of Dipeptide-Based Catalysts for the Enantioselective Addition of Vinylzinc Reagents to Aldehydes

Christopher M. Sprout, Meaghan L. Richmond, and Christopher T. Seto*

Department of Chemistry, Brown University, 324 Brook Street Box H, Providence, Rhode Island

02912

Email: Christopher_Seto@brown.edu

Table of Contents

General Methods	S2
Conditions for ee determinations of allylic alcohols 7a-7c , 63-64 , and 67-75	S3
Dipeptide ligand yields from solid-phase synthesis	S10
Ligand screening % ee results	S13
¹ H and ¹³ C NMR spectra for compounds 61 , 62 , 54 , 76 , 16	S16
¹ H and ¹³ C NMR spectra for (<i>E</i>)-allylic alcohols 67-75	S21
References	S30

General Methods. NMR spectra were recorded on either a 300 or 400 MHz instrument using CDCl₃ as the solvent and TMS as the internal standard. Mass spectra were recorded using electrospray, electron impact, or fast atom bombardment ionization methods. Enantiomeric excess was determined using a HPLC with a Daicel Chiralcel OD-H column (0.46 cm i.d. × 25 cm) with UV detection at 219 nm. Optical rotations were recorded on a polarimeter at ambient temperature and at a wavelength of 589 nm (c = g/100 mL). Melting points are uncorrected. Solvents were of reagent grade and were used as received except for CH₂Cl₂, toluene, and Et₂O which were purified by passage through a column of activated alumina.¹

Solvents were removed by rotary evaporation at water aspirator pressure followed by further drying at approximately 0.3 mm Hg. Benzaldehyde, cyclohexanecarboxaldehyde, and all alkynes were distilled prior to use. 4-Chlorobenzaldehyde was recrystallized prior to use. 4-Bromobenzaldehyde, 4-methoxybenzaldehyde, 2-naphthylaldehyde, 1-naphthylaldehyde, and *trans*-cinnamaldehyde were purified by column chromatography. Oxime resin (0.9 mmol/g, 100-200 mesh) was purchased from Advanced ChemTech and was washed with DMF (3 × 3 mL) prior to use. Anhydrous procedures were conducted using oven dried or flame dried glassware and standard syringe and cannula transfer techniques. Alkenylzinc reactions were conducted under a nitrogen atmosphere without stirring. Reactions were monitored using TLC. TLC plates were visualized using either an ethanol solution of phosphomolybdic acid (PMA) or UV at 254 nm. Racemic standards for all allylic alcohols were synthesized and analyzed by HPLC or by ¹⁹F NMR analysis of a Mosher's ester derivative in order to confirm the peak assignments of the two enantiomers. The absolute configurations of allylic alcohols **67-75** are inferred based upon the supposition that the reactions catalyzed by the optimized ligand **54** give the same sense of stereochemical induction for the formation of these products as for formation of the allylic alcohol products **7a-c** and **63-64**.

Conditions for ee Determinations

(*R,E*)-(-)-4,4-Dimethyl-1-phenylpent-2-en-1-ol (**7a**)

$[\alpha]_D^{24} = -43$ ($c = 0.61$, CHCl_3) {lit.² $[\alpha]_D^{20} = +49.3$ ($c = 1.7$, CHCl_3 , for 98% ee (*S*)); 94% ee (*R*) by HPLC analysis (Chiralcel OD-H column eluted with hexanes:2-propanol (99:1) at 1.0 mL/min and detected at 219 nm), $t_R = 15.1$ min for (*R*) and $t_R = 24.0$ min for (*S*).

Figure S1. HPLC trace of 4,4-dimethyl-1-phenylpent-2-en-1-ol (**7a**).

(*R,E*)-(-)-3-Cyclopropyl-1-phenylprop-2-en-1-ol (**63**)

$[\alpha]_D^{26} = -71$ ($c = 1.0$, CHCl_3) {lit.³ $[\alpha]_D^{25} = +58.8$ ($c = 0.98$, CHCl_3 , for 94.5% ee (*S*)); 92% ee (*R*) by HPLC analysis (Chiralcel OD-H column eluted with hexanes:2-propanol (99:1) at 1.0 mL/min and detected at 219 nm), $t_R = 28.7$ min for (*R*) and $t_R = 40.2$ min for (*S*).

Figure S2. HPLC trace of 3-cyclopropyl-1-phenylprop-2-en-1-ol (**63**).

(*R,E*)-(-)-1-Phenylnon-2-en-1-ol (7c)

$[\alpha]_D^{25} = -27$ ($c = 1.0$, CHCl_3) {lit.² $[\alpha]_D^{20} = +34.4$ ($c = 2.0$, CHCl_3 , for 94% ee (*S*)); 92% ee (*R*) by HPLC analysis (Chiralcel OD-H column eluted with hexanes:2-propanol (99:1) at 1.0 mL/min and detected at 219 nm), $t_R = 18.3$ min for (*R*) and $t_R = 27.6$ min for (*S*).

Figure S3. HPLC trace of (*R,E*)-(-)-1-phenylnon-2-en-1-ol (**7c**).

(*R,E*)-(-)-1-Phenylhept-2-en-1-ol (7b)

$[\alpha]_D^{25} = -29$ ($c = 1.03$, CHCl_3) {lit.² $[\alpha]_D^{20} = +38.0$ ($c = 2.0$, CHCl_3 , for 96% ee (*S*)); 92% ee (*R*) by HPLC analysis (Chiralcel OD-H column eluted with hexanes:2-propanol (99:1) at 1.0 mL/min and detected at 219 nm), $t_R = 18.2$ min for (*R*) and $t_R = 26.4$ min for (*S*).

Figure S4. HPLC trace of (*R,E*)-(-)-1-phenylhept-2-en-1-ol (**7b**).

(*R,E*)-(-)-3-Cyclohexyl-1-phenylprop-2-en-1-ol (64)

$[\alpha]_D^{26} = -30$ ($c = 1.0$, CHCl_3) {lit.² $[\alpha]_D^{20} = +36.7$ ($c = 1.6$, CHCl_3 , for 95% ee (*S*)); 89% ee (*R*) by HPLC analysis (Chiralcel OD-H column eluted with hexanes:2-propanol (99:1) at 1.0 mL/min and detected at 219 nm), $t_R = 20.3$ min for (*R*) and $t_R = 30.3$ min for (*S*).

Figure S5. HPLC trace of (*R,E*)-(-)-3-cyclohexyl-1-phenylprop-2-en-1-ol (**64**).

(*R,E*)-(-)-(4-Chlorophenyl)-4,4-dimethylpent-2-en-1-ol (67)

$[\alpha]_D^{25} = -56$ ($c = 0.99$, CHCl_3); 95% ee by HPLC analysis (Chiralcel OD-H column eluted with hexanes at 1.0 mL/min and detected at 219 nm), $t_R = 61.5$ min for (*R*) and $t_R = 76.2$ min for (*S*).

Figure S6. HPLC trace of (*R,E*)-(-)-(4-chlorophenyl)-4,4-dimethylpent-2-en-1-ol (**67**).

(*R,E*)-(-)-(4-Bromophenyl)-4,4-dimethylpent-2-en-1-ol (68)

$[\alpha]_D^{25} = -57$ ($c = 2.24$, CHCl_3); 95 % ee by HPLC analysis (Chiralcel OD-H column eluted with hexanes at 1.0 mL/min and detected at 219 nm), $t_R = 67.1$ min for (*R*) and $t_R = 98.5$ min for (*S*).

Figure S7. HPLC trace of (*R,E*)-(-)-(4-bromophenyl)-4,4-dimethylpent-2-en-1-ol (**68**).

(*R,E*)-(-)-Methyl 4-(1-hydroxy-4,4-dimethylpent-2-enyl)benzoate (69)

$[\alpha]_D^{25} = -67$ ($c = 2.62$, CHCl_3); 94% ee by HPLC analysis (Chiralcel OD-H column eluted with hexanes:2-propanol (95:5) at 1.0 mL/min and detected at 219 nm), $t_R = 13.1$ min for (*S*) and $t_R = 14.9$ min for (*R*).

Figure S8. HPLC trace of (*R,E*)-(-)-methyl 4-(1-hydroxy-4,4-dimethylpent-2-enyl)benzoate (**69**).

(*R,E*)-(-)-1-(4-Methoxyphenyl)-4,4-dimethylpent-2-en-1-ol (71)

$[\alpha]_D^{25} = -21$ ($c = 0.99$, CHCl_3); 91% ee by HPLC analysis (Chiralcel OD-H column eluted with hexanes:2-propanol (99:1) at 1.0 mL/min and detected at 219 nm), $t_R = 24.6$ min for (*R*) and $t_R = 31.2$ min for (*S*).

Figure S9. HPLC trace of (*R,E*)-(-)-1-(4-methoxyphenyl)-4,4-dimethylpent-2-en-1-ol (71).

(*R,E*)-(-)-4,4-Dimethyl-1-(naphthalene-2-yl)pent-2-en-1-ol (72)

$[\alpha]_D^{25} = -52$ ($c = 1.00$, CHCl_3); 91% ee by HPLC analysis (Chiralcel OD-H column eluted with hexanes:2-propanol (99.5:0.5) at 1.0 mL/min and detected at 219 nm), $t_R = 68.7$ min for (*R*) and $t_R = 75.2$ min for (*S*).

Figure S10. HPLC trace of (*R,E*)-(-)-4,4-dimethyl-1-(naphthalene-2-yl)pent-2-en-1-ol (72).

(*S,E*)-(-)-6,6-Dimethyl-1-phenylhept-4-en-3-ol (73)

$[\alpha]_D^{24} = -7.5$ ($c = 0.96$, CHCl_3); 81% ee by HPLC analysis (Chiralcel OD-H column eluted with hexanes:2-propanol (99.5:0.5) at 1.0 mL/min and detected at 219 nm), $t_R = 22.5$ min for (*S*) and $t_R = 44.2$ min for (*R*).

Figure S11. HPLC trace of (*S,E*)-(-)-6,6-dimethyl-1-phenylhept-4-en-3-ol (**73**).

(*S,1E,4E*)-(-)-6,6-Dimethyl-1-phenylhepta-1,4-dien-3-ol (74)

$[\alpha]_D^{24} = +1.2$ ($c = 1.09$, CHCl_3); 78% ee by HPLC analysis (Chiralcel OD-H column eluted with hexanes:2-propanol (90:10) at 1.0 mL/min and detected at 219 nm), $t_R = 6.9$ min for (*S*) and $t_R = 10.9$ min for (*R*).

Figure S12. HPLC trace of (*S,1E,4E*)-(-)-6,6-dimethyl-1-phenylhepta-1,4-dien-3-ol (**74**).

(*R,E*)-(-)-4,4-Dimethyl-1-(naphthalene-1-yl)pent-2-en-1-ol (75)

$[\alpha]_D^{26} = +23$ ($c = 1.00$, CHCl_3); 59% ee by HPLC analysis (Chiralcel OD-H column eluted with hexanes:2-propanol (99:1) at 1.0 mL/min and detected at 219 nm), $t_R = 27.7$ min for (*S*) and $t_R = 43.9$ min for (*R*).

Figure S13. HPLC trace of (*R,E*)-(-)-4,4-dimethyl-1-(naphthalene-1-yl)pent-2-en-1-ol (**75**).

Chart S1. Solid-Phase Synthesis Yields of Ligands **6**, and **8-30**; Optimization of R².

*Recycled oxime resin was used to synthesize these ligands. **Synthesized using solution-phase.

Chart S2. Solid-Phase Synthesis Yields of Ligands **31-53**; Optimization of R³.

*Recycled oxime resin used to synthesize these ligands. **MeOH used to wash the ligand from the resin. ***Coupling reagent was 1.5 equiv HBTU/1.5 equiv HATU and used recycled oxime resin.

Chart S3. Solid-Phase Synthesis Yields of Ligands **54-59**; Optimization of R¹.

Table S1. Effect of Ligand R² Substituent on the Addition of Alkenylzinc to Benzaldehyde^{a,b}

Ligand	R ² = side chain of amino acid	% ee ^{c,d}	Ligand	R ² = side chain of amino acid	% ee ^{c,d}
8	Asn	83	19	Gly	55
9	Met	81	20	<i>t</i> -Leu	53
10	Ala	68	21	Tyr(Bn)	53
11	Orn(Cbz)	67	22	Ile	52
12	Trp	67	23	Glu(cHx) ^e	49
6	Val	67	24	Nle	46
13	Gln	65	25	Chg	43
14	Phe	64	26	Thr(Bn)	38
15	Ser(Bn)	63	27	Arg(Ts)	37
16	His(Bom) ^{e,f}	62	28	Asp(cHx) ^e	24
17	Leu	60	29	Lys(Cbz)	19
18	Cys(Bn)	59	30	Pro	9

^aExperiments performed in duplicate unless noted other wise. ^bList of abbreviations: Chg = cyclohexylglycine, cHx = cyclohexyl, Bom = benzyloxymethyl, Nle = norleucine, Orn = ornithine. ^c% ee measured by HPLC (Chiralcel OD-H). ^dAbsolute configuration was assigned as (*R*) by comparison of the optical rotation with the reported value. ^eBased on one experiment. ^fCompound was synthesized by solution-phase methods and tested in purified form.

Table S2. Effect of Ligand R³ Substituent on the Addition of Alkenylzinc to Benzaldehyde^{a,b}

Ligand	R ³ = side chain of amino acid	% ee ^{c,d}	Ligand	R ³ = side chain Of amino acid	% ee ^{c,d}
9	Cys(Tr)	81 (<i>R</i>)	42	Tyr(<i>t</i> Bu)	39 (<i>R</i>)
31	Glu(<i>t</i> Bu)	64 (<i>R</i>)	43	Trp	37 (<i>R</i>)
32	Chg	61 (<i>R</i>)	44	<i>t</i> -Leu	37 (<i>R</i>)
33	Orn(Cbz)	56 (<i>R</i>)	45	Arg(Mts)	34 (<i>R</i>)
34	Val	56 (<i>R</i>)	46	Ala	33 (<i>R</i>)
35	Ile	56 (<i>R</i>)	47	Pro	29 (<i>R</i>)
36	Lys(Boc)	55 (<i>R</i>)	48	Ser(<i>t</i> Bu)	26 (<i>R</i>)
37	Leu	50 (<i>R</i>)	49	Asp(<i>t</i> Bu)	23 (<i>R</i>)
38	Gln	47 (<i>R</i>)	50	Thr(<i>t</i> Bu)	21 (<i>R</i>)
39	Met	47 (<i>R</i>)	51	His(Boc)	12 (<i>R</i>)
40	Nle	44 (<i>R</i>)	52	Asn(Xan)	NR
41	Phe	40 (<i>R</i>)	53	Gly	6 (<i>S</i>)

^aAll experiments performed in duplicate. ^bList of abbreviations: Chg = cyclohexylglycine, Mts = mesitylene-2-sulfonyl, Nle = norleucine, Orn = ornithine, Xan = xanthyl, NR = no reaction. ^c% ee measured by HPLC (Chiralcel OD-H). ^dAbsolute configuration was assigned by comparison of the known HPLC retention times for each enantiomer.

Table S3. Effect of ligand R¹ Substituent on the Addition of Alkenylzinc to Benzaldehyde^a

Ligand	R ¹	% ee ^b
54	-CH ₂ CH ₃	86
55	-CH(CH ₃) ₂	85
56	-CH ₃	82
57	-(CH ₂) ₅ -	81
9	-(CH ₂) ₂ O(CH ₂) ₂ -	81
58	-(CH ₂) ₃ CH ₃	81
59	-(CH ₂) ₄ -	72

^aAll experiments performed in duplicate. ^b% ee of (*R*)-allylic alcohols as measured by HPLC (Chiralcel OD-H).

Figure S14. ^1H NMR spectrum for carbamate **61**.

Figure S15. ^{13}C NMR spectrum for carbamate **61**.

Figure S16. ¹H NMR spectrum for amine **62**.

Figure S17. ¹³C NMR spectrum for amine **62**.

Figure S18. ^1H NMR spectrum for ligand **54**.

Figure S19. ^{13}C NMR spectrum for ligand **54**.

Figure S20. ^1H NMR spectrum for carbamate **76**.

Figure S21. ^{13}C NMR spectrum for carbamate **76**.

Figure S22. ^1H NMR spectrum for ligand **16**.

Figure S23. ^{13}C NMR spectrum for ligand **16**.

Figure S24. ^1H NMR spectrum for alcohol **67**.

Figure S25. ^{13}C NMR spectrum for alcohol **67**.

Figure S26. ^1H NMR spectrum for alcohol **68**.

Figure S27. ^{13}C NMR spectrum for alcohol **68**.

Figure S28. ¹H NMR spectrum for alcohol **69**.

Figure S29. ¹³C NMR spectrum for alcohol **69**.

Figure S30. ¹H NMR spectrum for alcohol **70**.

Figure S31. ¹³C NMR spectrum for alcohol **70**.

Figure S32. ¹H NMR spectrum for alcohol **71**.

Figure S33. ¹³C NMR spectrum for alcohol **71**.

Figure S34. ¹H NMR spectrum for alcohol **72**.

Figure S35. ¹³C NMR spectrum for alcohol **72**.

Figure S36. ^1H NMR spectrum for alcohol **73**.

Figure S37. ^{13}C NMR spectrum for alcohol **73**.

Figure S38. ^1H NMR spectrum for alcohol **74**.

Figure S39. ^{13}C NMR spectrum for alcohol **74**.

Figure S40. ^1H NMR spectrum for alcohol **75**.

Figure S41. ^{13}C NMR spectrum for alcohol **75**.

References

(1) Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J.

Organometallics **1996**, *15*, 1518.

(2) Oppolzer, W.; Radinov, R. N. *Helv. Chim. Acta* **1992**, *75*, 170.

(3) Chen, Y. K.; Lurain, A. E.; Walsh, P. J. *J. Am. Chem. Soc.* **2002**, *124*, 12225.