Supporting Information

A Superior Method for the Reduction of Secondary Phosphine Oxides

Carl A. Busacca*, Jon C. Lorenz, Nelu Grinberg, Nizar Haddad, Matt Hrapchak, Bachir Latli, Heewon Lee, Paul Sabila, Anjan Saha, Max Sarvestani, Sherry Shen, Richard Varsolona, Xudong Wei, and Chris H. Senanayake

*Department of Chemical Development, Boehringer-Ingelheim
Pharmaceuticals, Inc. 900 Ridgebury Rd., Ridgefield, CT 06877*

General

Symmetric secondary phosphines oxides (SPO's) which were commercially available were **1**, **3**, **17**, and **19**. Symmetric secondary phosphines oxides (SPO's) which were not commercially available were prepared by the method of Hays¹ by adding 1 eq. diethylphosphite to 3.3 eq. Grignard at ~0°C and warming to ambient temperature. Compounds **4**, **5**, **7**, and **11** were prepared by the known methods.² Compounds **13** and **14** were also prepared by the known procedures.³ Representative examples are shown below for the synthesis of SPO's **8**, **12** and **17**. Non-symmetric SPO's were prepared by the addition of 1 eq. ethylphenylphosphinate to 2.2 eq. Grignard at ~0°C and warming to ambient temperature, essentially following the procedure of Emmick.⁴ We found that commercial sources of ethylphenylphosphinate were inferior, and the distilled reagent was best prepared from phenylphosphinic acid by the Hewitt reaction shortly before use.⁵ Representative examples are shown below for the synthesis of SPO's **15** and **16**. Only SPO **8**, and secondary phosphines **29-30** have not been previously reported and their synthesis and characterization are shown below. SPO's **10** and **20** and phosphines **31** and **41** have been previously reported,⁶ yet we show alternate preparations of these materials. The Grignard intermediate for synthesis of **20** was prepared by the known procedure and should NOT be prepared from Mg metal.⁷ ¹H NMR were recorded at 400 or 500 MHz, ³¹P NMR were recorded at 161 or 201 MHz, ¹⁹F NMR were recorded at 376 MHz, and ¹³C NMR were recorded at 100, 125, or 150 MHz on Bruker Avance spectrometers.

¹ (a) Hays, H.R. *J. Org. Chem.* **1968**, 33, 3690. (b) Harger, M.J.P.; Westlake, S. *Tetrahedron* **1982**, 38(10), 1511.

² (a) Tsvetkov, E.N.; Shvets, A.A.; Korolev, B.A.; Bondarenko, N.A.; Goncharova, L.V.; Osmolovskaya, L.A. *J. Gen. Chem. USSR* (Eng. Trans.) **1992**, 62, 1833. (b) Kapoor, P.N.; Pathak, D.D.; Gaur, G.; Kutty, M. *J. Organomet. Chem.* **1984**, 276(2), 167. (c) Busacca, C.A.; Grossbach, D.; Campbell, S.J.; Dong, Y.; Eriksson, M.C.; Harris, R.E.; Jones, P.J.; Kim, J.-Y.; Lorenz, J.C.; McKellop, K.B.; O'Brien, E.M.; Qiu, F.; Simpson, R.D.; Smith, L.; So, R.C.; Spinelli, E.M.; Vitous, J.; Zavattaro, C. *J. Org. Chem.* **2004**, 69(16), 5187. (d) Harger, M.J.P.; *J. Chem. Soc. Perkin Trans. 2* **1980**, 154.

³ (a) Kapoor, P.N.; Venzani, L.M. *Helv. Chim. Acta* **1977**, 60(8), 2824. (b) Wife, R.L.; Oort, A.B.v.; Doorn, J.A.v.; Leeuwen, P.W.N.M. *Synthesis* **1983**, 71.

⁴ (a) Emmick, T.L.; Letsinger, R.L. *J. Am. Chem. Soc.* **1968**, 90(13), 3459. (b) Shioji, K.; Tsukimoto, S.; Tanaka, H.; Okuma, K. *Chem. Lett.* **2003**, 32(7), 604.

⁵ (a) Hewitt, D.G. *Aust. J. Chem.* **1979**, 32, 463. (b) Afarinkia, K.; Yu, H.-W. *Tetrahedron Lett.* **2003**, 44(4), 781.

⁶ (a) Unruh, J.D.; Segmuller, B.E.; Chapa, G.R.; Pryor, K.E. U.S. Pat. 5,567,856, 1996. (b) Rajanbabu, T.V.; Casalnuovo, A.L.; Ayers, T.A.; Warren, T.H. *J. Am. Chem. Soc.* **1994**, 116(22), 9869.

⁷ (a) Leazer, J.L.; Cvetovich, R.; Tsay, F.-R.; Dolling, U.; Vickery, T.; Bachert, D. *J. Org. Chem.* **2003**, 68(9), 3695. (b) Yakelis, N.A.; Bergman, R.G. *Organometallics* **2005**, 24(14), 3579.

GC analyses were performed on a HP 5890 GCMS. H₂ was detected by TCD (250°C detector temperature) with a splitless injection using argon as carrier gas, using an Agilent CarbonPLOT column, 320µm x 30m, film thickness: 1.5 µm, 1 mL/min. flow. Isobutane was detected by FID (280°C detector temperature) with a split injection (50:1) using helium as a carrier gas, with a J+W DB-1 column, 250µm x 30m, film thickness: 1.0 µm, 1 mL/min. flow. In each analysis, the column was held at 35°C for 3 minutes, then ramped to 80°C at 5°C/minute, with a total run time of 12 minutes. Retention times were 2.88 min. for H₂ and 2.57 min. for isobutane. GC experiments were performed by mixing 1.0 mmol of both the SPO and reducing agent in a 10 mL flask for 1 minute, then sampling the headspace in the flask via syringe. All reagents were used as received unless stated otherwise. Digital photographs of glassware setups are found at the end of the Supporting Information.

General Procedure A: Symmetric Secondary Phosphine Oxides.

Di-n-butyl phosphine oxide 17.

A 250 mL RB flask equipped with an addition funnel was evacuated/Ar filled (3X), then charged with 50 mL 2M n-BuMgCl/THF (100 mmol, 3.3 eq.), and the solution cooled to 0°C under Ar. A solution of 3.90 mL diethylphosphite (30.3 mmol, 1.0 eq.) in 10 mL THF was then added dropwise over 15 minutes. The mixture was aged 15 minutes at 0°C, then the bath was removed, and the mixture stirred two hours at ambient temperature, then cooled again to 0°C. 75 mL 0.1N HCl was then added dropwise over 20 minutes, then 75 mL MTBE was added, and the mixture agitated well for 5 minutes. The upper organic phase was decanted from the gel and saved. To the remaining gel was added 100 mL CH₂Cl₂, and the mixture agitated well for 5 minutes. The resultant mixture was then filtered through a Celite pad, washing the pad with CH₂Cl₂. The filtrate phases were separated, and the organic phase combined with the first organic phase, dried (MgSO₄), and the solvents removed *in vacuo*. The residue was azeotroped with hexane (1 X 100 mL) causing a precipitate to form. This solid was then recrystallized from 25 mL boiling hexane to give 4.40 g **17** (90%) as colorless plates. M.p. 55-56°C; ¹H NMR (500 MHz, CDCl₃) δ: 6.90 (d, ¹J_{H-P} = 449 Hz, 1H), 1.82 (m, 4H), 1.62 (m, 4H), 1.47 (m, 4H), 0.94 (t, J = 7.4 Hz, 6H). ¹³C NMR (125 MHz, CDCl₃) δ: 28.39 (t, ¹J_{CP} = 91 Hz), 27.83 (t, ²J_{CP} = 65 Hz), 23.67 (t), 13.42 (q). ³¹P NMR (202 MHz, CDCl₃) δ: 36.09 (¹J_{PH} = 449 Hz). Anal. Calcd for C₈H₁₉PO: C, 59.24; H, 11.81; Found C, 58.84; H, 12.02;

Diphenyl-phosphine oxide 1.

¹H NMR (500 MHz, CDCl₃) δ: 8.09 (d, ¹J_{HP} = 482 Hz, 1H), 7.71 (m, 4H), 7.57 (m, 2H), 7.50 (m, 4H). ³¹P NMR (202 MHz, CDCl₃) δ: 21.72 (¹J_{PH} = 482 Hz).

Bis-(4-fluorophenyl)-phosphine oxide 2.

Prepared from 4-fluorophenylmagnesium bromide and diethylphosphite by General Procedure A and chromatographed on silica gel eluting with 70:30 EtOAc:Hexane to give the product (80%) as a colorless oil after drying under high vacuum. ^1H NMR (400 MHz, CDCl_3) δ : 8.05 (d, $^1J_{\text{HP}} = 486$ Hz, 1H), 7.69-7.62 (m, 4H), 7.19-7.13 (m, 4H). ^{31}P NMR (162 MHz, CDCl_3) δ : 18.86 ($^1J_{\text{PH}} = 488$ Hz). ^{19}F NMR (376 MHz, CDCl_3) δ : -105.00 Anal. Calcd for $\text{C}_{12}\text{H}_9\text{F}_2\text{OP}$: C, 60.51; H, 3.81;

Found C, 60.20; H, 4.01.

Bis-(4-chlorophenyl)-phosphine oxide 3.

^1H NMR (500 MHz, CDCl_3) δ : 8.08 (d, $^1J_{\text{HP}} = 486$ Hz, 1H), 7.62 (dd, $J = 13.5, 8.5$ Hz, 4H), 7.50 (dd, $J = 8.0, 1.5$ Hz, 4H). ^{31}P NMR (202 MHz, CDCl_3) δ : 19.06 ($^1J_{\text{PH}} = 486$ Hz).

Bis-(4-methylphenyl)-phosphine oxide 4.

Prepared from 4-methylphenylmagnesium bromide and diethylphosphite by General Procedure A and recrystallized from hot EtOAc to give the product (89%) as a colorless solid. ^1H NMR (500 MHz, CDCl_3) δ : 8.02 (d, $^1J_{\text{HP}} = 478$ Hz, 1H), 7.57 (dd, $J = 14, 8$ Hz, 4H), 7.28 (dd, $J = 8, 2$ Hz, 4H). ^{31}P NMR (202 MHz, CDCl_3) δ : 20.38 ($^1J_{\text{PH}} = 478$ Hz).

Bis-(3-fluorophenyl)-phosphine oxide 5.

Prepared from 3-fluorophenylmagnesium bromide and diethylphosphite by General Procedure A and chromatographed on silica gel eluting with 60% EtOAc/Hex to give the product (74%) as a colorless oil after drying under high vacuum. ^1H NMR (500 MHz, CDCl_3) δ : 8.04 (d, $^1J_{\text{HP}} = 486$ Hz, 1H), 7.55-7.48 (m, 3H), 7.43-7.38 (m, 2H), 7.30-7.27 (m, 2H). ^{31}P NMR (202 MHz, CDCl_3) δ : 18.54 ($^1J_{\text{PH}} = 487$ Hz).

Bis-(3-chlorophenyl)-phosphine oxide 6.

Prepared from 3-chlorophenylmagnesium bromide and diethylphosphite by General Procedure A and chromatographed on silica gel eluting with CH_2Cl_2 , then 5% $\text{MeOH}/\text{CH}_2\text{Cl}_2$ to give the product (74%) as a colorless oil after drying under high vacuum. ^1H NMR (500 MHz, CDCl_3) δ : 8.06(d, $^1J_{\text{HP}} = 493$ Hz, 1H), 7.68 (d, $J = 14$ Hz, 2H), 7.61-7.56 (m, 4H), 7.48 (dt, $J = 3.1, 7.6$ Hz, 2H). ^{31}P NMR (202 MHz, CDCl_3) δ : 18.92 ($^1J_{\text{PH}} = 493$ Hz). Anal. Calcd for $\text{C}_{12}\text{H}_9\text{Cl}_2\text{OP}$: C, 53.17; H, 3.35; Found C, 52.90; H, 3.54.

Bis-(3-fluoro-5-methylphenyl)-phosphine oxide 7.

A 4 neck 1L RBF fitted with mechanical stirrer, reflux condenser and addition funnel was evacuated/Ar filled (3 X), then charged with 12.9g $\text{Mg}(0)$ powder (0.529 mol, 3.3 eq.), then again evacuated/Ar filled (3 X). 180 mL dry Et_2O was then added *via* canula, and the mixture heated to a gentle reflux. To this was then added ~20% of a solution of 100g bromide (0.529 mol, 3.3 eq.) in 200 mL Et_2O . The mixture was refluxed 10 min., then a few crystals of I_2 were added, causing Grignard formation to begin. The remaining bromide solution was then added dropwise so as to maintain reflux. When the addition was complete, the mixture was then refluxed 2h, cooled to 0°C , then 20.6 mL diethylphosphite (160 mmol, 1 eq.) in 30 mL Et_2O was added dropwise over 15 minutes. The

mixture was then stirred 18h at RT, then cooled to 0°C and cautiously quenched with 600 mL 1N HCl. 400 mL EtOAc was then added, and the mixture stirred vigorously for 30 min., then the phases were separated. The aq. phase was reextracted with EtOAc (2 X 400 mL), and the combined EtOAc phases dried (MgSO_4) and the solvents removed *in vacuo* to give a yellow oil which crystallized on standing. The resultant solid was then triturated with ~200 mL hot 10:1 Hex: EtOAc and filtered and air dried to give 35.5 g **7** (83%) as a colorless solid. m.p. $76.9\text{--}79.6^\circ\text{C}$. ^1H NMR (500 MHz, CDCl_3) δ : 8.01 (d, $^1J_{\text{HP}} = 488$ Hz, 1H), 7.35 (d, $J = 14$ Hz, 2H), 7.17 (dd, $J = 14.6, 7.8$ Hz, 2H), 7.11 (d, $J = 9.5$ Hz, 2H), 2.44 (s, 6H). ^{13}C NMR (100 MHz, CDCl_3) δ : 162.84 (s, $^1J_{\text{CF}} = 249$ Hz, $^3J_{\text{CP}} = 19.1$ Hz), 142.29 (s, $^3J_{\text{CF}} = 14.8$ Hz, $^3J_{\text{CP}} = 7.3$ Hz), 133.11 (s, $^1J_{\text{CP}} = 100$ Hz, $^3J_{\text{CF}} = 4.4$ Hz), 127.09 (d, $^2J_{\text{CP}} = 10.6$ Hz, $^4J_{\text{CF}} = 2.7$ Hz), 120.83 (d, $^2J_{\text{CF}} = 20.8$ Hz, $^4J_{\text{CP}} = 2.2$ Hz), 114.56 (d, $^2J_{\text{CF}} = 22.4$ Hz, $^2J_{\text{CP}} = 12.9$ Hz), 21.45 (q). ^{31}P NMR (202 MHz, CDCl_3) δ : 19.36 ($^1J_{\text{PH}} = 488$ Hz). ^{19}F NMR (376 MHz, CDCl_3) δ : -111.79, -111.81. Anal Calcd for $\text{C}_{14}\text{H}_{13}\text{F}_2\text{OP}$: C, 63.16; H, 4.92. Found: C, 63.08; H, 4.85.

Bis-(3,5-difluoro-4-methoxyphenyl)-phosphine oxide 8.

Prepared from 3,5-difluoro-4-methoxy-phenylmagnesium bromide and diethylphosphite by General Procedure A. Chromatographed on silica gel eluting with 1:1 EtOAc :Hexane, then recrystallized from hot cyclohexane to give the product (80%) as a colorless solid. M.p. $67.0\text{--}67.5^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.97 (d, $^1J_{\text{HP}} = 496$ Hz, 1H), 7.28-7.20 (m, 4H), 4.11 (s, 6H). ^{13}C NMR (125 MHz, CDCl_3) δ : 155.39 (C3, ddd, $^1J_{\text{CF}} = 254$ Hz, $^3J = 21.4$ Hz, $^3J = 5.4$ Hz), 140.44 (C4, t, $^2J_{\text{CF}} = 11.2$ Hz), 124.23 (C1, dt, $^1J_{\text{CP}}$

= 106 Hz, $^3J_{\text{CF}} = 6.2$ Hz), 114.67 (C2, 10 lines), 61.57 (C5, t, $^4J_{\text{CF}} = 3.9$ Hz). ^{31}P NMR (202 MHz, CDCl_3) δ : 16.64 ($^1J_{\text{PH}} = 496$ Hz). ^{19}F NMR (376 MHz, CDCl_3) δ : -125.03 Anal. Calcd for $\text{C}_{14}\text{H}_{11}\text{F}_4\text{O}_3\text{P}$: C, 50.31; H, 3.32; Found C, 50.38; H, 3.14.

Bis-(3,5-dichlorophenyl)-phosphine oxide 9.

Prepared from 3,5-dichlorophenylmagnesium bromide and diethylphosphite by General Procedure A. The crude material was recrystallized from 1:1 EtOAc:Heptane to give the product (68%) as a colorless solid. M.p. 145-146 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.02 (d, $^1J_{\text{HP}} = 501$ Hz, 1H), 7.60 (m, 6H). ^{31}P NMR (162 MHz, CDCl_3) δ : 16.07 ($^1J_{\text{PH}} = 502$ Hz). Anal. Calcd for $\text{C}_{12}\text{H}_7\text{Cl}_4\text{OP}$: C, 42.39; H, 2.08; Found C, 42.00; H, 2.50.

Bis-(3,5-difluorophenyl)-phosphineoxide 10.

A 4-neck 500 mL jacketed flask fitted with a mechanical stirrer, addition funnel, thermocouple and inert gas inlet was evacuated/Ar filled (2X), then charged with 21.9 mL 3,5-difluorobromobenzene (36.6g, 190 mmol, 3.30 eq.) and 90 mL THF. The solution was cooled to -11°C by means of a digitally-controlled chiller (an intimate ice/salt mixture can also be used) under Ar. The addition funnel was then charged with 100 mL 2M i-PrMgCl/THF (200 mmol, 3.47 eq.) via canula under N_2 pressure. Dropwise Grignard addition was then started, and maintained at such a rate that the internal temperature did not exceed -5°C (with an ice/salt bath, it was necessary to install a new ice/salt bath to maintain the internal temperature <-5°C

after about 50% of the Grignard solution had been added). The addition required 30 minutes. At the completion of the addition, 15 mL THF was used to wash the addition funnel walls into the flask. The internal temperature was then raised to 0 °C, and maintained at this temperature for 1 hour. At this time an aliquot was quickly withdrawn and quenched into 4N HCl, then analyzed by HPLC. This showed only ~4% of the starting arylbromide (relative retention time, **RRT**, = **1.00**) remained, and 3,5-difluorobenzene was present (**RRT 0.87**).

The reaction mixture was cooled to -10°C, then the addition funnel was charged with 30 mL THF, followed by 7.16 mL 98% diethylphosphite (54.5 mmol, 1 eq.). This solution was then added dropwise over 25 minutes, causing the internal temperature to rise to 5°C. The addition funnel walls were again washed into the reaction flask with 15 mL THF. The cold bath was then removed and the reaction mixture was stirred at ambient temperature. After 1.5 hours at ambient temperature, HPLC analysis showed the formation of the desired product (**RRT** = **0.77**) was no longer increasing. The reaction mixture was then cooled to -5°C and the addition funnel charged with 100 mL 3N HCl. This acid solution was then added dropwise over 20 minutes, as the internal temperature rose to 10°C. 100 mL MTBE was then added, the bath was removed, and the mixture stirred well for 5 minutes. The phases were separated, and the aq. phase was then reextracted with MTBE (1 X 100 mL). The combined organic phases were dried (MgSO_4), and the solvents removed in vacuo. The residue was then dried under high vacuum (50°C/~1 mm) for 15 minutes to remove any fluoroarenes, to give 16 g of an oily solid.

This material was suspended in 50 mL boiling hexane, then EtOAc was added until a clear solution was just obtained (~10 mL). The resulting mixture was then allowed to cool slowly with stirring to room temperature. After one hour at ambient temperature, the resultant slurry was filtered, using ~25 mL hexane

to transfer the slurry and wash the cake. The solids were dried on the frit with N₂ blowing vigorously on them for 1 hour to give 12.7g (85%) of SPO **10** as a colorless solid. M.p. 88-90°C; ¹H NMR (400 MHz, CDCl₃) δ: 8.06 (d, ¹J_{HP} = 499 Hz, 1H), 7.29-7.25 (m, 2H), 7.24-7.21 (m, 2H), 7.06 (dt, *J* = 8.0, 2.0 Hz, 2H). ¹³C NMR (100 MHz, CDCl₃) δ: 163.31 (C3, ddd, ¹J_{CF} = 256 Hz, *J* = 21.1, 11.1 Hz) 134.24 (C1, dt, ¹J_{CP} = 101 Hz, ³J_{CF} = 7.0 Hz), 113.59 (C2, 7 lines), 108.86 (C4, t, ²J_{CF} = 25.2 Hz). ³¹P NMR (202 MHz, CDCl₃)δ: 16.52 (¹J_{PH} = 499 Hz). ¹⁹F NMR (376 MHz, CDCl₃)δ: -105.43 Anal. Calcd for C₁₂H₇F₄OP: C, 52.57; H, 2.57; Found C, 52.45; H, 2.50;

Bis-(2-methylphenyl)-phosphine oxide **11**.

Prepared from 2-methylphenylmagnesium bromide and diethylphosphite by General Procedure A. The crude product was triturated with hexane to give **11** (69%) as a colorless solid. M.p. 97-98°C; ¹H NMR (500 MHz, CDCl₃) δ: 8.21 (d, ¹J_{HP} = 476 Hz, 1H), 7.71 (dd, *J* = 15.2, 7.4 Hz, 2H), 7.47 (t, *J* = 7.5 Hz, 2H), 7.33 (t, *J* = 7.5 Hz, 2H), 7.25 (dd, *J* = 12.4, 7.1 Hz, 2H), 2.38 (s, 6H). ³¹P NMR (202 MHz, CDCl₃)δ: 17.82 (¹J_{PH} = 476 Hz).

Bis-(4-dimethylaminophenyl)-phosphine oxide **12**.

Prepared from 0.5M 4-dimethylaminophenylmagnesium bromide/THF and diethylphosphite by General Procedure A. The crude material was recrystallized from hot EtOAc to give the product (57%) as a colorless solid. M.p. 151-154°C; ¹H NMR (500 MHz, CDCl₃) δ: 7.96 (d, ¹J_{HP} = 472 Hz, 1H), 7.51 (m, 4H), 6.73 (m, 4H), 3.01 (s, 12H). ¹H NMR (400 MHz, ⁶d acetone) δ: 7.82 (d, ¹J_{HP} = 460 Hz, 1H), 7.47 (dd, *J* = 8.8, 12.8 Hz, 2H), 6.77 (dd, *J* = 2.4, 8.8 Hz, 2H), 3.03 (s, 12H). ³¹P NMR (202 MHz, CDCl₃)δ: 22.31 (¹J_{PH} = 472 Hz). ³¹P NMR (202 MHz, ⁶d acetone)δ: 18.12 (¹J_{PH} = 460 Hz). Anal. Calcd for C₁₆H₂₁N₂OP: C, 66.65; H, 7.34; N, 9.72; Found C, 66.49; H, 7.50; N, 9.45.

Bis-(4-methoxyphenyl)-phosphine oxide **13**.

Prepared from 4-methoxyphenylmagnesium bromide and diethylphosphite by General Procedure A. The crude material was chromatographed on silica gel eluting with 2% MeOH/CH₂Cl₂, then recrystallized from hot EtOAc to give the product (43%) as a colorless solid. ¹H NMR (500 MHz, CDCl₃) δ: 8.06(d, ¹J_{HP} = 478 Hz, 1H), 7.63 (dd, *J* = 13.2, 8.5 Hz, 4H), 7.02 (dd, *J* = 8.6, 1.7 Hz, 4H), 3.88 (s, 6H). ³¹P NMR (202 MHz, CDCl₃)δ: 21.01 (¹J_{PH} = 478 Hz).

Bis-(2-methoxyphenyl)-phosphine oxide 14.

Prepared from 2-methoxyphenylmagnesium bromide and diethylphosphite by General Procedure A. The crude product was slurried in hot Et₂O, filtered hot, and the product (12%, first crop only) crystallized from the filtrate as a colorless solid. M.p. 130-132°C; ¹H NMR (500 MHz, CDCl₃) δ: 8.28(d, ¹J_{HP} = 517 Hz, 1H), 7.70 (ddd, *J* = 9.0, 7.5, 1.6 Hz, 2H), 7.54 (t, *J* = 8.2 Hz, 2H), 7.09 (dt, *J* = 7.5, 1.1 Hz, 2H), 6.93 (dd, *J* = 8.2, 4.6 Hz, 2H), 3.79 (s, 6H). ³¹P NMR (202 MHz, CDCl₃)δ: 9.26 (¹J_{PH} = 518 Hz).

Dicyclohexylphosphine oxide 18.

Prepared from cyclohexylmagnesium bromide and diethylphosphite by General Procedure A. The crude product was chromatographed on silica gel eluting with EtOAc to give an oil which rapidly crystallized on standing to give the product (91%) as a colorless solid. M.p. 71-73°C; ¹H NMR (400 MHz, CDCl₃) δ: 6.32 (d, ¹J_{HP} = 432 Hz, 1H), 2.00-1.12 (m, 22H). ³¹P NMR (162 MHz, CDCl₃)δ: 49.84 (¹J_{PH} = 433 Hz). Anal. Calcd for C₁₂H₂₃OP: C, 67.26; H, 10.82; Found C, 67.20; H, 11.00.

Di-tert-butylphosphine oxide 19.

¹H NMR (500 MHz, CDCl₃) δ: 6.30 (d, ¹J_{HP} = 453 Hz, 1H), 1.33 (d, *J* = 16.3 Hz, 18H). ³¹P NMR (202 MHz, CDCl₃)δ: 73.53 (¹J_{PH} = 453 Hz).

Bis-(3,5-di-(trifluoromethyl)phenyl)-phosphine oxide 20.

A 4-neck 500 mL flask was fitted with an addition funnel, thermocouple, and inert gas inlet, then charged with 20.7 mL 3,5-bis(trifluoromethyl)bromobenzene (120 mmol, 3.3 eq.). The flask was then evacuated/Ar filled (2X), and then charged with 20 mL THF. The solution was cooled to -78°C under Ar, then the addition funnel was charged with 60 mL 2M i-PrMgCl/THF (untitrated, nominally 120 mmol, 3.3 eq.) via canula under N₂ pressure. The Grignard solution was then added dropwise over 15 minutes, causing the internal temperature to rise to -60°C. The bath was then removed and replaced by a ice/H₂O bath. The internal temperature rapidly rose to 11°C, then began to fall again. An aliquot was removed by syringe, quenched into 2N HCl and analyzed by HPLC (C18, gradient, H₂O to MeCN). 7% of the arylbromide (relative retention time, RRT = 1.00) still remained, and 1,3-bis(trifluoromethyl)benzene was present at RRT 0.93. The slurry was thus recooled to -78°C and an additional 4.2 mL 2M i-PrMgCl/THF (8.4 mmol, 7% of original charge) was added by syringe, and an ice/H₂O bath installed again as described before. HPLC analysis of an aliquot at ~0°C showed <1% of the arylbromide remained. The walls of the addition funnel were washed into the flask with 5 mL of THF. The mixture was then cooled to -78°C and the addition funnel was charged with 10 mL THF + 4.79 mL diethylphosphite (98%, 36.4 mmol, 1 eq.). This solution was then added dropwise over 10 minutes, causing

the internal temperature to rise to -50°C . The cold bath was then removed, and the reaction allowed to warm to ambient temperature. After 1 hour at ambient temperature, HPLC indicated the reaction was complete to **SPO 20**, with RRT 0.96. The reaction mixture was then cooled to 2°C and 60 mL 3N HCl was added by addition funnel over 15 minutes. 50 mL sat's NaCl was then added, followed by 100 mL EtOAc. The phases were separated and the aqueous phase reextracted with 100 mL EtOAc. The combined organic phases were dried with a large amount of MgSO_4 , filtered, and the solvents removed from the filtrate first under house vacuum, then under high vacuum ($50^{\circ}\text{C}/1\text{mm}$, 15 minutes) to remove the 1,3-bis(trifluoromethyl)benzene byproduct. The resulting orange oil (19g) was dissolved in 30 mL boiling MTBE and allowed to cool to ambient temperature under Ar, with stirring, causing a thick slurry to form. 50 mL hexane was slowly added to this slurry, the mixture stirred 10 minutes, and filtered through a medium-fritted filter funnel, with a vigorous stream of N_2 flowing over the filter cake. After 2h, 11.8g **20** (68%) was obtained as a colorless solid. M.p. $127\text{--}128^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 8.25 (d, $^1J_{\text{HP}} = 504$ Hz, 1H), 8.20 (d, $J = 7.1$ Hz, 4H), 8.06 (s, 2H). ^{13}C NMR (150 MHz, CDCl_3) δ : 133.22 (C3, dq, $^2J_{\text{CF}} = 34$ Hz, $^3J_{\text{CP}} = 13$ Hz), 133.15 (C1, d, $^1J_{\text{CP}} = 101$ Hz), 130.75 (C2, dq, $^2J_{\text{CP}} = 12$ Hz, $^3J_{\text{CF}} = 3$ Hz), 127.22 (C4, m), 122.49 (C5, q, $^1J_{\text{CF}} = 273$ Hz). ^{31}P NMR (202 MHz, CDCl_3) δ : 14.46 ($^1J_{\text{PH}} = 505$ Hz). ^{19}F NMR (376 MHz, CDCl_3) δ : -63.03. Anal. Calcd for $\text{C}_{16}\text{H}_7\text{F}_{12}\text{OP}$: C, 40.53; H, 1.49; Found C, 40.45; H, 1.45.

Bis-(2,4,6-trimethylphenyl)-phosphine oxide **21**.

Prepared from 2,4,6-trimethylphenylmagnesium bromide and diethylphosphite by General Procedure A. The crude material was recrystallized from hot EtOAc to give the product (80%) as a colorless solid. M.p. $148\text{--}149^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 8.56 (d, $^1J_{\text{HP}} = 482$ Hz, 1H), 6.87 (d, $J = 3.5$ Hz, 4H), 2.40 (s, 12H), 2.29 (s, 6H). ^{31}P NMR (202 MHz, CDCl_3) δ : 11.91 ($^1J_{\text{PH}} = 480$ Hz). Anal. Calcd for $\text{C}_{18}\text{H}_{23}\text{OP}$: C, 75.50; H, 8.10; Found C, 75.22; H, 8.01.

General Procedure B: Non-Symmetric Secondary Phosphine Oxides.

Phenylisopropyl phosphine oxide **15**.

A 250 mL 3-neck RBF fitted with an addition funnel was evacuated/Ar filled (3X), then charged with 36.7 mL 2M *i*-PrMgCl/THF (73.4 mmol, 2.2 eq.) and 25 mL THF, then cooled to -10°C under Ar in an ice/salt bath. A solution of 5.00 mL ethylphenylphosphinate (33.3 mmol, 1 eq.) in 20 mL THF was then added dropwise via addition funnel over 20 minutes. The bath was then removed, the mixture stirred 1 hour at 23°C , then cooled to 0°C and quenched by the dropwise addition of 35 mL 1N HCl over 15 minutes. 50 mL Et_2O was added, agitated 5 minutes, then the upper organic phase was decanted (and saved) from a gel-like lower phase. To the lower phase was then added 100 mL additional 1N HCl, and it was extracted with CH_2Cl_2 (1 X 100 mL). The combined organics were dried (MgSO_4), and the solvents removed first under house vacuum ($35^{\circ}\text{C}/60$ mm), then high vacuum ($35^{\circ}\text{C}/1$ mm) to give 5.35g product **15** (96%) as a colorless oil. ^1H NMR (400 MHz, ^6d acetone) δ : 7.85-7.74 (m, 2H), 7.68-7.55 (m, 3H), 7.25 (dd, $^1J_{\text{HP}} = 455$ Hz, $^3J_{\text{HH}} = 2.4$ Hz, 1H), 2.22-2.10 (m, 1H), 1.59 (dd, $J = 17.1$, 7.1 Hz, 3H), 1.06 (dd, $J = 17.1$, 7.1 Hz, 3H). ^{13}C NMR (100 MHz, ^6d acetone) δ : 133.12 (d, $J_{\text{CP}} = 2$ Hz), 132.07 (s, $^1J_{\text{CP}} = 64$ Hz), 131.35 (d, $J_{\text{CP}} = 11$ Hz), 129.66 (d, $J_{\text{CP}} = 11$ Hz), 29.24 (d, $^1J_{\text{CP}} = 68$ Hz), 15.77 (q), 14.79 (q). ^{31}P NMR (162 MHz, ^6d acetone) δ : 35.25 ($^1J_{\text{PH}} = 455$ Hz). Anal. Calcd for $\text{C}_9\text{H}_{13}\text{OP}$: C, 64.28; H, 7.79; Found C, 63.90; H, 8.05.

Phenyl(*t*-butyl) phosphine oxide 16.

Prepared by General Procedure B using 1M *t*-BuMgCl/THF. The crude product was distilled (b.p. 160 °C/2mm) to give an oil which rapidly crystallized to give the product (78%) as a colorless solid. M.p. 49-51°C; ¹H NMR (400 MHz, CDCl₃) δ: 7.68-7.62 (m, 2H), 7.58-7.50 (m, 3H), 7.00 (d, ¹J_{HP} = 453 Hz, 1H), 1.12 (d, *J* = 16 Hz, 9H). ³¹P NMR (162 MHz, CDCl₃) δ: 47.63 (¹J_{PH} = 454 Hz). Anal. Calcd for C₁₀H₁₅OP: C, 65.92; H, 8.30; Found C, 65.75; H, 8.55.

General Procedure C: Non-Cryogenic Reduction of Secondary Phosphine Oxides.

Bis-(*p*-tolyl)phosphine 25.

See photographs in Figures 1-3. A 4-neck jacketed flask (Figure 1, reactor #1) equipped with a bottom drain valve containing a 24/40 joint, mechanical stirrer, addition funnel, and an argon sparge line was evacuated/Ar filled (3X), then charged with 45 mL 1M DIBAL-H/THF (45.0 mmol, 3 eq.) through the addition funnel. 10 mL THF was then used to wash the walls of

the addition funnel into the flask (Figure 1). The addition funnel was then charged with a solution of 3.45g SPO 4 (15.0 mmol, 1 eq.) in 40 mL THF. This solution was then added dropwise over 15 minutes (CAUTION: gas evolution). 10 mL THF was then used to wash the walls of the addition funnel into the flask. The resulting mixture was then stirred 20 minutes at ambient temperature, then 100 mL MTBE was added dropwise via addition funnel over 10 minutes. After 10 minutes, the sparge line was inserted beneath the solution surface and a slow argon flow was started and maintained throughout all subsequent operations. After sparging for 10 minutes, the mixture was cooled to an internal temperature of 5°C. The addition funnel was then charged with 50 mL 2N NaOH. This solution was then added dropwise via addition funnel over 20 minutes. 15 mL sat'd aq. NaCl was then added via addition funnel over 10 minutes. After 5 minutes the agitation was stopped, and the lower aqueous phase was drained to waste through the bottom drain valve. 50 mL 5% (w/v) NaCl in H₂O was then added dropwise via addition funnel over 10 minutes, and the lower aqueous phase again discarded to waste as described above.

The ground glass joint of the bottom drain valve was then mated to an Air-free© filter containing 10g MgSO₄ (Figure 2), which was in turn mated to a 4-neck receiver RBF (reactor #2) equipped with a stir bar, an addition funnel, a septum inlet, and a valve connected to a double manifold (Figure 3). While the reactor bottom drain valve was still closed, the Air-free© filter/receiver RBF assembly was inerted by evacuation/Ar filling (3X), then the reactor drain valve was opened, and the reaction mixture was dropped into the Air-free© filter. House vacuum was then gently applied from the valve on the receiver RBF, filtering the solution into the receiver. 25 mL MTBE was then added dropwise via addition funnel to the first reactor over 10 minutes, and it was degassed by Ar sparging for a full 10 minutes, then this solution was dropped into the filter as a wash of the solids. An aliquot of the filtrate was removed by syringe through the septum inlet and added to an inerted screw-cap NMR tube containing a small amount of ⁸d THF. ³¹P NMR showed a single species with δ -43.02 ppm, ¹J_{PH} 217 Hz, the desired secondary phosphine 25. The addition funnel on the receiver was then charged with 40 mL 1M BH₃/THF (40 mmol, 2.7 eq.). This borane solution was then added over 20 minutes (CAUTION: gas evolution). After 15 minutes, an aliquot was removed for ³¹P NMR as described above, showing a single species at δ -2.80 ppm, the desired

phosphine borane complex. The solution was then cooled to 5°C, then 70 mL 1N HCl was added dropwise via addition funnel, the phases were separated, and the aqueous phase reextracted with 50 mL MTBE. The combined organics were washed with sat'd NaHCO₃ (1 X 100 mL), dried (MgSO₄), and the solvents removed in vacuo to give 4.0g of a colorless oil. This was chromatographed through a short column of silica gel eluting with 4:1 Hex:EtOAc to give 2.75g of the phosphine borane complex (80%) as a colorless oil which crystallized on standing in the freezer. ¹H NMR (500 MHz, C₆D₆) δ: 7.59 (dd, *J* = 11, 7.6 Hz, 4H), 7.29 (d, *J* = 7.7 Hz, 4H), 6.29 (dq, ¹*J*_{HP} = 378 Hz, ³*J*_{HH} = 6.8 Hz, 1H), 2.42 (s, 6H). ¹H NMR (500 MHz, ⁸d THF) δ: 7.62 (m, 4H), 7.30 (d, *J* = 7.8 Hz, 4H), 6.31 (dq, *J* = 379 Hz, 7 Hz, 1H), 2.39 (s, 6H). ¹³C NMR (150 MHz, ⁸d THF) δ: 142.82 (s, ⁴*J*_{CP} = 3.8 Hz), 133.85 (d, *J*_{CP} = 15.5 Hz), 130.58 (d, *J*_{CP} = 16.9 Hz), 125.18 (¹*J*_{CP} = 93.0 Hz), 21.61 (q). ³¹P NMR (202 MHz, C₆D₆) δ: -0.46 (d, ¹*J*_{PB} = 59 Hz, ¹*J*_{PH} = 380 Hz). ³¹P NMR (202 MHz, ⁸d THF) δ: -1.13 (¹*J*_{PB} = 51 Hz, ¹*J*_{PH} = 371 Hz).

Diphenylphosphine 22.

Prepared by General Procedure C with 3 eq. DIBAL-H/THF to give the product (86%) as a colorless oil. ¹H NMR (500 MHz, C₆D₆) δ: 7.50-7.47 (m, 4H), 7.14-7.12 (m, 6H), 5.32 (d, ¹*J*_{HP} = 215 Hz, 1H). ³¹P NMR (202 MHz, C₆D₆) δ: -27.08 (¹*J*_{PH} = 215 Hz).

Bis(4-fluorophenyl)-phosphine 23.

Prepared by General Procedure C with 3 eq. DIBAL-H/THF to give the product (90%) as a colorless oil. ¹H NMR (500 MHz, C₆D₆) δ: 6.97 (m, 4H), 6.61 (t, *J* = 8.4 Hz, 4H), 4.90 (¹*J*_{HP} = 216 Hz, 1H). ¹H NMR (400 MHz, ⁸d THF) δ: 7.46-7.37 (m, 4H), 7.02-6.92 (m, 4H). ¹⁹F NMR (376 MHz, ⁸d THF) δ: -115.72. ³¹P NMR (202 MHz, C₆D₆) δ: -30.65 (¹*J*_{PH} = 218 Hz). ³¹P NMR (162 MHz, ⁸d THF) δ: -46.22 (¹*J*_{PH} = 217 Hz).

Bis(4-chlorophenyl)-phosphine 24.

Prepared by General Procedure C with 3 eq. DIBAL-H/THF to give the product (83%) as a colorless oil. ¹H NMR (500 MHz, C₆D₆) δ: 7.10-7.04 (m, 8H), 4.99 (d, ¹*J*_{HP} = 217 Hz, 1H). ¹H NMR (500 MHz, ⁸d THF) δ: 7.38 (m, 4H), 7.27 (d, *J* = 10 Hz, 4H), 5.14 (d, ¹*J*_{HP} = 216 Hz, 1H). ³¹P NMR (202 MHz, C₆D₆) δ: -30.15 (¹*J*_{PH} = 217 Hz). ³¹P NMR (202 MHz, ⁸d THF) δ: -45.58 (¹*J*_{PH} = 219 Hz).

Bis(3-fluorophenyl)-phosphine 26.

Prepared by General Procedure C with 3 eq. DIBAL-H/THF to give the product (89%) as a colorless oil. ^1H NMR (500 MHz, C_6D_6) δ : 6.98 (t, $J = 7.5$ Hz, 2H), 6.89 (t, $J = 7.5$ Hz, 2H), 6.73 (m, 2H), 6.63 (dt, $J = 8.6, 2.2$ Hz, 2H), 4.90 (d, $^1J_{\text{HP}} = 218$ Hz, 1H). ^1H NMR (500 MHz, ^8d THF) δ : 7.54-7.27 (m, 6H), 7.19 (t, $J = 8.0$ Hz, 2H), 5.39 (d, $J = 218$ Hz, 1H). ^{19}F NMR (376 MHz, ^8d THF) δ : -115.35. ^{31}P NMR (202 MHz, C_6D_6) δ : -27.10 ($^1J_{\text{PH}} = 219$ Hz). ^{31}P NMR (202 MHz, ^8d THF) δ : -42.30 ($^1J_{\text{PH}} = 217$ Hz).

Bis(3-chlorophenyl)-phosphine 27.

Prepared by General Procedure C with 3 eq. DIBAL-H/THF to give the product (81%) as a colorless oil. ^1H NMR (500 MHz, C_6D_6) δ : 7.44 (m, 2H), 7.07-7.02 (m, 4H), 6.77 (m, 2H), 4.95 (d, $^1J_{\text{HP}} = 219$ Hz, 1H). ^1H NMR (500 MHz, ^8d THF) δ : 7.57-7.26 (m, 8H), 5.29 (d, $^1J_{\text{HP}} = 220$ Hz, 1H). ^{31}P NMR (202 MHz, C_6D_6) δ : -27.15 ($^1J_{\text{PH}} = 219$ Hz). ^{31}P NMR (202 MHz, ^8d THF) δ : -42.46 ($^1J_{\text{PH}} = 221$ Hz).

Bis-(3-fluoro-5-methyl-phenyl)-phosphine. 28.

Prepared by General Procedure C with 3 eq. DIBAL-H/THF to give the product (82%) as a light yellow oil. ^1H NMR (500 MHz, C_6D_6) δ : 7.07 (t, $J = 7.3$ Hz, 2H), 7.00 (d, $J = 7.8$ Hz, 2H), 6.63 (d, $J = 9.6$ Hz, 2H), 5.11 (d, $^1J_{\text{HP}} = 218$ Hz, 1H). ^1H NMR (400 MHz, CDCl_3) δ : 7.10-7.08 (d, $J = 7.8$ Hz, 2H), 6.98-6.94 (t, $J = 7.93$ Hz, 2H), 6.86-6.83 (d, $J = 9.6$ Hz, 2H), 5.19 (d, $^1J_{\text{HP}} = 220$ Hz, 1H), 2.30 (s, 6H). ^{19}F NMR (376 MHz, CDCl_3) δ : -113.98. ^{31}P NMR (202 MHz, C_6D_6) δ : -26.37 ($^1J_{\text{PH}} = 217$ Hz). ^{31}P NMR (162 MHz, CDCl_3) δ : -39.66 ($^1J_{\text{PH}} = 219$ Hz).

Bis-[3,5-difluoro-4-methoxyphenyl]-phosphine 29.

Prepared by General Procedure C with 3 eq. DIBAL-H to give the product (90%) as a colorless oil. ^1H NMR (500 MHz, C_6D_6) δ : 6.81 (m, 4H), 4.80 (d, $^1J_{\text{HP}} = 220$ Hz, 1H), 3.66 (s, 6H). ^1H NMR (500 MHz, THF) δ : 7.08 (t, $J = 7.0$ Hz, 4H), 5.15 (d, $^1J_{\text{HP}} = 222$ Hz, 1H), 3.90 (s, 6H). ^{13}C NMR (100 MHz, THF) δ : 155.81 (dt, $^1J_{\text{CF}} = 260$ Hz, $J = 7.0$ Hz), 129.38 (t, $J = 7.0$ Hz), 117.74 ($\underline{\text{CH}}$, dt, $J = 28, 7.0$ Hz), 112.50 (multiple lines), 61.40 (q). ^{19}F NMR (376 MHz, THF) δ : -127.28; ^{31}P NMR (202

MHz, C₆D₆) δ : -25.71 ($^1J_{\text{PH}} = 219$ Hz). ^{31}P NMR (202 MHz, THF) δ : -39.00, $^1J_{\text{PH}} = 225$ Hz.

Bis-[2-methylphenyl]-phosphine 32.

Prepared by General Procedure C with 3 eq. DIBAL-H to give the product (80%) as a colorless oil which crystallized on standing. ^1H NMR (500 MHz, C₆D₆) δ : 7.38 (t, $J = 7.0$ Hz, 2H), 7.16 (t, $J = 7.5$ Hz, 2H), 7.09 (dd, $J = 7.3, 3.3$ Hz, 2H), 7.03 (t, $J = 7.4$ Hz, 2H), 5.21 ($^1J_{\text{HP}} = 219$ Hz, 1H). ^1H NMR (500 MHz, ^8d THF) δ : 7.28-7.00 (m, 8H), 5.13 (d, $^1J_{\text{HP}} = 219$ Hz, 1H), 2.42 (s, 6H). ^{31}P NMR (202 MHz, C₆D₆) δ : -44.64 ($^1J_{\text{PH}} = 219$ Hz). ^{31}P NMR (202 MHz, ^8d THF) δ : -59.07, $^1J_{\text{PH}} = 222$ Hz.

Bis-[4-dimethylaminophenyl]-phosphine 33.

Prepared by General Procedure C using 5 eq. DIBAL-H to give the product (92%) as a colorless solid. ^1H NMR (500 MHz, C₆D₆) δ : 7.69 (t, $J = 7.6$ Hz, 4H), 6.63 (d, $J = 8.6$ Hz, 4H), 5.63 (d, $^1J_{\text{HP}} = 212$ Hz, 1H), 2.55 (s, 12H). ^1H NMR (500 MHz, ^8d THF) δ : 7.32 (m, 4H), 6.70 (d, $J = 8$ Hz, 4H), 5.08 (d, $^1J_{\text{HP}} = 200$ Hz, 1H), 3.12 (s, 6H). ^{31}P NMR (202 MHz, C₆D₆) δ : -31.97 ($^1J_{\text{PH}} = 212$ Hz). ^{31}P NMR (202 MHz, ^8d THF) δ : -45.77, $^1J_{\text{PH}} = 198$ Hz.

Bis-[4-methoxyphenyl]-phosphine 34.

Prepared by General Procedure C using 5 eq. DIBAL-H to give the product (91%) as a colorless solid. ^1H NMR (500 MHz, C₆D₆) δ : 7.27 (t, $J = 7.6$ Hz, 4H), 6.58 (d, $J = 8.4$ Hz, 4H), 5.18 (d, $^1J_{\text{HP}} = 214$ Hz, 1H), 3.10 (s, 6H). ^1H NMR (500 MHz, THF) δ : 7.41 (m, 4H), 6.89 (d, $J = 7.9$ Hz, 4H), 5.16 (d, $^1J_{\text{HP}} = 214$ Hz, 1H), 3.79 (s, 6H). ^{31}P NMR (202 MHz, C₆D₆) δ : -31.26 ($^1J_{\text{PH}} = 214$ Hz). ^{31}P NMR (202 MHz, THF) δ : -45.00, $^1J_{\text{PH}} = 214$ Hz.

Bis-[2-methoxyphenyl]-phosphine 35.

Prepared by General Procedure C using 5 eq. DIBAL-H to give the product (75%) as a colorless solid. ^1H NMR (500 MHz, ^{10}d Et₂O) δ : 7.28 (t, $J = 7.7$ Hz, 2H), 7.22 (t, $J = 7.0$ Hz, 2H), 6.85 (m, 4H), 5.23 (d, $^1J_{\text{HP}} = 224$ Hz, 1H), 3.82 (s, 6H). ^{31}P NMR (202 MHz, ^{10}d Et₂O) δ : -74.21, $^1J_{\text{PH}} = 224$ Hz.

Phenylisopropyl phosphine 36.

Prepared by General Procedure C with 3 eq. DIBAL-H to give the product (90%) as a colorless oil. ^1H NMR (500 MHz, C_6D_6) δ : 7.52-7.49 (m, 2H), 7.20-7.18 (m, 3H), 4.11 (dd, $^1J_{\text{HP}} = 205$ Hz, $^3J_{\text{HH}} = 6.0$ Hz, 1H), 2.02 (m, 1H), 1.14-1.07 (m, 6 lines, 6H). ^1H NMR (500 MHz, ^8d THF) δ : 7.57-7.20 (m, 2H), 7.37-7.31 (m, 3H), 4.02 (dd, $J = 6$, 202 Hz, 1H), 2.13 (m, 1H), 1.10-1.09 (m, 8 lines, 6H). ^{31}P NMR (202 MHz, C_6D_6) δ : -11.97 ($^1J_{\text{PH}} = 203$ Hz). ^{31}P NMR (202 MHz, ^8d THF) δ : -25.68, $^1J_{\text{PH}} = 201$ Hz.

Phenyl-t-butyl phosphine 37.

Prepared by General Procedure C using 3 eq. DIBAL-H to give the product (86%) as a colorless oil. ^1H NMR (500 MHz, C_6D_6) δ : 7.51 (m, 2H), 7.20 (m, 3H), 4.05 (d, $^1J_{\text{HP}} = 207$ Hz, 1H), 1.14 (d, $J = 12$ Hz, 9H). ^{31}P NMR (202 MHz, C_6D_6) δ : 7.90 (4 lines, $^1J_{\text{PH}} = 206$ Hz).

Di-n-butyl phosphine 38.

Prepared by General Procedure C using 3 eq. DIBAL-H to give the product (85%) as a colorless oil. ^1H NMR (500 MHz, C_6D_6) δ : 3.25 (dpent, $^1J_{\text{HP}} = 192$ Hz, $^3J_{\text{HH}} = 6.9$ Hz, 1H), 1.64-1.37 (m, 12H), 0.96 (t, $J = 7.3$ Hz, 6H). ^{31}P NMR (202 MHz, C_6D_6) δ : -55.98 ($^1J_{\text{PH}} = 192$ Hz).

Dicyclohexyl phosphine 39.

Prepared by General Procedure C using 3 eq. DIBAL-H to give the product (88%) as a colorless oil. ^1H NMR (500 MHz, C_6D_6) δ : 3.05 (dt, $^1J_{\text{HP}} = 194$ Hz, $^3J_{\text{HH}} = 5.8$ Hz, 1H), 1.92-1.67 (m, 11H), 1.50-1.26 (m, 11H). ^{31}P NMR (202 MHz, C_6D_6) δ : -14.41 ($^1J_{\text{PH}} = 194$ Hz).

Di-t-butyl phosphine 40.

Prepared by General Procedure C using 3 eq. DIBAL-H to give the product (87%) as a colorless oil. ^1H NMR (500 MHz, C_6D_6) δ : 3.28 (d, $^1J_{\text{HP}} = 199$ Hz, 1H), 1.29 (d, $J = 11.3$ Hz, 18H). ^{31}P NMR (202 MHz, C_6D_6) δ : 33.22 ($^1J_{\text{PH}} = 200$ Hz).

Bis-[3,5-di-(trifluoromethyl)phenyl]-phosphine 41.

A 250 mL 4-neck flask was equipped with an argon sparge line, addition funnel, thermocouple (through a septum), and inert gas valve. The flask was charged with 5.00g SPO **20** (10.5 mmol, 1 eq.) and 25 mL cyclohexane. To this slurry, stirring under Ar at ambient temperature, was then added 21.1 mL 1M DIBAL-H/cyclohexane (21.1 mmol, 2 eq.) dropwise via addition funnel over 5 minutes. A clear, dark solution was formed, and the internal temperature rose to 32.4°C, then began to fall. 15 minutes after the completion of the addition, 9 mL MTBE was added via addition funnel over 5 minutes. The resulting mixture was stirred a further 15 minutes, then cooled to -78 °C under Ar. The addition funnel was then

charged with 40 mL 2N NaOH, and this was added dropwise over 10 minutes, causing a thick slurry (ice) to form. The cold bath was then removed, and the mixture allowed to warm to room temperature. Agitation was stopped, and a 15 cm long needle attached to a 60 mL syringe was then inserted through the septum, the tip of the needle was pressed against the bottom of the flask, and the lower aqueous phase was withdrawn and discarded. 20 mL H_2O in a small (separate) flask was degassed by bubbling Ar through it for 10 minutes, then added to the reaction flask by syringe. The mixture was agitated for 5 minutes, then agitation was stopped and the lower aqueous phase withdrawn by syringe and discarded as before.

An Airfree© filter (see Figure 2) was charged first with 18g silica gel, then 14g MgSO_4 was added. These solids were then wetted by flowing ~100 mL 5:1 (v/v) cyclohexane:MTBE through them under gravity. The filter thus prepared was then fitted with a rubber septum and placed atop a second 4-neck 250 mL flask (see Figure 3) that was equipped with a short-path distillation head to a chilled (-78 °C) 250 mL collection flask, and an inert gas/vacuum valve. The filter/receiver flask assembly was inerted by flowing argon through it for ~5 minutes. The organic solution containing the secondary phosphine was then withdrawn by syringe and injected into the filter through the septum. Gentle vacuum was applied from the valve on the receiver flask, drawing the organic solution through the solids. At this time, 40 mL 5:1 (v/v) cyclohexane:MTBE was added to the reaction flask by syringe, and degassed by bubbling argon through it for 10 minutes. This solution was then withdrawn by syringe and injected into the filter, through the septum, as a wash of the solids. The filtered solution in the receiver flask was then subjected to distillation (40°C/60 mm) until minimal stirrable volume was achieved. The resulting viscous oil was then placed under high vacuum (23°C/1mm) for ~15 minutes, causing crystallization of the product to occur. 3.4g of phosphine **41** (71%) was thus obtained as a light yellow solid. ^1H NMR (500 MHz, C_6D_6) δ : 7.67 (s, 2H), 7.54 (d, $J = 6.2$ Hz, 4H), 4.63 (d, $^1J_{\text{HP}} = 224$ Hz, 1H). ^1H NMR (500 MHz, CDCl_3) δ : 7.80 (m, 6H), 5.39 (d, $^1J_{\text{HP}} = 222$ Hz). ^{13}C NMR (125 MHz, CDCl_3) δ : 136.38 (d, $J = 16$ Hz), 133.73 (d, $J = 18$ Hz), 132.49 (dq, $J = 5.8, 34$ Hz), 123.41 (m), 123.00 (q, $^1J_{\text{CF}} = 273$ Hz). ^{31}P NMR (202 MHz, C_6D_6) δ : -29.02 ($^1J_{\text{PH}} = 223$ Hz). ^{31}P NMR (202 MHz, CDCl_3) δ : -40.43 ($^1J_{\text{PH}} = 229$ Hz); ^{19}F NMR (376 MHz, CDCl_3) δ : -63.40.

General Procedure D: Cryogenic Reductions of Secondary Phosphine Oxides.

Bis-(3,5-difluorophenyl)-phosphine 31.

NOTE: This reduction can also be carried out at 0°C with identical results.

A jacketed 4-neck 500 mL RBF with a bottom drain valve (similar to Figure 1) was fitted with a mechanical stirrer, addition funnel and thermocouple, then evacuated/Ar filled (2X), then charged with 5.00g bis(3,5-difluorophenyl)-phosphineoxide **10** (18.2 mmol, 1.0 eq.) and 50 mL THF. The solution was cooled by means of a digitally-controlled chiller set at -30°C under Ar. The addition funnel was then charged with 54.7 mL 1M DIBAL-H/THF (54.7 mmol, 3 eq.) via syringe. When the internal temperature reached -25°C, dropwise addition of the DIBAL-H solution was started so that the internal temperature did not exceed -20°C, requiring 30 minutes for the addition. The internal temperature was then adjusted to -20°C and maintained at this temperature for 1 hour. The chiller was then set to -30°C and the addition funnel was charged with 50 mL MTBE. When the internal temperature reached -25°C, dropwise addition of MTBE was started, maintaining the internal temperature below -20°C, requiring 10 minutes. The chiller was then set to -35°C. After 5 minutes, the addition funnel was charged with 100 mL 2N NaOH, and this solution was then added dropwise, maintaining the internal temperature below -20°C: **NOTE:** The initial addition is very exothermic and the addition must be done slowly. The complete addition required 40 minutes. The addition funnel was then charged with 20 mL sat'd aq. NaCl, and this was added dropwise over 5 minutes.

The chiller was then set at 22°C. When the internal temperature had stabilized, the agitation was stopped and the lower aqueous phase was dropped to waste through the bottom drain valve. Agitation was restarted and the addition funnel was then charged with 50 mL 5% aq. NaCl solution. This solution was then added dropwise over 10 minutes. Agitation was again stopped and the lower aqueous phase dropped to waste. The ground glass joint of the bottom drain valve was then mated to an Air-free© filter containing 15g MgSO₄ (Figure 2), on top of 10g silica gel, which was in turn mated to a 4-neck receiver RBF (reactor #2) equipped with a stir bar, a short path distillation head with receiver, a septum inlet, and a valve connected to a double manifold (similar to Figure 3). While the reactor bottom drain valve was still closed, the Air-free© filter/receiver RBF assembly was inerted by evacuation/Ar filling (3X), then the reactor drain valve was opened, and the reaction mixture was dropped slowly into the Air-free© filter. House vacuum was then gently applied from the valve on the receiver RBF, slowly filtering the solution into the receiver. 25 mL MTBE was then added dropwise via addition funnel to the first reactor, and it was degassed by Ar sparging for a full 10 minutes, then this solution was dropped into the filter as a wash of the solids.

The filtrate solution was then distilled using house vacuum at ~40°C until a semi-solid mixture was obtained. High vacuum (~1 mm) was then applied at ambient temperature for 1 hour to give 3.76g secondary phosphine **31** (80%) as a colorless solid. ¹H NMR (500 MHz, C₆D₆) δ: 6.72-6.69 (m, 4H), 6.46-6.42 (dt, J = 8.9, 2.1 Hz, 2H), 4.68 (d, ¹J_{HP} = 220 Hz, 1H). ¹H NMR (400 MHz, THF) δ: 7.37-7.28 (m, 4H), 7.20-7.10 (m, 2H), 5.50 (d, ¹J_{HP} = 224 Hz, 1H). ¹³C NMR (100 MHz, THF) δ: 164.14 (C3, ddd, ¹J_{CF} = 252 Hz, J = 12.1, 9.1 Hz), 139.68 (C1, dd, J = 14.1, 7.0 Hz), 117.39 (C2, 6 lines), 105.35 (C4, t, ²J_{CF} = 25.2 Hz). ³¹P NMR (202 MHz, C₆D₆) δ: -24.94 (¹J_{PH} = 221 Hz). ³¹P NMR (162 MHz, THF) δ: -39.76, ¹J_{PH} = 224 Hz; ¹⁹F NMR (376 MHz, THF) δ: -111.73.

Bis-(3,5-dichlorophenyl)-phosphine 30.

Prepared by General Procedure D to give the product (90%) as a colorless solid. ^1H NMR (500 MHz, C_6D_6) δ : 7.05 (dd, $J = 6.6, 1.8$ Hz, 4H), 7.02 (t, $J = 1.8$ Hz, 2H), 4.62 (d, $^1J_{\text{HP}} = 221$ Hz, 1H). ^1H NMR (500 MHz, ^8d THF) δ : 7.66-7.43 (m, 6H), 5.39 (d, $^1J_{\text{HP}} = 230$ Hz, 1H). ^{31}P NMR (202 MHz, C_6D_6) δ : -26.54 ($^1J_{\text{PH}} = 221$ Hz). ^{31}P NMR (202 MHz, ^8d THF) δ : -41.17, $^1J_{\text{PH}} = 233$ Hz).

General Procedure E: Triisobutylaluminum Reduction of Secondary Phosphine Oxides.

Bis-(2,4,6-trimethylphenyl)-phosphine 42.

A jacketed 4-neck 250 mL RBF with a bottom drain valve (similar to Figure 1) was fitted with a mechanical stirrer, addition funnel, inert gas line and thermocouple, and was then evacuated/Ar filled (2X), then charged with 572mg SPO **21** (2.00 mmol, 1 eq.), followed by 10 mL PhMe. To the resulting solution was then added 4.03 mL neat TIBA (16.0 mmol, 8 eq.) via syringe over ~5 minutes. The resulting clear solution was heated to 70°C and maintained there for 24 hours, then cooled to -78°C. The addition funnel was then charged with 30 mL 2N NaOH, and this basic solution was then added dropwise over ~ 10 minutes to the cold solution causing a slurry of ice to form. The cold bath was then removed, and the mixture allowed to warm to ambient temperature with good stirring. The lower aqueous

phase was then dropped to waste. 30 mL H₂O in a flask was degassed by bubbling Ar through it for 10 minutes, then it was transferred to the flask via syringe, and the mixture agitated for 5 minutes, then the phases were allowed to settle, and the lower aqueous phase was discarded as before. The 24/40 joint on the bottom drain valve was then mated to an Airfree©filter (see Figure 2) containing 8g MgSO₄ on top of 5g Si gel which was in turn connected to a receiver flask (see Figure 3) which was equipped with a short-path distillation head to a chilled receiver. The filter/receiver flask assembly was inerted, then the bottom drain valve of the reactor flask was slowly opened. Vacuum was applied to the receiver flask, drawing the organic solution slowly through the drying agent into the receiver flask. 25 mL MTBE was then charged to the reactor flask, and it was degassed with the Ar sparge line for 10 minutes, then dropped into the Airfree©filter as a wash, then vacuum again applied to the receiver flask. Short-path distillation of the solvents was then carried out at 40-50°C under house vacuum, until minimal stirrable volume was achieved. High vacuum (~1mm) was then applied for ~30 minutes, giving the product **42** as a colorless solid, 0.41g, 75%. ¹H NMR (500 MHz, C₆D₆) δ: 6.68 (s, 4H), 5.31 (d, ¹J_{HP} = 229 Hz, 1H), 2.26 (s, 12H), 2.07 (s, 6H). ³¹P NMR (202 MHz, C₆D₆)δ: -79.59, ¹J_{PH} = 229 Hz).

Photographs of Glassware Setups

Figure 1. Reactor #1 with Jointed Bottom Drain Valve

Figure 2. Chemglass Air-free © Filter
Containing MgSO_4 on frit

Figure 3. Reactor 1/Filter/Reactor 2 Assembly

(C₆H₅)₂POH -SPO 1

Current Data Parameters
NAME SPO 1
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050804
Time 10.43
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT CDCl₃
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 90.5
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300090 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

SPO 1

Current Data Parameters

NAME 12725c
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050811
Time 10.44
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 7298.2
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

SPO 2

Current Data Parameters
 NAME spo-2
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050804
 Time 2.27
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT CDCl3
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 57
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300091 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

(4-F-C₆H₄)₂POH -SPO 2

Current Data Parameters
NAME SPO 2
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050804
Time 10.58
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 4597.6
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters
SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

(4-Cl-C₆H₄)₂POH

Current Data Parameters
NAME SPO 3
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050803
Time 13.40
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT CDCl3
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 181
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300091 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

(4-Cl-C₆H₄)₂POH

Current Data Parameters
NAME SPO 3
EXPNO 4
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050803
Time 13.43
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 64
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 11585.2
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters
SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

(4-Me-C₆H₄)₂POH -SPO 4

Current Data Parameters
 NAME SPO 4
 EXPNO 4
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050804
 Time 9.25
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT CDCl₃
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 35.9
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1299876 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

(4-Me-C₆H₄)₂POH -SPO 4

Current Data Parameters

NAME SPO 4
EXPNO 5
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050804
Time 9.26
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 3251
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

(3-F1-C6H4)2POH

Current Data Parameters
 NAME 5991007
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050808
 Time 15.56
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT CDCl3
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 90.5
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300000 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

(3-F1-C6H4)2POH

Current Data Parameters

NAME 5991007
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050808
Time 15.59
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 5792.6
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

SPO-6

Current Data Parameters
 NAME 7248-145-spo-6
 EXPNO 5
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050805
 Time 9.20
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT CDCl3
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 40.3
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1299657 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

Current Data Parameters

NAME 7248-145-spo-6
EXPNO 7
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050805
Time 9.23
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 2298.8
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

3-F, 5-Me-C6H3

Current Data Parameters
NAME 7561-041-7
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050803
Time 11.08
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT CDCl3
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 287.4
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300091 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

3-F, 5-Me-C₆H₃

Current Data Parameters

NAME 7561-041-7
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050803
Time 11.11
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 10321.3
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

6039076z
F2-OMe-SPO, CDC13

Current Data Parameters
NAME 6039076z
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050803
Time 17.43
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT CDC13
NS 32
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 161.3
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300093 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

6039076z
F2-OMe-SPO, CDCl₃

Current Data Parameters

NAME 6039076z
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050803
Time 17.46
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl₃
NS 64
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 8192
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

3,5-Cl₂-C₆H₃

Current Data Parameters
 NAME 7561-041-9
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050803
 Time 11.19
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT CDCl₃
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 456.1
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300091 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

3,5-Cl₂-C₆H₃

Current Data Parameters
NAME 7561-041-9
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050803
Time 11.23
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl₃
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 10321.3
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters
SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

3,5-F₂-C₆H₃

Current Data Parameters
 NAME 7561-041-10
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050803
 Time 11.30
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT CDCl3
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 362
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300000 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

3,5-F₂-C₆H₃

Current Data Parameters
NAME 7561-041-10
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050803
Time 11.33
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl₃
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 11585.2
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters
SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

SPO-11

Current Data Parameters
 NAME 7248-145-spo-11
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050803
 Time 17.07
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT CDCl3
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 90.5
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300088 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

Current Data Parameters
 NAME 7248-145-spo-11
 EXPNO 3
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050803
 Time 17.10
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zgpg30
 TD 65536
 SOLVENT CDCl3
 NS 16
 DS 4
 SWH 80645.164 Hz
 FIDRES 1.230548 Hz
 AQ 0.4063732 sec
 RG 8192
 DW 6.200 usec
 DE 6.00 usec
 TE 300.0 K
 D1 2.00000000 sec
 d11 0.03000000 sec
 d12 0.00002000 sec

===== CHANNEL f1 =====
 NUC1 31P
 P1 13.80 usec
 PL1 -3.50 dB
 SFO1 202.4461871 MHz

===== CHANNEL f2 =====
 CPDPRG2 waltz16
 NUC2 1H
 PCPD2 70.00 usec
 PL2 -3.50 dB
 PL12 16.02 dB
 PL13 120.00 dB
 SFO2 500.1320005 MHz

F2 - Processing parameters
 SI 32768
 SF 202.4563100 MHz
 WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 1.40

4-NMe₂-C₆H₄

Current Data Parameters
 NAME 7057126
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050803
 Time 12.00
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT CDCl₃
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 90.5
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300094 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

4-NMe₂-C₆H₄

Current Data Parameters

NAME 7057126
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050803
Time 12.03
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl₃
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 6502
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

(4-MeO-C₅H₄)₂POH -SPO 13

Current Data Parameters
 NAME 7581057-58
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050810
 Time 12.18
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT CDCl₃
 NS 64
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 256
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300092 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

4557062x
4-OMe-SPO, CDCl₃

Current Data Parameters

NAME 4557062x
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050808
Time 14.04
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl₃
NS 32
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 6502
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

2-OMe-C₆H₄

Current Data Parameters
 NAME 7561-041-14
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050803
 Time 11.40
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT CDCl₃
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 406.4
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300091 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

2-OMe-C₆H₄

Current Data Parameters
NAME 7561-041-14
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050803
Time 11.43
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 10321.3
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters
SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

C6H5-P-i-Pr

Current Data Parameters
 NAME 7057098
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050803
 Time 12.10
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT CDCl3
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 50.8
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300092 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

C6H5-P-i-Pr

Current Data Parameters

NAME 7057098
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050803
Time 12.13
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 5792.6
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

7057020z3

Ph-t-Bu-POH, CDCl₃

7.69
7.68
7.67
7.66
7.58
7.58
7.57
7.57
7.52
7.52
7.51
7.50
7.50
7.49
7.27
6.58

1.25
1.25
1.17
1.14

16 (CDCl₃)

Current Data Parameters
NAME 7057020z3
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050803
Time 18.07
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT CDCl₃
NS 32
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 181
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300095 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

7057020z3
Ph-t-Bu-POH, CDCl₃

Current Data Parameters

NAME 7057020z3
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050803
Time 17.59
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl₃
NS 32
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 10321.3
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

7057140z
n-Bu₂POH, CDCl₃

Current Data Parameters
NAME 7057140z
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050805
Time 14.14
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT CDCl₃
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 11.3
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1298506 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

7057140z
n-Bu₂POH, CDCl₃

Current Data Parameters

NAME 7057140z
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050805
Time 14.09
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl₃
NS 32
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 1149.4
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

7057023z
Cy2POH, CDCl3

Current Data Parameters
NAME 7057023z
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050804
Time 18.10
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT CDCl₃
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 32
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300107 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

7057023z
Cy2POH, CDCl₃

Current Data Parameters

NAME 7057023z
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050804
Time 18.05
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl₃
NS 64
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 3649.1
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

7057022x
t-Bu2POH, CDCl3

Current Data Parameters
NAME 7057022x
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050805
Time 15.31
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT CDCl3
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 64
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300100 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

7057022x
t-Bu2POH, CDCl3

Current Data Parameters

NAME 7057022x
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050805
Time 15.25
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 32
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 5160.6
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

Current Data Parameters
NAME 7665-002-4
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050803
Time 10.27
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT CDCl3
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 574.7
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300093 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

(3,5-CF₃)₂ -SPO

14.46

Current Data Parameters
NAME 758101
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050815
Time 11.26
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 32
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 7298.2
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters
SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

100 50 0 -50 -100 -150 -200 ppm

7057-127

Current Data Parameters
 NAME 7057-127
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050725
 Time 19.09
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT CDCl3
 NS 128
 DS 8
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 143.7
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300096 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

7057-127

Current Data Parameters

NAME 7057-127
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050725
Time 19.20
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 128
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 10321.3
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

7.50
7.49
7.49
7.48
7.47
7.47
7.28
7.14
7.13
7.13
7.12
7.12

5.53

5.10

Current Data Parameters
NAME 75119991
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050805
Time 15.52
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT C6D6
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 90.5
DW 67.800 use
DE 4.50 use
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 use
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300000 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

75119991
Ph2PH, C6D6

22 (C₆D₆)

7.6 7.4 7.2 7.0 6.8 6.6 6.4 6.2 6.0 5.8 5.6 5.4 5.2 ppm

35.82

54.95

9.23

75119991
Ph2PH, C6D6

-27.08

Current Data Parameters
NAME 75119991
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050805
Time 15.45
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 32
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 5160.6
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters
SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

22 (C₆D₆)

Current Data Parameters

NAME 7511333x
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050810
Time 13.00
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT C6D6
NS 128
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 406.4
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====

NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters

SI 65536
SF 500.1300000 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

7511333x
4-F-DPP, C6D6

-30.68

23 (C₆D₆)

Current Data Parameters
NAME 7511333x
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050810
Time 12.39
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 256
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 11585.2
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters
SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

Current Data Parameters
 NAME 7662016
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050810
 Time 16.16
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT C6D6
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 90.5
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300000 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

(4-Cl-C₆H₄)₂PH

-30.15

Current Data Parameters
NAME 7662016
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050810
Time 16.19
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 10321.3
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters
SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

150 100 50 0 -50 -100 -150 -200 ppm

Current Data Parameters

NAME 7511555x
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050809
Time 17.29
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT CDCl3
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 50.8
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====

NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters

SI 65536
SF 500.1300000 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

7511555x
p-Tol2PH.BH3, CDCl3

-0.32
-0.61

Current Data Parameters
NAME 7511555x
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050809
Time 17.20
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 128
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 5160.6
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters
SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

7.10
7.09
7.09
7.04
7.02
7.01
6.88
6.88
6.87
6.87
6.85
6.85
6.84
6.84
6.78
6.78
6.77
6.76
6.75

5.25

4.81

5991010x
3-F-DPP, C6D6

26 (C₆D₆)

Current Data Parameters
NAME 5991010x
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050806
Time 16.27
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT C6D6
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 90.5
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300000 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

7.4 7.2 7.0 6.8 6.6 6.4 6.2 6.0 5.8 5.6 5.4 5.2 5.0 4.8 ppm

22.14

22.02

22.61

21.91

11.30

5991010x
3-F-DPP, C6D6

-27.06

26 (C₆D₆)

Current Data Parameters

NAME 5991010x
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050806
Time 16.29
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 32
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 8192
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

150 100 50 0 -50 -100 -150 -200 ppm

5474190x

3-Cl-DPP, C₆D₆

7.44
7.44
7.43
7.43
7.22
7.05
7.04
7.02
6.78
6.78
6.76
6.75

5.16

4.73

Current Data Parameters

NAME 5474190x
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050806
Time 16.03
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT C₆D₆
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 90.5
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====

NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters

SI 65536
SF 500.1300000 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

5474190x
3-Cl-DPP, C6D6

-27.15

Current Data Parameters
NAME 5474190x
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050806
Time 15.58
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 32
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 7298.2
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters
SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

150 100 50 0 -50 -100 -150 -200 ppm

Current Data Parameters
NAME 6039034x
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050806
Time 16.52
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT C6D6
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 161.3
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300000 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

6039034x
3-F, 5-Me-DPP, C6D6

-26.37

28 (C₆D₆)

Current Data Parameters

NAME 6039034x
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050806
Time 16.54
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 32
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 10321.3
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

150 100 50 0 -50 -100 -150 -200 ppm

Current Data Parameters
NAME 5991124x
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050808
Time 8.19
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT C6D6
NS 32
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 322.5
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300000 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

5991124x
F2-OMe-SPP, C6D6

-25.71

Current Data Parameters

NAME 5991124x
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050808
Time 8.14
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 64
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 9195.2
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

Current Data Parameters
 NAME 5474187x
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050806
 Time 14.21
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT C6D6
 NS 32
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 287.4
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300000 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

30 (C₆D₆)

Current Data Parameters

NAME 5474187x
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050806
Time 14.24
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 64
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 10321.3
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

Current Data Parameters
NAME 7248-147-sp
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050810
Time 15.25
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT C6D6
NS 128
DS 8
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 50.8
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300000 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

Bis-3,5-difluoroSP

-24.94

31 (C₆D₆)

Current Data Parameters

NAME 7248-147-sp
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050810
Time 15.28
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 6502
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

150 100 50 0 -50 -100 -150 -200 ppm

7.16
7.14
7.10
7.09
7.09
7.08
7.05
7.03
7.02

5.43

4.99

2.40

2-MeSP

32 (C₆D₆)

Current Data Parameters
NAME 7511789y
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050815
Time 17.15
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT C6D6
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 90.5
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300000 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

7.5 7.0 6.5 6.0 5.5 5.0 4.5 4.0 3.5 3.0 2.5 ppm

12.97
13.00
11.71
14.36

6.80

41.10

7511789y
2-MeSP, C6D6

-44.65

32 (C₆D₆)

Current Data Parameters

NAME 7511789y
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050815
Time 17.21
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 8192
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

150 100 50 0 -50 -100 -150 -200 ppm

Current Data Parameters
 NAME 7662017
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050810
 Time 18.15
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT C6D6
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 90.5
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300000 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

(NMe₂-C₆H₄)₂PH

-31.97

Current Data Parameters

NAME 7662017
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050810
Time 18.19
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 11585.2
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

150 100 50 0 -50 -100 -150 -200 ppm

MeOPhSP

Current Data Parameters
 NAME 7248-148-4
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050812
 Time 17.30
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT C6D6
 NS 128
 DS 8
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 456.1
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1301182 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

MeOPhSP

-31.26

Current Data Parameters

NAME 7248-148-4
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050812
Time 17.33
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 13004
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

7.25
7.24
7.21
7.21
7.18
6.88
6.87
6.86
6.84

5.46

5.01

4.61

3.93
3.87
3.80
3.80

35 ($^{10}\text{d Et}_2\text{O}$)

Current Data Parameters

NAME 7511011z8
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050816
Time 16.09
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT DEE
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 14.3
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====

NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters

SI 65536
SF 500.1299824 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

24.97

23.31

6.36

38.60

7511011z8
2-OMeSP, EE

-74.21

35 ($^{10}\text{d Et}_2\text{O}$)

Current Data Parameters

NAME 7511011z8
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050816
Time 16.06
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT DEE
NS 32
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 7298.2
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 ^{31}P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 ^1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4563100 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

100 50 0 -50 -100 -150 -200 ppm

Current Data Parameters
 NAME 7248-146sp3
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050810
 Time 16.30
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT C6D6
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 45.3
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300000 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

PhiPrSP

-12.18

Current Data Parameters

NAME 7248-146sp3
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050810
Time 16.33
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 5792.6
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

150 100 50 0 -50 -100 -150 -200 ppm

Current Data Parameters
NAME 7511666a
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050809
Time 12.39
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT C6D6
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 143.7
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300000 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

7511666a
Ph-t-Bu-PH, C6D6

8.20
7.60

37 (C₆D₆)

Current Data Parameters
NAME 7511666a
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050809
Time 12.33
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 64
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 13004
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters
SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

Current Data Parameters
 NAME 7248-147-bu2sp
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050811
 Time 16.18
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT C6D6
 NS 128
 DS 8
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 90.5
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300000 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

Bis-nBuSP

-55.81

38 (C₆D₆)

Current Data Parameters

NAME 7248-147-bu2sp
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050811
Time 16.21
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 16
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 7298.2
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

150 100 50 0 -50 -100 -150 -200 ppm

3.25
3.23
3.13

2.87
2.86
2.85

1.92
1.90
1.80
1.80
1.79
1.77
1.76
1.70
1.67
1.52
1.50
1.49
1.47
1.34
1.32
1.31
1.28
1.27
1.26
1.24

7511777a
Cy2PH, C6D6

39 (C₆D₆)

Current Data Parameters
NAME 7511777a
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050809
Time 8.41
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT C6D6
NS 32
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 456.1
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters
SI 65536
SF 500.1300000 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

7511777b
Cy2PH, C6D6

-14.41

Current Data Parameters

NAME 7511777b
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050809
Time 9.37
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 1024
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 10321.3
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

Current Data Parameters
 NAME 7511888b
 EXPNO 3
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050809
 Time 7.55
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT C6D6
 NS 16
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 362
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300000 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

7511888b
t-Bu₂-PH, C₆D₆

33.22

40 (C₆D₆)

Current Data Parameters
NAME 7511888b
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050809
Time 7.50
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 64
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 14596.5
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters
SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

Current Data Parameters
 NAME 7511068x
 EXPNO 3
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050806
 Time 12.48
 INSTRUM spect
 PROBHD 5 mm BBI 1H-BB
 PULPROG zg10
 TD 32768
 SOLVENT C6D6
 NS 32
 DS 4
 SWH 7374.631 Hz
 FIDRES 0.225056 Hz
 AQ 2.2217205 sec
 RG 256
 DW 67.800 usec
 DE 4.50 usec
 TE 303.0 K
 D1 2.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.40 usec
 PL1 -3.50 dB
 SFO1 500.1334009 MHz

F2 - Processing parameters
 SI 65536
 SF 500.1300000 MHz
 WDW EM
 SSB 0
 LB 0.05 Hz
 GB 0
 PC 1.00

7511068x
(CF₃)₄-DPP, C₆D₆

-29.02

41 (C₆D₆)

Current Data Parameters

NAME 7511068x
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050806
Time 12.43
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C₆D₆
NS 32
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 13004
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

150 100 50 0 -50 -100 -150 -200 ppm

Mes2 SP

Current Data Parameters

NAME 7381-179-3
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050812
Time 17.56
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zg10
TD 32768
SOLVENT C6D6
NS 16
DS 4
SWH 7374.631 Hz
FIDRES 0.225056 Hz
AQ 2.2217205 sec
RG 90.5
DW 67.800 usec
DE 4.50 usec
TE 303.0 K
D1 2.00000000 sec

===== CHANNEL f1 =====

NUC1 1H
P1 7.40 usec
PL1 -3.50 dB
SFO1 500.1334009 MHz

F2 - Processing parameters

SI 65536
SF 500.1300634 MHz
WDW EM
SSB 0
LB 0.05 Hz
GB 0
PC 1.00

Mes2 SP

-79.57

Current Data Parameters

NAME 7381-179-3
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters

Date_ 20050812
Time 18.07
INSTRUM spect
PROBHD 5 mm BBI 1H-BB
PULPROG zgpg30
TD 65536
SOLVENT C6D6
NS 128
DS 4
SWH 80645.164 Hz
FIDRES 1.230548 Hz
AQ 0.4063732 sec
RG 14596.5
DW 6.200 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====

NUC1 31P
P1 13.80 usec
PL1 -3.50 dB
SFO1 202.4461871 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16
NUC2 1H
PCPD2 70.00 usec
PL2 -3.50 dB
PL12 16.02 dB
PL13 120.00 dB
SFO2 500.1320005 MHz

F2 - Processing parameters

SI 32768
SF 202.4536060 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

150 100 50 0 -50 -100 -150 -200 ppm