A statistical evaluation of the origin of cypress domes in Florida

Eron R. Raines*

School of Geography, Environment and Earth Science, Victoria University of Wellington, Wellington 6012, New Zealand
School of Earth and Environmental Sciences, University of Wollongong, Wollongong, NSW 2522, Australia

ARTICLE INFO

Article history:

Received 3 December 2019

Received in revised form 5 March 2020

Accepted 6 March 2020

Available online 13 March 2020

Keywords:

Cypress domes

Sinkholes

Big Cypress National Preserve

Ecologic drill

ABSTRACT

Two competing explanations exist for the origin of one type of karstic landform found in Florida called the cypress dome. One explanation relies on complex ecological feedbacks stemming from nutrient cycling suggesting biota contribute more significantly to processes of landscape evolution in Florida than anywhere else in the world. The second explanation is that the landforms are sinkholes that completely preclude the biological explanation while fitting more parsimoniously with the surrounding geological narrative. This work puts forward geostatistical analyses and a model linking the landforms to sinkholes, thus bolstering the geological explanation for origin of the landform. Satellite imagery of sinkholes occurring in limestone from locations spanning the planet was analyzed. Measurements of globally distributed limestone sinkhole surface areas are best characterized by an exponential distribution indicating sinkhole formation is robust to starting conditions (i.e., climate, tectonics). This observation is supported by an analysis of sinkhole geometry and geospatial dispersion. This demonstrates the geospatial parameters space for globally distributed groups of sinkholes forming in limestone are statistically indistinguishable despite sinkhole formation in different climates, tectonic regimes, and at different times. Employing this observation as a tool, sinkholes are directly compared to the cypress domes in Florida and are found to be statistically indistinguishable. From the striking similarity in spatial parameter spaces in conjunction with the geologic history of the area, it is interpreted that these landforms originate through geologic, *not* biologic, processes.

© 2020 Elsevier B.V. All rights reserved.

1. Introduction

In the shallow limestones of South Florida are cypress domes: depressional, bowl-like landforms that occur in spatial patterns corresponding with the distribution of vegetation (Fig. 1) (Vernon, 1947; Watts et al., 2014). Vegetation in cypress domes is tallest in the center and decrease in height radially, which is what gives cypress domes their dome-like appearance and their name.

The biological explanation for the origin of cypress domes is that vegetation once clustered in dense pockets where nutrients were the highest (Cohen et al., 2015). Vegetation acidified the soil to acquire these nutrients, which overtime dissolved the shallow limestone creating the bowl-like landforms we see today (Chamberlin et al., 2018; Dong et al., 2019; Zhang et al., 2019). This biological explanation is predicated on cypress domes not having originated as sinkholes (Chamberlin et al., 2018; Cohen et al., 2015; Dong et al., 2018; Dong et al., 2019; Quintero and Cohen, 2019; Zhang et al., 2019). However, limestone sinkholes do co-occur with the cypress domes and are known to create similar depressional landforms that similarly affect

the distribution of vegetation (Antonić et al., 2001; Bátori et al., 2014; Bishop, 1967; Fleming and Van Alstine, 1999; Kobal et al., 2015; Kruse et al., 2006; Metz, 2011; Petuch and Roberts, 2007; Schneider, 1973; Sinclair, 1982). Because there are sinkholes present where there are cypress domes indicates that the conditions conducive for sinkhole development have at one time existed.

The sinkhole explanation comes from observations made of the local geology. The top-most limestone layer in the area is the Tamiami formation. This formation is 15–30 m thick and it stopped forming some 2.54 Mya, giving an upper age limit on the sinkholes (Petuch and Roberts, 2007). Fluctuations in sea level over the past 2.54 Mya introduced coastal depositional environments to South Florida with the potential to have completely filled the sinkholes with sediments (Lane et al., 2011; Miller et al., 2005; Shinn et al., 1996; van Hengstum et al., 2011).

This paper develops the sinkhole explanation by statistically evaluating sinkholes and cypress domes starting from the assumptions that the conditions necessary for sinkhole formation were once present, that changes in sea level occur, and that a fluctuating sea level would have introduced the area to sufficiently high sedimentation rates resulting from introduced oceanic depositional processes (Lane et al., 2011; Miller et al., 2005; Schneider, 1973). All three assumptions have some measure of observational support.

* School of Geography, Environment and Earth Science, Victoria University of Wellington, Wellington 6012, New Zealand.

E-mail address: eron.raines@vuw.ac.nz.

Fig. 1. Elements A and B taken with permission from Watts et al. (2014). (A) Cypress domes studied are located within Big Cypress National Preserve, which occupies 295,000 ha in southern Florida, USA. Cypress domes are also found throughout Florida. (B) The dome-shaped profiles of pond cypress (*Taxodium distichum*) are distinctive in an otherwise flat landscape. (C) This Google Earth image (Google, Inc., Mountain View, CA USA) shows a 3 km² area of Cypress domes taken from the middle of the Preserve. (D–F) Show a 3 km² Google Earth image taken of sinkholes outside Livno, Canton 10, FB&H; near Vaughn, New Mexico, USA; and just outside Tan-Tan, Morocco, respectively. Vegetation differs because of the climate between C–F, which causes some visual differences between the images. However, the size, shape, and distribution of landforms within each image is not immediately dissimilar. This is striking because the landform within each image is of different age, formed in a different climate, and has been exposed to different tectonic regimes.

2. Study area

The cypress domes are located within Big Cypress National Preserve (295,500 km²) on the shallow, porous Tamiami group of limestones found in South Florida (Figs. 1 and 2). This group of limestones are 15–30 m thick and the youngest formed some 2.54 Mya (Petuch and Roberts, 2007). The cypress domes are geographically situated proximal to the Gulf of Mexico and the Atlantic Ocean with an average elevation of 4 m above mean sea level. Groundwater levels in porous limestone near the ocean rises and falls in direct response to sea level fluctuations (Fig. 1) (Brown et al., 2018; Gulley et al., 2013). Over the past 2.54 Mya, sea level has ranged from −100 to +50 m relative to current mean sea level (Miller et al., 2005). Lowering of sea level in the geologic past drastically reduced groundwater levels in South Florida creating conditions conducive for sinkhole formation (LeGrand, 1973; Tihansky, 1999). The subsequent rise in mean sea level introduced significant oceanic depositional processes ultimately responsible for the formation of what is currently the state of Florida (Petuch and Roberts, 2007). It is well established that these depositional processes fill sinkholes submerged by sea water throughout the Caribbean and Florida (Brown et al., 2013; Lane et al., 2011; Shinn et al., 1996; van Hengstum et al., 2011).

3. Methods

3.1. Constructing a geospatial database

Sinkholes forming in limestones were first positively identified in groups (groups = 23; total landforms = 2861) occurring throughout the planet (Duque et al., 2010; Harrington, 1957; Neal and Johnson, 2002; Reznichenko, 2012; Wanaïm et al., 2014; Žebre and Stepišnik, 2015). The central-most location, surface area, and perimeter of each sinkhole was measured from satellite imagery using ArcGIS. The same measurements were made on cypress domes (groups = 14; total landforms = 1502). Measurements of both landforms were then characterized using regression analyses, Average Nearest Neighbor Distances, and circularity indices (Table 1; see supplemental information: S.I. 1).

3.2. Analysis of data

Regression analyses were performed on rank-ordered area measurements made for each group to determine which scaling law, exponential or power, best fit the data (see supplemental information: S.I. 1). Because the number of parameters were the same for each regression, the best fit could be determined simply by the amount of variance explained (i.e., the R^2) by the model. In all comparisons the exponential distribution best described the scaling of the areas measured within each group.

Using Kolmogorov-Smirnov tests (KS tests) in a stepwise fashion, statistics generated for the Average Nearest Neighbor Distances and circularity indices between groups of sinkholes and the cypress domes (see supplemental information: S.I. 2) were compared. The result of the KS test (i.e., reject or fail to reject the null hypothesis that the distributions are the same) was tallied for every stepwise comparison between groups from which was generated a contingency table of count data (Table 2). To interpret the contingency table, we simplified it using Fischer's Exact Test (see supplemental information: S.I. 2). In all comparisons made using KS tests, the proportion of reject and fail to reject outcomes for each group were statistically the same (Table 3). This indicates that the size, shape, and spatial distribution of sinkholes can be used as a numerical fingerprint to positively and blindly identify a group of sinkholes as sinkholes. Using this approach, all groups of cypress domes were identified through their numerical fingerprint as sinkholes.

4. Results

Regression analysis of the surface area of sinkholes, the part that is open at the land's surface, indicated they are best described by an exponential distribution as opposed to a power-law distribution (Fig. 3), which matches past work done in the area (Quintero and Cohen, 2019; Watts et al., 2014). In studies on how landforms develop, exponential or power-law distributions are used to infer whether the genetic mechanisms responsible for a landform are most strongly affected by scale-specific or scale-free processes, respectively, and how starting conditions affected the landform's emergence as well as how ongoing processes facilitated the formation of the landform. (Kéfi et al., 2010; Scanlon et al., 2007; Yizhaq et al., 2017).

Fig. 2. Study sites span the planet with landforms (i.e., cypress domes or sinkholes) forming in spatiotemporally independent limestone outcroppings. These limestone outcroppings formed under different conditions, at different times, experience different climates, and are subjected to varying tectonic regimes.

Table 1

Compilation of geospatial parameters from the Cypress domes and sinkholes from around the world. **NNR's** > 1 indicate landforms within groups are spatially over-dispersed. **Z-scores** > 2 indicate the observed NNR is outside the 95% confidence interval. The **inverse cumulative distribution** of landform areas informs on if the landform's size is controlled by global processes (power-law) or local processes (exponential). For all sites the exponential distribution proved the best fit. **Circularity indices** range from -1 to 1 and quantify how far a shape deviates from being perfectly circular (i.e. 0).

Site	Geospatial analysis	Average nearest neighbor distances		Inverse cumulative distribution		Circularity indices	# of landforms	Latitude	Longitude
		Type (reference)	NNR	Z-score	Exponential distribution (r^2)				
Arizona, USA	Sinkholes (Neal and Johnson, 2002)	1.32	3.61	0.96	0.67	−0.04	33	34.79816	−110.588307
Castile and Leon, ESP	Sinkholes (Duque et al., 2010)	1.14	3.31	0.99	0.78	−0.05	117	42.762724	−4.130129
		1.16	4.03	0.98	0.67	−0.06	134	42.773045	−4.099769
Dakhla-Oued Ed-Dahab, MAR	Sinkholes (Wanaim et al., 2014)	1.22	2.55	0.96	0.59	−0.07	37	28.364945	−11.217788
		1.11	2.38	0.94	0.81	−0.07	128	28.1789	−11.337115
		1.21	5.47	0.93	0.85	−0.07	173	28.18979	−11.372473
		1.22	5.69	0.94	0.84	−0.08	180	28.249229	−11.349301
		1.27	8.12	0.92	0.89	−0.07	234	28.208624	−11.344012
Dinaric Alps, BIH	Sinkholes (Žebre and Stepišnik, 2015)	1.18	3.51	0.97	0.83	−0.06	92	44.247948	16.637697
		1.24	5.26	0.99	0.84	−0.05	95	44.145592	16.834408
		1.34	6.81	0.97	0.75	−0.04	102	44.237247	16.686295
		1.19	4.70	0.97	0.63	−0.07	126	44.230562	16.705928
		1.38	8.69	0.97	0.76	−0.05	129	44.198204	16.757825
		1.38	9.04	0.92	0.90	−0.04	131	44.22539	16.72594
		1.34	8.03	0.92	0.79	−0.04	142	44.212138	16.742403
		1.30	7.82	0.99	0.71	−0.06	154	44.244872	16.659978
		1.24	6.49	0.99	0.75	−0.05	166	44.217632	16.699369
		1.40	12.61	0.98	0.74	−0.04	234	44.221264	16.6799
New Mexico, USA	Sinkholes (Harrington, 1957)	1.23	2.74	0.94	0.87	−0.05	32	34.429422	−105.094851
		1.28	4.99	0.95	0.79	−0.05	73	34.452401	−105.122597
		1.37	6.58	0.97	0.85	−0.07	78	34.458729	−105.175213
		1.36	5.44	0.97	0.91	−0.08	128	34.387464	−105.138188
Timaru, NZ	Sinkholes (Reznichenko, 2012)	1.08	1.98	0.98	0.85	−0.04	140	−44.443376	170.969235
Florida, USA	Bowl-like landforms (Watts et al., 2014)	1.25	4.35	0.98	0.71	−0.06	49	26.10008	−81.093944
		1.49	8.41	0.98	0.71	−0.04	62	25.981117	−81.22647
		1.54	8.78	0.96	0.69	−0.04	62	25.939322	−80.951138
		1.16	3.98	0.95	0.65	−0.04	72	26.093673	−81.17228
		1.36	8.18	0.94	0.68	−0.04	75	25.981367	−81.205292
		1.36	7.47	0.90	0.67	−0.03	76	25.949934	−80.927787
		1.29	6.29	0.97	0.76	−0.05	79	25.893357	−81.137975
		1.42	8.59	0.99	0.79	−0.04	86	−81.072767	25.91517
		1.37	8.50	0.97	0.74	−0.03	102	26.027782	−81.001431
		1.45	8.86	0.94	0.75	−0.04	102	26.0187	−81.160097
		1.46	11.06	0.97	0.63	−0.03	140	25.92956	−81.204068
		1.31	9.13	0.89	0.80	−0.05	167	25.93031	−81.17643
		1.24	6.98	0.95	0.79	−0.05	177	25.991799	−81.078071
		1.33	10.69	0.91	0.83	−0.05	253	25.993593	−81.05631

Sinkholes within the globally distributed groups were found to be significantly over-dispersed (Table 1; Fig. 4). The shape of sinkholes determined using the circularity index (see supplemental information: S.I. 1) is consistent between all global sites (Tables 1 and 3). This consistency emerges despite differences in sinkhole age, climate, limestone chemistry, and tectonic regime, which means that the shape of sinkhole is unaffected by these geomorphic drivers.

4.1. Cypress domes

The analysis of cypress domes revealed that they, like sinkholes, are spatially over-dispersed (Table 1). The statistics from this observation matches the results of others who support the biologic explanation, indicating that the results from both studies are comparable (Watts et al., 2014). The shape of cypress domes, determined using

Table 2

Counts of the results from stepwise Kolmogorov-Smirnov tests of geospatial parameters. Fail to reject and reject results from the Kolmogorov-Smirnov tests are indicated by $p < 0.05$ and $p > 0.05$, respectively.

Test ID	Samples compared		Description	Counts of NND KS-test results		Counts of circularity KS-test results	
	Sample 1	Sample 2		$p < 0.05$	$p > 0.05$	$p < 0.05$	$p > 0.05$
A	Sinkhole subsample	Sinkhole subsample	Groups of sinkholes compared within the same site	39	71	35	75
B	All sinkholes	Sinkhole subsample	Groups from all sinkhole sites compared to groups at one sinkhole site	49	71	40	80
C	Cypress domes	Cypress domes	Groups of bowl-like depressions compared within the same site	19	36	21	34
D	Cypress domes	Sinkhole subsample	Groups of bowl-like depressions compared to groups at one sinkhole site	13	32	12	33
E	Cypress domes	All sinkholes	All bowl-like landform groups compared to all sinkhole groups	94	148	72	170

Table 3

Comparison of the KS-test results using Fisher's Exact Test: a proportions test used to determine if the $p < 0.05$ to $p > 0.05$ count ratio is the same for the Test ID's compared. The Upper and Lower Limits are determined from the Odds Ratio. When 1 is contained within the Limits the two Test ID's are said to be the same. This is assumed a significant result with p -value's > 0.05 . For all Test ID's, and thus all sites, the $p < 0.05$ to $p > 0.05$ count ratios for both the NND and Circularity parameters are statistically the same meaning the sites are statistically indistinguishable.

Test ID's compared	NND				Circularity			
	Odds ratio	Lower limit	Upper limit	p-Value	Odds ratio	Lower limit	Upper limit	p-Value
A vs B	0.59	0.26	1.30	0.21	0.73	0.31	1.64	0.46
A vs C	1.30	0.51	3.35	0.67	1.69	0.67	4.42	0.29
A vs D	0.74	0.32	1.66	0.46	0.78	0.33	1.78	0.57
A vs E	0.64	0.29	1.33	0.24	0.86	0.38	1.83	0.73
B vs C	0.77	0.37	1.56	0.50	1.23	0.60	2.51	0.61
B vs D	0.80	0.45	1.40	0.42	0.93	0.52	1.68	0.89
B vs E	0.92	0.58	1.48	0.73	0.85	0.52	1.40	0.55
C vs D	0.96	0.46	1.99	1.00	1.32	0.63	2.74	0.49
C vs E	0.83	0.42	1.59	0.64	1.46	0.75	2.79	0.26
D vs E	0.87	0.52	1.41	0.56	1.10	0.65	1.84	0.71

the circularity index (see supplemental information: S.I. 1), is consistent between all sites (Tables 1 and 3). Analysis of the surface area of these landforms revealed that their size distribution is best mathematically described by an exponential distribution as opposed to a power-law distribution (Fig. 3). This result also matches the work done by Watts et al. (2014), which indicates that the same geomorphic phenomenon they measured was also measured here with the same results and indicates that their methods were effectively duplicated.

Both the globally distributed sinkholes and cypress domes are spatially over-dispersed, have similar circularity indices, and exhibit exponential scaling of their surface area distributions (Table 1). Comparison of the over-dispersion and circularity metrics between groups indicates that the sinkholes and the cypress domes are statistically indistinguishable (Table 3).

5. Discussion

Collecting sediments from cypress domes is an approach that could be used to document depositional processes and provide a means for directly testing the sinkhole hypothesis. However, the study setting is remote and geophysically locating the covered sinkholes remains a serious challenge. The South Florida study area is characterized by a shallow subsurface rich in marl: a carbonate clay that strongly attenuates ground penetrating radar signals. Resistivity imaging would work, however, it requires some guidance from GPR, which did not work (Carbonel et al., 2014; Kruse et al., 2006).

Collecting sediment cores deep enough to begin to account for the past 2.54 Mya would require a professional drill rig, which would be nearly impossible to navigate to the sinkhole center – assuming the center of the sinkholes could be found using geophysics to begin with. Sediments collected by hand can only penetrate 2–3 m and thus only inform on very recent processes (Zhang et al., 2019). With the inherent field difficulties, geospatial analysis of satellite imagery provided the most effective method to test the biological and geological explanations.

Previous geospatial analyses conducted on satellite imagery by Watts et al. (2014) were interpreted to support the biological explanation and were subsequently used as evidence that the cypress domes did not originate as sinkholes (Chamberlin et al., 2018; Dong et al., 2018; Dong et al., 2019; Quintero and Cohen, 2019; Watts et al., 2014; Zhang et al., 2019). Such geospatial analyses are also commonly used to study sinkholes (de Carvalho et al., 2014; Šegina et al., 2018; Theilen-Willige, 2018; Theilen-Willige et al., 2014). Using satellite imagery, a geospatial analysis was conducted to characterize both the cypress domes and sinkholes located across the planet (Fig. 2). This characterization provided a numeric fingerprint to statistically evaluate cypress domes and sinkholes. The cypress dome analysis yielded results almost identical to Watts et al. (2014), which indicated their methods had been successfully replicated. The predictive ability of the numeric fingerprints was tested, and it was found that both cypress domes and sinkholes could be positively identified using them (Tables 2 and 3). The next step was to test for the similarity between cypress domes and sinkholes. Because each could be positively identified through

Fig. 3. The x-axis is figure area in m^2 . The y-axis is the proportion of the feature areas less than the maximal area measured at a given site (see SI). The yellow circles are digitized landforms. The area of the yellow circles was measured for a group to determine what scaling best described the landform's area distribution. The regression analysis was performed on a rank-ordered data set normalized by the largest area value within that group.

Fig. 4. Nearest Neighbor Ratios are used to quantify the relative spatial orientation of objects in two dimensions. The four squares on the left side are actual observations (Table 1), whereas the two squares on the right are of a random distribution.

their numeric fingerprints, the numeric fingerprint for cypress domes and sinkholes were statistically tested for similarity and they were found to be identical (Table 3, Fig. 3).

Regression analyses of the size distributions of both the sinkhole groups and groups of cypress domes revealed that both are exponentially distributed (Table 1). The exponential distribution of cypress dome sizes was interpreted as evidence for the presence of a scale-specific formation mechanism responsible for the emergence of the landform, which was attributed to biological processes affected by spatial feedbacks (Quintero and Cohen, 2019; Watts et al., 2014). These workers posited that if cypress domes had originated as sinkholes, they would exhibit power-law scaling. Their reasoning was that power-law distributions would evidence the absence of inhibitory feedbacks that these workers posited (without references) would be the case for sinkholes. However, as the geospatial analysis presented here shows, size distributions of sinkholes forming in limestone anywhere on Earth are exponentially distributed (Table 1).

6. Conclusion

This study began as an attempt to validate the biologic explanation by rigorously testing the sinkhole origin hypothesis. It was initially believed that, given the number of recent observations supporting the biologic explanation (Chamberlin et al., 2018; Cohen et al., 2015; Dong et al., 2018; Dong et al., 2019; Quintero and Cohen, 2019; Watts et al., 2014; Zhang et al., 2019), this explanation would prevail. However, employing the same techniques used to first establish the biologic explanation, it is shown here that these cypress domes are statistically indistinguishable from sinkholes. The observed exponential scaling of the cypress domes' area distributions was used in support of the biologic explanation to discredit a sinkhole origin (Quintero and Cohen, 2019; Watts et al., 2014). The rationale was that sinkholes should exhibit power-law scaling. However, this assumption had not been tested at the same spatial scales that the cypress domes were analyzed. This work shows that, at these specific spatial scales, the size distributions of sinkholes are also best described by an exponential distribution that more closely linked the cypress domes to sinkholes. Statistical evaluation of the geospatial parameters further revealed that cypress domes and sinkholes were indistinguishable. Because these geospatial parameters can be used to robustly identify both a cypress dome as a cypress

dome and a sinkhole as a sinkhole, it is hard to reconcile them as different types of landforms if, using this approach, a cypress dome is identified as a sinkhole. Given the geologic history of South Florida and the multiple statistical observations linking cypress domes to sinkholes, it is concluded here that the sinkhole explanation best describes the formation of cypress domes.

The success of the sinkhole explanation does not, of course, argue for the absence of biological processes acting on the limestones of South Florida. It is well accepted that plants do spatially cluster and dissolve rock where the relative abundance of ecologically important nutrients is highest (Hahm et al., 2014; Landeweert et al., 2001). Therefore, it is not suggested here that plants act any differently in South Florida, just that these processes are not responsible for creating landforms.

Author contributions

E.R.R. collected and tabulated data, did all analyses, and wrote the manuscript.

Declaration of competing interest

The author declares that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The Everglades Depth Estimation Network (EDEN) project and the US Geological Survey provided the regional elevation data used. The journal *Earth Surface Processes and Landforms* (Elsevier) provided permissions to reprint elements A and B of Fig. 1 from Watts et al. (2014). Jonathan Phillips, Scott Lecce, Augusto Auler, Claire Lukens, and an anonymous reviewer provided much helpful feedback and discussion that has improved this manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.geomorph.2020.107161>.

References

- Antonić, O., Hatic, D., Pernar, R., 2001. DEM-based depth in sink as an environmental estimator. *Ecol. Model.* 138 (1–3), 247–254.
- Bátori, Z., Farkas, T., Erdős, L., Tölgyesi, C., Körmöczy, L., Vojtkó, A., 2014. A comparison of the vegetation of forested and non-forested solution dolines in Hungary: a preliminary study. *Biologia* 69 (10), 1339–1348.
- Bishop, E., 1967. A Tentative Classification of Lake Shorelines in Florida Lakes. Florida Board of Conservation, Division of Water Resources, Tallahassee, FL.
- Brown, A.L., Reinhardt, E.G., van Hengstum, P.J., Pilarczyk, J.E., 2013. A coastal Yucatan sinkhole records intense hurricane events. *J. Coast. Res.* 30 (2), 418–428.
- Brown, A.L., Martin, J.B., Kamenov, G.D., Ezell, J.E., Sreaton, E.J., Gulley, J., Spellman, P., 2018. Trace metal cycling in karst aquifers subject to periodic river water intrusion. *Chemical Geology*.
- Carbonel, D., Rodríguez, V., Gutiérrez, F., McCalpin, J.P., Linares, R., Roqué, C., Zarroca, M., Guerrero, J., Sasowsky, I., 2014. Evaluation of trenching, ground penetrating radar (GPR) and electrical resistivity tomography (ERT) for sinkhole characterization. *Earth Surf. Process. Landf.* 39 (2), 214–227.
- de Carvalho, O., Guimarães, R., Montgomery, D., Gillespie, A., Trancoso Gomes, R., de Souza Martins, É., Silva, N., 2014. Karst depression detection using ASTER, ALOS/PRISM and SRTM-derived digital elevation models in the Bambuí Group, Brazil. *Remote Sensing* 6 (1), 330–351.
- Chamberlin, C.A., Bianchi, T.S., Brown, A.L., Cohen, M.J., Dong, X., Flint, M.K., Martin, J.B., McLaughlin, D.L., Murray, A.B., Pain, A., 2018. Mass balance implies Holocene development of a low-relief karst patterned landscape. *Chemical Geology*.
- Cohen, M., Quintero, C., Ward, N., Raines, E., Brown, A., Martin, J., Bianchi, T., McLaughlin, D., Osborne, T., Heffernan, J., 2015. An ecological drill: biogeomorphic pattern evolution in a low-relief carbonate landscape. *Proceedings AGU Fall Meeting Abstracts*.
- Dong, X., Cohen, M.J., Martin, J.B., McLaughlin, D.L., Murray, A.B., Ward, N.D., Flint, M.K., Heffernan, J.B., 2018. Ecohydrologic processes and soil thickness feedbacks control limestone-weathering rates in a karst landscape. *Chemical Geology*.
- Dong, X., Murray, A.B., Heffernan, J.B., 2019. Ecohydrologic feedbacks controlling sizes of cypress wetlands in a patterned karst landscape. *Earth Surf. Process. Landf.* 44 (5), 1178–1191.
- Duque, J.F.M., García, J.C., Urquí, L.C., 2010. Organización de información geomorfológica orientada a la ordenación y gestión de espacios naturales. El caso de Covalagua y Las Tuerces (Palencia, España). *Boletín de la Real Sociedad Española de Historia Natural. Sección Geológica* 104 (1), 71–92.
- Fleming, G.P., Van Alstine, N.E., 1999. Plant communities and floristic features of sinkhole ponds and seepage wetlands in southeastern Augusta County, Virginia. *Banisteria* 13, 67–94.
- Gulley, J., Martin, J., Moore, P., Murphy, J., 2013. Formation of phreatic caves in an eogenetic karst aquifer by CO₂ enrichment at lower water tables and subsequent flooding by sea level rise. *Earth Surface Processes and Landforms* 38 (11), 1210–1224.
- Hahm, W.J., Riebe, C.S., Lukens, C.E., Araki, S., 2014. Bedrock composition regulates mountain ecosystems and landscape evolution. *Proc. Natl. Acad. Sci.* 111 (9), 3338–3343.
- Harrington, E.R., 1957. Sinkholes, bottomless lakes, and the Pecos River. *The Scientific Monthly* 84 (6), 302–308.
- van Hengstum, P.J., Scott, D.B., Gröcke, D.R., Charette, M.A., 2011. Sea level controls sedimentation and environments in coastal caves and sinkholes. *Mar. Geol.* 286 (1–4), 35–50.
- Kéfi, S., Eppinga, M.B., de Ruiter, P.C., Rietkerk, M., 2010. Bistability and regular spatial patterns in arid ecosystems. *Theor. Ecol.* 3 (4), 257–269.
- Kobal, M., Bertonecelj, I., Pirotti, F., Daksobler, I., Kutnar, L., 2015. Using Lidar data to analyse sinkhole characteristics relevant for understory vegetation under forest cover—case study of a high karst area in the Dinaric Mountains. *PloS one* 10 (3), e0122070.
- Kruse, S., Grasmueck, M., Weiss, M., Viggiano, D., 2006. Sinkhole structure imaging in covered karst terrain. *Geophysical Research Letters* 33 (16).
- Landeweert, R., Hoffland, E., Finlay, R.D., Kuyper, T.W., van Breemen, N., 2001. Linking plants to rocks: ectomycorrhizal fungi mobilize nutrients from minerals. *Trends Ecol. Evol.* 16 (5), 248–254.
- Lane, P., Donnelly, J.P., Woodruff, J.D., Hawkes, A.D., 2011. A decadal-resolved paleohurricane record archived in the late Holocene sediments of a Florida sinkhole. *Marine Geology* 287 (1–4), 14–30.
- LeGrand, H., 1973. Hydrological and ecological problems of karst regions: hydrological actions on limestone regions cause distinctive ecological problems. *Science* 179 (4076), 859–864.
- Metz, P.A., 2011. Factors that Influence the Hydrologic Recovery of Wetlands in the Northern Tampa Bay Area, Florida. US Department of the Interior, US Geological Survey.
- Miller, K.G., Kominz, M.A., Browning, J.V., Wright, J.D., Mountain, G.S., Katz, M.E., Sugarman, P.J., Cramer, B.S., Christie-Blick, N., Pekar, S.F., 2005. The Phanerozoic record of global sea-level change. *Science* 310 (5752), 1293–1298.
- Neal, J.T., Johnson, K.S., 2002. McCauley Sinks: a compound breccia pipe in evaporite karst, Holbrook basin, Arizona, USA. *Carbonates and Evaporites* 17 (2), 98.
- Petuch, E.J., Roberts, C., 2007. *The Geology of the Everglades and Adjacent Areas*. CRC Press.
- Quintero, C.J., Cohen, M.J., 2019. Scale-dependent patterning of wetland depressions in a low-relief karst landscape. *J. Geophys. Res.* Earth 124 (8), 2101–2117.
- Reznichenko, N., 2012. *A Field Guide to the Geology of the Castle Hill Basin*.
- Scanlon, T.M., Caylor, K.K., Levin, S.A., Rodríguez-Iturbe, I., 2007. Positive feedbacks promote power-law clustering of Kalahari vegetation. *Nature* 449 (7159), 209–212.
- Schneider, D.L., 1973. *The Limnology of a Meromictic Sinkhole Lake in South Florida*.
- Šegina, E., Benac, Č., Rubinić, J., Knez, M., 2018. Morphometric analyses of dolines—the problem of delineation and calculation of basic parameters. *Acta Carsologica* 47 (1).
- Shinn, E.A., Reich, C.D., Locker, S.D., Hine, A.C., 1996. A giant sediment trap in the Florida Keys. *Journal of Coastal Research* 953–959.
- Sinclair, W.C., 1982. Sinkhole Development Resulting from Ground-water Withdrawal in the Tampa Area, Florida. US Geological Survey, Water Resources Division.
- Theilen-Willige, B., 2018. Detection of karst features in the black hills area in South Dakota/Wyoming, USA, based on evaluations of remote sensing data. *Geosciences* 8 (6), 192.
- Theilen-Willige, B., Malek, H., Charif, A., El Bchari, F., Chaibi, M., 2014. Remote sensing and GIS contribution to the investigation of karst landscapes in NW-Morocco. *Geosciences* 4 (2), 50–72.
- Tihansky, A.B., 1999. Sinkholes, west-central Florida: land subsidence in the United States. *US Geological Survey Circular* 1182, 121–140.
- Vernon, R.O., 1947. Cypress domes. *Science* 105 (2717), 97–99.
- Wanaim, A., Belfoul, M.A., Bouzid, S., Faik, F., 2014. Speleogenesis and evidences tectonic into Lakhssas karst system (Western Anti-Atlas, Morocco). *Open Journal of Geology* 4 (7), 314.
- Watts, A.C., Watts, D.L., Cohen, M.J., Heffernan, J.B., McLaughlin, D.L., Martin, J.B., Kaplan, D.A., Osborne, T.Z., Kobziar, L.N., 2014. Evidence of biogeomorphic patterning in a low-relief karst landscape. *Earth Surf. Process. Landf.* 39 (15), 2027–2037.
- Yizhaq, H., Ish-Shalom, C., Raz, E., Ashkenazy, Y., 2017. Scale-free distribution of Dead Sea sinkholes: observations and modeling. *Geophys. Res. Lett.* 44 (10), 4944–4952.
- Žebre, M., Stepišnik, U., 2015. Glaciokarst landforms and processes of the southern Dinaric Alps. *Earth Surf. Process. Landf.* 40 (11), 1493–1505.
- Zhang, X., Bianchi, T.S., Cohen, M.J., Martin, J.B., Quintero, C.J., Brown, A.L., Ares, A.M., Heffernan, J.B., Ward, N., Osborne, T.Z., 2019. Initiation and development of wetlands in southern Florida karst landscape associated with accumulation of organic matter and vegetation evolution. *J. Geophys. Res. Biogeo.* 124 (6), 1604–1617.