Achieving High Selectivity and Facile Displacement with a New Thiol Auxiliary for Boron Mediated Aldol Reactions

Sandra Fanjul, Alison N. Hulme,* John W. White

School of Chemistry, The University of Edinburgh, West Mains Road, Edinburgh, EH9 3JJ, UK.

Supporting Information (Spectra)

Preparation of thiol auxiliary 1	S7-S16
Acylation and <i>anti</i> aldol reaction	S17-S25
Displacement reactions	S26-S35

(1*S*,2*R*)-2-(*N*-Benzyl-*N*-mesitylenesulfonylamino)-1-phenylpropane-1-thiol 1

2'-(*N*-Benzyl-*N*-mesitylenesulfonylamino)-1'-phenylpropyl (1'*S*,2*S*,2'*R*,3*S*)-2,5-dimethyl-3-hydroxy-thiopentanoate **7c**

2'-(*N*-Benzyl-*N*-mesitylenesulfonylamino)-1'-phenylpropyl (1'*S*,2*R*,2'*R*,3*R*)-2,5-dimethyl-3-hydroxy-thiopentanoate
minor diastereomer

2'-(*N*-Benzyl-*N*-mesitylenesulfonylamino)-1'-phenylpropyl (1'*S*,2*S*,2'*R*,3*R*)-2,4-dimethyl-3-hydroxythiolpent-4-eneoate **7b**

2'-(*N*-Benzyl-*N*-mesitylenesulfonylamino)-1'-phenylpropyl (1'*S*,2*S*,2'*R*,3*R*)-3-hydroxy-2-methyl-3-phenylthiolpropionate **7e**