Acid-catalyzed rearrangement of diaryl hydrazides for the efficient synthesis of functionalized 2,2'-diamino-1,1'-biaryls.

Young-Kwan Lim, Jin-Woo Jung and Cheon-Gyu Cho*

Department of Chemistry, Hanyang University, Seoul 133-791, Korea.

Center for Bioactive Molecular Hybrids, Yonsei University, Seoul 120-747, Korea.

ccho@hanyang.ac.kr

Supporting Information

General Methods. ¹H NMR spectra were recorded at 400MHz and ¹³C NMR spectra at 100MHz, with either TMS (δ = 0) or the signal for residual CHCl₃ in the CDCl₃ solvent (7.24) as internal standards. *J* values are reported in Hz. High resolution mass spectra were measured by using FAB method. Flash column chromatography was performed with Kieselgel 60 Art 9385(230-400 mesh). All solvents used were purified according to standard procedures.

Table of Contents

Copy of ¹ H/ ¹³ C NMR/IR spectra of 1a	S2 – S4
Copy of ¹ H/ ¹³ C NMR/IR spectra of 3a – 3k	S5 – S37
Copy of ¹ H/ ¹³ C NMR/IR spectra of 4a – 4k	S38 – S67
Copy of ¹ H/ ¹³ C NMR/IR spectra of 6a, 6b, 9a-9c	S68 – S82
Copy of ¹ H/ ¹³ C NMR/IR spectra of 7b, 10b, 10c	S83 – S91

Mode =

2004-03-17

Res = 4 20 scans/minute

Mode =

Res = 4 20 scans/minute

2004-02-1

Mode =

Res = 4 20 scans/minute

2004-02-1

Mode =

04-03-0

Res = 4 20 scans/minute

Mode =

2004-02-1

Res = 4 20 scans/minute

Mode =

04-03-0

Res = 4 19 scans/minute

Mode =

04-03-0

Res = 4 20 scans/minute

Mode =

2004-02-23

Res = 4 19 scans/minute

Mode =

04-02-23

Res = 4 20 scans/minute

Mode =

2004-02-23

Res = 4 19 scans/minute

Mode =

2004-02-23

Res = 4 20 scans/minute

Mode =

2004-02-23

Res = 4 19 scans/minute

Mode =

2004-02-23

Res = 4 19 scans/minute

Mode =

04-03-0

Res = 4 19 scans/minute

Mode =

04-03-0

Res = 4 20 scans/minute

Mode =

04-03-0

Res = 4 19 scans/minute

Mode =

04-03-0

Res = 4 19 scans/minute

Mode =

04-03-0

Res = 4 19 scans/minute

Mode =

04-03-0

Res = 4 19 scans/minute

Mode =

04-03-0

Res = 4 19 scans/minute