Ag-CATALYZED ASYMMETRIC MANNICH REACTIONS OF ENOL ETHERS AND ARYL, ALKYL, ALKENYL AND ALKYNYL IMINES

Nathan S. Josephsohn, Marc L. Snapper* and Amir H. Hoveyda*

Department of Chemistry, Merkert Chemistry Center, Boston College
Chestnut Hill, Massachusetts 02467

SUPPORTING INFORMATION

General. Infrared (IR) spectra were recorded on a Perkin Elmer 781 spectrophotometer, $\nu_{\max}$ in cm^{-1} . Bands are characterized as broad (br), strong (s), medium (m), or weak (w). ^1H NMR spectra were recorded on a Varian Unity INOVA 400 (400 MHz) spectrometer. Chemical shifts are reported in ppm from tetramethylsilane with the solvent resonance resulting from incomplete deuteration as the internal standard (CDCl_3 : 7.26). Data are reported as follows: chemical shift, integration, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, br = broad, m = multiplet), and coupling constants in Hz. ^{13}C NMR spectra were recorded on a Varian Unity INOVA 400 (100 MHz) with complete proton decoupling. Chemical shifts are reported in ppm from tetramethylsilane with the solvent resonance as the internal standard (CDCl_3 : 77.16). Enantiomeric ratios were determined by high-performance liquid chromatography (HPLC) by a Shimadzu chromatograph and chiralpak AD, OD and OJ columns (4.6 mm x 250 mm chiral column by Chiral Technologies) in comparison with authentic racemic materials. Elemental analyses were performed by Robertson Microlit Laboratories (Madison, New Jersey). All high-resolution mass spectrometry was performed at Boston College on a Micromass LCT ESI-MS (positive mode). Optical rotation values were recorded on a Rudolph Research Analytical Autopol IV polarimeter.

Unless otherwise stated, all reactions conducted under open atmosphere in 16 x 150 mm borosilicate test tubes. All liquid reagents were handled with a Fisherbrand Finnpiptette repeating pipetter. J.T. Baker reagent grade THF was used without purification. Fisherbrand HPLC grade 2-Propanol was used without purification. Unless otherwise stated silver acetate (99%), iodobenzene diacetate, DIBAL-H, LAH, NaCNBH_3 , formaldehyde (30% v/v in H_2O), Boc_2O , *p*-anisidine, *o*-anisidine 1-phenyl-1-(trimethylsilyloxy)ethylene (**3a**) and all commercially available aldehydes (for imines) were purchased from Aldrich and used without further purification.

o-Anisidyl imines,¹ 5-oxo-pentanoic acid methyl ester,² trimethyl[(1-methylethenyl)oxy]-silane³ (**3b**) and ligand **1**⁴ were prepared according to literature procedure.

¹ a) Saito, S; Hatanaka, K.; Yamamoto, H. *Org. Lett.* **2000**, 2, 1891-1894. (b) Adrian, Jr. J. C.; Barkin, J. L.; Hassib, L. *Tetrahedron Lett.* **1999**, 40, 2457-2460.

² Mioskowski, P. C. C.; Falk, J. R. *Tetrahedron Lett.* **1989**, 30, 2545-2548.

³ Walshe, N. D. A.; Goodwin, G. B. T.; Smith, G. C.; Woodward, F. E. *Org. Synth.* **1986**, 65, 1-5.

⁴ Josephsohn, N. J.; Snapper, M. L.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2003**, 125, 1418-1419.

Representative procedure for the Ag-catalyzed asymmetric Mannich reaction:

Ligand **1** (18 mg, 36 μ mol) and AgOAc (6.0 mg, 36 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 1 mL of THF and allowed to stir for 5 min at 22 °C. Imine **2d** (310 mg, 1.20 mmol) was added immediately followed by 100 μ L *i*-PrOH (1.3 mmol). The test tube was capped with a septum, and allowed to stir in a cold room at 4 °C for 5 min. Enol ether **3a** (0.5 mL, 2.4 mmol) was added and the reaction mixture was allowed to stir at 4 °C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:Et₂O in most cases 4:1 hexanes: EtOAc when enol ether **3b** is used), to deliver 396 mg (1.05 mmol, 88% yield) of the desired product **4d** as a yellow-orange solid.

(-)-3-(2-Methoxyphenylamino)-1,3-diphenylpropan-1-one (4aa);

A modified procedure was used. Ligand **1** (6.0 mg, 12 μ mol) and AgOAc (2.0 mg, 12 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.2 mL of THF and allowed to stir for 5 min at 22 °C. Imine **2a** (85 mg, 0.4 mmol) was added immediately followed by 33 μ L *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir in a cold bath at -10 °C for 5 min. Enol ether **3a** (160 μ L, 0.8 mmol) was added and the reaction was allowed to stir at -10 °C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:Et₂O) to deliver 72 mg (0.22 mmol, 54%) of the desired product **4aa** as a yellow solid. IR (neat, NaCl): 3410 (w), 3063 (w), 1686 (s), 1599 (s), 1512 (s), 1456 (s), 1357 (m), 1258 (m), 1227 (s), 1128 (m), 1028 (m), 732 (s), 700 (s). ¹H NMR (CDCl₃, 400 MHz): δ 7.92-7.90 (2H, d, *J* = 7.32 Hz), 7.56-7.20 (8H, m), 6.76-6.60 (3H, m), 6.45 (1H, dd, *J* = 7.7, 1.5 Hz), 5.06 (1H, apparent triplet, *J* = 6.4 Hz), 5.00 (1H, br), 3.84 (3H, s), 3.50 (2H, apparent doublet, *J* = 5.9 Hz). ¹³C NMR (CDCl₃, 100 MHz): δ 198.0, 147.2, 146.4, 143.3, 137.0, 133.4, 128.9, 128.8, 128.3, 127.4, 126.6, 121.2, 117.0, 111.5, 109.6, 55.7, 54.5, 46.8. HRMS (ES) Calcd for C₂₂H₂₁NO₂Na: 354.1465; Found: 354.1470. $[\alpha]_D^{20} = -36.2^\circ$ (*c* = 0.347, CHCl₃) for a 94% *ee* sample.

The optical purity of **4aa** was established by chiral HPLC analysis (OD column, 9/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-4-(2-Methoxyphenylamino)-4-phenylbutan-2-one (4ab); A modified procedure was used. Ligand **1** (6.0 mg, 12 μ mol) and AgOAc (2.0 mg, 12 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.2 mL of THF and allowed to stir for 5 min at 22 $^{\circ}$ C. Imine **2a** (85 mg, 0.4 mmol) was added immediately followed by 33 μ L *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir in a cold bath at -10 $^{\circ}$ C for 5 min. Enol ether **3b** (130 μ L, 0.80 mmol) was added and the reaction was allowed to stir at -10 $^{\circ}$ C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:EtOAc) to deliver 70 mg (0.26 mmol, 65%) of the desired product **4ab** as a clear colorless oil. IR (neat, NaCl): 3413 (w), 2926 (w), 1715 (s), 1602 (m), 1515 (s), 1453 (m), 1359 (m), 1228 (s), 1028 (m), 741 (m), 697 (m). ¹H NMR (CDCl₃, 400 MHz): δ 7.36-7.19 (5H, m), 6.75-6.60 (3H, m), 6.42 (1H, dd, *J* = 7.7, 1.3 Hz), 4.92-4.85 (2H, m), 3.85 (3H, s), 2.95 (2H, ddd, *J* = 16.1, 7.5, 5.9 Hz), 2.10 (3H, s). ¹³C NMR (CDCl₃, 100 MHz): δ 207.1, 147.3, 143.0, 136.9, 128.1, 127.7, 126.7, 121.5, 117.4, 111.7, 109.8, 55.8, 54.5, 52.1, 31.0. HRMS (ES) Calcd for C₁₇H₁₉NO₂Na: 292.1320; Found: 292.1313. $[\alpha]_D^{20}$ = -38.2 $^{\circ}$ (*c* = 0.427, CHCl₃) for a 92% *ee* sample.

The optical purity of **4ab** was established by chiral HPLC analysis (OD column, 9/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-3-(4-Methoxyphenyl)-3-(2-methoxyphenylamino)-1-phenylpropan-1-one (4ba);

A modified procedure was used. Ligand **1** (18 mg, 36 μ mol) and AgOAc (6.0 mg, 36 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 1 mL of toluene and allowed to stir for 5 min at 22 °C. Imine **2b** (320 mg, 1.20 mmol) was added immediately followed by 100 μ L *i*-PrOH (1.30 mmol). The test tube was capped with a septum, and allowed to stir in a cold bath at 23 °C for 5 min. Enol ether **3a** (0.5 mL, 2.4 mmol) was added and the reaction was allowed to stir at 23 °C

for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:Et₂O) to deliver 88 mg (0.24mmol, 61%) of the desired product **4ba** as a white solid. IR (neat, NaCl): 1681 (m), 1600 (m), 1513 (s), 1457 (m), 1246 (m), 1227 (m), 1028 (m), 736 (m). ¹H NMR (CDCl₃, 400 MHz): δ 7.92 (2H, dd, *J* = 8.3, 0.7 Hz), 7.58-7.36 (5H, m), 6.87-6.48 (6H, m), 5.04 (1H, apparent doublet, *J* = 6.4 Hz), 4.97 (1H, br), 3.86 (3H, s), 3.77 (3H, s), 3.51 (2H, apparent doublet, *J* = 6.4 Hz). ¹³C NMR (CDCl₃, 100 MHz): δ 198.2, 158.9, 147.2, 137.0, 136.9, 135.2, 133.6, 128.7, 128.3, 127.7, 121.2, 117.0, 114.3, 111.5, 109.6, 55.6, 55.3, 53.9, 46.8. HRMS (ES) Calcd. for C₂₃H₂₀NO₃Na: 384.1582; Found: 384.1576. Anal Calcd. for C₁₈H₁₇NO₂: C, 76.43, H, 6.41; Found: C, 77.36, H, 6.48. $[\alpha]_D^{20} = -33.5^\circ$ (*c* = 0.967, CHCl₃) for a 96% *ee* sample.

The optical purity of **4ba** was established by chiral HPLC analysis (OD column, 9/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-4-(4-Methoxyphenyl)-4-(2-methoxyphenylamino)-butan-2-one (4bb);

A modified procedure was used. Ligand **1** (10 mg, 20 μ mol) and AgOAc (3.0 mg, 20 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.4 mL of toluene and allowed to stir for 5 min at 22 °C. Imine **2b** (100 mg, 0.40 mmol) was added

immediately followed by 33 μ L *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir in a cold room at 4 °C for 5 min. Enol ether **3b** (130 μ L, 0.80 mmol) was added and the reaction was allowed to stir at 4 °C for 18 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH_2Cl_2 (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH_2Cl_2) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH_2Cl_2 and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:EtOAc) to deliver 85 mg (0.29 mmol, 71%) of the desired product **4bb** as a light yellow oil. IR (neat, NaCl): 1712 (m), 1606 (m), 1507 (s), 1457 (m), 1240 (s), 1227 (m), 1184 (m), 1184 (m), 1028 (m), 730 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.28 (2H, d, J = 8.6 Hz), 6.85 (2H, dd, J = 8.1, 0.7 Hz), 6.77-6.61 (3H, m), 6.45 (1H, d, J = 7.9 Hz), 4.84, (1H, apparent doublet, J = 6.8 Hz), 4.79 (1H, br), 3.86 (3H, s), 3.77 (3H, s), 2.94 (2H, ddd, J = 15.7, 7.1, 6.2 Hz), 2.10 (3H, s). ^{13}C NMR (CDCl_3 , 100 MHz): δ 207.0, 158.9, 147.1, 136.8, 134.8, 127.5, 121.2, 117.0, 114.2, 111.4, 109.6, 55.6, 55.3, 53.6, 51.8, 30.8. HRMS (ES) Calcd for $\text{C}_{18}\text{H}_{21}\text{NO}_3\text{Na}$: 322.1418; Found: 322.1419. $[\alpha]_{\text{D}}^{20} = -17.3^\circ$ (c = 2.35, CHCl_3) for a 90% *ee* sample.

The optical purity of **4bb** was established by chiral HPLC analysis (OD column, 9/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-3-(2-Methoxyphenylamino)-3-(4-nitrophenyl)-1-phenylpropan-1-one (4ca); A

modified procedure was used. Ligand **1** (18 mg, 36 μ mol) and AgOAc (6.0 mg, 36 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 1 mL of THF and allowed to stir for 5 min at 22 °C. Imine **2c** (310 mg, 1.20 mmol) was added immediately followed by 100 μ L *i*-PrOH (1.30 mmol). The test tube was capped with a septum, and allowed to stir in a cold bath at -5 °C for 5 min. Enol ether **3a** (0.5 mL, 2.4 mmol) was added and the reaction was allowed to stir at -5 °C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH_2Cl_2 (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH_2Cl_2) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH_2Cl_2 and separated as stated above. The

combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:Et₂O) to deliver 396 mg (1.1 mmol, 88%) of the desired product **4ca** as a orange solid. IR (neat, NaCl): 1684 (m), 1596 (m), 1515 (s), 1453 (m), 1340 (s), 1228 (m), 741 (m). ¹H NMR (CDCl₃, 400 MHz): δ 8.16 (2H, d, *J* = 8.6 Hz), 7.92 (2H, dd, *J* = 7.1, 1.8 Hz), 7.65-7.44 (5H, m), 6.80-6.65 (3H, m), 6.35 (1H, dd, *J* = 7.3, 1.8 Hz), 5.18 (1H, apparent doublet, *J* = 6.2 Hz), 5.13 (1H, s), 3.87 (3H, s), 3.57 (2H, ddd, *J* = 16.7, 6.8, 5.7 Hz). ¹³C NMR (CDCl₃, 100 MHz): δ 196.6, 151.1, 147.3, 147.2, 136.6, 136.2, 133.7, 128.9, 128.2, 127.6, 124.1, 121.2, 117.8, 111.4, 109.8, 55.6, 53.9, 46.2. HRMS (ES) Calcd for C₂₂H₂₀N₂O₄Na: 399.1312; Found: 399.1321. [α]_D²⁰ = -55.0° (*c* = 0.58, CHCl₃) for a 92% *ee* sample.

The optical purity of **4ca** was established by chiral HPLC analysis (OD column, 1/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-4-(2-Methoxyphenylamino)-4-(4-nitrophenyl)-butan-2-one (4cb); A modified procedure was used. Ligand **1** (10 mg, 20 μmol) and AgOAc (3.0 mg, 20 μmol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.4 mL of THF and allowed to stir for 5 min at 22 °C.

Imine **2c** (105 mg, 0.40 mmol) was added immediately followed by 33 μL *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir in a cold bath at -10 °C for 5 min. Enol ether **3b** (130 μL, 0.80 mmol) was added and the reaction was allowed to stir at -10 °C for 16 h. The reaction was quenched upon

addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:2 hexanes:EtOAc) to deliver 114 mg (0.36 mmol, 91%) of the desired product **4cb** as a orange solid. IR (neat, NaCl): 1713 (m), 1601 (m), 1514 (s), 1452 (w), 1346 (s), 1228 (m), 1022 (w), 742 (m). ¹H NMR (CDCl₃, 400 MHz): δ 8.15 (2H, dd, *J* = 6.8, 2.0 Hz), 7.55 (2H, dd, *J* = 6.8, 2.0 Hz), 6.79-6.64 (3H, m), 6.29 (1H, dd, *J* = 7.2, 2.4 Hz), 4.99-4.96 (2H, m), 3.88 (3H, s), 3.00 (2H, ddd, *J* = 16.7, 7.0, 5.9 Hz), 2.15 (3H, s). ¹³C NMR (CDCl₃, 100 MHz): δ 205.6, 150.8, 147.3, 147.1, 136.0, 127.5, 124.1, 121.2, 117.9, 111.4, 109.8, 55.6, 53.6, 51.2,

30.7. HRMS (ES) Calcd for $C_{17}H_{18}NO_2NaCl$: 326.0927; Found: 326.0924. $[\alpha]_D^{20} = -38.0^\circ$ ($c = 0.93$, $CHCl_3$) for a 94% *ee* sample.

The optical purity of **4cb** was established by chiral HPLC analysis (OJ column, 9/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-3-(4-Chlorophenyl)-3-(2-methoxyphenylamino)-1-phenylpropan-1-one (4da);

The representative procedure was followed. The crude product was purified by silica gel chromatography (4:2 hexanes:Et₂O) to deliver 370 mg (1.0 mmol, 83%) of the desired product **4da** as an orange solid. IR (neat, NaCl): 1678 (s), 1599 (s), 1515 (s), 1448 (m), 1255 (m), 1225 (s), 1032 (m), 737 (s). ¹H NMR (CDCl₃, 400 MHz): δ 7.90 (2H, d, $J = 8.0$ Hz), 7.58-7.25 (7H, m), 6.77-6.62 (3H, m), 6.40 (1H, d, $J = 7.7$ Hz), 5.05-5.03 (2H, m), 3.85 (3H, s), 3.49 (2H, ddd, $J = 16.7, 7.0, 5.7$ Hz). ¹³C NMR (CDCl₃, 100 MHz): δ 197.6, 147.1, 141.8, 136.8, 136.6, 133.5, 133.0, 129.0, 128.8, 128.3, 128.0, 121.2, 117.3, 111.5, 109.6, 55.6, 53.9, 46.6. HRMS (ES) Calcd for $C_{22}H_{20}NO_2NaCl$: 388.1070; Found: 388.1080. $[\alpha]_D^{20} = -53.7^\circ$ ($c = 0.947$, $CHCl_3$) for a 96% *ee* sample.

The optical purity of **4da** was established by chiral HPLC analysis (OD column, 94/6 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-4-(4-Chlorophenyl)-4-(2-methoxyphenylamino)-butan-2-one (4db); A modified procedure was used. Ligand **1** (10 mg, 20 μ mol) and AgOAc (3.0 mg, 20 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.4 mL of THF and allowed to stir for 5 min at 22 $^\circ$ C. Imine **2d** (100 mg, 0.40 mmol)

was added immediately followed by 33 μ L *i*-PrOH (0.44 mmol). The test tube was capped with a septum, and allowed to stir in a cold bath at -10 °C for 5 min. Enol ether **3b** (130 μ L, 0.80 mmol) was added and the reaction was allowed to stir at -10 °C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH_2Cl_2 (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH_2Cl_2) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH_2Cl_2 and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:2 hexanes:EtOAc) to deliver 96 mg (0.32 mmol, 79%) of the desired product **4db** as a clear colorless oil. IR (neat, NaCl): 1712 (s), 1607 (m), 1509 (s), 1607 (m), 1226 (s), 1029 (m), 740 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.30-7.24 (4H, m), 6.76-6.61 (3H, m), 6.35 (1H, dd, J = 7.7, 1.6 Hz), 4.85-4.82 (2H, m), 3.85 (3H, s), 2.92 (2H, ddd, J = 16.1, 7.1, 5.5 Hz), 2.10 (3H, s). ^{13}C NMR (CDCl_3 , 100 MHz): δ 206.4, 147.1, 141.5, 136.5, 133.0, 129.0, 127.9, 121.2, 117.4, 111.4, 109.7, 55.6, 53.6, 51.7, 30.8. HRMS (ES) Calcd for $\text{C}_{17}\text{H}_{18}\text{NO}_2\text{NaCl}$: 326.0923; Found: 326.0925. $[\alpha]_{\text{D}}^{20}$ = -36.6° (c = 0.970, CHCl_3) for a 90% *ee* sample.

The optical purity of **4db** was established by chiral HPLC analysis (OD column, 9/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-3-(2-Methoxyphenylamino)-3-(3-nitrophenyl)-1-phenylpropan-1-one (4ea); A

modified procedure was used. Ligand **1** (6.0 mg, 12 μ mol) and AgOAc (2.0 mg, 12 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.2 mL of THF and allowed to stir for 5 min at 22 °C. Imine **2e** (115 mg, 0.40 mmol) was added immediately followed by 33 μ L *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir in a cold bath at -10 °C for 5 min. Enol ether **3a** (160 μ L, 0.80 mmol) was added and the reaction was allowed to stir at -10 °C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH_2Cl_2 (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH_2Cl_2) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH_2Cl_2 and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel

chromatography (4:2 hexanes:Et₂O) to deliver 162 mg (0.15 mmol, 97%) of the desired product **4ea** as a orange solid. IR (neat, NaCl): 1683 (s), 1597 (m), 1529 (s), 1449 (m), 1344 (s), 1226 (m), 738 (s). ¹H NMR (CDCl₃, 400 MHz): δ 8.33 (1H, s), 8.07 (1H, dd, *J* = 8.1, 1.5 Hz), 7.92 (2H, dd, *J* = 8.1, 1.3 Hz), 7.82 (1H, d, *J* = 7.7 Hz), 7.60-7.44 (4H, m), 6.79-6.65 (3H, m), 6.39 (1H, dd, *J* = 7.5, 1.8 Hz), 5.17 (1H, apparent doublet, *J* = 6.2 Hz), 5.12 (1H, br), 3.88 (3H, s), 3.57 (2H, ddd, *J* = 16.7, 6.8, 5.9 Hz). ¹³C NMR (CDCl₃, 100 MHz): δ 197.0, 148.8, 147.2, 145.8, 136.6, 136.2, 133.7, 133.3, 129.8, 128.9, 128.3, 122.6, 121.6, 121.2, 117.8, 111.5, 109.8, 55.7, 53.9, 46.4. HRMS (ES) Calcd for C₂₂H₂₀N₂O₄Na: 399.1316; Found: 399.1321. Anal Calcd for C₂₂H₂₀N₂O₄: C, 70.20, H, 5.36; Found: C, 70.15, H, 5.20. [α]_D²⁰ = -61.8° (*c* = 0.630, CHCl₃) for a 86% *ee* sample.

The optical purity of **4ea** was established by chiral HPLC analysis (OD column, 75/25 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-4-(2-Methoxyphenylamino)-4-(3-nitrophenyl)-butan-2-one (4eb); A modified procedure was used. Ligand **1** (6.0 mg, 12 μmol) and AgOAc (2.0 mg, 12 μmol) were

weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.4 mL of THF and allowed to stir for 5 min at 22 °C. Imine **2e** (115 mg, 0.4 mmol) was added immediately followed by 33 μL *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir in a cold bath at -5 °C for 5 min. Enol ether **3b** (130 μL, 0.80 mmol) was added and the reaction mixture was allowed to stir at -5 °C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:2 hexanes:EtOAc) to deliver 121 mg (0.380 mmol, 96%) of the desired product **4eb** as a yellow oil. IR (neat, NaCl): 1712 (m), 1600 (s), 1513 (s), 1463 (m), 1345 (s), 1221 (m), 742 (s). ¹H NMR (CDCl₃, 400 MHz): δ 8.25 (1H, s), 8.10-8.07 (1H, m), 7.74 (1H, d, *J* = 7.7 Hz), 7.48 (1H, t, *J* = 7.9 Hz), 6.79-6.64 (3H, m), 6.35 (1H, dd, *J* = 7.3, 1.8 Hz), 5.10-4.97 (2H, m), 3.89 (3H, s), 3.02 (2H, ddd, *J* = 16.7, 7.1, 5.7 Hz), 2.16 (3H, s). ¹³C NMR (CDCl₃, 100 MHz): δ 205.7, 148.8, 147.2, 145.5, 136.0, 133.1, 129.8, 122.6, 121.5, 121.2, 117.9, 111.5, 109.8, 55.7, 53.6,

51.4, 30.7. HRMS (ES) Calcd for $C_{17}H_{18}N_2O_4Na$: 337.1162; Found: 337.1164. $[\alpha]_D^{20} = -42.3^\circ$ ($c = 0.68$, $CHCl_3$) for a 86% *ee* sample.

The optical purity of **4eb** was established by chiral HPLC analysis (OD column, 75/25 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-3-(2-Methoxyphenylamino)-3-naphthalen-2-yl-1-phenylpropan-1-one (4fa); A

modified procedure was used. Ligand **1** (6.0 mg, 12 μ mol) and AgOAc (2.0 mg, 12 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.4 mL of THF and allowed to stir for 5 min at 22 $^\circ$ C. Imine **2f** (105 mg, 0.40 mmol) was added immediately followed by 33 μ L *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir in a cold bath at -5 $^\circ$ C for 5 min. Enol ether **3a** (160 μ L, 0.8 mmol) was added and the reaction mixture was allowed to stir at -5 $^\circ$ C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH_2Cl_2 (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH_2Cl_2) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH_2Cl_2 and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:EtOAc) to deliver 135 mg (0.35 mmol, 88%) of the desired product **4fa** as a white solid. IR (neat, NaCl): 3413 (w), 3057 (w), 1684 (s), 1602 (s), 1517 (s), 1453 (m), 1222 (s), 1028 (m), 747 (s). 1H NMR ($CDCl_3$, 400 MHz): δ 8.10-7.43 (12, m), 6.80-6.62 (3H, m), 6.53 (1H, dd, $J = 7.7, 1.5$ Hz), 5.25 (1H, apparent triplet, $J = 6.4$ Hz), 5.11 (1H, br), 3.89 (3H, s), 3.61 (2H, apparent doublet, $J = 6.4$ Hz). ^{13}C NMR ($CDCl_3$, 100 MHz): δ 197.9, 147.1, 140.8, 137.0, 136.9, 133.6, 133.4, 133.0, 128.8, 128.6, 128.3, 128.0, 127.8, 126.1, 125.8, 125.3, 124.8, 121.2, 117.1, 111.6, 109.6, 55.6, 54.7, 46.8. HRMS (ES) Calcd for $C_{26}H_{23}NO_2Na$: 404.1632; Found: 404.1626. $[\alpha]_D^{20} = -53.57^\circ$ ($c = 1.37$, $CHCl_3$) for a 96% *ee* sample.

The optical purity of **4fa** was established by chiral HPLC analysis (OD column, 1/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-4-(2-Methoxyphenylamino)-4-naphthalen-2-yl-butan-2-one (**4fb**); A modified procedure was used. Ligand **1** (6.0 mg, 12 μ mol) and AgOAc (2.0 mg, 12 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.4 mL of THF and allowed to stir for 5 min at 22 $^{\circ}$ C. Imine **2f** (105 mg, 0.40 mmol) was added immediately followed by 33 μ L *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir in a cold bath at -5 $^{\circ}$ C for 5 min. Enol ether **3b** (130 μ L, 0.80 mmol) was added and the reaction mixture was allowed to stir at -5 $^{\circ}$ C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:2 hexanes:EtOAc) to deliver 120 mg (0.38 mmol, 94%) of the desired product **4fb** as a white solid. IR (neat, NaCl): 3412 (w), 3049 (w), 1714 (s), 1599 (s), 1515 (s), 1460 (m), 1218 (m), 1128 (m), 1031 (m), 735 (s). ¹H NMR (CDCl₃, 400 MHz): δ 7.85-7.81 (4H, m), 7.53-7.44 (3H, m), 6.79 (1H, dd, *J* = 7.2, 1.1 Hz), 6.73-6.62 (2H, m), 6.51 (1H, dd, *J* = 7.2, 1.4 Hz), 5.06 (1H, dd, *J* = 6.8, 6.4 Hz), 5.01 (1H, s), 3.91 (3H, s), 3.05 (2H, ddd, *J* = 14.0, 7.1, 6.0 Hz), 2.13 (3H, s). ¹³C NMR (CDCl₃, 100 MHz): δ 206.7, 147.1, 140.4, 136.8, 133.6, 132.9, 128.8, 128.0, 127.8, 126.2, 125.9, 125.2, 124.5, 121.2, 117.2, 111.5, 109.6, 55.6, 54.4, 51.8, 30.8. HRMS (ES) Calcd for C₂₁H₂₁NO₂Na: 342.1464; Found: 342.1470. $[\alpha]_D^{20}$ = -55.5 $^{\circ}$ (*c* = 0.40, CHCl₃) for a 92% *ee* sample.

The optical purity of **4fb** was established by chiral HPLC analysis (OD column, 9/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-3-(2-Bromophenyl)-3-(2-methoxyphenylamino)-1-phenylpropan-1-one (4ga); A

modified procedure was used. Ligand **1** (10 mg, 20 μ mol) and AgOAc (3.0 mg, 20 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.4 mL of toluene and allowed to stir for 5 min at 23 $^{\circ}$ C. Imine **2g** (115 mg, 0.40 mmol) was added immediately followed by 33 μ L *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir 23 $^{\circ}$ C for 5 min. Enol ether **3a** (160 μ L, 0.80 mmol) was added and the reaction mixture was allowed to stir at 23 $^{\circ}$ C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (8:2 hexanes:Et₂O) to deliver 64 mg (0.63 mmol, 46%) of the desired product **4ga** as a white solid. IR (neat, NaCl): 3410 (w), 1686 (s), 1599 (s), 1512 (s), 1456 (m), 1258 (m), 1221 (s), 1022 (s), 737 (s). ¹H NMR (CDCl₃, 400 MHz): δ 7.99 (2H, d, *J* = 7.1 Hz), 7.58-7.43 (5H, m), 7.21 (1H, dt, *J* = 7.5, 1.3 Hz), 7.09 (1H, dt, *J* = 7.7, 1.6 Hz), 6.76-6.59 (3H, m), 6.22 (1H, dd, *J* = 7.7, 1.4 Hz), 5.38 (1H, br), 5.32 (1H, dd, *J* = 8.9, 3.5 Hz), 3.88 (3H, s), 3.58 (1H, dd, *J* = 15.7, 3.7 Hz), 3.37 (1H, dd, *J* = 15.7, 8.8 Hz). ¹³C NMR (CDCl₃, 100 MHz): δ 198.1, 147.2, 141.4, 136.8, 135.4, 133.5, 133.2, 129.0, 128.8, 128.5, 128.4, 128.2, 122.9, 121.2, 117.2, 111.5, 109.6, 55.7, 54.0, 44.4. HRMS (ES) Calcd for C₂₂H₂₀NO₂NaBr: 432.0583; Found: 432.0575. [α]_D²⁰ = -89.1 $^{\circ}$ (*c* = 1.85, CHCl₃) for a 76% *ee* sample.

The optical purity of **4ga** was established by chiral HPLC analysis (OD column, 9/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-4-(2-Bromophenyl)-4-(2-methoxyphenylamino)-butan-2-one (4gb); A modified

procedure was used. Ligand **1** (10 mg, 20 μ mol) and AgOAc (3.0 mg, 20 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.4 mL of THF and allowed to stir for 5 min at 22 $^{\circ}$ C. Imine **2g** (115 mg, 0.40 mmol) was added immediately followed by 33 μ L *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir in a cold room at 4 $^{\circ}$ C for 5 min. Enol ether **3b** (130 μ L, 0.80 mmol) was added and the reaction mixture was allowed to stir at 4

$^{\circ}$ C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH_2Cl_2 (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH_2Cl_2) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH_2Cl_2 and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:2 hexanes:EtOAc) to deliver 97 mg (0.28 mmol, 70%) of the desired product **4gb** as a clear colorless oil. IR (neat, NaCl): 3414 (m), 2928 (w), 1713 (s), 1601 (m), 1514 (s), 1452 (s), 1352 (m), 1228 (s), 1022 (m), 735 (s). ^1H NMR (CDCl_3 , 400 MHz): δ 7.56 (1H, dd, J = 6.9, 0.4 Hz), 7.41 (1H, dd, J = 7.8, 1.6 Hz), 7.22 (1H, t, J = 7.5 Hz), 7.09 (1H, t, J = 7.5 Hz), 6.77-6.61 (3H, m), 6.22, (1H, dd, J = 7.6, 1.5 Hz), 5.19 (1H, dd, J = 9.5, 3.6 Hz), 5.10-5.00 (1H, br), 3.89 (3H, s), 3.03 (1H, dd, J = 15.6, 3.6 Hz), 2.76 (1H, dd, J = 15.6, 9.5 Hz), 2.21 (3H, s). ^{13}C NMR (CDCl_3 , 100 MHz): δ 206.9, 147.2, 140.9, 136.1, 133.2, 129.0, 128.2, 127.9, 122.9, 121.3, 117.5, 111.5, 109.6, 55.7, 53.8, 49.8, 30.0. HRMS (ES) Calcd for $\text{C}_{17}\text{H}_{18}\text{NO}_2\text{NaBr}$: 370.0410; Found: 370.0419. Anal Calcd for $\text{C}_{17}\text{H}_{18}\text{BrNO}_2$: C, 58.63, H, 5.21; Found: C, 59.03, H, 5.00. $[\alpha]_D^{20}$ = -89.5° (c = 0.62, CHCl_3) for a 80% *ee* sample.

The optical purity of **4gb** was established by chiral HPLC analysis (OD column, 9/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-3-Furan-2-yl-3-(2-methoxyphenylamino)-1-phenylpropan-1-one (4ha); A modified procedure was used. Ligand **1** (6.0 mg, 12 μ mol) and AgOAc (2.0 mg, 12 μ mol) were weighed into a 16 x 150 mm test tube. The contents

were dissolved in 0.4 mL of THF and allowed to stir for 5 min at 22 °C. Imine **2h** (80 mg, 0.4 mmol) was added immediately followed by 33 μ L *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir in a cold room at 4 °C for 5 min.

Enol ether **3a** (160 μ L, 0.80 mmol) was added and the reaction mixture was allowed to stir at 4 °C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:Et₂O) to deliver 108 mg (0.33 mmol, 84%) of the desired product **4ha** as a clear colorless oil. IR (neat, NaCl): 3402 (w), 1690 (s), 1608 (s), 1513 (s), 1457 (m), 1249 (m), 1224 (m), 1029 (m), 739 (s). ¹H NMR (CDCl₃, 400 MHz): δ 7.95 (2H, d, *J* = 7.5 Hz), 7.58-7.32 (4H, m), 6.87-6.68 (4H, m), 6.28-6.23 (2H, m), 5.30 (1H, apparent triplet, *J* = 6.4 Hz), 4.86 (1H, br), 3.84 (3H, s), 3.61 (2H, ddd, *J* = 16.7, 6.6, 6.0 Hz). ¹³C NMR (CDCl₃, 100 MHz): δ 197.8, 155.1, 147.3, 141.7, 136.9, 136.5, 133.3, 128.7, 128.2, 121.3, 117.5, 111.3, 110.4, 109.8, 106.5, 55.6, 48.3, 42.9. HRMS (ES) Calcd for C₂₀H₁₉NO₃Na: 344.1259; Found: 344.1263. $[\alpha]_D^{20}$ = +18.1° (*c* = 3.77, CHCl₃) for a 84% *ee* sample.

The optical purity of **4h** was established by chiral HPLC analysis (OD column, 9/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-4-Furan-2-yl-4-(2-methoxyphenylamino)-butan-2-one (4hb); A modified procedure was used. Ligand **1** (6.0 mg, 12 μ mol) and AgOAc (2.0 mg, 12 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.4 mL of THF and allowed to stir for 5 min at 22 °C. Imine **2h** (80 mg, 0.4 mmol) was added immediately followed by 33 μ L *i*-PrOH (0.45mmol). The test tube was capped with a septum, and allowed to stir in a cold room at 4 °C for 5 min. Enol ether **3b** (130 μ L, 0.80 mmol) was added and the reaction mixture was allowed to stir at 4 °C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:2 hexanes:Et₂O) to deliver 81 mg (0.31 mmol, 78%) of the desired product **4hb** as a clear colorless oil. IR (neat, NaCl): 3402 (m), 2941 (w), 1713 (s), 1601 (s), 1514 (s), 1458 (s), 1365 (m), 1252 (s), 1228 (s), 1128 (s), 1034 (s), 742 (s). ¹H NMR (CDCl₃, 400 MHz): δ 7.32 (1H, s), 6.85-6.68 (4H, m), 6.27 (1H, dd, *J* = 3.2, 2.0 Hz), 6.19 (1H, d, *J* = 3.2 Hz), 5.06 (1H, apparent triplet, *J* = 6.4), 4.74 (1H, s), 3.84 (3H, s), 3.04 (2H, d, *J* = 6.4 Hz), 2.15 (3H, s). ¹³C NMR (CDCl₃, 100 MHz): δ 206.5, 154.9, 147.3, 141.8, 136.3, 121.3, 117.6, 111.3, 110.4, 109.9, 106.4, 55.6, 48.0, 47.8, 30.7. HRMS (ES) Calcd for C₁₅H₁₇NO₃Na: 282.1105; Found: 282.1106. [α]_D²⁰ = +44.5° (*c* = 2.10, CHCl₃) for a 88% *ee* sample.

The optical purity of **4hb** was established by chiral HPLC analysis (OD column, 8/2 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-3-(2-Methoxyphenylamino)-1,5-diphenylpent-4-en-1-one (4ia); The representative procedure was followed. The crude product was purified by silica gel chromatography (4:1 hexanes:Et₂O) to deliver 218 mg (0.6 mmol, 51%) of the desired product **4ia** as a white solid. IR (neat, NaCl): 3408 (w), 1683 (s), 1602 (s), 1513 (s), 1451 (s), 1224 (s), 1029 (m), 966 (m), 739 (s), 689 (s). ¹H NMR (CDCl₃, 400 MHz): δ 7.96 (2H, dd, *J* = 7.1, 1.3 Hz), 7.58-7.16 (8H, m), 6.85-6.61 (5H, m), 6.31 (1H, dd, *J* = 15.9, 6.0 Hz), 4.76-4.71 (2H, m), 3.85 (3H, s), 3.42 (2H, ddd, *J* = 16.3, 6.6, 5.7 Hz). ¹³C NMR (CDCl₃, 100 MHz): δ 198.2, 147.3, 137.1, 136.9, 133.4, 130.7, 128.8, 128.6, 128.3, 127.6, 126.6, 125.4, 124.7, 121.4, 117.3, 111.5, 109.8, 55.6, 52.0, 44.5. HRMS (ES) Calcd for C₂₄H₂₃NO₂Na: 380.1628; Found: 380.1626. Anal Calcd for C₂₄H₂₃NO₂: C, 80.64, H, 6.49; Found: C, 80.58, H, 6.50. [α]_D²⁰ = +1.4° (*c* = 0.97, CHCl₃) for a 90% *ee* sample.

The optical purity of **4ia** was established by chiral HPLC analysis (OD column, 8/2 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-4-(2-Methoxyphenylamino)-6-phenylhex-5-en-2-one (4ib); The representative procedure was followed. The crude product was purified by silica gel chromatography (4:1 hexanes:EtOAc) to deliver 273 mg (0.92 mmol, 77%) of the desired product **4ib** as a clear colorless oil. IR (neat, NaCl): 3408 (w), 1709 (s), 1595 (s), 1513 (s), 1457 (m), 1218 (s), 1029 (m), 966 (m), 746 (s). ¹H NMR (CDCl₃, 400 MHz): δ 7.33-7.17 (5H, m), 6.84-6.55 (5H, m), 6.20 (1H, dd, *J* = 16.0, 6.0 Hz),

4.60-4.51 (2H, m), 3.85 (3H, s), 2.86 (2H, ddd, $J = 16.1, 6.8, 6.6$ Hz), 2.18 (3H, s). ^{13}C NMR (CDCl_3 , 100 MHz): δ 206.9, 147.2, 136.8, 136.7, 130.8, 130.5, 128.6, 127.7, 126.6, 121.4, 117.3, 111.4, 109.8, 55.6, 51.7, 49.5, 30.9. HRMS (ES) Calcd for $\text{C}_{19}\text{H}_{21}\text{NO}_2\text{Na}$: 318.1471; Found: 318.1470. $[\alpha]_{\text{D}}^{20} = +31.9^\circ$ ($c = 0.446$, CHCl_3) for a 89% *ee* sample.

The optical purity of **4ib** was established by chiral HPLC analysis (OD column, 7/3 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-5-(4-Methoxyphenyl)-3-(2-methoxyphenylamino)-1-phenylpent-4-en-1-one (**4ja**);

A modified procedure was used. Ligand **1** (18 mg, 36 μmol) and AgOAc (6.0 mg, 36 μmol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 1 mL of toluene and allowed to stir for 5 min at 22 $^\circ\text{C}$. Imine **2j** (320 mg, 1.20 mmol) was added immediately followed by 100 μL *i*-PrOH (1.30 mmol). The test tube was capped with a septum, and allowed to stir at 22 $^\circ\text{C}$ for 5 min. Enol ether **3a** (0.5 mL, 2.4 mmol) was added and the reaction mixture was allowed to stir at 22 $^\circ\text{C}$ for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH_2Cl_2 (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH_2Cl_2) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH_2Cl_2 and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes: Et_2O) to deliver 343 mg (0.89 mmol, 74%) of the desired product **4ja** as a white solid. IR (neat, NaCl): 3408 (w), 1677 (m), 1602 (m), 1507 (s), 1457 (m), 1249 (s), 1218 (m), 1174 (m), 1029 (m), 733 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.95 (2H, d, $J = 7.1$ Hz), 7.57-7.22 (5H, m), 6.84-6.55 (7H, m), 6.16 (1H, dd, $J = 16.0, 6.0$ Hz), 4.70-4.67 (2H, m), 3.84 (3H, s), 3.77 (3H, s), 3.40 (2H, ddd, $J = 15.9, 6.6, 5.5$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 198.3, 159.3, 147.2, 137.2, 136.8, 133.4, 130.2, 129.7, 128.8, 128.5, 128.3, 127.7, 121.4, 117.2, 114.0, 111.5, 109.7, 55.6, 55.4, 52.0, 44.6. HRMS (ES) Calcd for $\text{C}_{25}\text{H}_{25}\text{NO}_3\text{Na}$: 410.1726; Found: 410.1726. $[\alpha]_{\text{D}}^{20} = +10.5^\circ$ ($c = 0.740$, CHCl_3) for a 96% *ee* sample.

The optical purity of **4ja** was established by chiral HPLC analysis (OD column, 8/2 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-6-(4-Methoxyphenyl)-4-(2-methoxyphenylamino)-hex-5-en-2-one (4jb); A modified procedure was used. Ligand **1** (10 mg, 20 μ mol) and AgOAc (3.0 mg, 20 μ mol)

were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.4 mL of toluene and allowed to stir for 5 min at 22 °C. Imine **2j** (110 mg, 0.40 mmol) was added immediately followed by 33 μ L *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir at 22 °C for 5 min. Enol ether **3b** (130 μ L, 0.80 mmol) was added and the reaction mixture was allowed to stir at 22 °C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:EtOAc) to deliver 62 mg (0.2 mmol, 47%) of the desired product **4jb** as a white solid. IR (neat, NaCl): 3404 (w), 1711 (m), 1605 (s), 1512 (s), 1456 (m), 1252 (s), 1177 (m), 1022 (m), 737 (m). ¹H NMR (CDCl₃, 400 MHz): δ 7.25 (2H, dd, *J* = 7.7, 0.4 Hz), 6.84-6.50 (7H, m), 6.05 (1H, dd, *J* = 15.9, 6.2 Hz), 4.59-4.47 (2H, m), 3.84 (3H, s), 3.77 (3H, s), 2.84 (2H, ddd, *J* = 16.1, 6.6, 6.0 Hz), 2.17 (3H, s). ¹³C NMR (CDCl₃, 100 MHz): δ 207.0, 159.3, 147.2, 136.8, 130.2, 129.6, 128.2, 127.7, 121.4, 117.2, 114.1, 111.4, 109.7, 55.6, 55.4, 51.8, 49.6, 30.9. HRMS (ES) Calcd for C₂₀H₂₃NO₃Na: 348.1580; Found: 348.1576. [α]_D²⁰ = +27.4° (*c* = 0.667, CHCl₃) for a 90% *ee* sample.

The optical purity of **4jb** was established by chiral HPLC analysis (OD column, 9/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-3-(2-Methoxyphenylamino)-5-(4-nitrophenyl)-1-phenylpent-4-en-1-one (4ka); A modified procedure was used. Ligand **1** (6.0 mg, 12 μ mol) and AgOAc (2.0 mg, 12 μ mol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.4 mL of THF and allowed to stir for 5 min at 22 $^{\circ}$ C. Imine **2k** (115 mg, 0.40 mmol) was added immediately followed by 33 μ L *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir in a cold bath at -5 $^{\circ}$ C for 5 min. Enol ether **3a** (160 μ L, 0.80 mmol) was added and the reaction mixture was allowed to stir at -5 $^{\circ}$ C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:Et₂O) to deliver 160 mg (0.40 mmol, 99%) of the desired product **4ka** as a orange solid. IR (neat, NaCl): 3402 (w), 1683 (m), 1595 (s), 1513 (s), 1457 (w), 1344 (s), 1224 (m), 746 (m). ¹H NMR (CDCl₃, 400 MHz): δ 8.11 (2H, d, *J* = 8.8 Hz), 7.97 (2H, d, *J* = 7.1 Hz), 7.61-7.41 (4H, m), 6.87-6.68 (5H, m), 6.54 (2H, dd, *J* = 16.0, 5.5 Hz) 4.81-4.77 (2H, m), 3.87 (3H, s), 3.47 (2H, ddd, *J* = 16.8, 6.8, 5.5 Hz). ¹³C NMR (CDCl₃, 100 MHz): δ 197.7, 147.1, 146.8, 143.3, 136.8, 136.4, 135.9, 133.5, 128.8, 128.6, 128.2, 127.0, 123.9, 121.3, 117.4, 111.2, 109.8, 55.5, 51.5, 44.1. HRMS (ES) Calcd for C₂₄H₂₂N₂O₄Na: 425.1458; Found: 425.1477. Anal Calcd for C₂₄H₂₂N₂O₄: C, 71.63, H, 5.51; Found: C, 71.74, H, 5.54. [α]_D²⁰ = +7.4 $^{\circ}$ (*c* = 2.29, CHCl₃) for a 92% *ee* sample.

HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:Et₂O) to deliver 160 mg (0.40 mmol, 99%) of the desired product **4ka** as a orange solid. IR (neat, NaCl): 3402 (w), 1683 (m), 1595 (s), 1513 (s), 1457 (w), 1344 (s), 1224 (m), 746 (m). ¹H NMR (CDCl₃, 400 MHz): δ 8.11 (2H, d, *J* = 8.8 Hz), 7.97 (2H, d, *J* = 7.1 Hz), 7.61-7.41 (4H, m), 6.87-6.68 (5H, m), 6.54 (2H, dd, *J* = 16.0, 5.5 Hz) 4.81-4.77 (2H, m), 3.87 (3H, s), 3.47 (2H, ddd, *J* = 16.8, 6.8, 5.5 Hz). ¹³C NMR (CDCl₃, 100 MHz): δ 197.7, 147.1, 146.8, 143.3, 136.8, 136.4, 135.9, 133.5, 128.8, 128.6, 128.2, 127.0, 123.9, 121.3, 117.4, 111.2, 109.8, 55.5, 51.5, 44.1. HRMS (ES) Calcd for C₂₄H₂₂N₂O₄Na: 425.1458; Found: 425.1477. Anal Calcd for C₂₄H₂₂N₂O₄: C, 71.63, H, 5.51; Found: C, 71.74, H, 5.54. [α]_D²⁰ = +7.4 $^{\circ}$ (*c* = 2.29, CHCl₃) for a 92% *ee* sample.

The optical purity of **4ka** was established by chiral HPLC analysis (OD column, 75/25 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-4-(2-Methoxyphenylamino)-6-(4-nitrophenyl)-hex-5-en-2-one (4kb); A modified procedure was used. Ligand **1** (6.0 mg, 12 μ mol) and AgOAc (2.0 mg, 12 μ mol) were

weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.4 mL of THF and allowed to stir for 5 min at 22 °C. Imine **2k** (115 mg, 0.40 mmol) was added immediately followed by 33 μ L *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir in a cold bath at -5 °C for 5 min. Enol ether **3b** (130 μ L, 0.80 mmol) was added and the reaction mixture was allowed to stir at -5 °C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous

HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:Et₂O) to deliver 101 mg (0.30 mmol, 74%) of the desired product **4kb** as a orange solid. IR (neat, NaCl): 3402 (w), 1715 (m), 1595 (m), 1513 (s), 1344 (s), 1224 (m), 1023 (w), 739 (m). ¹H NMR (CDCl₃, 400 MHz): δ 8.13 (2H, dd, *J* = 6.9, 1.8 Hz), 7.44 (2H, dd, *J* = 6.9, 1.7 Hz), 6.86-6.63 (5H, m), 6.44 (1H, dd, *J* = 15.9, 5.7 Hz), 4.62-4.55 (2H, m), 3.87 (3H, s), 2.91 (2H, ddd, *J* = 16.5, 6.4, 5.8 Hz), 2.21 (3H, s). ¹³C NMR (CDCl₃, 100 MHz): δ 206.4, 147.1, 147.0, 143.3, 136.4, 135.6, 128.7, 127.1, 124.0, 121.3, 117.6, 111.2, 109.8, 55.6, 51.4, 49.1, 30.8. HRMS (ES) Calcd for C₁₉H₂₀N₂O₄Na: 363.1311; Found: 363.1321. Anal Calcd for C₁₉H₂₀N₂O₄: C, 67.05, H, 5.92; Found: C, 67.39, H, 5.71. $[\alpha]_D^{20}$ = +30.9° (*c* = 1.67, CHCl₃) for a 90% *ee* sample.

The optical purity of **4kb** was established by chiral HPLC analysis (OD column, 75/25 hexanes / *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-3-(2-Methoxyphenylamino)-1,5-diphenylpent-4-yn-1-one (4la); The representative procedure was followed. The crude product was purified by silica gel chromatography (4:1 hexanes:Et₂O) to deliver 395 mg (1.10 mmol, 93%) of the desired product **4la** as a white solid. IR (neat, NaCl): 3389 (w), 3062 (w), 1690 (s), 1602 (s), 1513 (s), 1451 (s), 1249 (s), 1224 (s), 1123 (m), 1035 (m), 909 (m), 758 (s), 689 (s). ¹H NMR (CDCl₃, 400 MHz): δ 8.01 (2H, d, *J* = 7.1 Hz), 7.60-4.45 (3H, m), 7.32-7.21 (5H, m), 6.98-6.91 (2H, m), 6.82-6.74 (2H, m), 5.07 (1H, apparent triplet, *J* = 6.4 Hz), 4.82 (1H, br), 3.84 (3H, s), 3.60 (2H, ddd, *J* = 16.4, 6.6, 6.0 Hz). ¹³C NMR (CDCl₃, 100 MHz): δ 197.1, 147.7, 137.0, 136.1, 133.4, 131.8, 128.8, 128.3, 128.2, 127.6, 122.9, 121.3, 118.1, 112.1, 109.9, 89.3, 83.3, 55.6, 44.3, 42.5. HRMS (ES) Calcd for C₂₄H₂₁NO₂Na: 378.1472; Found: 378.1470. [α]_D²⁰ = +82.0° (*c* = 0.62, CHCl₃) for a 92% *ee* sample.

The optical purity of **4la** was established by chiral HPLC analysis (OD column, 7/3 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-4-(2-Methoxyphenylamino)-6-phenylhex-5-yn-2-one (4lb); A modified procedure was used. Ligand **1** (6.0 mg, 12 μmol) and AgOAc (2.0 mg, 12 μmol) were weighed into a 16 x 150 mm test tube. The contents were dissolved in 0.4 mL of THF and allowed to stir for 5 min at 22 °C. Imine **2l** (95 mg, 0.4 mmol) was added immediately followed by 33 μL *i*-PrOH (0.45 mmol). The test tube was capped with a septum, and allowed to stir in a cold bath at -5 °C for 5 min.

Enol ether **3b** (130 μ L, 0.80 mmol) was added and the reaction mixture was allowed to stir at -5 $^{\circ}$ C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH_2Cl_2 (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH_2Cl_2) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH_2Cl_2 and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:EtOAc) to deliver 107 mg (0.37 mmol, 91% yield) of the desired product **4b** as a light orange solid. IR (neat, NaCl): 3383 (w), 1715 (m), 1602 (m), 1507 (s), 1457 (m), 1243 (m), 1224 (m), 1123 (m), 1029 (m), 758 (m), 739 (m), 689 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.35-7.22 (5H, m), 6.90-6.72 (4H, m), 4.82 (1H, apparent triplet, J = 6.4 Hz), 4.67 (1H, br), 3.83 (3H, s), 3.03 (2H, ddd, J = 16.3, 6.4, 6.2 Hz), 2.24 (3H, s). ^{13}C NMR (CDCl_3 , 100 MHz): δ 205.8, 147.7, 136.0, 131.9, 128.3, 128.3, 122.9, 121.3, 118.3, 112.2, 109.9, 89.0, 83.2, 55.6, 49.1, 42.3, 31.0. HRMS (ES) Calcd for $\text{C}_{19}\text{H}_{19}\text{NO}_2\text{Na}$: 316.1309; Found: 316.1313. $[\alpha]_{\text{D}}^{20}$ = +101 $^{\circ}$ (c = 1.12, CHCl_3) for a 88% *ee* sample.

The optical purity of **4b** was established by chiral HPLC analysis (OD column, 75/25 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-4-(2-Methoxyphenylamino)-tetradecan-2-one(7m); To a 16 x 150 mm test tube charged with 250 μ L (1.2 mmol) of undecylic aldehyde (**m**) cooled to -

5 $^{\circ}$ C was added 135 μ L (1.2 mmol) of *o*-anisidine. The solution was allowed to stir at that temperature for 5 min upon which time it turned cloudy. THF was then added (1.0 mL) and the solution was allowed to warm to 22 $^{\circ}$ C and ligand **1** (31 mg, 60 μ mol) and AgOAc (10 mg, 60 μ mol) were added and the solution was allowed to stir at that temperature for 5 min. The solution was then cooled to 4 $^{\circ}$ C in a cold room and enol ether **3b** (400 μ L, 2.40 mmol) was added and the reaction mixture was allowed to stir at 4 $^{\circ}$ C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH_2Cl_2 (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH_2Cl_2) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH_2Cl_2 and separated as stated

above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:Et₂O) to deliver 240 mg (0.72 mmol, 60%) of the desired product **4m** as a clear colorless oil. IR (neat, NaCl): 3422 (w), 2917 (s), 2855 (s), 1710 (m), 1597 (m), 1512 (m), 1455 (m), 1353 (w), 1217 (m), 1030 (w). ¹H NMR (CDCl₃, 400 MHz): δ 6.85 (1H, ddd, *J* = 7.6, 1.3, 1.1 Hz), 6.77 (1H, dd, *J* = 7.8, 1.1 Hz), 6.67-6.61 (2H, m), 4.18 (1H, s), 3.84 (1H, partially covered), 3.83 (3H, s), 2.72 (1H, dd, *J* = 16.3, 5.1 Hz), 2.57 (1H, dd, *J* = 16.2, 6.9 Hz), 2.14 (3H, s), 1.59-1.25 (18H, m), 0.88 (3H, t, *J* = 6.6 Hz). ¹³C NMR (CDCl₃, 100 MHz): δ 208.2, 147.0, 137.2, 121.5, 116.4, 110.3, 109.8, 55.5, 49.4, 48.6, 35.4, 32.0, 31.0, 29.7, 29.7, 29.6, 29.4, 26.3, 22.8, 14.3. HRMS (ES) Calcd for C₂₁H₃₆NO₂: 334.2748; Found: 224.2746. [α]_D²⁰ = +0.187 ° (*c* = 2.14, CHCl₃) for a 92% *ee* sample.

The optical purity of **4m** was established by chiral HPLC analysis (OD column, 95/5 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-4-Cyclohexyl-4-(2-methoxyphenylamino)-butan-2-one (7n); In a 16 x 150 mm test tube charged with 145 μ L (1.20 mmol) of cyclohexane carboxaldehyde (**n**) cooled to -5 °C in a cold bath was added 135 μ L (1.20 mmol) of *o*-anisidine and the solution was

allowed to stir at that temperature for 5 min upon which time it turned cloudy. THF was then added (1.0 mL) and the solution was allowed to warm to 22 °C and ligand **1** (30.5 mg, 60 μ mol) and AgOAc (10 mg, 60 μ mol) were added and the solution was allowed to stir at that temperature for 5 min. The solution was then cooled to 4 °C in a cold room and enol ether **3b** (400 μ L, 2.40 mmol) was added and the reaction mixture was allowed to stir at 4 °C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:Et₂O) to deliver 176 mg (0.64 mmol, 53%) of the desired product **4n** as a clear colorless oil. IR (neat, NaCl): 3422 (w), 2929 (s), 2855 (m), 1716 (s), 1603 (s), 1518 (s), 1455 (m), 1257 (m), 1217 (m), 1121 (w), 1030 (m). ¹H NMR (CDCl₃, 400 MHz): δ 6.84 (1H, dddd, *J* = 8.1, 1.3, 0.9, 0.5 Hz), 6.75 (1H, dd, *J* = 7.9, 1.5 Hz), 6.66-6.60 (2H, m), 4.27 (1H, s), 3.83 (3H, s), 3.78 (1H, q, *J* = 6.0 Hz), 2.62 (2H, dddd, *J* = 16.0, 9.2, 6.6, 5.7 Hz), 2.14 (3H, s),

1.86-1.82 (1H, s), 1.79-1.72 (2H, m), 1.68-1.64 (2H, m), 1.61-1.53 (1H, m), 1.27-1.0 (5H, m). ^{13}C NMR (CDCl_3 , 100 MHz): δ 208.3, 146.9, 137.5, 121.4, 116.1, 110.3, 109.8, 55.6, 54.0, 46.2, 42.1, 30.7, 29.5, 29.3, 26.6, 26.4. HRMS (ES) Calcd for $\text{C}_{17}\text{H}_{25}\text{NO}_2\text{Na}$: 298.1787; Found: 298.1783. $[\alpha]_{\text{D}}^{20} = +7.1^\circ$ ($c = 2.4$, CHCl_3) for a 94% *ee* sample.

The optical purity of **4n** was established by chiral HPLC analysis (OD column, 95/5 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below

(-)-4-(2-Methoxyphenylamino)-6-methylheptan-2-one (7o); In a 16 x 150 mm test tube charged with 35 μL (0.40 mmol) of isovaleraldehyde (**o**) cooled to -5°C in a cold

bath was added 45 μL (0.40 mmol) of *o*-anisidine and the solution was allowed to stir at that temperature for 5 min upon which time it turned cloudy. THF was then added (0.5 mL) and the solution was allowed to warm to 22°C upon which time ligand **1** (10 mg, 0.02 mmol) and AgOAc (3.0 mg, 0.02 mmol) were added and the solution was allowed to stir at that temperature for 5 min. The solution was then cooled to 4°C in a cold room and enol ether **3b** (130 μL , 0.80 mmol) was added and the reaction mixture was allowed to stir at 4°C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH_2Cl_2 (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH_2Cl_2) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH_2Cl_2 and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes: Et_2O) to deliver 41 mg (0.16 mmol, 41%) of the desired product **4o** as a clear colorless oil. IR (neat, NaCl): 3402 (w), 2953 (m), 1707 (s), 1595 (s), 1514 (s), 1458 (s), 1246 (m), 1221 (m), 1028 (m), 735 (s). ^1H NMR (CDCl_3 , 400 MHz): δ 6.85 (1H, dt, $J = 7.8, 1.4$ Hz), 6.76 (1H, dd, $J = 7.3, 1.1$ Hz), 6.66-6.62 (2H, m), 4.13 (1H, s), 3.97-3.90 (1H, m), 3.83 (3H, s), 2.72 (1H, dd, $J = 16.3, 4.7$ Hz), 2.55 (1H, dd, $J = 16.3, 7.1$ Hz), 2.13 (3H, s), 1.80-1.71 (1H, m), 1.54-1.34 (2H, m), 0.92 (6H, dd, $J = 11.4, 6.8$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 208.3, 147.0, 137.2, 121.5, 116.4, 110.1, 109.8, 55.6, 48.9, 57.4, 45.0, 31.1, 25.1, 23.3, 22.3. HRMS (ES) Calcd for $\text{C}_{15}\text{H}_{23}\text{NO}_2\text{Na}$: 272.1624; Found: 272.1626. $[\alpha]_{\text{D}}^{20} = -7.0$ ($c = 1.2$, CHCl_3) for a 94% *ee* sample.

The optical purity of **4o** was established by chiral HPLC analysis (OD column, 95/5 hexanes / *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(-)-[1-(4-Chlorophenyl)-3-oxo-3-phenylpropyl]-carbamic acid *tert*-butyl ester (10**);**

To a solution of $\text{PhI}(\text{OAc})_2$ (4.20 g, 13.1 mmol) in 20 mL of MeOH and 1 mL of AcOH at 22 °C was added **4da** (1.2 g, 3.3 mmol, in 2.0 mL MeOH) slowly at 22 °C over 30 min by a pressure equalizing addition funnel. Upon complete addition the reaction was allowed to stir at 22 °C for an additional 30 min upon which time 15 mL of a 10% aq. HCl solution (wt %) was added. The mixture was allowed to stir for 30 minutes at which time 15 mL of a 10% aq. $\text{Na}_2\text{S}_2\text{O}_3$ solution (wt %) was added and stirring was allowed to continue for an additional 30 min. The solution was made basic by the addition of Na_2CO_3 upon which the solution turns dark red, and Boc_2O was added (2.90 g, 13.1 mmol) followed by 5 mL of CH_2Cl_2 . The mixture was allowed to stir at 22 °C for 12 h. The layers were separated and the aqueous layer was washed with 2 x 50 mL of CH_2Cl_2 . The combined organic layers were washed with brine (1 x 20 mL), dried over MgSO_4 and concentrated. The product was recrystallized from hot hexanes yield 935 mg (2.60 mmol, 79%) of **10** as a white solid. IR (neat, NaCl): 3340 (w), 2356 (m), 1690 (s), 1489 (m), 1440 (w), 1360 (m), 1244 (w), 1165 (m), 1092 (m), 1006 (m), 750 (w). ^1H NMR (CDCl_3 , 400 MHz): δ 8.86 (2H, d, $J = 7.5$ Hz), 7.53 (1H, t, $J = 7.3$ Hz), 7.43-7.39 (2H, m), 7.28-7.22 (4H, m), 5.66 (1H, br), 5.20 (1H, br), 3.63-3.60 (1H, m), 3.39 (1H, dd, $J = 16.8, 5.9$ Hz), 1.39 (9H, s). ^{13}C NMR (CDCl_3 , 100 MHz): δ 197.9, 155.3, 140.5, 136.7, 133.6, 133.0, 128.8, 128.7, 128.2, 127.9, 79.9, 50.9, 44.0, 28.4. HRMS (ES) Calcd for $\text{C}_{20}\text{H}_{22}\text{NO}_3\text{NaCl}$: 382.1185; Found: 382.1186. Anal Calcd for $\text{C}_{20}\text{H}_{22}\text{ClNO}_3$: C, 66.75, H, 6.16; Found: C, 66.71, H, 6.15. $[\alpha]_D^{20} = -4.8^\circ$ ($c = 1.9$, CHCl_3) for a >99% *ee* sample.

The optical purity of **4ba** was established by chiral HPLC analysis (AD column, 9/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-(3-Oxo-1-phenylethynylbutyl)-carbamic acid *tert*-butyl ester (11**);** To a solution of $\text{PhI}(\text{OAc})_2$ (0.55g, 1.7 mmol) in 5.0 mL of MeOH and 0.10 mL of AcOH at 22 °C was added **4b** (125 mg, 0.43 mmol, in 1.0 mL MeOH) slowly at 22 °C over 20 min by pipette. Upon complete addition the reaction was allowed to stir at 22 °C for an additional 30 min upon which time 5 mL of a 10% aq. HCl solution (wt %) was added. The mixture was allowed to stir for 30 minutes at which time 5 mL of a 10% aq. $\text{Na}_2\text{S}_2\text{O}_3$ solution (wt %) was added and stirring was allowed to continue for an additional 30 min. The solution was made basic by the addition of Na_2CO_3 upon which the solution turns dark red, and Boc_2O was added (370 mg, 1.7 mmol) followed by 2 mL of CH_2Cl_2 . The reaction was allowed to stir at 22 °C for 12 h. The layers were separated and the aqueous layer was washed with 2 x 25 mL of CH_2Cl_2 . The combined organic layers were washed with brine (1 x 10 mL), dried over MgSO_4 and concentrated. The product was purified by silica gel chromatography (3 : 2 hexanes : Et_2O) to yield 85 mg (0.3 mmol, 70%) of **11** as a yellow orange oil. IR (neat, NaCl): 3345 (s), 2974 (m), 1715 (s), 1501 (s), 1369 (s), 1249 (m), 1167 (s), 1048 (m), 765 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.36-7.25 (5H, m), 5.36 (1H, br), 4.95 (1H, br), 2.90 (2H, ddd, J = 16.8, 6.2, 4.8 Hz), 2.19 (3H, s), 1.43 (9H, s). ^{13}C NMR (CDCl_3 , 100 MHz): δ 205.8, 154.8, 131.8, 128.5, 128.3, 122.6, 87.6, 83.1, 80.2, 48.7, 39.7, 30.7, 28.4. HRMS (ES) Calcd for $\text{C}_{17}\text{H}_{21}\text{NO}_3\text{Na}$: 310.1423; Found: 310.1419. $[\alpha]_{\text{D}}^{20}$ = +48.0° (c = 0.767, CHCl_3) for a 88% *ee* sample.

(-)-5-(2-Methoxyphenylamino)-7-oxo-7-phenylheptanoic acid methyl ester (8**);** In a 16 x 150 mm test tube charged with 156 μL (1.20 mmol) of 5-oxo-pentanoic acid methyl ester⁵ cooled to -5 °C in a cold bath was added 135 μL (1.20 mmol) of *o*-anisidine and the solution was allowed to stir at this temperature for 5 min upon which time it turned cloudy. THF was added (1.0 mL) and the solution was allowed to warm to 22 °C at which time ligand **1** (31 mg, 60 μmol) and AgOAc (10 mg, 60 μmol) were added and the solution was allowed to stir at this temperature for 5 min. The solution was cooled to 4 °C in a cold room and enol ether **3a** (0.5 mL, 2.4

⁵ Huckstep, J.; Taylor, R.J.K. *Synthesis* **1982**, 881-882.

mmol) was added and the mixture was allowed to stir at 4 °C for 16 h. The reaction was quenched upon addition of 2 mL of 10 % aqueous HCl followed by vigorous stirring for 5 min. CH₂Cl₂ (5 mL) was added and the mixture was allowed to stir vigorously for 1 min. The layers were allowed to separate, and the bottom layer (CH₂Cl₂) was removed with a pipette and placed into a round-bottom flask. The aqueous layer was washed with 2 x 5 mL portions of CH₂Cl₂ and separated as stated above. The combined organic layers were concentrated (without drying) and purified by silica gel chromatography (4:1 hexanes:Et₂O) to deliver 239 mg (0.67 mmol, 56%) of the desired product **8** as a light yellow oil. IR (neat, NaCl): 3408 (w), 2942 (m), 1734 (s), 1683 (s), 1602 (s), 1513 (s), 1451 (m), 1224 (s), 1180 (m), 1029 (m), 746 (s). ¹H NMR (CDCl₃, 400 MHz): δ 7.91 (2H, d, *J* = 7.7 Hz), 7.55 (1H, dd, *J* = 7.5, 7.1 Hz), 7.44 (2H, dd, *J* = 7.9, 7.5 Hz), 6.84 (1H, dd, *J* = 8.0, 7.3 Hz), 6.76 (1H, d, *J* = 8.0 Hz), 6.75-6.65 (2H, m), 4.30 (1H, s), 4.10-4.02 (1H, m), 3.83 (3H, s), 3.64 (3H, s), 3.28 (1H, dd, *J* = 16.8, 3.7 Hz), 3.13 (1H, dd, *J* = 16.8, 7.3 Hz), 2.40-2.30 (2H, m), 1.99-1.60 (4H, m). ¹³C NMR (CDCl₃, 100 MHz): δ 199.5, 173.9, 133.2, 128.7, 128.1, 121.4, 121.2, 116.6, 115.1, 110.3, 109.8, 55.5, 51.6, 49.2, 43.2, 34.8, 33.9, 21.8. HRMS (ES) Calcd for C₂₁H₂₆NO₄: 356.1845; Found: 356.1862. [α]_D²⁰ = -12.6 ° (*c* = 1.27, CHCl₃) for a >99% *ee* sample.

The optical purity of **8** was established by chiral HPLC analysis (AD column, 9/1 hexanes/ *i*-PrOH eluent, 1 mL/min and a lamp setting of 254 nm); chromatograms are shown below.

(+)-6-(2-Oxo-2-phenylethyl)-piperidin-2-one (15); To a solution of PhI(OAc)₂ (0.325, 1.00 mmol) in 5 mL of MeOH and 0.10 mL of AcOH at 22 °C was added **14** (90 mg, 0.25 mmol, in 1.0 mL MeOH) slowly at 22 °C over 20 min by pipette. Upon complete addition the reaction was allowed to stir at 22 °C for an additional 30 min upon which time 5 mL of a 10% aq. HCl solution (wt %) was added. The reaction was allowed to stir for 30 minutes after which time 5 mL of a 10% aq. Na₂S₂O₃ solution (wt %) was added and stirring was continued for an additional 30 min. The solution was then made basic by the addition of Na₂CO₃ at which time the solution turns dark red. CH₂Cl₂ was added (2 mL) and the mixture was allowed to stir at 22 °C for 12 h. The layers were then separated and the aqueous layer was washed with 2 x 25 mL of CH₂Cl₂. The combined organic layers were washed with brine (1 x 10 mL), dried over MgSO₄ and concentrated. The product was purified by silica gel chromatography (4:1 EtOAc:MeOH) to yield 44 mg (0.20 mmol, 80% yield) as a white solid. IR (neat, NaCl): 3254 (w), 1681 (s), 1662 (s), 1600 (s), 1370 (s), 1215 (m), 960 (m), 761 (m), 686 (m). ¹H

NMR (CDCl₃, 400 MHz): δ 7.92 (2H, d, J = 7.0 Hz), 7.61-7.45 (3H, m), 6.42 (1H, s), 4.09-4.03 (1H, m), 3.17 (2H, ddd, J = 17.7, 9.1, 3.7 Hz), 2.45-2.27 (2H, m), 2.02-1.97 (1H, m), 1.96-1.87 (1H, m), 1.84-1.74 (1H, m), 1.57-1.48 (1H, m). ¹³C NMR (CDCl₃, 100 MHz): δ 198.3, 171.9, 136.5, 133.9, 128.9, 128.1, 48.7, 45.5, 31.7, 28.6, 19.6. HRMS (ES) Calcd for C₁₃H₁₅NO₂Na: 240.1006; Found: 240.1000. $[\alpha]_D^{20}$ = +45.0° (c = 0.6000, CHCl₃).

(–)-Sedamine; In a 100 mL flask amide **15** (250 mg, (1.15 mmol) was dissolved in 20 mL of THF (distilled from Na-benzophenone ketyl) and cooled to –78 °C. A 1M solution of DIBAL-H in hexanes (5.75 mL, 5.75 mmol) was added slowly to the solution by syringe over 30 min. The solution was allowed to stir at –78 °C for an additional 30 min at which time it was warmed to –40 °C and 130 mg (3.4 mmol) of LAH was added directly. The reaction was allowed to stir at –40 °C for 30 min then warm to 22 °C and continued to stir for an additional 1 h. The reaction was then cooled to 0 °C and a 10% (aq.) solution (wt %) of sodium potassium tartrate was added drop wise until the vigorous gas evolution stopped. The solution was allowed to warm to 22 °C and 15 mL of 10% (aq.) solution of sodium potassium tartrate was added and the mixture was allowed to continue stirring for 30 min. The organic layer was separated and the aqueous layer was washed with 3 x 10 mL of CH₂Cl₂. The combined organic layers were dried over MgSO₄ and concentrated to give a light yellow solid. The analysis of the ¹H NMR spectrum of this solid shows a 16:1 diastereoselectivity of desired syn reduction to undesired anti reduction.

The resulting solid was then subjected to modified Borch reductive amination conditions⁶. The unpurified product was dissolved in 12 mL of CH₃CN at 22 °C and NaCNBH₃ (360 mg, 5.75 mmol) was added to the solution directly followed by 3 mL of 37% formaldehyde (aq.) solution (0.35 mol) and the reaction was allowed to stir for 5 min. AcOH (600 μ L) was added to the solution in 3 parts over 2 h at 40 min intervals. After each addition of AcOH the reaction was highly exothermic. The solution was allowed to stir at 22 °C for 12 h after which time 5 mL of 0.1M NaOH (aq.) solution was added and the mixture was allowed to continue stirring for 30 min followed by the addition of 5 mL of CH₂Cl₂. The organic layer was separated and aqueous layer was washed with 3 x 10 mL of CH₂Cl₂. The combined organic layers were dried over MgSO₄ and concentrated. The product was purified by column chromatography on neutral alumina (9 : 1 CH₂Cl₂ : MeOH) to provide 224 mg (1.0 mmol, 89% of (–)-sedamine as a white solid. IR (neat, NaCl): 3269 (br), 2936 (s), 2860 (m), 2785 (m), 1740 (m), 1683 (m), 1457 (m), 1375 (m), 1230 (m), 1060 (s), 1029 (s), 758 (m), 702 (s). ¹H NMR (CDCl₃, 400 MHz): δ 7.38-7.20 (5H, m), 6.22 (1H, br), 4.87 (1H, dd, J = 10.5, 2.4 Hz), 3.06-3.02 (1H, m), 2.85-2.81 (1H, m), 2.56-2.51 (1H, m), 2.47 (3H, s), 2.14-2.06 (1H, m), 1.74-1.71 (1H, m), 1.64-1.54 (2H, m), 1.48-1.44 (4H, m). ¹³C NMR (CDCl₃, 100 MHz): δ 145.8, 128.3, 127.1, 127.0, 125.7, 125.6, 74.7, 61.0, 51.4, 40.1, 39.9, 25.9, 22.5, 20.7. HRMS (ES) Calcd for C₁₄H₂₂NO: 220.1704; Found: 220.1701. $[\alpha]_D^{20}$ = –60.8° (c = 1.21,

⁶ Borch, R. F.; Hassid, A. I. *J. Org. Chem.* **1972**, 37, 2321-2323.

EtOH) for a 16:1 diastereomeric mixture. Literature, $[\alpha]_D^{20} = +87.0$, (c = 1.10, EtOH) for (+)-sedamine⁷. M.P. = 59.5-62 °C; literature, 59.0-61 °C.

⁷ Cossy, J.; Willis, C.; Bellosta, V.; BouzBouz, S. *J. Org. Chem.* **2002**, 67, 1982-1992.