Supporting Information

Air Stable Racemization Catalyst for Dynamic Kinetic Resolution of Alcohols at Room Temperature

Namdu Kim, Soo-Byung Ko, Min Serk Kwon, Mahn-Joo Kim,* and Jaiwook
Park*

*National Research Laboratory of Chirotechnology, Department of
Chemistry, Division of Molecular and Life Science, Pohang University
of Science and Technology, San 31 Hyojadong, Pohang, Kyeongbuk 790-
784, Korea.*

Synthesis of 3a.

A solution of $\eta^4-(C_4Ph_4CO)(CO)_3Ru$ (200 mg, 0.351 mmol) and benzyl chloride (44.4 mg, 0.351 mmol) in toluene (4 mL) was placed in a 50-mL flask equipped with a grease-free high-vacuum stopcock under argon. The mixture was heated at 110 °C under argon for 3 days. The reaction mixture was concentrated and purified by chromatography on a silica-gel column to give pale yellow solid **3a** (152 mg, 65% yield). m.p.: dec. 148-150 °C; 1H NMR ($CDCl_3$, 300MHz) (7.53 - 7.56 (m, 4H), 6.93 - 7.29 (m, 21H), 4.83 (s, 2H, CH₂); ^{13}C NMR ($CDCl_3$, 75MHz) (197.5 (CO),

136.0, 133.1, 132.4, 129.9, 129.8, 128.9, 128.7, 128.6, 128.5, 128.4, 128.1, 128.0, 101.9, 89.0, 76.2(CH₂); IR (cm⁻¹) 2036.7, 1989.4; Elemental analysis (%) calcd for C₃₈H₂₇ClO₃Ru: C 68.31; H 4.07; found: C 68.19; H 4.12.

Synthesis of **3b**

From η^4 -(C₄Ph₄CO)(CO)₃Ru (200 mg, 0.351 mmol) and benzyl bromide (60.1 mg, 0.351 mmol) pale yellow solid **3b** (120 mg, 48% yield) was obtained. m.p.: dec. 171-173 °C ; ¹H NMR (CDCl₃, 300MHz) δ 7.55 - 7.58 (m, 4H), 6.94 - 7.31 (m, 21H), 4.86 (s, 2H, CH₂); ¹³C NMR(CDCl₃, 75MHz) δ 198.0 (CO), 136.2, 133.2, 132.5, 129.9, 129.2, 129.1, 129.0, 128.8, 128.8, 128.4, 101.8, 88.9, 76.2; IR (cm⁻¹) 2035.1, 1977.1; Elemental analysis (%) calcd for C₃₈H₂₇BrO₃Ru: C 64.05; H 3.82; found: C 63.94; H 3.86.

Synthesis of **7**

A solution of *p*-chloromethylbenzoyl chloride (567 mg, 3.0 mmol) in dichloromethane (5.0 mL) was added dropwise to a suspension of hydroxymethyl polystyrene (1.77 g, 2.0 mmol; substitution: 1.13

mmol/g)) in dichloromethane (20 mL) at 0 °C. The mixture was stirred for 10 h at room temperature and filtered. The resulting solid was washed with water (20 mL), acetone (20 mL), and dichloromethane (20 mL), and dried under vacuum to give pale yellow solid **7** (2.08 g, 100%). IR (cm⁻¹) 1717.3 (See the spectra given below).

Synthesis of 4

In a 50-mL flask equipped with a grease-free high-vacuum stopcock, **7** (1.04 g, 1.0 mmol), η^4 -(C₄Ph₄CO)(CO)₃Ru (570 mg, 1.0 mmol), and dry toluene (10 mL) were placed under argon. The mixture was stirred at 150 °C for 5 days, and filtered. The resulting solid was washed with acetone (20 mL) and dichloromethane (20 mL), and dried under vacuum to give yellow solid **4** (1.27 g, 42.4 % yield). IR (cm⁻¹) 2042.7, 1992.7, 1719.0 (See the following spectra).

NMR Analysis for a Mixture of **3a**, K_3PO_4 , and 1-Phenylethanol

A mixture of **3a** (6.7 mg, 10 μmol), (*S*)-1-phenylethanol (30 μL , 0.25 mmol), K_3PO_4 (53 mg, 0.25 mmol), and toluene- d_8 (0.8 mL) in an NMR tube was shaken occasionally for 2 h at room temperature for the NMR analysis. See the following spectra.

^1H NMR spectrum

3a + toluene- d_8

3a + toluene- d_8 + 1-phenylethanol + K_3PO_4

^{13}C NMR spectrum

3a + toluene- d_8

3a + toluene- d_8 + 1-phenylethanol + K_3PO_4