**Alkyl- or Aryl-thiolation of Aryl Iodide via Cleavage of S-S bond of Disulfide
Compound by Nickel Catalyst and Zinc**

Nobukazu Taniguchi*

Department of Chemistry, Fukushima Medical University, Fukushima 960-1295 Japan

taniguti@fmu.ac.jp

SUPPORTING INFORMATION

TABLE OF CONTENTS

Contents	Page number
General procedure	S2
References for known compounds	S2~S4
Analytical data	S4~S9

General Procedure. All reactions were carried out under a nitrogen atmosphere. NMR spectra were recorded on a JEOL EX-270 spectrometer (270 MHz for ^1H , 67.5 MHz for ^{13}C). Chemical shifts are reported in δ ppm referenced to an internal tetramethylsilane standard for ^1H NMR and chloroform-*d* (δ 77.0) for ^{13}C NMR. IR spectra were measured by Perkin-Elmer Spectrum One FT-IR spectrometer.

References for known compounds:

Diphenyl sulfide (3aa): (a) Zheng, N.; McWilliams, C.; Fleitz, F. J.; Armstrong III, J. D.; Volante, R. P. *J. Org. Chem.* **1998**, *63*, 9606–9607. (b) Migita, T.; Shimizu, T.; Asami, Y.; Shiobara, J.-i.; Kato, Y.; Kosugi, M. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 1385–1389.

Phenyl 4-tolyl sulfide (3ab): Migita, T.; Shimizu, T.; Asami, Y.; Shiobara, J.-i.; Kato, Y.; Kosugi, M. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 1385–1389.

Methyl phenyl sulfide (3ac): Rossi, R. A.; Palacios, S. M. *J. Org. Chem.* **1981**, *46*, 5300–5304.

***n*-Butyl phenyl sulfide (3ad):** (a) Zheng, N.; McWilliams, C.; Fleitz, F. J.; Armstrong III, J. D.; Volante, R. P. *J. Org. Chem.* **1998**, *63*, 9606–9607. (b) Migita, T.; Shimizu, T.; Asami, Y.; Shiobara, J.-i.; Kato, Y.; Kosugi, M. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 1385–1389. (c) Bates, C. G.; Gujadhur, R. K.; Venkataraman, D. *Org. Lett.* **2002**, *4*, 2803–2806.

Phenyl *iso*-propyl sulfide (3ae): Cogolli, P.; Maiolo, F.; Testaferri, L.; Tingoli, M.; Tiecco, M. *J. Org.*

Chem. **1979**, *44*, 2642–2646.

***n*-Butyl 2-tolyl sulfide (3bd)**: (a) Zheng, N.; McWilliams, C.; Fleitz, F. J.; Armstrong III, J. D.; Volante, R. P. *J. Org. Chem.* **1998**, *63*, 9606–9607. (b) Migita, T.; Shimizu, T.; Asami, Y.; Shiobara, J.-i.; Kato, Y.; Kosugi, M. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 1385–1389. (c) Bates, C. G.; Gujadhur, R. K.; Venkataraman, D. *Org. Lett.* **2002**, *4*, 2803–2806.

***n*-Butyl 2-methoxyphenyl sulfide (3cd)**: Bates, C. G.; Gujadhur, R. K.; Venkataraman, D. *Org. Lett.* **2002**, *4*, 2803–2806.

***n*-Butyl 4-tolyl sulfide (3fd)**: Bates, C. G.; Gujadhur, R. K.; Venkataraman, D. *Org. Lett.* **2002**, *4*, 2803–2806.

***n*-Butyl 4-methoxyphenyl sulfide (3gd)**: (a) Zheng, N.; McWilliams, C.; Fleitz, F. J.; Armstrong III, J. D.; Volante, R. P. *J. Org. Chem.* **1998**, *63*, 9606–9607. (b) Migita, T.; Shimizu, T.; Asami, Y.; Shiobara, J.-i.; Kato, Y.; Kosugi, M. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 1385–1389. (c) Bates, C. G.; Gujadhur, R. K.; Venkataraman, D. *Org. Lett.* **2002**, *4*, 2803–2806.

***n*-Butyl 4-bromophenyl sulfide (3hd)**: Bates, C. G.; Gujadhur, R. K.; Venkataraman, D. *Org. Lett.* **2002**, *4*, 2803–2806.

Phenyl 2-tolyl sulfide (3ba): (a) Bunnett, J. F.; Creary, X. *J. Org. Chem.* **1974**, *39*, 3173–3174. (b) Kosugi, M.; Ogata, T.; Terada, M.; Sano, H.; Migita, T. *Bull. Chem. Soc. Jpn.* **1985**, *58*, 3657–3658. (c) H.

Cristau, J.; Chabaud, B.; Chene, A.; Christol, H. *Synthesis*, **1981**, 892–895.

The reaction conditions and results are summarized in Table 2. A typical procedure is shown in the experimental section.

To a mixture of NiBr₂ (6.6 mg, 0.03 mmol), bpy (4.7 mg, 0.03 mmol), zinc (dust) (39.2 mg, 0.6 mmol) and DMF (0.5 mL) were added iodobenzene (**1a**) (61.2 mg, 0.3 mmol) and diphenyl disulfide (**2a**) (32.8 mg, 0.15 mmol), and the mixture was stirred at 110 °C for 18 h. After evaporation of the solvent, the residue was dissolved in Et₂O, and the resulting mixture was filtered through Celite pad. The solution was washed with H₂O and saturated sodium chloride and dried over anhydrous magnesium sulfate. Chromatography on silica gel (Hexane) gave diphenyl sulfide (**3aa**) (50.1 mg, 90%). ¹H NMR (CDCl₃) δ 7.23–7.36 (m, 10H); ¹³C NMR (CDCl₃) δ 127.0, 129.1, 131.0, 135.7; IR (neat) 1579, 1475, 1438 cm⁻¹; Anal. Calcd for C₁₂H₁₀S C, 77.37; H, 5.41. Found: C, 77.24; H, 5.53.

3ab: ¹H NMR (CDCl₃) δ 2.34 (s, 3H), 7.17–7.33 (m, 9H); ¹³C NMR (CDCl₃) δ 21.1, 126.3, 127.1, 127.5, 129.0, 129.1, 129.7, 130.0, 132.2; IR (neat) 3058, 1581, 1491, 1476 cm⁻¹; Anal. Calcd for C₁₃H₁₂S C, 77.95; H, 6.04. Found: C, 77.69; H, 6.22.

3ac: ¹H NMR (CDCl₃): δ 2.48 (s, 3H), 7.10–7.16 (m, 1H), 7.24–7.31 (m, 4H); ¹³C NMR (CDCl₃) δ 15.9, 125.0, 126.7, 128.8, 138.4; Anal. Calcd for C₇H₈S C, 67.69; H, 6.49. Found: C, 67.68; H, 6.51.

3ad: ¹H NMR (CDCl₃): δ 0.92 (t, *J* = 7.2 Hz, 3H), 1.40–1.48 (m, 2H), 1.53–1.66 (m, 2H), 2.91 (t, *J* = 7.4

Hz, 2H), 7.12–7.34 (m, 5H); ^{13}C NMR (CDCl_3) δ 13.6, 21.9, 31.2, 33.3, 125.6, 128.7, 128.8, 137.0; IR (neat) 2957, 2929, 1584, 1480 cm^{-1} ; Anal. Calcd for $\text{C}_{10}\text{H}_{14}\text{S}$ C, 72.23; H, 8.49. Found: C, 72.21; H, 8.50.

3ae: ^1H NMR (CDCl_3): δ 1.29 (d, $J = 6.6$ Hz, 6H), 3.32–3.42 (m, 1H), 7.21–7.31 (m, 3H), 7.37–7.42 (2H, m); ^{13}C NMR (CDCl_3) δ 23.1, 38.2, 126.6, 128.7, 131.9, 135.5; IR (neat) 2929, 1583, 1479 cm^{-1} ; Anal. Calcd for $\text{C}_9\text{H}_{12}\text{S}$ C, 71.00; H, 7.94. Found: C, 70.97; H, 7.99.

The reaction conditions and results are summarized in Table 3. A typical procedure is shown in the experimental section.

To a mixture of NiBr_2 (6.6 mg, 0.03 mmol), bpy (4.7 mg, 0.03 mmol), zinc (dust) (39.2 mg, 0.6 mmol) and DMF (0.5 mL) were added iodobenzene (**1a**) (61.2 mg, 0.3 mmol) and di-*n*-butyl disulfide (**2d**) (26.8 mg, 0.15 mmol), and the mixture was stirred at 110 °C for 48 h. After evaporation of the solvent, the residue was dissolved in Et_2O , and the resulting mixture was filtered through Celite pad. The solution was washed with H_2O and saturated sodium chloride and dried over anhydrous magnesium sulfate. Chromatography on silica gel (Hexane) gave *n*-butyl phenyl sulfide (**3ad**) (40.1 mg, 81 %).

3bd: ^1H NMR (CDCl_3) δ 0.93 (t, $J = 7.2$ Hz, 3H), 1.42–1.53 (m, 2H), 1.53–1.68 (m, 2H), 2.36 (s, 3H), 2.89 (t, $J = 7.4$ Hz, 2H), 7.06–7.26 (m, 4H); ^{13}C NMR (CDCl_3) δ 13.6, 20.3, 22.0, 31.0, 32.5, 125.2, 126.2, 127.3, 129.9, 136.3, 137.2; IR (neat) 3059, 2957, 1582, 1468 cm^{-1} ; Anal. Calcd for $\text{C}_{11}\text{H}_{16}\text{S}$ C, 73.27; H, 8.49. Found: C, 73.39; H, 8.75.

3cd: ^1H NMR (CDCl_3) δ 0.92 (t, $J = 7.3$ Hz, 3H), 1.42–1.54 (m, 2H), 1.59–1.67 (m, 2H), 2.88 (t, $J = 7.3$ Hz, 2H), 3.89 (s, 3H), 6.83 (d, $J = 8.2$ Hz, 1H), 6.91 (dd, $J = 7.6$ and 7.9 Hz, 1H), 7.15 (dd, $J = 7.9$ and 8.2 Hz, 1H), 7.24 (d, $J = 7.6$ Hz, 1H); ^{13}C NMR (CDCl_3) δ 13.6, 22.0, 30.9, 31.5, 55.7, 110.0, 120.9, 125.3, 126.6, 128.7, 157.0; IR (neat) 3059, 2957, 1582, 1468 cm^{-1} ; Anal. Calcd for $\text{C}_{11}\text{H}_{16}\text{OS}$ C, 67.30; H, 8.22. Found: C, 67.35; H, 8.18.

3dd: ^1H NMR (CDCl_3) δ 0.88 (t, $J = 7.2$ Hz, 3H), 1.36–1.45 (m, 2H), 1.47–1.60 (m, 2H), 2.73 (t, $J = 7.2$ Hz, 2H), 4.32 (br, 2H), 6.65–6.71 (m, 2H), 7.09 (dd, $J = 7.9$ and 8.2 Hz, 1H), 7.36 (d, $J = 7.6$ Hz, 1H); ^{13}C NMR (CDCl_3) δ 13.6, 21.8, 31.7, 34.5, 114.7, 118.4, 129.3, 133.5, 135.5, 148.0; IR (neat) 3450, 3358, 1605, 1478 cm^{-1} ; Anal. Calcd for $\text{C}_{10}\text{H}_{15}\text{NS}$ C, 66.25; H, 8.34; N, 7.73. Found: C, 66.12; H, 8.38; N, 7.75.

3ed: ^1H NMR (CDCl_3) δ 0.88 (t, $J = 7.2$ Hz, 3H), 1.34–1.43 (m, 2H), 1.48–1.56 (m, 2H), 2.68 (t, $J = 7.2$ Hz, 2H), 6.75 (br, 1H), 6.85 (dd, $J = 8.5$ and 7.6 Hz, 1H), 6.97 (d, $J = 8.2$ Hz, 1H), 7.25 (dd, $J = 8.2$ and 8.5 Hz, 1H), 7.45 (d, $J = 7.6$ Hz, 1H); ^{13}C NMR (CDCl_3) δ 13.5, 21.7, 31.7, 36.5, 114.6, 119.2, 120.6, 130.9, 135.9, 156.9; IR (CHCl_3) 3400, 2960, 2930, 1471 cm^{-1} ; Anal. Calcd for $\text{C}_{10}\text{H}_{14}\text{OS}$ C, 65.89; H, 7.74. Found: C, 65.89; H, 7.87.

3fd: ^1H NMR (CDCl_3) δ 0.90 (t, $J = 7.3$ Hz, 3H), 1.36–1.49 (m, 2H), 1.54–1.66 (m, 2H), 2.31 (s, 3H), 2.87 (t, $J = 7.3$ Hz, 2H), 7.08 (d, $J = 8.6$ Hz, 2H), 7.24 (d, $J = 8.6$ Hz, 2H); ^{13}C NMR (67 MHz, CDCl_3) δ 13.6, 20.9, 21.9, 31.3, 34.0, 129.3, 129.7, 133.1, 135.8; IR (neat) 2957, 2927, 1492 cm^{-1} ; Anal. Calcd for $\text{C}_{11}\text{H}_{16}\text{S}$ C, 73.27; H, 8.94. Found: C, 73.25; H, 8.97.

3gd: ^1H NMR (CDCl_3) δ 0.89 (t, $J = 7.2$ Hz, 3H), 1.37–1.45 (m, 2H), 1.51–1.59 (m, 2H), 2.81 (t, $J = 7.2$ Hz, 2H), 3.79 (s, 3H), 6.83 (d, $J = 8.6$ Hz, 2H), 7.33 (d, $J = 8.6$ Hz, 2H); ^{13}C NMR (CDCl_3) δ 13.6, 21.7, 31.4, 35.4, 55.2, 114.4, 127.0, 132.8, 158.7; IR (neat) 2957, 2930, 1593, 1493 cm^{-1} ; Anal. Calcd for $\text{C}_{11}\text{H}_{16}\text{OS}$ C, 67.30; H, 8.22. Found: C, 67.31; H, 8.20.

3hd: ^1H NMR (CDCl_3) δ 0.92 (t, $J = 7.3$ Hz, 3H), 1.39–1.47 (m, 2H), 1.54–1.64 (m, 2H), 2.89 (t, $J = 7.3$ Hz, 2H), 7.17 (d, $J = 8.6$ Hz, 2H), 7.38 (d, $J = 8.6$ Hz, 2H); ^{13}C NMR (CDCl_3) δ 13.5, 21.9, 31.0, 33.3, 119.3, 128.7, 130.8, 131.8; IR (neat) 2957, 2928, 1473 cm^{-1} ; Anal. Calcd for $\text{C}_{10}\text{H}_{13}\text{SBr}$ C, 48.99; H, 5.34. Found: C, 48.87; H, 5.37.

3id: ^1H NMR (CDCl_3) δ 0.91 (t, $J = 7.2$ Hz, 3H), 1.39–1.47 (m, 2H), 1.54–1.64 (m, 2H), 2.88 (t, $J = 7.2$ Hz, 2H), 7.23 (s, 4H); ^{13}C NMR (CDCl_3) δ 13.5, 21.8, 31.1, 33.5, 128.9, 130.2, 131.5, 135.5; IR (neat) 2957, 2928, 1476 cm^{-1} ; Anal. Calcd for $\text{C}_{10}\text{H}_{13}\text{OSCl}$ C, 59.84; H, 6.53. Found: C, 59.97; H, 6.71.

3jd: ^1H NMR (CDCl_3) δ 0.94 (t, $J = 7.2$ Hz, 3H), 1.43–1.54 (m, 2H), 1.61–1.70 (m, 2H), 2.97 (t, $J = 7.4$ Hz, 2H), 7.34 (d, $J = 8.2$ Hz, 2H), 7.49 (d, $J = 8.2$ Hz, 2H); ^{13}C NMR (CDCl_3) δ 13.5, 21.9, 30.8, 32.1, 125.4, 125.5, 125.6, 127.1, 142.8; IR (neat) 2961, 1607 cm^{-1} ; Anal. Calcd for $\text{C}_{11}\text{H}_{13}\text{F}_3\text{S}$ C, 56.39; H, 5.59. Found: C, 56.66; H, 5.68.

3kd: ^1H NMR (CDCl_3) δ 0.88 (t, $J = 7.2$ Hz, 3H), 1.38–1.46 (m, 2H), 1.49–1.60 (m, 2H), 2.76 (t, $J = 7.2$ Hz, 2H), 3.67 (br, 2H), 6.61 (d, $J = 8.6$ Hz, 2H), 7.23 (d, $J = 8.6$ Hz, 2H); ^{13}C NMR (CDCl_3) δ 13.6, 21.8, 31.5, 36.0, 115.5, 123.9, 133.6, 145.7; IR (neat) 3019, 2975, 1593, 1494 cm^{-1} ; Anal. Calcd for $\text{C}_{10}\text{H}_{15}\text{NS}$ C, 77.10; H, 8.10; N, 14.80. Found: C, 77.10; H, 8.10; N, 14.80.

66.25; H, 8.34. Found: C, 66.08; H, 8.49.

3ld: ^1H NMR (CDCl_3) δ 0.89 (t, $J = 7.3$ Hz, 3H), 1.36–1.44 (m, 2H), 1.53–1.59 (m, 2H), 2.81 (t, $J = 7.3$ Hz, 2H), 5.08 (br, 1H), 6.76 (d, $J = 8.6$ Hz, 2H), 7.28 (d, $J = 8.6$ Hz, 2H); ^{13}C NMR (CDCl_3) δ 13.6, 21.7, 31.3, 35.5, 115.9, 133.1, 138.4, 154.6; IR (neat) 3019, 2961, 1494 cm^{-1} ; Anal. Calcd for $\text{C}_{10}\text{H}_{14}\text{OS}$ C, 65.89; H, 7.74. Found: C, 65.87; H, 7.78.

3md: ^1H NMR (CDCl_3) δ 0.95 (t, $J = 7.3$ Hz, 3H), 1.44–1.53 (m, 2H), 1.67–1.72 (m, 2H), 3.00 (t, $J = 7.3$ Hz, 2H), 7.31 (d, $J = 8.2$ Hz, 2H), 7.99 (d, $J = 8.2$ Hz, 2H); ^{13}C NMR (CDCl_3) δ 13.5, 22.0, 30.7, 31.6, 126.1, 130.5, 137.9, 145.9, 174.8; IR (CHCl_3) 3019, 2962, 1689, 1593 cm^{-1} ; Anal. Calcd for $\text{C}_{11}\text{H}_{14}\text{O}_2\text{S}$ C, 62.83; H, 6.71. Found: C, 62.70; H, 6.78.

3nd: ^1H NMR (CDCl_3) δ 0.95 (t, $J = 7.2$ Hz, 3H), 1.43–1.51 (m, 2H), 1.63–1.70 (m, 2H), 2.98 (t, $J = 7.2$ Hz, 2H), 3.89 (s, 3H), 7.28 (d, $J = 8.5$ Hz, 2H), 7.92 (d, $J = 8.5$ Hz, 2H); ^{13}C NMR (CDCl_3) δ 13.6, 22.0, 30.8, 31.8, 52.0, 126.2, 126.5, 129.8, 144.5, 166.8; IR (neat) 2955, 2930, 1720, 1595 cm^{-1} ; Anal. Calcd for $\text{C}_{12}\text{H}_{16}\text{O}_2\text{S}$ C, 64.25; H, 7.19. Found: C, 64.23; H, 7.20.

3od: ^1H NMR (CDCl_3) δ 0.95 (t, $J = 7.3$ Hz, 3H), 1.43–1.52 (m, 2H), 1.63–1.72 (m, 2H), 2.56 (s, 3H), 2.99 (t, $J = 7.3$ Hz, 2H), 7.28 (d, $J = 8.4$ Hz, 2H), 7.84 (d, $J = 8.4$ Hz, 2H); ^{13}C NMR (CDCl_3) δ 13.5, 21.9, 26.3, 30.7, 31.6, 126.2, 128.6, 133.6, 144.9, 197.1; IR (neat) 2958, 2929, 1679, 1589 cm^{-1} ; Anal. Calcd for $\text{C}_{12}\text{H}_{16}\text{OS}$ C, 69.19; H, 7.74. Found: C, 69.00; H, 7.83.

3pd: ^1H NMR (CDCl_3) δ 0.94 (t, $J = 7.4$ Hz, 3H), 1.39–1.52 (m, 2H), 1.62–1.73 (m, 2H), 2.79 (s, 3H),

2.99 (t, $J = 7.4$ Hz, 2H), 6.45 (d, $J = 15.6$ Hz, 2H), 6.72 (d, $J = 15.6$ Hz, 2H), 7.15–7.31 (m, 5H); ^{13}C NMR (CDCl_3) δ 13.6, 21.9, 31.5, 32.3, 125.3, 125.4, 126.6, 126.7, 128.6, 137.2; IR (neat) 2956, 2928, 1597, 1569 cm^{-1} ; Anal. Calcd for $\text{C}_{12}\text{H}_{16}\text{S}$ C, 74.94; H, 8.39. Found: C, 74.89; H, 8.42.

The reaction of 2-iodotoluene (**1b**) with $(\text{PhS})_2\text{Ni}(\text{II})(\text{bpy})_2$

To a mixture of $(\text{PhS})_2\text{Ni}(\text{II})(\text{bpy})_2$ (88.4 mg, 0.15 mmol), zinc (dust) (39.2 mg, 0.6 mmol) and DMF (0.5 mL) were added 2-iodotoluene (**1b**) (65.4 mg, 0.3 mmol), and the mixture was stirred at 110 °C for 18 h. After evaporation of the solvent, the residue was dissolved in Et_2O , and the resulting mixture was filtered through Celite pad. The solution was washed with H_2O and saturated sodium chloride and dried over anhydrous magnesium sulfate. Chromatography on silica gel (Hexane) gave Phenyl 2-tolyl sulfide (**3ba**) (55.3 mg, 92 %). ^1H NMR (270 MHz, CDCl_3) δ 2.37 (s, 3H), 7.13–7.33 (m, 8H), 7.49 (d, $J = 8.5$ Hz, 1H); ^{13}C NMR (67 MHz, CDCl_3) δ 20.5, 126.3, 126.7, 127.5, 127.9, 129.0, 129.1, 129.2, 129.6, 130.5, 133.0; IR (neat) 3059, 1582, 1476 cm^{-1} ; Anal. Calcd for $\text{C}_{13}\text{H}_{12}\text{S}$ C, 77.95; H, 6.04. Found: C, 77.98; H, 6.11.