Supplementary Information

“Alkynes as Synthetic Equivalents to Stabilized Wittig Reagents: Intra- and Intermolecular Carbonyl Olefinations Catalyzed by Ag(I), BF₃ and HBF₄”

Jong Uk Rhee and Michael J. Krische*

University of Texas at Austin

Department of Chemistry and Biochemistry

Austin, TX 78712

USA

Table of Contents

1. General Procedure.....	S2
2. Preparation of Substrate.....	S3
3. Representative intramolecular alkyne-carbonyl metathesis.....	S23
4. Representative intermolecular alkyne-carbonyl metathesis.....	S43
5. References.....	S54

1. Experimental Section

1. 1. General Procedure: All reactions were run under an atmosphere of argon, unless otherwise indicated. Anhydrous solvents were transferred by an oven-dried syringe, and flasks were flame-dried and cooled under a stream of nitrogen. All solvents were purchased from Fisher Co. and fresh distilled from calcium hydride (dichloromethane and dichloroethane) or from calcium hydride (THF and diethyl ether). Other chemicals were purchased from Aldrich Chemical Co. or Across Co. and used without further purification, unless otherwise noted.

Analytical thin-layer chromatography (TLC) was carried out using 0.2 mm commercial silica gel plates (EM Science precoated 60 F₂₅₄). Proton NMR spectra were recorded on Varian Gemini 300 (300 MHz), Varian Mercury 400 (400 MHz), and Varian Inova 500 (500 MHz). Chemical shifts are reported in delta (δ) units, parts per million (ppm) down field from TMS, and Hertz (Hz) is used for the coupling constants. ¹³C NMR spectra were recorded with Varian Gemini 300 (75 MHz), Varian Mercury 400 (100 MHz), and Varian Inova 500 (125 MHz) spectrometer. Chemical shifts are reported in delta (δ) units, parts per million (ppm) relative to the central line of CDCl₃ (δ = 77.00 ppm) unless otherwise mentioned. ¹³C NMR spectra were routinely run with broad band decoupling. The multiplicities are expressed like following: s = singlet, d = doublet, t = triplet, q = quartet. Infrared spectra were obtained on Perkin-Elmer 1600 Infrared Spectrometer and are reported in reciprocal centimeters (cm⁻¹). The relative intensities of IR spectra are reported as follows: br = broad, s = strong, m = medium, w = weak. High resolution mass spectra were measured by the University of Texas at Austin Instrumental Service. All melting points were determined by Thomas-Hoover capillary melting point apparatus and are uncorrected.

2. Preparation of Substrate

Preparation of *N*-But-2-ynyl-4-methyl-*N*-(2-oxo-ethyl)-benzenesulfonamide (1a)

A solution of *N*-allyl-*N*-but-2-ynyl-4-methyl-benzenesulfonamide (9.30 mmol, 2.45 g) in dichloromethane (125 mL) was purged with O₃ at -78 °C until no more starting material was observed by TLC analysis (20 min.). The excess O₃ in the solution was removed by purging of oxygen and argon at -78 °C for 5 min, respectively. To the mixture was added one equivalent of triphenyl phosphine (9.3 mmol, 2.44 g) at -78 °C, and then stirred it at -78 °C to rt under argon atmosphere over night. After all solvent was removed on a rotary evaporator, the crude mixture was purified by column chromatography eluting with hexane:EtOAc = 3:1. The isolated product was obtained as pale brown oil (1.81 g) and the isolated % yield is 73%.

1a

Pale brown oil. Column; hexane:EtOAc = 3:1. R_f = 29 (hexane:EtOAc = 2:1). ¹H NMR (CDCl₃, 300 MHz): δ 9.65 (t, J = 1.2 Hz, 1H), 7.70 (dt, J = 8.4, 1.8 Hz, 2H), 7.33 (d, J = 8.7 Hz, 2H), 4.08 (q, J = 2.4 Hz, 2H), 3.90 (d, J = 1.5 Hz, 2H), 2.42 (s, 3H), 1.59 (t, J = 2.1 Hz, 3H). ¹³C NMR (CDCl₃, 75 MHz): δ 197.6, 143.9, 134.6, 129.4, 127.4, 82.8, 71.0, 55.7, 39.1, 21.2, 2.9. IR (NaCl, neat): 3456 *Br s*, 3062*m*, 2923*m*, 2846*m*, 2728*w*, 1813*w*, 1732*s*, 1597*s*, 1572*w*, 1490*s*, 1442*s*, 1403*s*, 1349*s*, 1257*m*, 1162*s*, 1108*s*, 1008*m*, 928*m*, 871*m*, 815*m*, 758*m*, 709*m*, 692*m*, 660*m* cm⁻¹. HRMS (CI) m/z calcd for C₁₃H₁₆N₁O₅S₁ [M^+ + 1] 266.0851, found 266.0854.

Preparation of *N*-(3-Cyclopropyl-prop-2-ynyl)-4-methyl-*N*-(2-oxo-ethyl)-benzene-sulfonamide (2a)

A solution of *N*-allyl-*N*-(3-cyclopropyl-prop-2-ynyl)-4-methyl-benzenesulfonamide (5.87 mmol, 1.70 g) in dichloromethane (80 mL) was purged with O₃ at -78 °C until no more starting material was observed by TLC analysis (25 min.). The excess O₃ in the solution was removed by purging of oxygen and argon at -78 °C for 5 min, respectively. To the mixture was added one equivalent of triphenyl phosphine (5.87 mmol, 1.54 g) at -78 °C, and then stirred it at -78 °C to rt under argon atmosphere over night. After all solvent was removed on a rotary evaporator, the crude mixture was purified by column chromatography eluting with hexane:EtOAc = 3:1. The isolated product was obtained as colorless oil (1.26 g) and the isolated % yield is 74%.

2a

Colorless oil. Column; hexane:EtOAc = 3:1. R_f = 0.34 (hexane:EtOAc = 2:1). ¹H NMR (CDCl₃, 400 MHz): δ 9.66 (dd, J = 2.4, 1.6 Hz, 1H), 7.69 (d, J = 8.8 Hz, 2H), 7.35 (d, J = 8.4 Hz, 2H), 4.09 (d, J = 2.0 Hz, 2H), 3.87 (s, 2H), 2.45 (s, 3H), 0.98-1.05 (m, 1H), 0.63-0.68 (m, 2H), 0.38-0.42 (m, 2H). ¹³C NMR (CDCl₃, 100 MHz): δ 197.8, 144.1, 134.8, 129.7, 127.6, 90.6, 67.1, 55.9, 39.4, 21.5, 7.8, -1.0. IR (NaCl, neat): 3458 *Br s*, 3010*m*, 2938*m*, 2924*m*, 2863*m*, 2725*w*, 1921*w*, 1733*s*, 1597*m*, 1494*m*, 1442*m*, 1402*m*, 1348*s*, 1307*s*, 1292*m*, 1267*w*, 1162*s*, 1108*s*, 1090*s*, 1036*s*, 998*s*, 930*s*, 878*s*, 815*s*, 737*s*, 708*s*, 658*s*, 545*s* cm⁻¹. HRMS (CI) m/z calcd for C₁₅H₁₈N₁O₃S₁ [M^+ + 1] 292.1007, found 292.0997.

Preparation of 4-Methyl-N-(2-oxo-ethyl)-N-(3-phenyl-prop-2-ynyl)-benzenesulfonamide (3a)

A solution of *N*-allyl-4-methyl-*N*-(3-phenyl-prop-2-ynyl)-benzenesulfonamide (56.37 mmol, 18.34 g) in dichloromethane (400 mL) was purged with O₃ at -78 °C until no more starting material was observed by TLC analysis (10 min.). The excess O₃ in the solution was removed by purging of oxygen and argon at -78 °C for 5 min, respectively. To the mixture was added one equivalent of triphenyl phosphine (56.37 mmol, 14.78 g) at -78 °C, and then stirred it at -78 °C to rt under argon atmosphere over night. After all solvent was removed on a rotary evaporator, the crude mixture was purified by column chromatography eluting with hexane:EtOAc = 3:1. The isolated product was obtained as pale yellow solid (16.08 g) and the isolated % yield is 87%.

3a

Pale yellow solid. Column; hexane:EtOAc = 3:1. R_f = 0.18 (hexane:EtOAc = 3:1). Mp: 80-82 °C. ¹H NMR (CDCl₃, 300 MHz): δ 9.70 (t, J = 1.5 Hz, 1H), 7.74 (d, J = 8.4 Hz, 2H), 7.21-7.31 (m, 6H), 7.12 (dd, J = 7.5, 1.2 Hz, 2H), 4.38 (s, 2H), 3.92 (s, 2H), 2.35 (s, 3H). ¹³C NMR (CDCl₃, 75 MHz): δ 197.4, 144.2, 134.7, 131.4, 129.8, 128.7, 128.1, 127.6, 121.5, 86.5, 80.9, 56.0, 39.6, 21.4. IR (NaCl, neat): 3479 *Br s*, 3031*w*, 2921*s*, 2856*m*, 2732*w*, 1923*s*, 1732*s*, 1598*s*, 1494*s*, 1442*s*, 1402*s*, 1347*s*, 1257*m*, 1213*w*, 1160*s*, 1090*s*, 1001*s*, 929*s*, 872*m*, 816*s*, 751*s*, 710*s*, 681*s*, 658*s* cm⁻¹. HRMS (CI) m/z calcd for C₁₈H₁₈N₁O₃S₁ [M^+ + 1] 328.1007, found 328.1008.

Preparation of 4-Methyl-*N*-(2-oxo-ethyl)-*N*-(4-phenyl-but-3-ynyl)-benzenesulfonamide (4a)

A solution of *N*-allyl-4-methyl-*N*-(4-phenyl-but-3-ynyl)-benzenesulfonamide (3.06 mmol, 1.04 g) in dichloromethane (30 mL) was purged with O₃ at -78 °C until no more starting material was observed by TLC analysis (25 min.). The excess O₃ in the solution was removed by purging of oxygen and argon at -78 °C for 5 min, respectively. To the mixture was added one equivalent of triphenyl phosphine (3.06 mmol, 0.803 g) at -78 °C, and then stirred it at -78 °C to rt under argon atmosphere over night. After all solvent was removed on a rotary evaporator, the crude mixture was purified by column chromatography eluting with hexane:EtOAc = 4:1 to 3:1. The isolated product was obtained as yellow oil (329 mg) and the isolated % yield is 32%.

4a

Yellow oil. Column; hexane:EtOAc = 4:1 to 3:1. R_f = 0.25 (hexane:EtOAc = 3:1). ¹H NMR (CDCl₃, 400 MHz): δ 9.67 (s, 1H), 7.71 (d, J = 8.0 Hz, 2H), 7.26-7.35 (m, 7H), 4.01 (s, 2H), 3.43 (t, J = 6.8 Hz, 2H), 2.70 (t, J = 6.8 Hz, 2H), 2.39 (s, 3H). ¹³C NMR (CDCl₃, 100 MHz): δ 198.9, 143.9, 135.5, 131.3, 129.8, 128.1, 128.0, 127.1, 122.8, 86.1, 82.9, 57.9, 48.6, 21.3, 20.4. IR (NaCl, neat): 3480 *Br s*, 3029_w, 2920_s, 2859_m, 2731_w, 1920_s, 1730_s, 1599_s, 1490_s, 1441_s, 1401_s, 1349_s, 1257_m, 1214_w, 1158_s, 1093_s, 1004_s, 931_s, 870_m, 815_s, 750_s, 711_s, 680_s, 659_s cm⁻¹. HRMS (CI) m/z calcd for C₁₉H₂₀N₁O₃S₁ [M^+ + 1] 342.1164, found 342.1163.

Preparation of (2R)-5-Oxo-1-(3-phenyl-prop-2-ynyl)-pyrrolidine-2-carbaldehyde (5a)

To a stirred solution of oxalyl chloride (7.92 mmol, 1.01 g, 0.69 mL) in dichloromethane (16 mL) was added dropwise DMSO (15.69 mL, 1.23 g, 1.11 mL, dried from calcium hydride and stored over 4 Å MS) at -78 °C. After stirring at -78 °C for 15 min, a solution of (5R)-5-hydroxymethyl-1-(3-phenyl-prop-2-ynyl)-pyrrolidin-2-one (5.23 mmol, 1.21 g) in dichloromethane (8 mL) was added dropwise at -78 °C. The resulting mixture was stirred at -78 °C for 15 min, and then added triethylamine (20.92 mmol, 2.12 g, 2.92 mL) at -78 °C, and then the temperature increased to 0 °C and continued stirring for 1.5 h. After the reaction was quenched with saturated sodium bicarbonate solution, extracted it with diethyl ether. Combined organic was washed with H₂O and brine solution, dried over MgSO₄, filtered, and concentrated on a rotary evaporator to give crude mixture. The crude mixture was purified by column chromatography eluting with hexane:EtOAc = 1:3. Yellow oil was obtained as the product (879 mg) and the isolated yield is 74%.

5a

Yellow oil. Column; hexane:EtOAc = 1:3. R_f = 0.25 (hexane:EtOAc = 1:5). ¹H NMR (CDCl₃, 400 MHz): δ 9.74 (d, J = 2.4 Hz, 1H), 7.38-7.41 (m, 2H), 7.27-7.34 (m, 3H), 4.62 (d, J = 18.0 Hz, 1H), 4.42 (ddd, J = 9.6, 4.4, 2.0 Hz, 1H), 4.29 (d, J = 17.6 Hz, 1H), 2.49 (dd, J = 8.8, 2.8 Hz, 1H), 2.47 (d, J = 8.4 Hz, 1H), 2.29-2.41 (m, 1H), 2.09 (dddd, J = 10.0, 5.6, 4.4, 4.0 Hz, 1H). ¹³C NMR (CDCl₃, 100 MHz): δ 198.5, 174.5, 131.7, 128.7, 128.3, 122.0, 85.3, 82.4, 64.8, 32.5, 29.2, 19.3. IR (NaCl, neat): 3359 *Br s*, 2929_w, 1731_m, 1672_s, 1598_w, 1490_m, 1442_m, 1417_m, 1353_m, 1190_w, 1157_w, 1070_m, 968_w, 825_w, 758_s, 692_s cm⁻¹. HRMS (CI) m/z calcd for C₁₄H₁₄N₁O₂ [M^+ + 1] 228.1025, found 228.1026.

Preparation of (3-Phenyl-prop-2-ynyloxy)-acetaldehyde (6a)

A solution of (3-allyloxy-prop-1-ynyl)-benzene (22.3 mmol, 3.84 g) in dichloromethane (150 mL) was purged with O₃ at -78 °C until no more starting material was observed by TLC analysis (25 min.). The excess O₃ in the solution was removed by purging of oxygen and argon at -78 °C for 5 min, respectively. To the mixture was added one equivalent of triphenyl phosphine (22.3 mmol, 5.85 g) at -78 °C, and then stirred it at -78 °C to rt under argon atmosphere over night. After all solvent was removed on a rotary evaporator, the crude mixture was purified by column chromatography eluting with hexane:EtOAc = 5:1 to 3:1. Pale yellow oil was obtained as the product (3.594 g) and the isolated yield is 93%. The ¹H and ¹³C NMR spectra were agreed with the reported value.¹

6a

Pale yellow oil. Column; hexane:EtOAc = 5:1 to 3:1. R_f = 0.16 (hexane:EtOAc = 3:1). ¹H NMR (CDCl₃, 400 MHz): δ 9.73 (Br s, 1H), 7.42-7.43 (m, 2H), 7.26-7.31 (m, 3H), 4.48 (t, J = 3.2 Hz, 2H), 4.21 (Br s 2H). ¹³C NMR (CDCl₃, 100 MHz): δ 199.7, 131.5, 128.5, 128.1, 121.8, 87.3, 83.6, 74.5, 59.2.

Preparation of 2-(2-Oxo-ethyl)-2-(3-phenyl-prop-2-ynyl)-malonic acid diethyl ester (7a)²

A solution of 2-allyl-2-(3-phenyl-prop-2-ynyl)-malonic acid diethyl ester (70.0 mmol, 22.0 g) in dichloromethane (500 mL) was purged with O₃ at -78 °C until no more starting material was observed by TLC analysis (1 h 50 min.). The excess O₃ in the solution was removed by purging of oxygen and argon at -78 °C for 5 min, respectively. To the mixture was added one equivalent of triphenyl phosphine (70.0 mmol, 18.36 g) at -78 °C, and then stirred it at -78 °C to rt under argon atmosphere over night. After all solvent was removed on a rotary evaporator, the crude mixture was purified by column chromatography eluting with hexane:EtOAc = 7:1. Yellow oil was obtained as the product (15.08 g) and the isolated yield is 68%. The ¹H and ¹³C NMR spectra were agreed with the reported value.²

7a

Pale yellow oil. Column; hexane:EtOAc = 9:1. *R_f* = 0.22 (hexane:EtOAc = 7:1). ¹H NMR (CDCl₃, 300 MHz): δ 9.80 (s, 1H), 7.34-7.37 (m, 2H), 7.26-7.30 (m, 3H), 4.25 (qd, *J* = 7.2, 1.8 Hz, 4H), 3.28 (d, *J* = 1.2 Hz, 2H), 3.19 (s, 2H), 1.27 (dd, *J* = 7.2 Hz, 6H). ¹³C NMR (CDCl₃, 75 MHz): δ 198.8, 168.9, 131.6, 128.2, 128.2, 122.8, 84.1, 83.9, 62.2, 54.4, 46.2, 24.8, 13.9. MS (CI) *m/z* calcd for C₁₈H₂₁O₅ [*M*⁺ + 1] 317, found 317.

Preparation of 2-(3-Phenyl-prop-2-ynyloxy)-benzaldehyde (8a)³

To a solution of K₂CO₃ (10.5 mmol, 1.45 g) in DMF (10 mL) was added salicylaldehyde (10.0 mmol, 1.22 g, 1.07 mL) at rt. The reaction mixture was stirred at rt for 1 h, and followed by the addition of (3-bromo-prop-1-ynyl)-benzene (10.5 mmol, 2.05 g) at rt. The resulting mixture was further stirred at rt for 1 h, and diluted with diethyl ether. The organic was washed with H₂O and brine solution, dried over MgSO₄, filtered, and concentrated on a rotary evaporator to give crude mixture. The desired product was isolated by column chromatography eluting with hexane:EtOAc = 9:1. White solid was obtained and the isolated yield was 92% (2.04 g). The ¹H and ¹³C NMR spectra were agreed with the reported value.³

8a

Whit sold. Column; hexane:EtOAc = 9:1. *R*_f = 0.39 (hexane:EtOAc = 6:1). ¹H NMR (CDCl₃, 300 MHz): δ 10.53 (d, *J* = 0.3 Hz, 1H), 7.88 (dd, *J* = 8.1, 1.8 Hz, 1H), 7.58 (app td, *J* = 7.5, 1.8 Hz, 1H), 7.41-7.44 (m, 2H), 7.24-7.35 (m, 3H), 7.21 (d, *J* = 8.4 Hz, 1H), 7.09 (t, *J* = 7.2 Hz, 1H), 5.05 (s, 2H). ¹³C NMR (CDCl₃, 75 MHz): δ 189.7, 160.1, 135.7, 131.8, 128.9, 128.5, 128.3, 125.5, 121.9, 121.5, 88.1, 82.9, 57.3

Preparation of 2-But-2-ynyloxy-benzaldehyde(9a)⁴

To a solution of K_2CO_3 (11 mmol, 1.52 g) in DMF (10 mL) was added salicylaldehyde (10.0 mmol, 1.22 g, 1.07 mL) at rt. The reaction mixture was stirred at rt for 1 h, and followed by the addition of 1-bromo-but-3-ynr (10.5 mmol, 1.40 g, 0.95 mL) in DMF (5 mL) at rt. The resulting mixture was further stirred at rt for 70 h, and diluted with diethyl ether. The organic was washed with H_2O and brine solution, dried over MgSO_4 , filtered, and concentrated on a rotary evaporator to give yellow oil (1.70 g, 98% yield). The crude product was analytically pure (>95% based on ^1H NMR spectroscopy) and used without further purification.

9a

Yellow liquid. Column; hexane:EtOAc = 5:1. R_f = 0.50 (hexane:EtOAc = 5:1). ^1H NMR (CDCl_3 , 300 MHz): δ 10.48 (s, 1H), 7.84 (dd, J = 7.5, 1.8 Hz, 1H), 7.55 (app td, J = 78.7, 1.8 Hz, 1H), 7.11 (d, J = 8.4 Hz, 1H), 7.05 (dd, J = 7.5 Hz, 1H), 4.78 (q, J = 2.1 Hz, 2H), 1.85 (t, J = 2.4 Hz, 3H). ^{13}C NMR (CDCl_3 , 75 MHz): δ 189.6, 160.0, 135.6, 128.2, 125.3, 121.1, 113.2, 84.6, 73.2, 56.9, 3.5.

Preparation of 2-Methyl-2-(4-phenyl-but-3-ynyl)-cyclopentane-1,3-dione (10a)⁵

To a mixture of 2-but-3-ynyl-2-methyl-cyclopentane-1,3-dione (9.50 mmol, 1.56 g) and iodobenzene (10.45 mmol, 2.13 g, 1.16 mL) were added Pd(PPh₃)₂Cl₂ (19 μmol, 13.3 mg), CuI (38 μmol, 7.2 mg), and piperidine (19 mmol, 1.62 g, 1.88 mL) at rt. After the mixture was stirred at 40 °C for 21 h, all solvent was removed on a rotary evaporator. The crude mixture was subject to purification by column chromatography eluting with hexane:EtOAc = 8:1. Brown oil was obtained as the product and the isolated yield is 69% (1.565 g), and the ¹H and ¹³C NMR spectra were agreed with the reported value.⁵

10a

Brown oil. Column; hexane:EtOAc = 8:1. R_f = 0.23 (hexane:EtOAc = 7:1). ¹H NMR (CDCl₃, 400 MHz): δ 7.31-7.36 (m, 2H), 7.26-7.30 (m, 3H), 2.67-2.82 (m, 4H), 2.41 (t, J = 7.2 Hz, 2H), 2.09 (t, J = 6.8 Hz, 2H), 1.15 (s, 1H). ¹³C NMR (CDCl₃, 100 MHz): δ 216.1, 131.2, 128.3, 128.0, 123.0, 88.8, 83.1, 55.6, 34.9, 33.2, 21.6, 15.3. IR (NaCl, neat): 3056_w, 2964_w, 2926_w, 2868_w, 1763_m, 1720_s, 1637_w, 1597_w, 1571_w, 1490_m, 1451_m, 1419_m, 1373_m, 1341_w, 1284_m, 1154_w, 1071_m, 993_w, 915_w, 758_s, 732_m, 693_s cm⁻¹. HRMS (CI) m/z calcd for C₁₆H₁₇O₂ [M^+ + 1] 241.1229, found 241.1223.

3. Representative intramolecular alkyne-carbonyl metathesis

To a solution of *N*-But-2-ynyl-4-methyl-*N*-(2-oxo-ethyl)-benzenesulfonamide (**1a** , 0.4 mmol, 106 mg) in dichloroethane (4 mL) was added a solution of AgSbF₆ (0.5 M in THF, 10 mol%, 40 μ mol, 80 μ L) at rt. The mixture was refluxed under argon atmosphere for 12 h, and all starting material was consumed by TLC analysis. After all solvent was removed on a rotary evaporator, the crude mixture was purified by column chromatography eluting with hexane:EtOAc = 3:1. The product was isolated as pale yellow solid (64 mg) and the isolated yield is 60%.

1-[1-(Toluene-4-sulfonyl)-2,5-dihydro-1*H*-pyrrol-3-yl]-ethanone (1b)

1b

Pale yellow solid. Column; hexane:EtOAc = 3:1. R_f = 0.29 (hexane:EtOAc = 3:1). Mp: 156-158 °C. ¹H NMR (CDCl₃, 300 MHz): δ 7.72 (d, J = 8.1 Hz, 2H), 7.34 (d, J = 8.1 Hz, 2H), 6.52 (app t, J = 1.8 Hz, 1H), 4.34-4.37 (m, 2H), 4.23-4.26 (m, 2H), 2.43 (s, 3H), 2.27 (s, 3H). ¹³C NMR (CDCl₃, 75 MHz): δ 193.7, 143.8, 140.3, 135.6, 133.5, 129.8, 127.4, 55.7, 53.2, 26.2, 21.4. IR (NaCl, neat): 3072 w , 2918 w , 2855 w , 1671 s , 1636 m , 1596 m , 1457 m , 1384 m , 1339 s , 1304 m , 1282 m , 1259 m , 1186 w , 1161 s , 1098 m , 1067 m , 1016 w , 997 m , 948 w , 834 m , 818 m , 710 m , 669 s , 630 s , 590 s , 549 s cm⁻¹. HRMS (CI) m/z calcd for C₁₃H₁₆N₁O₃S₁ [M^+ + 1] 266.0851, found 266.0862.

Cyclopropyl-[1-(toluene-4-sulfonyl)-2,5-dihydro-1H-pyrrol-3-yl]-methanone (2b)

2b

White solid. Column; hexane:EtOAc = 4:1. R_f = 0.19 (hexane:EtOAc = 3:1). Mp: 130-132 °C. ^1H NMR (CDCl_3 , 300 MHz): δ 7.74 (d, J = 8.1 Hz, 2H), 7.33 (t, J = 7.8 Hz, 2H), 6.61 (app t, J = 2.1 Hz, 1H), 4.36-4.41 (m, 2H), 4.26-4.30 (m, 2H), 2.43 (s, 3H), 2.21 (app tt, J = 4.5, 3.3 Hz, 1H), 1.07 (app td, J = 7.5, 4.5 Hz, 2H), 0.92 (app td, J = 7.5, 3.6 Hz, 2H). ^{13}C NMR (CDCl_3 , 75 MHz): δ 195.7, 143.8, 140.6, 133.2, 133.7, 129.9, 127.5, 55.9, 53.6, 21.5, 17.4, 11.5. IR (NaCl, neat): 3087 w , 2999 w , 2924 w , 2854 w , 1657 s , 1628 w , 1597 w , 1447 w , 1412 s , 1346 s cm^{-1} . HRMS (CI) m/z calcd for $\text{C}_{15}\text{H}_{18}\text{N}_1\text{O}_3\text{S}_1$ [$\text{M}^+ + 1$] 292.1007, found 292.1005.

Phenyl-[1-(toluene-4-sulfonyl)-2,5-dihydro-1H-pyrrol-3-yl]-methanone (3b)

3b

Whit solid. Column; hexane:EtOAc = 4:1. R_f = 0.47 (hexane:EtOAc = 2:1). Mp: 134-136 °C. ^1H NMR (CDCl_3 , 300 MHz): δ 7.76 (d, J = 7.8 Hz, 2H), 7.66 (d, J = 7.8 Hz, 2H), 7.56 (t, J = 7.8 Hz, 1H), 7.42 (t, J = 7.8 Hz, 2H), 7.35 (d, J = 7.8 Hz, 2H), 6.36 (s, 1H), 4.40-4.44 (m, 4H), 2.43 (s, 3H). ^{13}C NMR (CDCl_3 , 75 MHz): δ 190.5, 143.85, 138.7, 137.4, 137.0, 133.5, 132.7, 129.9, 128.6, 128.5, 127.4, 55.6, 54.2, 21.4. IR (NaCl, neat): 2923w, 2853w, 1633s, 1595m, 1575m, 1494w, 1447m, 1367w, 1341s, 1306m, 1290m, 1249w, 1188m, 1163s, 1112s, 1079w, 1018w, 989m, 938w, 924w, 837w, 814m, 780w, 700m, 675s cm^{-1} . HRMS (CI) m/z calcd for $\text{C}_{18}\text{H}_{18}\text{N}_1\text{O}_3\text{S}_1$ [$\text{M}^+ + 1$] 328.1007, found 328.1013.

Phenyl-[1-(toluene-4-sulfonyl)-1,2,3,6-tetrahydro-pyridin-4-yl]-methanone (4b)

4b

White solid. Column; hexane:EtOAc = 4:1. R_f = 0.17 (hexane:EtOAc = 4:1). Mp: 143-145 °C. ^1H NMR (CDCl_3 , 300 MHz): δ 7.70 (d, J = 8.1 Hz, 2H), 7.58-7.61 (m, 2H), 7.53 (tt, J = 7.2, 1.5 Hz, 1H), 7.41 (t, J = 7.2 Hz, 2H), 7.35 (d, J = 7.8 Hz, 2H), 6.40 (app tt, J = 3.0, 1.8 Hz, 1H), 3.83 (app q, J = 3.0 Hz, 2H), 3.29 (t, J = 5.7 Hz, 2H), 2.62-2.66 (m, 2H), 2.43 (s, 3H). ^{13}C NMR (CDCl_3 , 75 MHz): δ 195.7, 143.9, 137.3, 136.4, 136.3, 132.9, 132.0, 129.7, 129.0, 128.2, 127.6, 44.8, 42.6, 24.3, 21.5. IR (NaCl, neat): 3049 w , 2960 w , 2918 w , 2866 w , 2822 w , 1642 s , 1594 s , 1577 m , 1492 m , 1448 m , 1432 s , 1388 m , 1380 m , 1341 s , 1309 s , 1285 m , 1202 w , 1163 s , 1121 m , 1094 m , 1052 w , 1019 w , 985 w , 970 m , 942 m , 879 w , 840 w , 817 m , 800 w , 767 m , 728 m , 702 s , 694 s , 671 s , 647 s , 603 m , 567 m , 549 s cm^{-1} . HRMS (CI) m/z calcd for $\text{C}_{19}\text{H}_{20}\text{N}_1\text{O}_3\text{S}_1$ [$\text{M}^+ + 1$] 342.1164, found 342.1155.

(7a*R*)-6-Benzoyl-1,2,5,7a-tetrahydro-pyrrolizin-3-one (5b)

5b

Yellow oil. Column; hexane:EtOAc = 1:4. R_f = 0.22 (hexane:EtOAc = 1:5). ^1H NMR (CDCl_3 , 300 MHz): δ 7.76 (dt, J = 7.2, 1.2 Hz, 2H), 7.59 (tt, J = 7.2, 1.5 Hz, 1H), 7.48 (tt, J = 7.8, 1.5 Hz, 2H), 6.58 (dd, J = 3.9, 1.8 Hz, 1H), 4.88-4.96 (m, 1H), 4.78 (ddd, J = 15.9, 3.9, 1.5 Hz, 1H), 4.08 (ddd, J = 15.6, 3.9, 2.1 Hz, 1H), 2.71 (ddd, J = 15.9, 12.3, 8.1 Hz, 1H), 2.48-2.48 (m, 1H), 2.41 (ddd, J = 16.2, 9.0, 0.6 Hz, 1H), 2.21 (app tt, J = 12.3, 9.0 Hz, 1H). ^{13}C NMR (CDCl_3 , 75 MHz): δ 193.2, 177.7, 141.8, 141.3, 137.4, 132.7, 128.7, 128.5, 68.1, 49.5, 33.4, 28.7. IR (NaCl, neat): 3047 *Br s*, 3063*w*, 2978*w*, 2929*m*, 2873*m*, 1759*m*, 1694*s*, 1643*s*, 1606*s*, 1577*m*, 1501*w*, 1447*m*, 1400*s*, 1354*s*, 1306*w*, 1275*s*, 1238*s*, 1215*m*, 1168*m*, 1108*w*, 1076*w*, 1054*w*, 1026*w*, 1000*w*, 912*m*, 889*w*, 860*w*, 790*w*, 761*m*, 712*s*, 656*s* cm^{-1} . HRMS (CI) m/z calcd for $\text{C}_{14}\text{H}_{14}\text{N}_1\text{O}_2$ [$\text{M}^+ + 1$] 228.1025, found 228.1024.

(2,5-Dihydro-furan-3-yl)-phenyl-methanone (6b)

6b

White solid. Column; hexane:EtOAc = 4:1. R_f = 0.41 (hexane:EtOAc = 3:1). Mp: 48-50 °C. ^1H NMR (CDCl_3 , 400 MHz): δ 7.78-7.80 (m, 2H), 7.58 (tt, J = 7.2, 1.2 Hz, 1 H), 7.47 (t, J = 8.0 Hz, 2 H), 7.41 (app pent, J = 4.0, 1 H), 4.99-5.02 (m, 2H), 4.91-4.94 (m, 2H). ^{13}C NMR (CDCl_3 , 100 MHz): δ 190.7, 139.8, 139.4, 137.8, 132.5, 128.6, 128.4, 76.6, 75.1. IR (NaCl, neat): 3063w, 2855m, 2251m, 1639s, 1598m, 1577m, 1447m, 1365m, 1295s, 1243m, 1180w, 1150m, 1072m, 1009w, 909s, 844w, 830w, 779w, 731s, 702s, 668m, 648m cm^{-1} . HRMS (CI) m/z calcd for $\text{C}_{11}\text{H}_{11}\text{O}_2$ [$\text{M}^+ + 1$] 175.0759, found 175.0764.

3-Benzoyl-cyclopent-3-ene-1,1-dicarboxylic acid diethyl ester (7 b)

7b

Pale yellow oil. Column; hexane:EtOAc = 20:1. R_f = 0.23 (hexane:EtOAc = 10:1). ^1H NMR (CDCl_3 , 300 MHz): δ 7.71-7.74 (m, 2H), 7.54 (tt, J = 7.2, 2.1 Hz, 1H), 7.43 (t, J = 7.8 Hz, 2H), 6.38 (app t, J = 2.1 Hz, 1H), 4.24 (q, J = 7.2 Hz, 4H), 3.44 (dd, J = 3.9, 1.8 Hz, 2H), 3.30 (dd, J = 4.5, 2.1 Hz, 2H), 1.28 (t, J = 7.2 Hz, 6H). ^{13}C NMR (CDCl_3 , 75 MHz): δ 192.8, 171.4, 142.0, 141.2, 138.2, 132.1, 128.8, 128.2, 61.8, 58.1, 41.8, 39.4, 14.0. IR (NaCl, neat): 3062 m , 2982 s , 2937 m , 1731 s , 1644 s , 1598 m , 1578 m , 1465 m , 1447 s , 1390 m , 1366 s , 1360 s , 1256 s , 1183 s , 1097 s , 1014 s , 938 m , 924 m , 862 m , 780 m , 708 s , 670 m cm^{-1} . HRMS (CI) m/z calcd for $\text{C}_{18}\text{H}_{21}\text{O}_5$ [$\text{M}^+ + 1$] 317.1389, found 317.1376.

(2H-1-Benzopyran-3-yl)-phenyl-methanone (8b)

8b

Orange-yellow oil. Column; hexane:EtOAc = 8:1. R_f = 0.49 (hexane:EtOAc = 7:1). ^1H NMR (CDCl_3 , 300 MHz): δ 7.70-7.72 (m, 2H), 7.57 (t, J = 7.8 Hz, 1H), 7.47 (t, J = 7.5 Hz, 2H), 7.24 (td, J = 9.3, 1.5 Hz, 1H), 4.06-4.10 (m, 2H), 6.87-6.93 (m, 3H), 5.15 (s, 2H). ^{13}C NMR (CDCl_3 , 75 MHz): δ 194.0, 155.4, 137.4, 137.0, 132.5, 131.9, 129.7, 129.3, 128.9, 128.4, 121.8, 120.9, 116.3, 65.1. IR (NaCl, neat): 3058 m , 2805 m , 1626 s , 1604 s , 1568 s , 1483 m , 1450 m , 1448 m , 1386 m , 1344 s , 1322 m , 1294 m , 1224 s , 1179 w , 1139 m , 1102 m , 1074 w , 1019 w , 942 w , 884 w , 849 w , 830 w , 757 m , 739 w , 708 m , 669 w cm^{-1} . HRMS (CI) m/z calcd for $\text{C}_{16}\text{H}_{13}\text{O}_2$ [$\text{M}^+ + 1$] 237.0916, found 237.0918.

1-(2*H*-1-Benzopyran-3-yl)-ethanone (9b)

9b

Pale yellow solid. Column; hexane:EtOAc = 40:1. R_f = 0.28 (hexane:EtOAc = 7:1). Mp: 84-86 °C. ^1H NMR (CDCl_3 , 400 MHz): δ 7.31 (s, 1H), 7.26 (app td, J = 6.8, 1.6 Hz, 1H), 7.17 (dd, J = 7.6, 1.6 Hz, 1H), 7.41 (app td, J = 7.6, 1.2 Hz, 1H), 6.86 (d, J = 8.4 Hz, 1H), 5.01 (d, J = 1.6 Hz, 2H), 2.41 (s, 3H). ^{13}C NMR (CDCl_3 , 100 MHz): δ 196.0, 155.5, 134.0, 132.4, 130.7, 129.1, 121.8, 120.7, 116.3, 64.2, 25.0. IR (NaCl, neat): 3057 w , 2922 m , 2853 m , 1734 m , 1706 m , 1656 s , 1604 s , 1569 m , 1484 m , 1459 m , 1392 m , 1360 m , 1337 m , 1314 m , 1291 s , 1238 m , 1211 s , 1151 w , 1120 m , 1067 w , 1032 w , 1005 w , 933 w , 886 w , 854 w , 755 s cm^{-1} . HRMS (CI) m/z calcd for $\text{C}_{11}\text{H}_{11}\text{O}_2$ [$\text{M}^+ + 1$] 175.0759, found 175.0765.

4-Benzoyl-6a-methyl-3,5,6,6a-tetrahydro-2H-pentalen-1-one (10b)

10b

Colorless oil. Column; hexane:EtOAc = 4:1. R_f = 0.34 (hexane:EtOAc = 4:1). ^1H NMR (CDCl_3 , 300 MHz): δ 7.70-7.73(m, 2H), 7.56 (tt, J = 7.5, 1.2 Hz, 1H), 7.46 (app tt, J = 6.9, 1.5 Hz, 2H), 2.95-3.15 (m, 2H), 2.70-2.78 (m, 1H), 2.29-2.57 (m, 3H), 2.02-2.18 (m, 1H), 1.94 (ddd, J = 13.2, 7.2, 1.8 Hz, 1H), 1.37 (s, 3H). ^{13}C NMR (CDCl_3 , 75 MHz): δ 216.7, 193.4, 162.6, 162.6, 139.4, 134.3, 132.3, 128.5, 128.4, 64.0, 40.2, 34.5, 32.5, 23.6, 20.7. IR (NaCl, neat): 3060w, 2963m, 2927m, 2865m, 1741s, 1638s, 1598m, 1577m, 1447s, 1371m, 1342s, 1272s, 1217w, 1176w, 1141w, 1066m, 876m, 793w, 724m, 697m cm^{-1} . HRMS (CI) m/z calcd for $\text{C}_{16}\text{H}_{17}\text{O}_2$ [$\text{M}^+ + 1$] 241.1229, found 241.1222.

4. Representative intermolecular alkyne-carbonyl metathesis

To a solution of 1-phenylpropyne (1.0 mmol, 102 mg, 110 μ L) and isobutyraldehyde (3.0 mmol, 216 mg, 271 μ L) in dichloroethane (4 mL) was added a solution of AgSbF₆ (0.5 M in THF, 10 mol%, 100 μ mol, 200 μ L) at rt. The mixture was refluxed under argon atmosphere for 16 h, and all starting material was consumed by TLC analysis. After all solvent was removed on a rotary evaporator, the crude mixture was purified by column chromatography eluting with hexane:EtOAc = 40:1. The product was isolated as yellow oil (124 mg) and the isolated yield is 71%.

(E)-2,4-Dimethyl-1-phenyl-pent-2-en-1-one (11b)⁶

11b

Yellow oil. Column; hexane:EtOAc = 40:1. R_f = 0.29 (hexane:EtOAc = 30:1). ¹H NMR (CDCl₃, 300 MHz): δ 7.63 (app dt, J = 6.9, 1.2 Hz, 2H), 7.49 (tt, J = 7.2, 1.2 Hz, 1H), 7.41 (app tt, J = 7.8, 1.2 Hz, 2H), 6.10 (dq, J = 9.6, 1.5 Hz, 1H), 2.78 (dq, J = 9.3, 6.9 Hz, 1H), 1.98 (d, J = 1.2 Hz, 3H), 1.03 (d, J = 6.9, 6H). ¹³C NMR (CDCl₃, 75 MHz): δ 199.3, 153.2, 138.7, 134.0, 131.3, 129.3, 128.0, 28.4, 21.9, 12.33. IR (NaCl, neat): 3057 w , 2962 s , 2924 m , 2871 m , 1729 w , 1648 s , 1598 w , 1576 w , 1466 w , 1446 w , 1388 w , 1364 w , 1318 m , 1271 m , 1177 w , 1110 w , 1018 m , 978 w , 894 w , 789 w , 712 m , 647 m cm⁻¹. HRMS (CI) m/z calcd for C₁₃H₁₇O₁ [M^+ + 1] 189.1279, found 189.1275.

(E)-2,5-Dimethyl-1-phenyl-hex-2-en-1-one (12b)⁷

12b

Yellow oil. Column; hexane:EtOAc = 40:1. R_f = 0.33 (hexane:EtOAc = 20:1). ^1H NMR (CDCl_3 , 500 MHz): δ 7.62 (app dt, J = 8.5, 1.5 Hz, 2 H), 7.50 (tt, J = 7.5, 1.5 Hz, 1H), 7.41 (app tt, J = 8.0, 1.5 Hz, 2H), 6.31 (tq, J = 7.6, 1.0 Hz, 1H), 2.17 (app td, J = 7.0, 0.5 Hz, 2H), 1.96 (s, 3H), 1.74 (app hep, J = 7.0 Hz, 1H), 0.93 (d, J = 7.0 Hz, 6H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 199.1, 145.8, 138.9, 137.1, 131.3, 129.3, 128.0, 38.2, 28.4, 22.6, 12.6. IR (NaCl, neat): 3060w, 2956s, 2869s, 1714w, 1689w, 1648s, 1597m, 1577w, 1465m, 1446m, 1385m, 1366m, 1315m, 1284s, 1263m, 1228m, 1170w, 1064w, 1004m, 902w, 780w, 705s, 636m cm^{-1} . HRMS (CI) m/z calcd for $\text{C}_{14}\text{H}_{19}\text{O}_1$ [$\text{M}^+ + 1$] 203.1436, found 203.1440.

(E)-3-(4-Nitro-phenyl)-1-phenyl-propenone (13b)

13b

Yellow oil. Column; hexane:EtOAc = 15:1. R_f = 0.25 (hexane:EtOAc = 10:1). ^1H NMR (CDCl_3 , 500 MHz): δ 8.25 (app dt, J = 9.0, 2.5 Hz, 2H), 7.78-7.70 (m, 2H), 7.56-7.59 (m, 3H), 7.49 (t, J = 7.5 Hz, 2H), 7.16 (s, 1H), 2.27 (d, J = 1.5 Hz, 3H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 198.4, 147.0, 142.1, 140.0, 138.0, 137.4, 132.2, 130.1, 129.4, 128.3, 123.5, 14.7. IR (NaCl, neat): 3061 w , 2931 w , 2849 w , 2927 w , 1648 s , 1596 s , 1517 s , 1446 m , 1344 s , 1315 m , 1262 s , 1180 w , 1146 w , 1109 m , 1012 s , 911 w , 852 m , 736 m , 712 m cm^{-1} . HRMS (CI) m/z calcd for $\text{C}_{16}\text{H}_{14}\text{N}_1\text{O}_3$ [$\text{M}^+ + 1$] 268.0974, found 268.0961.

Table 1: COSY Data for 13b

Proton No	¹ H δ (ppm) (multiplet, J (Hz), number of proton) ^{a, b}	COSY Correlation ^c
H-1	2.27 (d, <i>J</i> = 1.5 Hz, 3H)	H-8
H-5	7.78-7.70 (m, 2H)	H-6, H-7
H-6	7.49 (t, <i>J</i> = 7.5 Hz, 2H)	H-5, H-7, H-10, H-11
H-8	7.16 (s, 1H)	H-1, H-10, H-11
H-11	8.25 (app dt, <i>J</i> = 9.0, 2.5 Hz, 2H)	H-8, H-11

^aRecorded at 500 MHz. ^bAssignments based on HMQC, HMBC, NOSEY, COSY, and nOe data. ^cOnly those correlations which could be unambiguously assigned are recorded.

Table 2: NOSEY Data for 13b

Proton No	¹ H δ (ppm) (multiplet, J (Hz), number of proton) ^{a, b}	NOSEY Correlation ^c
H-1	2.27 (d, <i>J</i> = 1.5 Hz, 3H)	H-5, H-8, H-10
H-5	7.78-7.70 (m, 2H)	H-1, H-6, H-8
H-6	7.49 (t, <i>J</i> = 7.5 Hz, 2H)	H-5
H-8	7.16 (s, 1H)	H-1, H-5, H-10
H-10	7.56-7.59(m, 2H)	H-1, H-8, H-11
H-11	8.25 (app dt, <i>J</i> = 9.0, 2.5 Hz, 2H)	H-10

^aRecorded at 500 MHz. ^bAssignments based on NOSEY and COSY data. ^cOnly those correlations which could be unambiguously assigned are recorded.

COSY NMR

NOESY NMR

(E)-3-Cyclohex-3-enyl-2-methyl-1-phenyl-propenone (14b)

14b

Pale yellow oil. Column; hexane:EtOAc = 80:1. R_f = 0.40 (hexane:EtOAc = 20:1). ^1H NMR (CDCl_3 , 400 MHz): δ 7.64 (d, J = 8.4 Hz, 2H), 7.49 (t, J = 6.8 Hz, 1H), 7.40 (t, J = 7.6 Hz, 2H), 6.18 (d, J = 9.6 Hz, 1H), 5.68 (app t, J = 15.2 Hz, 2H), 2.71-2.79 (m, 1H), 2.05-2.17 (m, 3H), 2.00 (s, 3H), 1.74-1.86 (m, 2H), 1.39-1.49 (m, 1H). ^{13}C NMR (CDCl_3 , 100 MHz): δ 199.0, 150.5, 138.5, 134.9, 131.3, 129.3, 127.9, 126.9, 125.3, 33.9, 30.2, 27.6, 24.2, 12.4. IR (NaCl, neat): 3059 w , 3023 s , 2919 s , 2854 m , 2835 s , 1687 m , 1650 s , 1597 m , 1577 m , 1446 s , 1436 s , 1383 m , 1349 m , 1316 s , 1299 s , 1284 s , 1272 s , 1245 m , 1208 s , 1177 m , 1135 m , 1074 m , 1032 m , 1012 s , 958 w , 940 m , 902 m , 790 m , 742 m , 733 m , 712 s , 655 s , 644 s cm^{-1} . HRMS (CI) m/z calcd for $\text{C}_{16}\text{H}_{19}\text{O}_1$ [$\text{M}^+ + 1$] 227.1436, found 227.1435.

4. Reference

1. Zhao, L.; Lu, X. *Angew. Chem. Int. Ed.* **2002**, *41*, 4341
2. Chatani, N.; Morimoto, T.; Fukumoto, Y.; Murai, S. *J. Am. Chem. Soc.*, **1998**, *120*, 5335
3. Boger, D. L.; Corbett, W. L. *J. Org. Chem.*, **1993**, *58*, 2068
4. Bashiardes, G.; Safir, I.; Barbot, F.; Laduranty, J. *Tetrahedron Lett.*, **2004**, 1567
5. Balog, A.; Geib, S. V.; Curran, D. P. *J. Org. Chem.*, **1995**, *60*, 345
6. Tsuda, T.; Kiyoi, T.; Saegusa, T. *J. Org. Chem.*, **1990**, *55*, 2554
7. Miranda, P. O.; Díaz, D. D.; Padrón, J. I.; Ramírez, M. A.; Martín, V. S. *J. Org. Chem.*, **2005**, *43*, 1613