Amberlite IRA900N₃ as a new catalyst for the azidation of α,β -unsaturated ketones under solvent-free conditions

Luca Castrica, Francesco Fringuelli,* Luisa Gregoli, Ferdinando Pizzo and Luigi Vaccaro*

CEMIN - “Centro di Eccellenza Materiali Innovativi Nanostrutturati” Dipartimento di Chimica,
Laboratorio di Chimica Organica, Università di Perugia Via Elce di Sotto, 8 I-06123, Perugia, Italia

Fax: (+39) 075 5855560

E-mail: luigi@unipg.it; pizzo@unipg.it

CONTENTS

page S2	Experimental Section
page S3-S14	Full characterization data (¹ H NMR, ¹³ C NMR, IR, GC-EIMS, R _f) for compounds 2a-l
page S14-S39	Copies of the ¹ H and ¹³ C NMR for compounds 2a-l

Experimental Section

All chemicals were purchased and used without any further purification. GC-EIMS analyses were carried out with 70 eV electron energy. All ^1H NMR and ^{13}C NMR spectra were recorded at 200 MHz or 400 MHz, and at 50.3 or 100.6 MHz respectively, using a convenient deuterated solvent (reported in the characterization charts) and the residual peak as internal standard, or TMS in the case of CDCl_3 . IR spectra were recorded with FT-IR instrument, using CHCl_3 as solvent. Thin Layer Chromatography analyses were performed on silica gel on aluminum plates and UV and/or KMnO_4 were used as revealing systems. Column chromatographies were performed by using silica gel 230-400 mesh and eluting as reported in the following characterization charts. Amberlite IRA900F was purchased from Aldrich as "Fluoride on polymer support". Trimethylsilyl azide was purchased from Acros (97% pure, d: 0.860 Kg/L)

β -azido ketones **2a**¹, **2c**², **2f**³, **2g**³, **2i**^{4, 5}, **2j**⁵ and **2k**⁵ are known compounds, products **2b**, **2d**, **2e**, **2h**, and **2l** are new compounds.

Characterization data and copies of the ^1H and ^{13}C NMR spectra the compounds **2a-l** are reported below.

References

- 1 Taylor, S. M.; Zalatan, D. N.; Lerchner, A. M.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2005**, *127*, 1313-1317.
- 2 Davies, A. J.; Donald, A. S. R.; Marks, R. E. *J. Chem. Soc. (C)* **1967**, *21*, 109-112.
- 3 Chung, B. Y.; Park, Y. S.; Cho, I. S.; Hyun, B. C. *Bull. Korean Chem. Soc.* **1988**, *9*, 269-70.
- 4 Xu, L.-W.; Xia, C.-G.; Li, J.-W. Zhou, S.-L. *Tetrahedron Lett.* **2004**, *45*, 1219-1221.
- 5 Guerin, D. J.; Horstmann, T. E.; Miller, S. J. *Org. Lett.* **1999**, *1*, 1107-1109.

Chem. Name	4-azido-4-methylpentan-2-one (2g)			
Lit. Ref.	Bull. Korean Chem. Soc. 1988, 9, 269-70			
<div> </div> <p style="text-align: center;">Mol. Wt.: 141</p>				
<p>METHOD:</p> <p>In a screw capped vial equipped with a magnetic stirrer 4-methylpent-3-en-2-one (1g) (0.109 g, 1.0 mmol), Amberlite IRA900F (0.096 g, 0.25 mmol, 2.6 mmol/g), and azidotrimethylsilane (0.207 mL, 1.5 mmol) were consecutively added and the resulting mixture was left under vigorous stirring at 60 °C. After 3 h, ethyl acetate was added, the catalyst was recovered by filtration and the organic solvent was evaporated under vacuum to give pure 4-azido-4-methylpentan-2-one (2g) as an oil (82 % yield, 0.116 g).</p>				
Mol Formula	C ₆ H ₁₁ N ₃ O	m.p./b.p.	Oil	
Tlc - R _f (eluent)	0.26 (EtOAc/petroleum ether 1/9)			
FT-IR (CHCl ₃ , cm ⁻¹): 2104, 1724, 1464, 1368.				
Elemental Analysis: C, 51.05; H, 7.85; N, 29.77; found C, 51.15; H, 7.82; N, 29.62.				
¹ H NMR 400 MHz CDCl ₃	δ value	No. H	Mult.	j value/Hz
	1.36	6	s	
	2.17	3	s	
	2.57	2	s	
¹³ C NMR (100.6 MHz, CDCl ₃) δ : 26.0, 31.7, 53.1, 59.7, 205.8.				
GC-EIMS (m/z, %): 141 (M ⁺ , <1), 99 (14), 56 (53), 43 (100).				

Chem. Name	3-azido-cyclopentan-1-one (2i)			
Lit. Ref.	Tetrahedron Lett. 2004, 45, 1219-1221; Org. Lett. 1999, 1, 1107-1109			
<div> <p>1i</p><p>2i Mol. Wt.:125</p></div>				
METHOD:				
In a screw capped vial equipped with a magnetic stirrer cyclopent-2-en-1-one (1i) (0.084 g, 1.0 mmol), Amberlite IRA900F (0.096 g, 0.25 mmol, 2.6 mmol/g), and azidotrimethylsilane (0.414 mL, 3.0 mmol) were consecutively added and the resulting mixture was left under vigorous stirring at 30°C. After 4 h, ethyl acetate was added, the catalyst was recovered by filtration and the organic solvent was evaporated under vacuum and the reaction mixture was separated by silica gel column chromatography (8/2 CHCl ₃ /petroleum ether ; silica/sample 30/1) to give pure 3-azido-cyclopentan-1-one (2i) as an oil (70 % yield, 0.088 g).				
Mol Formula	C ₅ H ₇ N ₃ O	m.p./b.p.	Oil	
Tlc - R _f (eluent)	0.35 (CHCl ₃ /petroleum ether 8/2)			
FT-IR (CHCl ₃ , cm ⁻¹): 2111, 1742, 1607, 1452, 1402.				
Elemental Analysis: Calcd. C, 47.99; H, 5.64; N, 33.58; found C, 48.30, H, 5.65; N, 33.46.				
¹ H NMR 200 MHz CDCl ₃	δ value	No. H	Mult.	j value/Hz
	2.13-2.32	5	<i>m</i>	
	2.44	1	<i>dd</i>	18.3, 6.3
	4.29	1	<i>m</i>	
¹³ C NMR (100.6 MHz, CDCl ₃) δ : 28.8, 35.7, 44.0, 58.8, 214.9.				
GC-EIMS (m/z, %): 125 (M ⁺ , 29), 82 (10), 68 (46), 55 (100), 43 (12).

4-azido-heptan-2-one (**2a**)

4-azidononan-2-one (2b)

3-azido-1-(2',6',6'-trimethylcyclohex-1'-en-1'-yl)butan-1-one (2d)

2-(1'-azido-1'-methylethyl)-5-methylcyclohexanone (2f)

2-(1'-azido-1'-methylethyl)-5-methylcyclohexanone (2f)

4-azido-4-methylpentan-2-one (2g)

4-azido-4-phenylbutan-2-one (2h)

ppm

7.2606

3.8970
3.8806
3.8596

2.6907
2.6684
2.6193
2.5970
2.4652
2.4229
2.3934
2.3541
2.3295
2.3038
2.0970
2.0508
1.8711
1.8238
1.7854
1.7599
1.7395
1.7176
1.6786

0.0193

ppm
214.872

3-azido-cyclopentan-1-one (2i)

79.793
77.661
77.024
76.387
59.172
58.854
46.326
45.303
43.992
38.134
37.582
35.677
29.803
29.424
29.159
28.805
23.730
23.170
20.164

ppm
200
180
160
140
120
100
80
60
40
20
0

3-azido-cyclohexan-1-one (2j)

3-azido-4,4-dimethylcyclohexan-1-one (2k)

ppm

7.2592

3.5146
3.4909
3.4687
3.4451
2.7129
2.6898
2.6383
2.6153
2.5280
2.4820
2.4530
2.4075
2.3637
2.3339
2.3143
2.2842
1.8515
1.8212
1.7823
1.7515
1.7214
1.5722
1.5399
1.5269
1.4963
1.4698
1.4576
1.4249
1.1288
1.0556
0.9552
0.8639
0.1059
-0.0351

3-azidocycloheptan-2-one (2I)