**An Efficient Synthesis of 4-Halo-5-hydroxyfuran-2(5*H*)-ones via
the Sequential Halolactonization and γ -Hydroxylation of 4-Aryl-2,3-
alkadienoic Acids**

Shengming Ma*, Bin Wu, and Zhangjie Shi

*State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic
Chemistry, Chinese Academy of Sciences, 354 Fenglin Lu, Shanghai 200032, P. R.*

China

[*masm@mail.sioc.ac.cn*](mailto:masm@mail.sioc.ac.cn)

Supporting Information

Table of Contents

Analytical data for compounds 2	S2
Analytical data for compounds 4	S8
¹ H NMR and ¹³ C NMR spectra of those compounds	S13

Experimental Section

Starting Materials. 2,3-Allenic acids were prepared according to the known method by treatment of the acid chlorides with ethyl 2-(triphenylphosphorylidene)propionate and subsequent hydrolysis of the 2,3-allenic esters with 1.5 equiv of NaOH.¹

General procedure for 3-methyl-4-iodo-5-phenyl-5-hydroxyl-2(5*H*)-furanone **2a from the optically active salt of 2-methyl-4-phenylbuta-2,3-dienoic acid and (L)-(-)-cinchonidine (+)-**1aa**:**² A mixture of 2-methyl-4-phenyl-2,3-butadienoic acid **1a** (5.22g, 30 mmol) and (L)-(-)-cinchonidine (4.416g, 15 mmol) in 25 ml ethyl acetate was stirred at room temperature overnight to afford a salt, which could be readily recrystallized in ethyl acetate / methanol (15:1) for 4 times to afford 2.50 g (35.5%) of the optically active (+)-**1aa**: $[\alpha]_D^{20} = 94.6^\circ$ (c = 1, EtOH).

A mixture of the optically active salt (+)-**1aa** (468 mg, 1.0 mmol) and I₂ (280 mg, 1.1 mmol) in 15 ml THF and 5 ml DMF was stirred at room temperature overnight. Then the mixture was diluted with diethyl ether, washed subsequently with 5% Na₂S₂O₃, brine, and dried over Na₂SO₄. After evaporation, the residue was purified via flash chromatography on silica gel with petroleum ether / diethyl ether (3.5:1) as the eluent to afford 265 mg (84%) of **2a**. The structure of **2a** was unambiguously determined by the X-ray single crystal diffraction study.

The following compounds were prepared according to Procedure A.

Procedure A.

(1) Synthesis of 3-methyl-4-iodo-5-phenyl-5-hydroxyl-2(5*H*)-furanone (2a**).** A

mixture of 2-methyl-4-phenyl-2,3-butadienoic acid **1a** (86 mg, 0.494 mmol), LiOAc·2H₂O (63 mg, 0.62 mmol), and I₂ (250 mg, 0.98 mmol) in THF (2 mL) was stirred at room temperature for 3 h. Then DMF (1 mL) was added, and the mixture was stirred at 40°C for 36 h with O₂ (1 atm). Then the mixture was diluted with diethyl ether, washed subsequently with 5% Na₂S₂O₃, brine, and dried over Na₂SO₄. After evaporation, the residue was purified via flash chromatography on silica gel with petroleum ether / diethyl ether (3.5:1) as the eluent to afford 136 mg (87%) of **2a**: solid, mp 150-152 °C (acetone/*n*-hexane); ¹H NMR (300 MHz, CD₃COCD₃) δ 7.60-7.38 (m, 5 H), 7.25-7.15 (bs, 1 H), 1.94 (s, 3 H); ¹³C NMR (75.4 MHz, CD₃COCD₃) δ 169.9, 137.9, 137.1, 130.2, 129.3, 127.3, 127.2, 107.2, 13.2; MS (*m/z*) 316 (M⁺, 4.65), 67 (100); IR (KBr) 3282, 1742, 1732, 1643 cm⁻¹; Anal. Calcd for C₁₁H₉IO₃: C, 41.80; H, 2.87. Found: C, 41.83; H, 2.62.

(2) 3-(*n*-Propyl)-4-iodo-5-phenyl-5-hydroxyl-2(5H)-furanone (2b): The reaction of **1b** (101 mg, 0.50 mmol), I₂ (257 mg, 1.01 mmol), and LiOAc·2H₂O (65 mg, 0.64 mmol) afforded 143 mg (83%) of **2b**: liquid; ¹H NMR (300 MHz, CD₃COCD₃) δ 7.62-7.35 (m, 5 H), 7.22 (s, 1 H), 2.34 (t, *J* = 7.7 Hz, 2 H), 1.73-1.52 (m, 2 H), 0.96 (t, *J* = 7.4 Hz, 3 H); ¹³C NMR (75.4 MHz, CD₃COCD₃) δ 169.0, 139.6, 137.3, 129.5, 128.6, 127.3, 126.5, 106.4, 29.3, 20.7, 13.5; MS (*m/z*) 344 (M⁺, 4.14), 95 (100); IR (neat) 3364, 2961, 1744, 1637, 955, 867 cm⁻¹; HRMS calcd for C₁₃H₁₃IO₃ 343.9909.

Found 343.9931.

(3) 3-Benzyl-4-iodo-5-phenyl-5-hydroxy-2(5H)-furanone (2c): The reaction of **1c** (100 mg, 0.40 mmol), I_2 (124 mg, 0.49 mmol), and $LiOAc \cdot 2H_2O$ (55 mg, 0.54 mmol) afforded 113 mg (72%) of **2c**: solid, mp 151-155 °C (acetone/*n*-hexane); 1H NMR (300 MHz, CD_3COCD_3) δ 7.60-7.18 (m, 10 H), 3.72 (s, 2 H), 2.86 (s, 1 H); ^{13}C NMR (75.4 MHz, CD_3COCD_3) δ 170.0, 139.9, 138.4, 138.1, 130.7, 130.1, 129.9, 129.7, 129.4, 128.2, 127.6, 107.8, 34.3; MS (m/z) 392 (M^+ , 1.16), 105 (100); IR (KBr) 3234, 1724, 1642 cm^{-1} ; Anal. Calcd for $C_{17}H_{13}IO_3$: C, 52.06; H, 3.34. Found: C, 52.39; H, 3.48.

(4) 3-(*n*-Butyl)-4-iodo-5-(4'-methoxyphenyl)-5-hydroxy-2(5H)-furanone (2d): The reaction of **1d** (187 mg, 0.76 mmol), I_2 (388 mg, 1.53 mmol), and $LiOAc \cdot 2H_2O$ (93 mg, 0.91 mmol) afforded 160 mg (54%) of **2d**: liquid; 1H NMR (300 MHz, CD_3COCD_3) δ 7.39 (d, $J = 6.9$ Hz, 2 H), 6.87 (d, $J = 6.9$ Hz, 2 H), 5.20-4.30 (bs, 1 H), 3.79 (s, 3 H), 2.31 (t, $J = 7.7$ Hz, 2 H), 1.60-1.21 (m, 4 H), 0.91 (t, $J = 7.4$ Hz, 3 H); ^{13}C NMR (75.4 MHz, CD_3COCD_3) δ 169.8, 161.8, 140.5, 130.0, 129.0, 128.1, 115.2, 107.7, 56.0, 30.4, 28.3, 23.4, 14.5; MS (m/z) 388 (M^+ , 17.57), 135 (100); IR

(neat) 3360, 1744, 1638, 1609, 931 cm^{-1} ; HRMS calcd for $\text{C}_{15}\text{H}_{17}\text{IO}_4$ 388.0172. Found 388.0172.

(5) 3-(*n*-Butyl)-4-iodo-5-(4'-fluorophenyl)-5-hydroxyl-2(5*H*)-furanone (2e): The reaction of **1e** (117 mg, 0.50 mmol), I_2 (159 mg, 0.63 mmol), and $\text{LiOAc} \cdot 2\text{H}_2\text{O}$ (64 mg, 0.63 mmol) afforded 138 mg (73%) of **2e**: liquid; ^1H NMR (300 MHz, CD_3COCD_3) δ 7.40 (dd, $J = 5.0$ and 8.9 Hz, 2 H), 7.00 (t, $J = 8.6$ Hz, 2 H), 5.70-4.50 (bs, 1 H), 2.26 (t, $J = 7.7$ Hz, 2 H), 1.55-1.39 (m, 2 H), 1.39-1.18 (m, 2 H), 0.86 (t, $J = 7.4$ Hz, 3 H); ^{13}C NMR (75.4 MHz, CD_3COCD_3) δ 169.7, 164.4 (d, $J = 246.3$ Hz), 140.9, 134.5 (d, $J = 3.4$ Hz), 129.8 (d, $J = 8.7$ Hz), 127.6, 116.3 (d, $J = 21.9$ Hz), 107.0, 30.3, 28.1, 23.3, 14.5; ^{19}F NMR (282 MHz, CD_3COCD_3) δ -114.36; MS (m/z) 376 (M^+ , 2.25), 123 (100); IR (neat) 3389, 1732, 1624, 1605, 1235 cm^{-1} ; HRMS calcd for $\text{C}_{14}\text{H}_{14}\text{FIO}_3$ 375.9972. Found 376.0008.

(6) 3-(*n*-Butyl)-4-iodo-5-(4'-*i*-propylphenyl)-5-hydroxyl-2(5*H*)-furanone (2f): The reaction of **1f** (129 mg, 0.50 mmol), I_2 (155 mg, 0.61 mmol), and $\text{LiOAc} \cdot 2\text{H}_2\text{O}$ (64 mg, 0.63 mmol) afforded 139 mg (70%) of **2f**: solid, mp 113-114 $^\circ\text{C}$ (acetone/*n*-hexane); ^1H NMR (300 MHz, CD_3COCD_3) δ 7.44 (d, $J = 8.7$ Hz, 2 H), 7.31 (d, $J =$

9 Hz, 2 H), 7.25-7.10 (bs, 1 H), 3.05-2.85 (m, 1 H), 2.37 (t, $J = 7.4$ Hz, 2 H), 1.68-1.50 (m, 2 H), 1.50-1.33 (m, 2 H), 1.24 (d, $J = 6.9$ Hz, 6 H), 0.94 (t, $J = 7.2$ Hz, 3 H); ^{13}C NMR (75.4 MHz, CD_3COCD_3) δ 170.0, 151.4, 140.8, 135.8, 128.0, 127.8, 127.7, 107.6, 35.0, 30.5, 28.3, 24.7, 23.5, 14.6; MS (m/z) 400 (M^+ , 24.14), 147 (100); IR (KBr) 3380, 1733, 1632, 1220, cm^{-1} ; Anal. Calcd for $\text{C}_{17}\text{H}_{21}\text{IO}_3$: C, 51.01; H, 5.29. Found: C, 50.89; H, 5.28.

(7) 3-(*n*-Butyl)-4-iodo-5-(4'-methylphenyl)-5-hydroxyl-2(5*H*)-furanone (2g): The reaction of **1g** (115 mg, 0.50 mmol), I_2 (156 mg, 0.61 mmol), and $\text{LiOAc} \cdot 2\text{H}_2\text{O}$ (61 mg, 0.60 mmol) afforded 135 mg (73%) of **2g**: solid, mp 111-112 °C (acetone/*n*-hexane); ^1H NMR (300 MHz, CD_3COCD_3) δ 7.39 (d, $J = 8.4$ Hz, 2 H), 7.24 (d, $J = 7.8$ Hz, 2 H), 7.20-7.10 (bs, 1 H), 2.45-2.25 (m, 5 H), 1.62-1.48 (m, 2 H), 1.48-1.26 (m, 2 H), 0.94 (t, $J = 7.4$ Hz, 3 H); ^{13}C NMR (75.4 MHz, CD_3COCD_3) δ 169.9, 140.6, 140.3, 135.3, 130.1, 128.1, 127.4, 107.4, 30.3, 28.0, 23.3, 21.6, 14.5; MS (m/z) 372 (M^+ , 5.89), 119 (100); IR (KBr) 3374, 1720, 1627, 1209 cm^{-1} ; Anal. Calcd for $\text{C}_{15}\text{H}_{17}\text{IO}_3$: C, 48.41; H, 4.60. Found: C, 48.35; H, 4.53.

(8) 3-Methyl-4-iodo-5-(1'-naphthyl)-5-hydroxyl-2(5*H*)-furanone (2h): The reaction

of **1h** (88 mg, 0.39 mmol), I₂ (177 mg, 0.70 mmol), and LiOAc·2H₂O (70 mg, 0.69 mmol) afforded 112 mg (78%) of **2h**: solid, mp 158-160 °C (acetone/*n*-hexane); ¹H NMR (300 MHz, CD₃COCD₃) δ 8.60-8.30 (bs, 1 H), 8.10-7.85 (m, 3 H), 7.70-7.50 (m, 3 H), 7.45 (bs, 1 H), 2.09 (s, 3 H); ¹³C NMR (75.4 MHz, CD₃COCD₃) δ 169.7, 138.9, 138.2, 135.7, 133.2, 132.2, 131.9, 130.0, 127.6, 127.0, 126.9, 125.7, 108.7, 13.8; MS (*m/z*) 366 (M⁺, 31.19), 67 (100); IR (KBr) 3289, 1747, 1732, 1643, 1287 cm⁻¹; Anal. Calcd for C₁₅H₁₁IO₃: C, 49.20; H, 3.03. Found: C, 49.22; H, 2.72.

(9) 3-(*n*-Propyl)-4-iodo-5-(1'-naphthyl)-5-hydroxyl-2(5H)-furanone (2i): The reaction of **1i** (100 mg, 0.397 mmol), I₂ (177 mg, 0.70 mmol), and LiOAc·2H₂O (75 mg, 0.74 mmol) afforded 111 mg (71%) of **2i**: solid, mp 162-164 °C (acetone/*n*-hexane); ¹H NMR (300 MHz, CD₃COCD₃) δ 8.60-8.40 (bs, 1 H), 8.10-7.90 (m, 3 H), 7.67-7.10 (m, 4 H), 2.49 (t, *J* = 7.4 Hz, 2 H), 1.80-1.60 (m, 2 H), 1.03 (t, *J* = 7.2 Hz, 3 H); ¹³C NMR (75.4 MHz, CD₃COCD₃) δ 168.5, 141.3, 137.3, 134.8, 132.2, 131.4, 130.9, 129.2, 127.3, 126.6, 126.1, 125.9, 124.8, 20.7, 13.7; MS (*m/z*) 394 (M⁺, 36.78), 155 (100); IR (KBr) 3345, 1701, 1634, 1236 cm⁻¹; Anal. Calcd for C₁₇H₁₅IO₃: C, 51.80; H, 3.84. Found: C, 51.86; H, 3.66.

The following compounds were prepared according to Procedure B.

Procedure B.

(1) Synthesis of 3-methyl-4-chloro-5-phenyl-5-hydroxyl-2(5H)-furanone (4a). A solution of **1a** (139 mg, 0.8 mmol) and CuCl₂ (432 mg, 3.2 mmol) in acetone / H₂O (2:1, 4.5 mL) was stirred at 65°C for 4 h (in the case of CuCl₂) or 2 h (in the case of CuBr₂). The mixture was then diluted with diethyl ether, washed subsequently with diluted HCl, saturated NaHCO₃ and brine, and dried over Na₂SO₄. After evaporation, the residue was treated with LiOAc·2H₂O (44 mg, 0.43 mmol) in THF / DMF (2:1, 3 mL) at 40°C with O₂ (1 atm) for 8.5 h. Then the mixture was diluted with diethyl ether, washed with brine, and dried over Na₂SO₄. After filtration and evaporation, the residue was purified via flash chromatography on silica gel with hexane / ethyl acetate (10:1) as the eluent to afford 138 mg (77%) of **4a**: solid, mp 104-105 °C (diethyl ether/*n*-hexane); ¹H NMR (300 MHz, CD₃COCD₃) δ 7.62-7.51 (m, 2 H), 7.51-7.38 (m, 3 H), 7.32 (s, 1 H), 1.91 (s, 3 H); ¹³C NMR (75.4 MHz, CD₃COCD₃) δ 169.9, 154.0, 137.3, 130.7, 129.6, 127.2, 126.9, 105.6, 9.2; MS (*m/z*) 226 (M⁺(³⁷Cl), 6.41), 224 (M⁺(³⁵Cl), 19.32), 105 (100); IR (KBr) 3283, 1741, 1722, 1666 cm⁻¹; Anal. Calcd for C₁₁H₉ClO₃ C, 58.81; H, 4.04. Found C, 59.07; H, 4.08.

(2) 3-Methyl-4-bromo-5-phenyl-5-hydroxyl-2(5H)-furanone (4b): The reaction of **1a** (87 mg, 0.50 mmol), CuBr₂ (448 mg, 2.0 mmol), and LiOAc·2H₂O (53 mg, 0.52 mmol) afforded 73 mg (54%) of **4b**: solid, mp 116-117 °C (acetone/*n*-hexane); ¹H NMR (300 MHz, CD₃COCD₃) δ 7.60-7.50 (m, 2 H), 7.50-7.40 (m, 3 H), 7.40-7.28

(bs, 1 H), 1.92 (s, 3 H); ^{13}C NMR (75.4 MHz, CD_3COCD_3) δ 170.0, 146.6, 137.6, 130.8, 130.7, 129.6, 127.4, 106.4, 10.8; MS (m/z) 270 ($\text{M}^+(\text{}^{81}\text{Br})$, 11.95), 268 ($\text{M}^+(\text{}^{79}\text{Br})$, 12.39), 105 (100); IR (KBr) 3284, 1747, 1735, 1660, cm^{-1} ; Anal. Calcd for $\text{C}_{11}\text{H}_9\text{BrO}_3$ C, 49.10; H, 3.37. Found C, 49.20; H, 3.27.

(3) 3-(*n*-Propyl)-4-chloro-5-phenyl-5-hydroxyl-2(5*H*)-furanone (4c): The reaction of **1b** (162 mg, 0.80 mmol), CuCl_2 (432 mg, 3.2 mmol), and $\text{LiOAc}\cdot 2\text{H}_2\text{O}$ (43 mg, 0.42 mmol) afforded 95 mg (47%) of **4c**: liquid; ^1H NMR (300 MHz, CD_3COCD_3) δ 7.60-7.52 (m, 2 H), 7.50-7.42 (m, 3 H), 3.50-3.00 (bs, 1 H), 2.35 (t, $J = 7.4$ Hz, 2 H), 1.70-1.54 (m, 2 H), 0.94 (t, $J = 7.4$ Hz, 3 H); ^{13}C NMR (75.4 MHz, CD_3COCD_3) δ 169.7, 154.6, 137.4, 130.7, 130.2, 129.7, 127.2, 105.5, 26.4, 21.3, 14.3; MS (m/z) 254 ($\text{M}^+(\text{}^{37}\text{Cl})$, 4.81), 252 ($\text{M}^+(\text{}^{35}\text{Cl})$, 14.53), 105 (100); IR (neat) 3353, 1748, 1661, 1222 cm^{-1} ; HRMS calcd for $\text{C}_{13}\text{H}_{13}^{35}\text{ClO}_3$ 252.0553. Found 252.0576.

(4) 3-(*n*-Propyl)-4-bromo-5-phenyl-5-hydroxyl-2(5*H*)-furanone (4d): The reaction of **1b** (122 mg, 0.60 mmol), CuBr_2 (532 mg, 2.38 mmol), and $\text{LiOAc}\cdot 2\text{H}_2\text{O}$ (32 mg, 0.31 mmol) afforded 89 mg (50%) of **4d**: liquid; ^1H NMR (300 MHz, CD_3COCD_3) δ

7.58-7.50 (m, 2 H), 7.50-7.42 (m, 3 H), 7.38-7.34 (bs, 1 H), 2.34 (t, $J = 7.5$ Hz, 2 H), 1.70-1.54 (m, 2 H), 0.95 (t, $J = 7.7$ Hz, 3 H); ^{13}C NMR (75.4 MHz, CD_3COCD_3) δ 169.8, 147.2, 137.7, 134.1, 130.6, 129.6, 127.3, 106.5, 27.8, 21.4, 14.3; MS (m/z) 298 ($\text{M}^+(\text{}^{81}\text{Br})$, 7.24), 296 ($\text{M}^+(\text{}^{79}\text{Br})$, 7.51), 95 (100); IR (neat) 3358, 1747, 1654, 1222 cm^{-1} ; HRMS calcd for $\text{C}_{13}\text{H}_{13}^{79}\text{BrO}_3$ 296.0048. Found 296.0077.

(5) 3-Benzyl-4-chloro-5-phenyl-5-hydroxyl-2(5H)-furanone (4e): The reaction of **1c** (151 mg, 0.60 mmol), CuCl_2 (327 mg, 2.42 mmol), and $\text{LiOAc} \cdot 2\text{H}_2\text{O}$ (60 mg, 0.59 mmol) afforded 143 mg (79%) of **4e**: solid, mp 136-138 $^\circ\text{C}$ (acetone/*n*-hexane); ^1H NMR (300 MHz, CD_3COCD_3) δ 7.58-7.51 (m, 3 H), 7.48-7.42 (m, 3 H), 7.36-7.30 (m, 4 H), 7.30-7.22 (m, 1 H), 3.72 (s, 2 H); ^{13}C NMR (75.4 MHz, CD_3COCD_3) δ 169.5, 155.1, 137.8, 137.3, 130.8, 129.9, 129.7, 129.5, 128.0, 127.2, 105.7; MS (m/z) 302 ($\text{M}^+(\text{}^{37}\text{Cl})$, 1.38), 300 ($\text{M}^+(\text{}^{35}\text{Cl})$, 4.01), 105 (100); IR (KBr) 3375, 1752, 1741, 1665, 1153 cm^{-1} ; Anal. Calcd for $\text{C}_{17}\text{H}_{13}\text{ClO}_3$ C, 67.89; H, 4.36. Found C, 67.82; H, 4.19.

(6) 3-Benzyl-4-bromo-5-phenyl-5-hydroxyl-2(5H)-furanone (4f): The reaction of **1c** (150 mg, 0.60 mmol), CuBr_2 (538 mg, 2.41 mmol), and $\text{LiOAc} \cdot 2\text{H}_2\text{O}$ (61 mg, 0.60

mmol) afforded 101 mg (49%) of **4f**: solid, mp 138-140 °C (acetone/*n*-hexane); ^1H NMR (300 MHz, CD_3COCD_3) δ 7.58-7.48 (m, 3 H), 7.48-7.40 (m, 3 H), 7.37-7.30 (m, 4 H), 7.30-7.21 (m, 1 H), 3.72 (s, 2 H); ^{13}C NMR (75.4 MHz, CD_3COCD_3) δ 169.8, 148.1, 137.8, 137.7, 133.5, 130.8, 130.0, 129.9, 129.8, 128.1, 127.4, 106.6, 31.9; MS (m/z) 346 ($\text{M}^+(\text{}^{81}\text{Br})$, 0.51), 344 ($\text{M}^+(\text{}^{79}\text{Br})$, 0.53), 105 (100); IR (KBr) 3382, 1745, 1735, 1657 cm^{-1} ; Anal. Calcd for $\text{C}_{17}\text{H}_{13}\text{BrO}_3$ C, 59.15; H, 3.80. Found C, 59.36; H, 3.77.

(7) 3-(*n*-Propyl)-4-chloro-5-(1'-naphthyl)-5-hydroxyl-2(5*H*)-furanone (4g**):** The reaction of **1i** (126 mg, 0.50 mmol), CuCl_2 (270 mg, 2.0 mmol), and $\text{LiOAc} \cdot 2\text{H}_2\text{O}$ (51 mg, 0.50 mmol) afforded 123 mg (81%) of **4g**: solid, mp 120-122 °C (acetone/*n*-hexane); ^1H NMR (300 MHz, CD_3COCD_3) δ 8.61 (bs, 1 H), 8.10-7.45 (m, 7 H), 2.44 (t, $J = 7.2$ Hz, 2 H), 1.78-1.60 (m, 2 H), 0.99 (t, $J = 7.4$ Hz, 3 H); ^{13}C NMR (75.4 MHz, CD_3COCD_3) δ 169.3, 154.3, 135.8, 132.9, 132.3, 131.9, 130.0, 127.5, 127.4, 127.1, 125.7, 107.3, 26.9, 21.4, 14.4; MS (m/z) 304 ($\text{M}^+(\text{}^{37}\text{Cl})$, 14.52), 302 ($\text{M}^+(\text{}^{35}\text{Cl})$, 43.05), 155 (100); IR (KBr) 3329, 1719, 1654 cm^{-1} ; Anal. Calcd for $\text{C}_{17}\text{H}_{15}\text{ClO}_3$ C, 67.44; H, 4.99. Found C, 67.21; H, 4.98.

(8) 3-(*n*-Propyl)-4-bromo-5-(1'-naphthyl)-5-hydroxyl-2(5*H*)-furanone (4h): The reaction of **1i** (126 mg, 0.50 mmol), CuBr₂ (450 mg, 2.0 mmol), and LiOAc·2H₂O (51 mg, 0.50 mmol) afforded 133 mg (77%) of **4h**: solid, mp 148-150 °C (acetone/*n*-hexane); ¹H NMR (300 MHz, CD₃COCD₃) δ 8.55 (bs, 1 H), 8.10-7.40 (m, 7 H), 2.45 (bs, 2 H), 1.80-1.60 (m, 2 H), 1.01 (t, *J* = 7.5 Hz, 3 H); ¹³C NMR (75.4 MHz, CD₃COCD₃) δ 169.6, 147.2, 135.8, 133.0, 132.4, 132.0, 130.1, 127.6, 127.23, 127.16, 125.8, 107.7, 28.3, 21.5, 14.6; MS (*m/z*) 348 (M⁺(⁸¹Br), 20.13), 346 (M⁺(⁷⁹Br), 20.65), 155 (100); IR (KBr) 3313, 1711, 1652, cm⁻¹; Anal. Calcd for C₁₇H₁₅BrO₃ C, 58.81; H, 4.35. Found C, 58.91; H, 4.24.

References:

- (1) *Allenenes in Organic Synthesis*; Schuster, H. F., Coppola, G. M., Eds.; Jone Wiley & Sons: New York, 1984.
- (2) Ma, S.; Wu, S. *Chem. Commun.* **2001**, 441.

wb-3-96-2-3-c

Archive directory: /export/home/masm/vnmrSYS/data
Sample directory:

Pulse Sequence: zgpg30
Solvent: Acetone
Ambient temperature
File: wb-3-96-2-3-c
Mercury-300BB "MERCURYvx300"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.000 sec
Width 18867.9 Hz
200 repetitions
OBSERVE C13, 75.4424690 MHz
DECOUPLE H1, 300.0310270 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 1.0 Hz
FT size 131072
Total time 0 min, 0 sec

3.722

2.861

2.085
2.084
2.058
2.057
2.051
2.048
2.043
2.042

wb-5-12-2-2-c

Archive directory: /export/home/masm/vnmrsvs/data
Sample directory:

Pulse Sequence: s2pu1

Solvent: Acetone
Ambient temperature
File: wb-5-12-2-2-c13
Mercury-300BB "DMC300"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.500 sec
Width 18867.9 Hz
11476 repetitions
OBSERVE C13, 75.4423844 MHz
DECOUPLE H1, 300.0310270 MHz
Power 36 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 1.0 Hz
FT size 65536
Total time 47 hr, 32 min, 47 sec

wb-5-27-2-c

Archive directory: /export/home/masm/vnmrsys/data
Sample directory:

Pulse Sequence: zgpg30
Solvent: Acetone
Ambient temperature
File: wb-5-27-2-c
Mercury-300BB "QNP300"

Relax. delay 3.000 sec
Pulse 45.0 degrees
Acq. time 3.000 sec
Width 18867.9 Hz
5984 repetitions
OBSERVE C13, 75.4423994 MHz
DECOUPLE H1, 300.0310270 MHz
Power 36 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 1.0 Hz
FT size 131072
Total time 107 hr, 43 min, 13 sec

2h

wb-5-51-c

Archive directory: /export/home/masm/vnmr/sys/data
Sample directory:

Pulse Sequence: s2pu1

Solvent: Acetone
Ambient temperature

File: wb-5-51-4-c
Mercury-300BB "OMC300"

Relax. delay 2.000 sec
Pulse 45.0 degrees
Acq. time 2.000 sec
Width 18867.9 Hz
684 repetitions
OBSERVE C13, 75.4423988 MHz
DECOUPLE H1, 300.0310270 MHz
Power 41 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 1.0 Hz
FT size 131072
Total time 14 hr, 26 min, 48 sec

wb-5-54-2-c

Archive directory: /export/home/masm/vnmrsys/data
Sample directory:

Pulse Sequence: s2pu1

Solvent: Acetone
Ambient temperature
File: wb-5-54-2-c
Mercury-300BB "DMC300"

Relax. delay 2.000 sec
Pulse 45.0 degrees
Acq. time 2.000 sec
Width 18867.9 Hz
1128 repetitions
OBSERVE C13, 75.4423962 MHz
DECOUPLE H1, 300.0310270 MHz
Power 41 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 1.0 Hz
FT size 131072
Total time 8 hr, 40 min, 5 sec

13C OBSERVE

Archive directory: /export/home/masm/vnmrsys/data
Sample directory:

Pulse Sequence: s2pu1