Supporting information

An Efficient Synthetic Route to Glyco-amino acid building blocks for Glycopeptide Synthesis

Mallesham Bejugam and Sabine L Flitsch*

s.flitsch@ed.ac.uk

School of Chemistry, Joseph Black Buildings, The University of Edinburgh, West
Mains Road, Edinburgh EH9 3JJ, United Kingdom

General considerations

The reducing sugars (GlcNAc, cellobiose, maltose, maltotriose), DIPEA, anhydrous DMSO and DMF were purchased from Aldrich, and lactose from Lancaster. Chitobiose octaacetate was purchased from Dextra labs and ammonium carbonate from BDH labs. Hydroxybenzotriazole (HOBt) was purchased from Acros chemicals and HBTU, Wang resin and all N- α -Fmoc-protected L-amino acids from Novabiochem. TLC plates (Silica gel 60F₂₅₄) were purchased from Merck and silica-gel (35-70 micron) from Fisher Scientific. Developed TLC plates were visualised under UV lamp and stained with p-anisaldehyde solution. Fmoc-SPPS was monitored by ninhydrin test. All microwave reactions were carried out in CEM Discover automated microwave instrument. The coupling and hydrolysis reactions were carried out under nitrogen atmosphere.

¹H NMR, ¹³C NMR, COSY and HSQC were recorded on Bruker arx250, dpx360, avo 600 instruments. Chemical shifts were relative to the deuterated solvent peak and were in parts per million (ppm). The mass spectra were recorded on HP1050 using electrospray ionisation and FAB technique. Melting points were determined on Gallenkamp melting point apparatus. LC-MS were run on Alliance HT WATERS

2790, using Phenomenex Luna 5 μ 2.1/250mm column (C18) and mobile phase was (0.1% TFA in H₂O, 0.1% TFA in Acetonitrile).

Experimental procedures:

General procedure for microwave-assisted Kochetkov amination reaction (2a-f):

The automated microwave tube was charged with reducing sugar (0.25g), ammonium carbonate (1.25g, 5fold excess) and anhydrous DMSO (0.8ml). The tube was sealed and placed under automated microwave at 40⁰C, 250psi pressure and 10 watts power for 90 minutes under cooling the vessel. The reaction tube was taken off and the reaction mixture was transferred into a round bottom flask. The reaction mixture was freeze dried for over night to remove excess of ammonia and DMSO to afford β -glycosyl amines (**2a-f**) as colourless hygroscopic solids in excellent yields. These amines were used without further purification.

Preparation of N- α -Cbz-L-Asn[GlcNAc]-OⁱBu (**5**)

5

To a stirred suspension of **2a** (0.17g, 0.77mmol) in anhydrous DMF (3ml) and NaHCO₃ (0.13g, 1.53mmol) was added N- α -Cbz-Asp(F)-OⁱBu (0.34g, 1.15mmol) in anhydrous DMF (3ml). The reaction mixture was stirred at room temperature for 2.5hrs. The solvent was removed under reduced pressure. The crude product was purified by flash chromatography using 12% MeOH in CHCl₃ as an eluent to afford **5** as a colourless solid (0.25g, 62%, melting point 196°-198°C). IR (KBr): 3295, 3091, 1747, 1703, 1654, 1296 cm⁻¹. ¹H NMR (600MHz, CD₃OD-d⁴) δ 7.35 (m, 5H), 5.14 (m, 2H), 4.97 (d, 1H, J= 9.7Hz), 4.61(m, 1H), 3.93 (d, 2H, J= 10.7Hz), 3.84 (dd, 1H, J= 1.6Hz, 10.6Hz), 3.76 (dd, 1H, J= 1.6, 10.6Hz), 3.67 (dd, 1H, J= 1.6Hz, 10.5Hz),

3.45 (m, 1H), 3.32 (m, 5H), 2.76 (dd, 1H, J= 5.7Hz, 17.0Hz), 2.70 (dd, 1H, J= 7.2Hz, 16.9Hz), 1.95 (m, 1H), 1.92 (s, 3H), 0.92 (d, 6H, J= 6.7Hz) ^{13}C NMR (600MHz, $\text{CD}_3\text{OD}-d^4$) δ 175.8, 174.2, 173.8, 159.6, 139.3, 130.7, 130.3, 130.2, 81.6, 81.0, 77.5, 73.8, 73.1, 69.9, 57.5, 57.3, 53.4, 39.7, 30.2, 24.1, 20.6. HRMS MS (FAB) calculated mass for $\text{C}_{24}\text{H}_{35}\text{N}_3\text{O}_{10}$, 526.2401 $[\text{M} + \text{H}]^+$ and was found at 526.2405 $[\text{M} + \text{H}]^+$

Preparation of N- α -Cbz-L-Asn[GlcNAc]-OH (6):

6

To a stirred solution of **5** (0.15g, 0.3mmol) in DMF (2ml) and phosphate buffer (pH 7.0, 2ml) was added 2M HCl to maintain the pH at 7.0, followed by L-cysteine (9mg) and papain (43mg). The reaction mixture was stirred at 35 $^{\circ}\text{C}$ for 8hrs and the pH was maintained at 7.0 by adding 0.2M NaOH solution. The stirring was continued for further 7hrs. The reaction mixture was filtered through a celite pad and was washed with water (3ml) and the filtrate pH was adjusted to 4.0 by adding 2M HCl and the reaction mixture was concentrated under reduced pressure. The crude product was purified by flash chromatography using 20% MeOH in CHCl_3 as an eluent to afford **6** as a white solid (85mg, 63 %, isolated yield). IR (KBr): 3294, 1704, 1683, 1650, 1272 cm^{-1} . ^1H NMR (360MHz, $\text{DMSO}-d^6$) δ 8.22 (d, 1H (NH), J= 9.0Hz), 7.83 (d, 1H(NH), J= 8.9Hz), 7.38 (m, 6H (ArHx5, NH), 4.80 (t, 1H, J= 9.3Hz), 4.32 (dd, 1H, J= 7.6Hz, 13.4Hz), 3.70 (dd, 1H, J= 7.6Hz, 13.5Hz), 3.55 (m, 1H), 3.23 (m, 4H), 2.64 (dd, 1H, J= 5.3Hz, 14.3Hz), 2.48 (dd, 1H, J= 5.3Hz, 14.3Hz), 1.77 (s, 3H). ^{13}C NMR (250MHz, D_2O) δ 175.2, 172.9, 172.3, 157.9, 136.2, 129.1, 129.0, 128.5, 79.3, 78.3, 74.4, 70.4, 68.2, 61.2, 55.3, 51.2, 38.4, 31.3, 22.7. MS (ESI) calculated mass for $\text{C}_{20}\text{H}_{27}\text{N}_3\text{O}_{10}$, m/z 469.31 and was found at 470.13 $[\text{M}+\text{H}]^+$, 492.14 $[\text{M}+\text{Na}]^+$

Preparation of N- α -Fmoc-L-Asn[GlcNAc]-O^tBu (**3a**)

3a

To a 25ml round-bottom flask was added N- α -Fmoc-L-aspartic acid α -tert butyl ester (0.49g, 1.2mmol), anhydrous DMF (3ml), HOBT (0.15g, 1.1mmol) and HBTU (1.13g, 3.0mmol). After stirring for 15 min, a solution of **2a** (0.22g, 1.0mmol) was added in anhydrous DMF (3ml). The reaction mixture was stirred at room temperature under nitrogen atmosphere for 24hrs. The solvent was removed under reduced pressure. The crude product was purified by flash chromatography using 13% MeOH in CHCl₃ as an eluent to afford **3a** as a white solid (0.54g, 88%, melting point 182°-184°C). IR (KBr): 3304, 2924, 1702, 1655, 1547, 1084 cm⁻¹. ¹H NMR (250MHz, CD₃OD-d⁴) δ 7.63 (d, 2H, J= 7.3Hz), 7.54 (d, 2H, J= 7.3Hz), 7.25 (t, 2H, J= 7.4Hz), 7.15 (t, 2H, J= 7.4Hz), 4.82 (d, 1H, J= 9.7Hz), 4.31(m, 2H), 4.09 (m, 2H) , 3.65 (m, 3H), 3.23 (m, 3H), 2.70 (m, 2H), 1.82 (s, 3H), 1.28 (s, 9H). ¹³C NMR (360 MHz, CD₃OD-d⁴) δ 172.6, 170.8, 170.1, 162.9, 156.9, 143.3, 140.7, 126.9, 126.3, 124.3, 119.0, 81.1, 78.4, 77.8, 74.3, 69.8, 66.1, 54.2, 50.8, 36.6, 35.0, 29.7, 26.3, 21.0. HRMS (FAB) calculated mass for C₃₁H₃₉N₃O₁₀, 614.2714[M + H]⁺ and was found 614.2712 [M+H]⁺.

Preparation of N- α -Fmoc-L-Asn[GlcNAc]-OH (**4a**)

4a

To a 50ml round-bottom flask was added **3a** (0.49g, 1.2mmol), anhydrous DCM (9ml), TFA (9ml) and anisole (0.9ml). The reaction mixture was stirred at room

temperature under nitrogen atmosphere for 4hrs. The reaction mixture was concentrated under reduced pressure and azeotroped with toluene ($2 \times 20\text{ml}$) and DCM ($2 \times 20\text{ml}$). The residue was triturated with ether to afford a white coloured solid. The crude product was purified by flash chromatography using 20% MeOH in CHCl_3 as an eluent to afford **4a** as a white solid (0.155g, 86%, melting point 240°C dec). IR (KBr): 3306, 3072, 1697, 1654, 1540, 1283, 1081 cm^{-1} . ^1H NMR (360MHz, $\text{D}_2\text{O} + \text{DMSO-d}^6$) δ 7.78 (d, 2H, $J = 7.3\text{Hz}$), 7.60 (d, 2H, $J = 7.3\text{Hz}$), 7.37 (t, 2H, $J = 7.5\text{Hz}$), 7.35 (t, 2H, $J = 7.5\text{Hz}$), 4.98 (d, 1H, $J = 9.7\text{Hz}$), 4.42 (t, 1H, $J = 6.1\text{Hz}$), 4.28 (t, 2H, $J = 6.1\text{Hz}$), 4.19 (m, 1H), 3.65 (m, 3H), 3.35 (m, 2H), 3.31 (d, 1H, $J = 6.2\text{Hz}$), 2.42 (m, 2H), 1.87 (s, 3H). ^{13}C NMR (360MHz, $\text{D}_2\text{O} + \text{DMSO-d}^6$) δ 178.2, 175.6, 174.4, 158.1, 144.9, 141.9, 129.2, 128.6, 126.3, 121.3, 79.2, 78.6, 75.2, 70.5, 67.5, 61.5, 55.3, 53.8, 47.9, 23.1. HRMS (FAB) calculated mass for $\text{C}_{27}\text{H}_{31}\text{N}_3\text{O}_{10}$, 558.2088 $[\text{M} + \text{H}]^+$ and was found at 558.2089 $[\text{M} + \text{H}]^+$

Preparation of N- α -Fmoc-L-Asn [Glc(β 1-4)Glc]-O^tBu (**3b**)

3b

To a 25ml round-bottom flask was added N- α -Fmoc-L-aspartic acid α -tert butyl ester (0.25g, 0.62mmol), anhydrous DMF (3ml), HOBt (83 mg, 0.61mmol) and HBTU (0.58g, 1.53mmol). After stirring for 15 min, was added a solution of **2b** (0.175g, 0.51mmol) in anhydrous DMF (3ml). The reaction mixture was stirred at room temperature under nitrogen atmosphere for 24hrs. The solvent was removed under reduced pressure. The crude product was purified by flash chromatography using 20% MeOH in CHCl_3 as an eluent to afford **3b** as a white solid (0.34g, 92%, melting point $176^\circ\text{--}178^\circ\text{C}$). IR (KBr): 3398, 1716, 1539, 1429 cm^{-1} . ^1H NMR (250MHz, $\text{CD}_3\text{OD-d}^4$) δ 7.7 (d, 2H, $J = 7.3\text{Hz}$), 7.60(d, 2H, $J = 7.3\text{Hz}$), 7.34 (t, 2H, $J = 7.0\text{Hz}$), 7.26 (t, 2H, $J =$

7.0Hz), 4.96 (d, 1H, J= 12.1Hz), 4.35 (t, 1H, J= 6.0Hz), 4.26 (m, 2H), 4.17 (t, 1H, J= 6.7Hz), 3.77 (ddd, 4H, J= 1.86Hz, 11.9Hz, 22.7Hz), 3.57 (dd, 1H, J= 4.5Hz, 11.2Hz), 3.45 (dd, 1H, J= 4.5Hz, 11.0Hz), 3.38 (m, 2H), 3.21(m, 5H), 2.76 (m, 2H), 1.35 (s, 9H) ^{13}C NMR (360MHz, $\text{CD}_3\text{OD}-d^4$) δ 171.3, 170.2, 156.5, 143.3, 140.6, 126.9, 126.3, 124.4, 119.0, 102.7, 81.3, 78.9, 78.4, 76.2, 75.8, 75.4, 72.9, 71.7, 69.4, 66.2, 60.5, 59.7, 50.7, 37.9, 36.6, 26.3. HRMS MS (FAB) calculated mass for $\text{C}_{35}\text{H}_{46}\text{N}_2\text{O}_{15}$, 735.2976 $[\text{M} + \text{H}]^+$ and was found at 735.2976 $[\text{M} + \text{H}]^+$

Preparation of N- α -Fmoc-L-Asn[Glc(β 1-4)Glc]-OH (**4b**)

4b

To a 50ml round-bottom flask was added **3b** (0.15g, 0.2mmol), anhydrous DCM (6ml), TFA (6ml) and anisole (0.6ml). The reaction mixture was stirred under nitrogen atmosphere at room temperature for 4hrs. The reaction mixture was concentrated under reduced pressure and azeotroped with toluene ($2 \times 20\text{ml}$) and DCM ($2 \times 20\text{ml}$). Finally the residue was triturated with ether to afford **4b** as a white coloured solid (0.11g, 80%, melting point $180^\circ\text{-}182^\circ\text{C}$). IR (KBr): 3367, 2919, 1699, 1541, 1163, 1078 cm^{-1} . ^1H NMR (360MHz, $\text{D}_2\text{O} + \text{DMSO}-d^6$) δ 7.71 (d, 2H, J= 7.5Hz), 7.51 (d, 2H, J= 7.5Hz), 7.31 (t, 2H, J= 7.3Hz), 7.23 (t, 2H, J= 7.3Hz), 4.75 (d, 1H, J= 9.2Hz), 4.35 (m, 3H), 4.11 (t, 2H, J= 6.2Hz), 3.72 (m, 2H), 3.63 (m, 2H), 3.55 (m, 4H), 3.35 (m, 5H) 2.85 (m, 2H). ^{13}C NMR (360MHz, $\text{D}_2\text{O} + \text{DMSO}-d^6$) δ 175.7, 174.1, 158.9, 145.2, 142.4, 129.8, 129.2, 126.8, 122.7, 104.2, 80.7, 79.8, 77.8, 77.6, 77.0, 76.5, 74.7, 73.0, 72.7, 71.0, 68.2, 62.1, 61.3, 52.3, 48.2, 42.9, 42.4, 39.4, 38.5. HRMS (FAB) calculated mass for $\text{C}_{31}\text{H}_{38}\text{N}_2\text{O}_{15}$, 679.2350 $[\text{M} + \text{H}]^+$ and was found at 679.2352 $[\text{M} + \text{H}]^+$.

Preparation of N- α -Fmoc-L-Asn[Gal(β 1-4)Glc]-O^tBu (**3c**)

3c

To a 25ml round-bottom flask was added N- α -Fmoc-L-aspartic acid α -tert butyl ester (0.23g, 0.56mmol), anhydrous DMF (3ml), HOBt (76mg, 0.56mmol) and HBTU (0.53g, 1.41mmol). After stirring for 15 min, was added a solution of **2c** (0.16g, 0.47mmol) in anhydrous DMF (3ml). The reaction mixture was stirred at room temperature under nitrogen atmosphere for 24hrs. The solvent was removed under reduced pressure. The crude product was purified by flash chromatography using 20% MeOH in CHCl₃ as an eluent to afford **3c** as a white solid (0.24g, 70 %, melting point 170°C dec). IR (KBr): 3397, 1698, 1539, 1428 cm⁻¹. ¹H NMR (360MHz, CD₃OD-d⁴) δ 7.87 (d, 2H, J= 7.5Hz), 7.71 (d, 2H, J= 7.5Hz), 7.43 (t, 2H, J= 7.4Hz), 7.38 (t, 2H, J= 7.4Hz), 4.93 (d, 1H, J= 9.2Hz), 4.51 (t, 1H, J= 6.7Hz), 4.40 (m, 2H), 4.28 (t, 1H, J= 6.7Hz), 3.84 (m, 5H), 3.55 (m, 6H), 3.34 (m, 2H), 2.79 (m, 2H), 1.49 (s, 9H). ¹³C NMR (360MHz, CD₃OD) δ 170.9, 170.1, 156.2, 143.3, 140.6, 127.0, 126.4, 124.5, 117.2, 103.2, 81.1, 78.9, 78.7, 76.2, 75.3, 75.2, 72.8, 71.6, 70.5, 68.3, 66.0 60.5, 59.8, 50.6, 37.9, 36.6, 26.3. HRMS MS (FAB) calculated mass for C₃₅H₄₆N₂O₁₅, 735.2976 [M + H]⁺ and was found at 735.2977 [M+H]⁺.

Preparation of N- α -Fmoc-L-Asn[Gal(β 1-4)Glc]-OH (**4c**)

4c

To a 50ml round-bottom flask was added **3c** (0.15g, 0.2mmol), anhydrous DCM (6ml), TFA (6ml) and anisole (0.6ml). The reaction mixture was stirred at room

temperature under nitrogen atmosphere for 4hrs. The reaction mixture was concentrated under reduced pressure and azeotroped with toluene (2 × 20ml) and DCM (2 × 20ml). Finally the residue was triturated with ether to afford **4c** as a white coloured solid (0.12g, 87%, 192°-194°C). IR (KBr): 3389, 2931, 1715, 1654, 1540, 1080 cm⁻¹. ¹H NMR (250MHz, DMSO-d⁶) δ 8.16 (d, 2H, J= 7.3Hz), 7.98 (d, 2H, J= 7.3Hz), 7.67 (t, 2H, J= 7.4Hz), 7.61 (t, 2H, J= 7.4Hz), 5.10 (t, 1H, J= 9.1Hz), 4.63 (m, 1H), 4.47 (m, 3H), 3.95 (m, 6H), 3.35 (m, 7H), 2.85 (m, 2H). HRMS (FAB) calculated mass for C₃₁H₃₈N₂O₁₅, 679.2350 [M + H]⁺ and was found at 679.2355 [M + H]⁺

Preparation of N-α-Fmoc-L-Asn[GlcNAc(β1-4)GlcNAc]-O^tBu (**3d**)

3d

To a 25ml round-bottom flask was added N-α-Fmoc-L-aspartic acid α-tert butyl ester (0.15g, 0.38mmol), anhydrous DMF (3ml), HOBt (50mg, 0.38mmol) and HBTU (0.21g, 0.57mmol). After stirring for 15 min, was added a solution of **2d** (0.16g, 0.47mmol) in anhydrous DMF (3ml) and DIPEA (67μl, 0.38mmol). The reaction mixture was stirred at room temperature under nitrogen atmosphere for 24hrs. The solvent was removed under reduced pressure. The crude product was purified by flash chromatography using 23% MeOH in chloroform as an eluent to afford **3d** as a white solid (0.106g, 70%). ¹H NMR (360MHz, DMSO-d⁶) δ 8.30 (d, 1H, J= 8.9Hz), 7.91 (d, 2H, J= 7.4Hz), 7.83 (d, 1H, J= 8.9Hz), 7.73 (d, 2H, J= 7.4Hz), 7.34 (t, 2H, J= 7.4Hz), 5.15 (d, 1H, J= 5.4Hz), 5.05 (d, 1H, J= 5.3Hz), 4.85 (t, 1H, J= 9.4Hz), 4.81 (d, 1H, J= 4.7Hz), 4.78 (t, 1H, J= 4.7Hz), 4.63 (t, 1H, J= 4.8Hz), 4.38 (m, 3H), 3.75 (m, 2H), 3.64 (m, 2H), 3.20 (d, 2H, J= 10.1Hz), 3.05 (m, 2H), 2.75 (m, 2H), 1.87 (s, 3H), 1.82 (s, 3H), 1.41 (s, 9H). ¹³C NMR (360MHz, DMSO-d⁶) δ 172.0, 171.1, 170.9,

170.6, 157.3, 145.2, 142.1, 129.1, 128.8, 128.5, 121.6, 104.3, 82.7, 80.6, 80.1, 78.4, 78.1, 75.4, 74.3, 72.1, 67.1, 62.4, 61.1, 58.9, 56.8, 52.2, 48.0, 38.4, 29.8, 24.5, 24.2. MS (ESI) calculated mass for $C_{39}H_{52}N_4O_{15}$, m/z 816.3 and was found at 817.1 $[M+H]^+$.

Preparation of N- α -Fmoc-L-Asn[GlcNAc(β 1-4)GlcNAc]-OH (4d**)**

4d

To a 25ml round-bottom flask was added **3d** (32mg, 0.04mmol), anhydrous DCM (2ml), TFA (2ml) and anisole (0.2ml). The reaction mixture was stirred at room temperature under nitrogen atmosphere for 4hrs. The reaction mixture was concentrated under reduced pressure and azeotroped with toluene (2×10 ml) and DCM (2×10 ml). Finally the residue was triturated with ether to afford **4d** as a white coloured solid (26mg, 89%). 1H NMR (360MHz, DMSO- d_6) δ 8.57 (d, 1H, $J=9.4$ Hz), 8.24 (d, 1H, $J=9.4$ Hz), 8.16 (bs, 1H, 7.88 (d, 2H, $J=7.3$ Hz), 7.74 (d, 1H, $J=9.2$ Hz), 7.71 (d, 2H, $J=7.3$ Hz), 7.46 (d, 1H, $J=8.32$ Hz), 7.41 (t, 2H, $J=7.42$ Hz), 7.32 (t, 2H, $J=7.10$ Hz), 4.83 (t, 1H, $J=9.3$ Hz), 4.25 (m, 4H), 3.76 (m, 2H), 3.52 (m, 6H), 3.22 (m, 2H), 3.13 (m, 3H), 2.83 (m, 2H), 1.85 (s, 3H), 1.81 (s, 3H). MS (ESI) calculated mass for $C_{35}H_{44}N_4O_{15}$, m/z 760.6 and was found at 783.6 $[M+Na]^+$

Preparation of N- α -Fmoc-L-Asn[Glc(α 1-4) Glc(α 1-4)Glc]-O t Bu (3e**)**

3e

To a 25ml round-bottom flask was added N- α -Fmoc-L-aspartic acid α -tert butyl ester (0.2g, 0.48mmol), anhydrous DMF (3ml), HOBt (64mg, 0.48mmol) and HBTU (0.27g, 0.71mmol). After stirring for 15 min, was added a solution of **2e** (0.12g, 0.24mmol) in anhydrous DMF (3ml). The reaction mixture was stirred at room temperature under nitrogen atmosphere for 24hrs. The solvent was removed under reduced pressure. The crude product was purified by flash chromatography using 25% MeOH in CHCl₃ as an eluent to afford **3e** as a white coloured crystalline solid (0.15g, 71%, melting point 126°-130°C). ¹H NMR (360MHz, CD₃OD-d⁴) δ 7.68 (d, 2H, J= 7.4Hz), 7.55 (d, 2H, J= 7.4Hz), 7.28 (t, 2H, J= 7.3Hz), 7.20 (t, 2H, J= 7.3Hz), 5.08 (m, 2H), 4.80 (d, 1H, J= 13.6Hz), 4.39 (t, 1H, J= 6.0Hz), 4.26 (m, 2H), 4.12 (t, 1H, J= 12.9 Hz), 3.74 (m, 5H), 3.59 (m, 5H), 3.45 (m, 5H), 3.22 (m, 3H), 2.75 (m, 2H), 1.35 (s, 9H). ¹³C NMR (360MHz, CD₃OD-d⁴) δ 171.2, 170.14, 156.45, 143.2, 140.5, 126.8, 126.2, 124.3, 118.9, 100.9, 100.7, 81.2, 79.3, 79.0, 78.9, 76.7, 76.1, 73.1, 72.9, 72.8, 72.2, 71.7, 71.4, 71.3, 69.5, 66.2, 60.7, 60.1, 50.6, 47.3, 36.9, 26.2. MS (ESI) calculated mass for C₄₁H₅₆N₂O₂₀, m/z 896.8 and was found at 897.4 [M+H]⁺, 919.3 [M+Na]⁺.

Preparation of N- α -Fmoc-L-Asn[Glc(α 1-4) Glc(α 1-4) Glc]-OH (**4e**)

4e

To a 25ml round-bottom flask was added **3e** (95mg, 0.11mmol), anhydrous DCM (4ml), TFA (4ml) and anisole (0.4ml). The reaction mixture was stirred at room temperature under nitrogen atmosphere for 4hrs. The reaction mixture was concentrated under reduced pressure and azeotroped with toluene (2 × 15ml) and

DCM (2 × 15ml). Finally the residue was triturated with ether to afford **4e** as a white coloured solid (79mg, 92%). IR (KBr): 3387, 2932, 1675, 1540, 1151 cm⁻¹. ¹H NMR (360MHz, CD₃OD-d⁴) δ 7.81(d, 2H, J= 7.4Hz), 7.68 (d, 2H, J= 7.4Hz), 7.41 (t, 2H, J= 7.3Hz), 7.33 (t, 2H, J= 7.3Hz), 5.20 (m, 2H), 4.83 (d, 1H, J= 10.6Hz), 4.62 (m, 4H), 4.32 (m, 3H), 3.84 (m, 5H), 3.71 (m, 4H), 3.56 (m, 5H), 3.35 (m, 4H), 2.87 (m, 2H). ¹³C NMR (360MHz, CD₃OD-d⁴) δ 172.9, 171.4, 156.5, 143.2, 140.6, 126.8, 126.3, 124.4, 119.0, 100.9, 100.6, 79.1, 77.0, 76.7, 76.2, 73.1, 73.0, 72.8, 72.3, 71.8, 71.5, 71.4, 69.5, 66.3, 65.0, 60.7, 60.1, 49.9, 47.1, 36.6. HRMS (FAB) calculated mass for C₃₇H₄₈N₂O₂₀, 841.2879 [M + H]⁺ and was found at 841.2866 [M + H]⁺.

Fmoc-SPPS of glycopeptide 7 and 8

Glycopeptides **7** and **8** were synthesised using standard Fmoc-SPPS protocol. Peptides were assembled on Wang resin (0.82mmol/gram), and the first amino acid was coupled using symmetrical anhydride method. The loading was calculated using UV detection of the Fmoc deprotection (0.56mmol/gram).

The remaining SPPS including coupling of glycosylated asparagine was performed using standard coupling conditions (Fmoc-amino acid (4eq), TBTU (4eq), HOBt (4eq), DIPEA (8eq), and DMF (2ml) for 4hrs followed Fmoc deprotection (20% piperidine in DMF for 1hr). The glyco-amino acids were incorporated into peptide sequence, using double coupling. Finally the resin and side chain protecting groups were removed by using a mixture (3ml) of TFA (94%), H₂O (2.5%), EDT (2.5%) and TIS (1%) for 4hrs, and the resin was filtered off and the filtrate was concentrated under reduced pressure followed by precipitation with cold ether. The glycopeptides **7** and **8** were dissolved separately in two round bottom flasks using (MeCN: H₂O: TFA (1:1:0.1%)) and were lyophilised to afford white solids, glycopeptides **7** (19mg, 67%) and **8** (17mg, 54%). These glyco peptides were analysed by LC-MS, using electrospray ionisation technique and HRMS (FAB) Calculated mass for

$\text{C}_{35}\text{H}_{57}\text{N}_{13}\text{O}_{16}\text{S}$ (glycopeptide **7**), 948.3845 $[\text{M} + \text{H}]^+$ and was found at 948.3849 $[\text{M} + \text{H}]^+$. Calculated mass for $\text{C}_{39}\text{H}_{64}\text{N}_{12}\text{O}_{21}\text{S}$ (glycopeptide **8**), 1069.4108 $[\text{M} + \text{H}]^+$ and was found at 1069.4106 $[\text{M} + \text{H}]^+$.

Source Data Parameters		F1 - Acquisition Parameters		F2 - Processing Parameters		F3 - Plot Parameters	
DATE	NOV4124103	DATE	20031128	DATE	20031128	DATE	20031128
TIME	17:59	TIME	17:59	TIME	17:59	TIME	17:59
PRGNO	1	PRGNO	1	PRGNO	1	PRGNO	1
EXPNO	10	EXPNO	10	EXPNO	10	EXPNO	10
PROCNO	1	PROCNO	1	PROCNO	1	PROCNO	1
EXPRES	10.00 Hz	EXPRES	10.00 Hz	EXPRES	10.00 Hz	EXPRES	10.00 Hz
PROBHD	5 mm QNP 1H/13	PROBHD	5 mm QNP 1H/13	PROBHD	5 mm QNP 1H/13	PROBHD	5 mm QNP 1H/13
NUC1	13C	NUC1	13C	NUC1	13C	NUC1	13C
NUC2	1H	NUC2	1H	NUC2	1H	NUC2	1H
NUC3	15N	NUC3	15N	NUC3	15N	NUC3	15N
NUC4	31P	NUC4	31P	NUC4	31P	NUC4	31P
NUC5	19F	NUC5	19F	NUC5	19F	NUC5	19F
NUC6	29Si	NUC6	29Si	NUC6	29Si	NUC6	29Si
NUC7	17O	NUC7	17O	NUC7	17O	NUC7	17O
NUC8	139La	NUC8	139La	NUC8	139La	NUC8	139La
NUC9	171Yb	NUC9	171Yb	NUC9	171Yb	NUC9	171Yb
NUC10	163Dy	NUC10	163Dy	NUC10	163Dy	NUC10	163Dy
NUC11	151Eu	NUC11	151Eu	NUC11	151Eu	NUC11	151Eu
NUC12	133Cs	NUC12	133Cs	NUC12	133Cs	NUC12	133Cs
NUC13	125Te	NUC13	125Te	NUC13	125Te	NUC13	125Te
NUC14	199Au	NUC14	199Au	NUC14	199Au	NUC14	199Au
NUC15	209Bi	NUC15	209Bi	NUC15	209Bi	NUC15	209Bi
NUC16	223Rn	NUC16	223Rn	NUC16	223Rn	NUC16	223Rn
NUC17	222Rn	NUC17	222Rn	NUC17	222Rn	NUC17	222Rn
NUC18	220Rn	NUC18	220Rn	NUC18	220Rn	NUC18	220Rn
NUC19	219Rn	NUC19	219Rn	NUC19	219Rn	NUC19	219Rn
NUC20	218Rn	NUC20	218Rn	NUC20	218Rn	NUC20	218Rn
NUC21	217Rn	NUC21	217Rn	NUC21	217Rn	NUC21	217Rn
NUC22	216Rn	NUC22	216Rn	NUC22	216Rn	NUC22	216Rn
NUC23	215Rn	NUC23	215Rn	NUC23	215Rn	NUC23	215Rn
NUC24	214Rn	NUC24	214Rn	NUC24	214Rn	NUC24	214Rn
NUC25	213Rn	NUC25	213Rn	NUC25	213Rn	NUC25	213Rn
NUC26	212Rn	NUC26	212Rn	NUC26	212Rn	NUC26	212Rn
NUC27	211Rn	NUC27	211Rn	NUC27	211Rn	NUC27	211Rn
NUC28	210Rn	NUC28	210Rn	NUC28	210Rn	NUC28	210Rn
NUC29	209Rn	NUC29	209Rn	NUC29	209Rn	NUC29	209Rn
NUC30	208Rn	NUC30	208Rn	NUC30	208Rn	NUC30	208Rn
NUC31	207Rn	NUC31	207Rn	NUC31	207Rn	NUC31	207Rn
NUC32	206Rn	NUC32	206Rn	NUC32	206Rn	NUC32	206Rn
NUC33	205Rn	NUC33	205Rn	NUC33	205Rn	NUC33	205Rn
NUC34	204Rn	NUC34	204Rn	NUC34	204Rn	NUC34	204Rn
NUC35	203Rn	NUC35	203Rn	NUC35	203Rn	NUC35	203Rn
NUC36	202Rn	NUC36	202Rn	NUC36	202Rn	NUC36	202Rn
NUC37	201Rn	NUC37	201Rn	NUC37	201Rn	NUC37	201Rn
NUC38	200Rn	NUC38	200Rn	NUC38	200Rn	NUC38	200Rn
NUC39	199Rn	NUC39	199Rn	NUC39	199Rn	NUC39	199Rn
NUC40	1

BM 051-013

Dec_111203, 00551220, 54704926, 469097, 16.0M, Contamin 5, 4x40K

Channel	Chemical Shift (ppm)	Intensity
1H	2.7075	1.00
1H	1.2581	1.00
1H	1.3460	1.00
1H	1.3760	1.00
1H	1.3762	1.00
1H	1.6716	1.00
1H	3.6599	1.00
1H	3.8705	1.00
1H	3.8707	1.00
1H	3.9301	1.00
1H	4.2044	1.00
1H	4.3203	1.00
1H	4.3205	1.00
1H	4.4831	1.00
1H	4.5810	1.00
1H	4.5812	1.00
1H	4.5813	1.00
1H	7.3335	1.00
1H	7.4005	1.00
1H	7.4771	1.00
1H	7.5292	1.00
1H	7.5300	1.00
1H	7.5293	1.00

Dec4_111203, BM051021P, Surfajugum, c1jstd, MeCN, Position: 6, Qp360

Current Data Parameters
NAME Dec4_111203
EXPNO 12
PROCNO 1
F2 - Acquisition Parameters
Date_ 20111111
Time 12.10
INSTRUM spect
PROBHD 5mm QNP 1H/13C
PULPROG zgpg30
PCPDPRG2 waltz16
SOLVENT MeCN
NS 4
DS 112
SW 23521.53 Hz
FIDRES 0.184565 Hz
AQ 1.494565 sec
RG 327.50
ZG 22.000 sec
ZS 8
TE 300.2 K
D1 1.000000 sec
d11 0.050000 sec
DELTA 0.050000 sec
===== CHANNEL f1 =====
NUC1 13C
P1 12.00 usec
PL1 0.00 dB
SFO1 90.5646055 MHz
===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
P2 12.00 usec
PL2 1.00 dB
SFO2 400.1460500 MHz
===== CHANNEL f3 =====
P3 553.96 usec
PL3 1.00 dB
SFO3 500.1354151 MHz
===== CHANNEL f4 =====
P4 1.40 usec
PL4 0.00 dB
SFO4 100.6281500 MHz
===== CHANNEL f5 =====
P5 1.40 usec
PL5 0.00 dB
SFO5 100.6281500 MHz
===== CHANNEL f6 =====
P6 1.40 usec
PL6 0.00 dB
SFO6 100.6281500 MHz
===== CHANNEL f7 =====
P7 1.40 usec
PL7 0.00 dB
SFO7 100.6281500 MHz
===== CHANNEL f8 =====
P8 1.40 usec
PL8 0.00 dB
SFO8 100.6281500 MHz
===== CHANNEL f9 =====
P9 1.40 usec
PL9 0.00 dB
SFO9 100.6281500 MHz
===== CHANNEL f10 =====
P10 1.40 usec
PL10 0.00 dB
SFO10 100.6281500 MHz
===== CHANNEL f11 =====
P11 1.40 usec
PL11 0.00 dB
SFO11 100.6281500 MHz
===== CHANNEL f12 =====
P12 1.40 usec
PL12 0.00 dB
SFO12 100.6281500 MHz
===== CHANNEL f13 =====
P13 1.40 usec
PL13 0.00 dB
SFO13 100.6281500 MHz
===== CHANNEL f14 =====
P14 1.40 usec
PL14 0.00 dB
SFO14 100.6281500 MHz
===== CHANNEL f15 =====
P15 1.40 usec
PL15 0.00 dB
SFO15 100.6281500 MHz
===== CHANNEL f16 =====
P16 1.40 usec
PL16 0.00 dB
SFO16 100.6281500 MHz
===== CHANNEL f17 =====
P17 1.40 usec
PL17 0.00 dB
SFO17 100.6281500 MHz
===== CHANNEL f18 =====
P18 1.40 usec
PL18 0.00 dB
SFO18 100.6281500 MHz
===== CHANNEL f19 =====
P19 1.40 usec
PL19 0.00 dB
SFO19 100.6281500 MHz
===== CHANNEL f20 =====
P20 1.40 usec
PL20 0.00 dB
SFO20 100.6281500 MHz
===== CHANNEL f21 =====
P21 1.40 usec
PL21 0.00 dB
SFO21 100.6281500 MHz
===== CHANNEL f22 =====
P22 1.40 usec
PL22 0.00 dB
SFO22 100.6281500 MHz
===== CHANNEL f23 =====
P23 1.40 usec
PL23 0.00 dB
SFO23 100.6281500 MHz
===== CHANNEL f24 =====
P24 1.40 usec
PL24 0.00 dB
SFO24 100.6281500 MHz
===== CHANNEL f25 =====
P25 1.40 usec
PL25 0.00 dB
SFO25 100.6281500 MHz
===== CHANNEL f26 =====
P26 1.40 usec
PL26 0.00 dB
SFO26 100.6281500 MHz
===== CHANNEL f27 =====
P27 1.40 usec
PL27 0.00 dB
SFO27 100.6281500 MHz
===== CHANNEL f28 =====
P28 1.40 usec
PL28 0.00 dB
SFO28 100.6281500 MHz
===== CHANNEL f29 =====
P29 1.40 usec
PL29 0.00 dB
SFO29 100.6281500 MHz
===== CHANNEL f30 =====
P30 1.40 usec
PL30 0.00 dB
SFO30 100.6281500 MHz
===== CHANNEL f31 =====
P31 1.40 usec
PL31 0.00 dB
SFO31 100.6281500 MHz
===== CHANNEL f32 =====
P32 1.40 usec
PL32 0.00 dB
SFO32 100.6281500 MHz
===== CHANNEL f33 =====
P33 1.40 usec
PL33 0.00 dB
SFO33 100.6281500 MHz
===== CHANNEL f34 =====
P34 1.40 usec
PL34 0.00 dB
SFO34 100.6281500 MHz
===== CHANNEL f35 =====
P35 1.40 usec
PL35 0.00 dB
SFO35 100.6281500 MHz
===== CHANNEL f36 =====
P36 1.40 usec
PL36 0.00 dB
SFO36 100.6281500 MHz
===== CHANNEL f37 =====
P37 1.40 usec
PL37 0.00 dB
SFO37 100.6281500 MHz
===== CHANNEL f38 =====
P38 1.40 usec
PL38 0.00 dB
SFO38 100.6281500 MHz
===== CHANNEL f39 =====
P39 1.40 usec
PL39 0.00 dB
SFO39 100.6281500 MHz
===== CHANNEL f40 =====
P40 1.40 usec
PL40 0.00 dB
SFO40 100.6281500 MHz
===== CHANNEL f41 =====
P41 1.40 usec
PL41 0.00 dB
SFO41 100.6281500 MHz
===== CHANNEL f42 =====
P42 1.40 usec
PL42 0.00 dB
SFO42 100.6281500 MHz
===== CHANNEL f43 =====
P43 1.40 usec
PL43 0.00 dB
SFO43 100.6281500 MHz
===== CHANNEL f44 =====
P44 1.40 usec
PL44 0.00 dB
SFO44 100.6281500 MHz
===== CHANNEL f45 =====
P45 1.40 usec
PL45 0.00 dB
SFO45 100.6281500 MHz
===== CHANNEL f46 =====
P46 1.40 usec
PL46 0.00 dB
SFO46 100.6281500 MHz
===== CHANNEL f47 =====
P47 1.40 usec
PL47 0.00 dB
SFO47 100.6281500 MHz
===== CHANNEL f48 =====
P48 1.40 usec
PL48 0.00 dB
SFO48 100.6281500 MHz
===== CHANNEL f49 =====
P49 1.40 usec
PL49 0.00 dB
SFO49 100.6281500 MHz
===== CHANNEL f50 =====
P50 1.40 usec
PL50 0.00 dB
SFO50 100.6281500 MHz
===== CHANNEL f51 =====
P51 1.40 usec
PL51 0.00 dB
SFO51 100.6281500 MHz
===== CHANNEL f52 =====
P52 1.40 usec
PL52 0.00 dB
SFO52 100.6281500 MHz
===== CHANNEL f53 =====
P53 1.40 usec
PL53 0.00 dB
SFO53 100.6281500 MHz
===== CHANNEL f54 =====
P54 1.40 usec
PL54 0.00 dB
SFO54 100.6281500 MHz
===== CHANNEL f55 =====
P55 1.40 usec
PL55 0.00 dB
SFO55 100.6281500 MHz
===== CHANNEL f56 =====
P56 1.40 usec
PL56 0.00 dB
SFO56 100.6281500 MHz
===== CHANNEL f57 =====
P57 1.40 usec
PL57 0.00 dB
SFO57 100.6281500 MHz
===== CHANNEL f58 =====
P58 1.40 usec
PL58 0.00 dB
SFO58 100.6281500 MHz
===== CHANNEL f59 =====
P59 1.40 usec
PL59 0.00 dB
SFO59 100.6281500 MHz
===== CHANNEL f60 =====
P60 1.40 usec
PL60 0.00 dB
SFO60 100.6281500 MHz
===== CHANNEL f61 =====
P61 1.40 usec
PL61 0.00 dB
SFO61 100.6281500 MHz
===== CHANNEL f62 =====
P62 1.40 usec
PL62 0.00 dB
SFO62 100.6281500 MHz
===== CHANNEL f63 =====
P63 1.40 usec
PL63 0.00 dB
SFO63 100.6281500 MHz
===== CHANNEL f64 =====
P64 1.40 usec
PL64 0.00 dB
SFO64 100.6281500 MHz
===== CHANNEL f65 =====
P65 1.40 usec
PL65 0.00 dB
SFO65 100.6281500 MHz
===== CHANNEL f66 =====
P66 1.40 usec
PL66 0.00 dB
SFO66 100.6281500 MHz
===== CHANNEL f67 =====
P67 1.40 usec
PL67 0.00 dB
SFO67 100.6281500 MHz
===== CHANNEL f68 =====
P68 1.40 usec
PL68 0.00 dB
SFO68 100.6281500 MHz
===== CHANNEL f69 =====
P69 1.40 usec
PL69 0.00 dB
SFO69 100.6281500 MHz
===== CHANNEL f70 =====
P70 1.40 usec
PL70 0.00 dB
SFO70 100.6281500 MHz
===== CHANNEL f71 =====
P71 1.40 usec
PL71 0.00 dB
SFO71 100.6281500 MHz
===== CHANNEL f72 =====
P72 1.40 usec
PL72 0.00 dB
SFO72 100.6281500 MHz
===== CHANNEL f73 =====
P73 1.40 usec
PL73 0.00 dB
SFO73 100.6281500 MHz
===== CHANNEL f74 =====
P74 1.40 usec
PL74 0.00 dB
SFO74 100.6281500 MHz
===== CHANNEL f75 =====
P75 1.40 usec
PL75 0.00 dB
SFO75 100.6281500 MHz
===== CHANNEL f76 =====
P76 1.40 usec
PL76 0.00 dB
SFO76 100.6281500 MHz
===== CHANNEL f77 =====
P77 1.40 usec
PL77 0.00 dB
SFO77 100.6281500 MHz
===== CHANNEL f78 =====
P78 1.40 usec
PL78 0.00 dB
SFO78 100.6281500 MHz
===== CHANNEL f79 =====
P79 1.40 usec
PL79 0.00 dB
SFO79 100.6281500 MHz
===== CHANNEL f80 =====
P80 1.40 usec
PL80 0.00 dB
SFO80 100.6281500 MHz
===== CHANNEL f81 =====
P81 1.40 usec
PL81 0.00 dB
SFO81 100.6281500 MHz
===== CHANNEL f82 =====
P82 1.40 usec
PL82 0.00 dB
SFO82 100.6281500 MHz
===== CHANNEL f83 =====
P83 1.40 usec
PL83 0.00 dB
SFO83 100.6281500 MHz
===== CHANNEL f84 =====
P84 1.40 usec
PL84 0.00 dB
SFO84 100.6281500 MHz
===== CHANNEL f85 =====
P85 1.40 usec
PL85 0.00 dB
SFO85 100.6281500 MHz
===== CHANNEL f86 =====
P86 1.40 usec
PL86 0.00 dB
SFO86 100.6281500 MHz
===== CHANNEL f87 =====
P87 1.40 usec
PL87 0.00 dB
SFO87 100.6281500 MHz
===== CHANNEL f88 =====
P88 1.40 usec
PL88 0.00 dB
SFO88 100.6281500 MHz
===== CHANNEL f89 =====
P89 1.40 usec
PL89 0.00 dB
SFO89 100.6281500 MHz
===== CHANNEL f90 =====
P90 1.40 usec
PL90 0.00 dB
SFO90 100.6281500 MHz
===== CHANNEL f91 =====
P91 1.40 usec
PL91 0.00 dB
SFO91 100.6281500 MHz
===== CHANNEL f92 =====
P92 1.40 usec
PL92 0.00 dB
SFO92 100.6281500 MHz
===== CHANNEL f93 =====
P93 1.40 usec
PL93 0.00 dB
SFO93 100.6281500 MHz
===== CHANNEL f94 =====
P94 1.40 usec
PL94 0.00 dB
SFO94 100.6281500 MHz
===== CHANNEL f95 =====
P95 1.40 usec
PL95 0.00 dB
SFO95 100.6281500 MHz
===== CHANNEL f96 =====
P96 1.40 usec
PL96 0.00 dB
SFO96 100.6281500 MHz
===== CHANNEL f97 =====
P97 1.40 usec
PL97 0.00 dB
SFO97 100.6281500 MHz
===== CHANNEL f98 =====
P98 1.40 usec
PL98 0.00 dB
SFO98 100.6281500 MHz
===== CHANNEL f99 =====
P99 1.40 usec
PL99 0.00 dB
SFO99 100.6281500 MHz
===== CHANNEL f100 =====
P100 1.40 usec
PL100 0.00 dB
SFO100 100.6281500 MHz

Decd_111210_0651023P_624P04epom_4pucay_4628_P0410n_6_04466

Peak	Chemical Shift (ppm)	Integration	Assignment
1	7.25	1.00	Aromatic H
2	6.85	1.00	Aromatic H
3	6.75	1.00	Aromatic H
4	6.65	1.00	Aromatic H
5	6.55	1.00	Aromatic H
6	6.45	1.00	Aromatic H
7	6.35	1.00	Aromatic H
8	6.25	1.00	Aromatic H
9	6.15	1.00	Aromatic H
10	6.05	1.00	Aromatic H
11	5.95	1.00	Aromatic H
12	5.85	1.00	Aromatic H
13	5.75	1.00	Aromatic H
14	5.65	1.00	Aromatic H
15	5.55	1.00	Aromatic H
16	5.45	1.00	Aromatic H
17	5.35	1.00	Aromatic H
18	5.25	1.00	Aromatic H
19	5.15	1.00	Aromatic H
20	5.05	1.00	Aromatic H
21	4.95	1.00	Aromatic H
22	4.85	1.00	Aromatic H
23	4.75	1.00	Aromatic H
24	4.65	1.00	Aromatic H
25	4.55	1.00	Aromatic H
26	4.45	1.00	Aromatic H
27	4.35	1.00	Aromatic H
28	4.25	1.00	Aromatic H
29	4.15	1.00	Aromatic H
30	4.05	1.00	Aromatic H
31	3.95	1.00	Aromatic H
32	3.85	1.00	Aromatic H
33	3.75	1.00	Aromatic H
34	3.65	1.00	Aromatic H
35	3.55	1.00	Aromatic H
36	3.45	1.00	Aromatic H
37	3.35	1.00	Aromatic H
38	3.25	1.00	Aromatic H
39	3.15	1.00	Aromatic H
40	3.05	1.00	Aromatic H
41	2.95	1.00	Aromatic H
42	2.85	1.00	Aromatic H
43	2.75	1.00	Aromatic H
44	2.65	1.00	Aromatic H
45	2.55	1.00	Aromatic H
46	2.45	1.00	Aromatic H
47	2.35	1.00	Aromatic H
48	2.25	1.00	Aromatic H
49	2.15	1.00	Aromatic H
50	2.05	1.00	Aromatic H
51	1.95	1.00	Aromatic H
52	1.85	1.00	Aromatic H
53	1.75	1.00	Aromatic H
54	1.65	1.00	Aromatic H
55	1.55	1.00	Aromatic H
56	1.45	1.00	Aromatic H
57	1.35	1.00	Aromatic H
58	1.25	1.00	Aromatic H
59	1.15	1.00	Aromatic H
60	1.05	1.00	Aromatic H
61	0.95	1.00	Aromatic H
62	0.85	1.00	Aromatic H
63	0.75	1.00	Aromatic H
64	0.65	1.00	Aromatic H
65	0.55	1.00	Aromatic H
66	0.45	1.00	Aromatic H
67	0.35	1.00	Aromatic H
68	0.25	1.00	Aromatic H
69	0.15	1.00	Aromatic H
70	0.05	1.00	Aromatic H

20-Jan-2004, 15:42:20
UV
Ar1
2.06e6

20-Jan-2004, 15:42:20
Scan ES+
6.30e5

DM-GLCNACNEW 116 (4.234) Cm (107:116)
355.3

Calculated mass 947.9 amu

Glycopeptide 7

20-Jan-2004, 16:11:40
Scan 55+
0.3064

BM-CELLOBIOSE-NEW 503 (18.302) Cm (501:510)
335.7

Calculated mass 1083.9 amu

Glycopeptide 8

20-Jan-2004, 16:11:40
Scan ES+
1.05e6

BM-CELLOBIOSE-NEW 115 (4.213) Cm (105:125)

100-181.1

Calculated mass 1068.9 amu

Glycopeptide 8