Supporting Information

An Enantioselective Total Synthesis of (+)-Ricciocarpin A

Ning-Wei Jan^a and Hsing-Jang Liu^{a,b*}

^a*Department of Chemistry, National Tsing Hua University, Hsinchu 30013, Taiwan, ROC*

^b*Institute of Chemistry, Academia Sinica, Nankang, Taipei 11529, Taiwan, ROC*

Compound 4:

To a solution of **3** (2.009 g, 7.97 mmol) in trimethyl orthoacetate (40 mL), was added 10 drops of propanoic acid. The flask was equipped with a Dean-Stark apparatus and a condenser. The mixture was heated at 165 °C for 48 h, during which time the methanol produced and most of the excess trimethyl orthoacetate were collected in the Dean-Stark apparatus. The residue was diluted with ether (40 mL), and the ethereal solution washed with 10% aqueous HCl (10 mL), saturated aqueous NaHCO₃ (10 mL), and brine. The organic phase was dried over MgSO₄, filtered and concentrated to give the crude product. Flash chromatography on silica gel (10% EtOAc in *n*-hexane) afforded iodo ester **4** (1.841 g, 91% yield based on 83% of consumed starting material) as a colorless oil.

$[\alpha]_D^{24} = -29.1$ ($c = 1.03$, CHCl₃)

IR (neat): $\nu_{\max}$ 2954, 2918, 1740, 1631, 1435 cm⁻¹

¹H-NMR (CDCl₃, 400 MHz): δ 6.29 (t, $J = 3.8$ Hz, 1H), 3.68 (s, 3H), 2.72-2.70 (m, 1H), 2.58 (dd, $J = 16.7, 3.2$ Hz, 1H), 2.36 (dd, $J = 16.7, 8.4$ Hz, 1H), 2.08-2.02 (m, 2H), 1.41 (ddd, $J = 13.6, 10.0, 6.6$ Hz, 1H), 1.23 (ddd, $J = 13.6, 5.8, 2.9$ Hz, 1H), 1.02 (s, 3H), 0.86 (s, 3H).

¹³C-NMR (CDCl₃, 100 MHz) : δ 173.4 (C), 137.6 (CH), 103.0 (C), 52.8 (CH), 51.8 (CH₃), 36.6 (CH₂), 34.6 (C), 29.5 (CH₂), 26.95 (CH₃), 26.90 (CH₂), 26.6 (CH₃).

HRMS : Calcd. for C₁₁H₁₇IO₂ : 308.0273; Found: 308.0267.

Anal. Calcd. for C₁₁H₁₇IO₂: C, 42.87; H, 5.56. Found: C, 42.64; H, 5.63.

Compound 6:

To a chilled (0 °C) suspension of lithium aluminum hydride (0.102 g, 2.69 mmol) in anhydrous ether (8 mL) was added slowly a solution of iodo ester **4** (0.415 g, 1.35 mmol) in ether (5 mL). The mixture was stirred for 1 h. This was followed by the sequential addition of 10% aqueous NaOH (0.1 mL), H₂O (0.1 mL) and again 10% aqueous NaOH (0.3 mL). The mixture was stirred vigorously until a white solid precipitated, filtered, and washed thoroughly with ether. The solvent was removed under reduced pressure to afford the crude product which was purified by flash column chromatography on silica gel eluting with 10% ethyl acetate in *n*-hexane to give alcohol **6** (0.356 g, 94% yield) as a colorless oil.

$[\alpha]_D^{24} = -43.6$ ($c = 1.58$, CHCl₃)

IR (neat): $\nu_{\max}$ 3334, 2919, 1632, 1471 cm⁻¹

¹H-NMR (CDCl₃, 400 MHz): δ 6.26 (t, $J = 3.6$ Hz, 1H), 3.80-3.68 (m, 2H), 2.09-2.04 (m, 3H), 1.89-1.80 (m, 1H), 1.75-1.66 (m, 1H), 1.48-1.40 (m, 2H), 1.26-1.20 (m, 1H), 0.99 (s, 3H), 0.93 (s, 3H).

¹³C-NMR (CDCl₃, 100 MHz) : δ 136.5 (CH), 104.2 (C), 62.5 (CH₂), 53.0 (CH), 35.2 (CH₂), 35.1 (C), 30.0 (CH₂), 27.7 (CH₃), 27.1 (CH₃+CH₂).

HRMS : calcd. for C₁₀H₁₇IO : 280.0324; found: 280.0315.

Compound 7:

To a chilled (0 °C) solution of alcohol **6** (0.210 g, 0.75 mmol) in CH₂Cl₂ (5 mL), was added Dess-Martin periodinane (15% by weight in CH₂Cl₂, 3.6 g, 1.27 mmol). The reaction mixture was warmed to room temperature and stirred for 4 h, at which time 10% aqueous Na₂S₂O₃ (10 mL) was added, and the resulting mixture was stirred until a clear solution was formed. The organic layer was washed with saturated aqueous sodium bicarbonate (10 mL), dried over MgSO₄, filtered, and concentrated to give the crude product. Purification by flash chromatography on silica gel (5% EtOAc in *n*-hexane) afforded aldehyde **7** (0.172 g, 82% yield) as a light yellow oil.

$[\alpha]_D^{24} = -35.2$ ($c = 1.90$, CHCl₃)

IR (neat): $\nu_{\max}$ 2920, 2717, 1727, 1633, 1472 cm⁻¹

¹H-NMR (CDCl₃, 400 MHz): δ 9.82 (t, $J = 1.8$ Hz, 1H), 6.33 (t, $J = 3.7$ Hz, 1H), 2.80-2.75 (m, 1H), 2.63 (ddd, $J = 17.8, 3.8, 1.5$ Hz, 1H), 2.56 (ddd, $J = 17.8, 7.3, 2.1$ Hz, 1H), 1.39 (ddd, $J = 13.7, 9.2, 6.9$ Hz, 1H), 1.28 (m, 1H), 1.03 (s, 3H), 0.82 (s, 3H).

¹³C-NMR (CDCl₃, 125 MHz) : δ 201.2 (C), 137.7 (CH), 102.7 (C), 50.9 (CH), 46.0 (CH₂), 34.5 (C), 30.0 (CH₂), 27.6 (CH₃), 26.8 (CH₂), 26.7 (CH₃).

HRMS : Calcd. for C₁₀H₁₅IO : 278.0168; Found: 278.0170.

Compounds 5 and 8:

To a chilled (-78 °C) solution of aldehyde **7** (0.156g, 0.56 mmol) in ether was added a solution of 3-lithiofuran in ether (prepared *in situ* by treatment of 3-bromofuran in ether with *n*-butyllithium at -78 °C; 0.24 M, 9.2 mL, 2.21 mmol) dropwise. The resulting reaction mixture was stirred at -60 °C for 3 h, at which time saturated aqueous ammonium chloride (10 mL) was added. The resulting mixture was extracted with EtOAc (15 mL x 4), and the combined organic phase was washed with brine, dried over MgSO₄, filtered and concentrated to give the crude product. Purification by flash chromatography on silica gel (2% EtOAc in *n*-hexane) yielded iodo alcohol **5** (0.058 g, 30% yield) followed by alcohol **8** (0.115 g, 59% yield), each as a colorless oil.

Compound 5:

$[\alpha]_D^{24} = -32.6$ ($c = 1.01$, CHCl₃)

IR (neat): $\nu_{\max}$ 3409, 2918, 1632, 1593, 1503 cm⁻¹

¹H-NMR (CDCl₃, 400 MHz): δ 7.42 (m, 1H), 7.38 (t, $J = 1.6$ Hz, 1H), 6.48 (dd, $J = 1.8, 0.9$ Hz, 1H), 6.31 (t, $J = 3.8$ Hz, 1H), 5.09 (m, 1H), 2.23 (m, 1H), 2.11-2.05 (m, 2H), 2.00 (ddd, $J = 14.5, 9.2, 4.3$ Hz, 1H), 1.86 (d, $J = 4.4$ Hz, 1H), 1.74 (ddd, $J = 14.5, 6.5, 4.8$ Hz, 1H), 1.50 (dt, $J = 13.6, 8.0$ Hz, 1H), 1.28 (m, 1H), 1.02 (s, 3H), 0.94 (s, 3H).

¹³C-NMR (CDCl₃, 100 MHz) : δ 143.2 (CH), 139.0 (CH), 137.2 (CH), 129.1 (C), 108.8 (CH), 103.1 (C), 66.2 (CH), 52.0 (CH), 40.9 (CH₂), 35.2 (C), 30.4 (CH₂), 27.23 (CH₃), 27.19 (CH₃).

HRMS : Calcd. for C₁₄H₁₉IO₂ : 346.0429; Found: 346.0428.

Compound 8:

$[\alpha]_D^{24} = -28.1$ ($c = 1.79$, CHCl₃)

IR (neat): $\nu_{\max}$ 3409, 2918, 1632, 1593, 1503 cm⁻¹

¹H-NMR (CDCl₃, 400 MHz): δ 7.41 (m, 1H), 7.38 (t, $J = 1.7$ Hz, 1H), 6.43 (dd, $J = 1.8, 0.8$ Hz, 1H), 6.26 (t, $J = 3.8$ Hz, 1H), 4.79 (m, 1H), 2.33 (m, 1H), 2.09-2.03 (m, 2H), 1.91 (ddd, $J = 15.1, 5.9, 4.6$ Hz, 1H), 1.84 (ddd, $J = 15.1, 9.2, 3.5$ Hz, 1H), 1.75 (d, $J = 6.4$ Hz, 1H), 1.48 (ddd, $J = 13.6, 10.0, 7.0$ Hz, 1H), 1.22 (m, 1H), 0.99 (s, 3H), 0.93 (s, 3H).

¹³C-NMR (CDCl₃, 100 MHz) : δ 143.4 (CH), 138.9 (CH), 136.4 (CH), 129.2 (C), 108.5 (CH), 105.4 (C), 66.7 (CH), 52.5 (CH), 40.9 (CH₂), 35.3 (C), 29.6 (CH₂), 27.7 (CH₃), 27.1 (CH₂), 26.8 (CH₃).

HRMS : Calcd. for C₁₄H₁₉IO₂ : 346.0429; Found: 346.0428.

Compound 9:

To a chilled (0 °C) mixture of alcohols **5** and **8** (0.100 g, 0.29 mmol) in CH₂Cl₂ (8 mL), was added Dess-Martin periodinane (15% by weight in CH₂Cl₂, 2.452 g, 0.87 mmol). The reaction mixture was warmed to room temperature and stirred for 6 h, at which time 10% aqueous Na₂S₂O₃ (8 mL) was added, and the resulting mixture was stirred until a clear solution was formed. The organic layer was washed with saturated aqueous sodium bicarbonate (8 mL), dried over MgSO₄, filtered, and concentrated to give the crude product. Purification by flash chromatography on silica gel (5% EtOAc in *n*-hexane) afforded ketone **9** (0.079 g, 79% yield) as a colorless oil.

$[\alpha]_D^{24} = -25.5$ ($c = 1.53$, CHCl₃)

IR (neat): $\nu_{\max}$ 2966, 2915, 1663, 1561 cm⁻¹

¹H-NMR (CDCl₃, 400 MHz): δ 8.07 (brs, 1H), 7.43 (m, 1H), 6.79 (dd, $J = 1.8, 0.7$ Hz, 1H), 6.31 (t, $J = 3.8$ Hz, 1H), 3.11-3.08 (m, 1H), 2.92 (dd, $J = 17.8, 3.3$ Hz, 1H), 2.85 (dd, $J = 17.8, 7.4$ Hz, 1H), 2.11-2.07 (m, 2H), 1.45 (ddd, $J = 13.7, 9.6, 6.6$ Hz, 1H), 1.29-1.20 (m, 1H), 1.03 (s, 3H), 0.79 (s, 3H).

¹³C-NMR (CDCl₃, 100 MHz): δ 193.1 (C), 146.8 (CH), 144.2 (CH), 137.4 (CH), 127.5 (C), 108.8 (CH), 104.3 (C), 50.6 (CH), 43.0 (CH₂), 34.7 (C), 30.1 (CH₂), 27.5 (CH₃), 27.0 (CH₂), 26.7 (CH₃).

HRMS : Calcd. for C₁₄H₁₇IO₂ : 344.0273; Found: 344.0288.

Compound 10:

Compound **5** (65.0 mg, 0.19 mmol) was dissolved in DMPU (0.7 mL) and anhydrous MeOH (0.3 mL). Et₃N (53 μ L, 0.38 mmol), PPh₃ (6.0 mg, 0.02 mmol) and Pd(OAc)₂ (2.7 mg, 0.01 mmol) were added. The mixture was purged with carbon monoxide for 10 min and then stirred under an atmosphere of carbon monoxide at 55 °C for 24 h. After cooling to room temperature, ether (15 mL) was added, and the resulting mixture was washed sequentially with water and brine. The solvent was evaporated, and the residue purified by column chromatography on silica gel (10% EtOAc in *n*-hexane) to provide the lactone **10** (41.0 mg, 89% yield) as a colorless oil.

$[\alpha]_D^{24} = +21.6$ ($c = 2.83$, CHCl₃)

IR (cast, CH₂Cl₂): $\nu_{\max}$ 2957, 2925, 1717, 1635, 1501 cm⁻¹

¹H-NMR (CDCl₃, 400 MHz): δ 7.41 (t, $J = 1.7$ Hz, 1H), 7.35 (dd, $J = 2.4, 1.5$ Hz, 1H), 7.22 (m, 1H), 6.28 (m, 1H), 5.54 (m, 1H), 2.26-2.20 (m, 3H), 2.08 (dt, $J = 13.7, 3.9$ Hz, 1H), 1.92 (ddd, $J = 13.8, 11.8, 4.7$ Hz, 1H), 1.40 (m, 2H), 0.94 (s, 3H), 0.78 (s, 3H).

¹³C-NMR (CDCl₃, 125 MHz) : δ 165.8 (C), 143.8 (CH), 142.3 (CH), 139.5 (CH), 127.0 (C), 125.5 (C), 108.0 (CH), 73.0 (CH), 38.6 (CH), 35.8 (CH₂), 30.7 (C), 28.4 (CH₃), 27.8 (CH₂), 23.8 (CH₂), 19.1 (CH₃).

HRMS : Calcd. for C₁₅H₁₈O₃ : 246.1256; Found: 246.1253.

(+)-Ricciocarpin A **1**

To a chilled (0 °C) solution of compound **10** (15.0 mg, 0.06 mmol) in anhydrous pyridine (1.5 mL) was added sodium borohydride (4.6 mg, 0.12 mmol) in one portion and the reaction mixture was allowed to warm to room temperature. After stirring for 6 h, the reaction mixture was diluted with EtOAc (15 mL) and washed sequentially with 1.0 *N* aqueous HCl and brine, dried over MgSO₄, filtered, and concentrated to give the crude product. After purification by flash chromatography on silica gel (5% EtOAc in *n*-hexane) (+)-ricciocarpin A (**1**) (11.5 mg, 76% yield) was obtained as a white solid.

$[\alpha]_{\text{D}}^{24} = +17.7$ ($c = 0.28$, CH₂Cl₂); natural **1**: $[\alpha]_{\text{D}}^{20} = +17.8$ ($c = 1.18$, CH₂Cl₂)¹

IR (cast, CH₂Cl₂): ν_{max} 2929, 2865, 1741, 1502 cm⁻¹

¹H-NMR (CDCl₃, 500 MHz): δ 7.43 (d, $J = 0.6$ Hz, 1H), 7.40 (t, $J = 1.7$ Hz, 1H), 6.39 (d, $J = 1.0$ Hz, 1H), 5.26 (dd, $J = 9.6, 4.5$ Hz, 1H), 2.39 (dt, $J = 12.3, 3.5$ Hz, 1H), 2.20-2.15 (m, 1H), 2.04 (dt, $J = 14.5, 9.3$ Hz, 1H), 1.91 (ddd, $J = 14.4, 6.8, 4.6$ Hz, 1H), 1.67-1.62 (m, 1H), 1.57-1.48 (m, 1H), 1.50-1.41 (m, 2H), 1.37-1.11 (m, 2H), 0.90 (s, 6H).

¹³C-NMR (CDCl₃, 100 MHz) : δ 175.2 (C), 143.6 (CH), 139.6 (CH), 124.8 (CH), 108.5 (CH), 71.7 (CH), 42.4 (CH), 40.5 (CH₂), 38.9 (CH), 33.7 (C), 29.9 (CH₂), 29.7 (CH₃), 27.2 (CH₂), 21.0 (CH₂), 18.5 (CH₃).

HRMS : Calcd. for C₁₅H₂₀O₃ : 248.1412; Found: 248.1411.

mp : 95-97 °C

Current Data Parameters
NAME PJII-iodoeter
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20041029
Time 11.48
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl3
NS 14
DS 2
SWH 4424.779 Hz
FIDRES 0.135034 Hz
AQ 3.7028339 sec
RG 512
DW 113.000 usec
DE 141.25 usec
TE 300.0 K
HL1 3 dB
P1 1.00000000 sec
SF01 400.1320059 MHz
NUCLEUS 1H

F2 - Processing parameters
SI 32768
SF 400.1300176 MHz
WDW EM
SSB 0
LB 0.01 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 22.00 cm
CY 12.50 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2 -0.500 ppm
F2P -200.07 Hz
PPMCM 0.50000 ppm/cm
HZCM 200.06502 Hz/cm

Current Data Parameters
 NAME PU11-8
 EXPNO 2
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20040810
 Time 22.09
 INSTRUM spect
 PROBHD 5 mm DUL 13C-1
 PULPROG zgpgdc
 TD 32768
 SOLVENT CDCl3
 NS 174
 DS 4
 SWH 22075.055 Hz
 FIDRES 0.673677 Hz
 AQ 0.7422452 sec
 RG 724.1
 DW 22.650 usec
 DE 6.00 usec
 TE 298.1 K
 D1 3.0000000 sec
 D11 0.03000000 sec
 MCREST 0.00000000 sec
 MCWAK 0.01500000 sec

===== CHANNEL f1 =====
 NUC1 13C
 P0 5.00 usec
 PL1 -2.00 dB
 SF01 100.6228372 MHz

===== CHANNEL f2 =====
 CPOPRG2 waltz16
 NUC2 1H
 PCPD2 95.00 usec
 PL2 -2.00 dB
 PL12 16.50 dB
 SF02 400.1316005 MHz

F2 - Processing parameters
 SI 32768
 SF 100.6127746 MHz
 WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 1.40

1D NMR plot parameters
 CX 22.00 cm
 CY 3.00 cm
 FIP 210.000 ppm
 F1 21128.68 Hz
 F2 -10.000 ppm
 F2 -1006.13 Hz

Current Data Parameters
NAME PUJI-125
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050316
Time_ 21.44
INSTRUM spect
PROBHD 5 mm DUL 13C-1
PULPROG zg30
TD 16384
SOLVENT CDCl3
NS 31
DS 2
SWH 4401.409 Hz
FIDRES 0.268641 Hz
AQ 1.8612725 sec
RG 181
DN 113.600 usec
DE 6.00 usec
TE 298.0 K
D1 1.00000000 sec
MCREST 0.00000000 sec
MCMRK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 10.30 usec
PL1 -2.00 dB
SFO1 400.1320173 MHz

F2 - Processing parameters
SI 16384
SF 400.1300165 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 24.00 cm
CY 15.00 cm
FIP 10.500 ppm
F1 4201.35 Hz
F2 -0.500 ppm
F3 -200.06 Hz
PRACH 0.45833 ppm/c
HZCN 183.39201 Hz/cm

Current Data Parameters
NAME PJII-125
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050316
Time 21.53
INSTRUM spect
PROBHD 5 mm DUL 13C-1
PULPROG zgpgdc
TO 32768
SOLVENT CDCl3
NS 6144
DS 4
SWH 22075.055 Hz
FIDRES 0.673677 Hz
AQ 0.7422452 sec
RG 1149.4
DW 22.650 usec
DE 6.00 usec
TE 298.0 K
D1 3.00000000 sec
D11 0.03000000 sec
MGEST 0.00000000 sec
NCHWK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 13C
P0 5.00 usec
PL1 -2.00 dB
SFO1 100.6228385 MHz

===== CHANNEL f2 =====
COPROG2 waltz16
NUC2 1H
PCPD2 95.00 usec
PL2 -2.00 dB
PL12 16.50 dB
SFO2 400.1316005 MHz

F2 - Processing parameters
SI 32768
SF 100.6127719 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 24.00 cm
CY 3.00 cm
FIP 210.000 ppm
F1 21128.68 Hz
F2 -10.000 ppm
F2 -1006.13 Hz
PRICH 9.16667 ppm/c
HZCM 922.283369 Hz/cm

Current Data Parameters
NAME PUJI-1165.m.
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050304
Time 16.09
INSTRUM spect
PROBHD 5 mm DUL 13C-1
PULPROG zg30
TD 16384
SOLVENT CDCl3
NS 13
DS 2
SKH 4401.409 Hz
FIDRES 0.268641 Hz
AQ 1.8612725 sec
RG 322.5
DM 113.600 usec
DE 6.00 usec
TE 295.4 K
D1 1.00000000 sec
NCREST 0.00000000 sec
HCMRK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 10.30 usec
PL1 -2.00 dB
SFO1 400.1320173 MHz

F2 - Processing parameters
SI 16384
SF 400.1300165 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 24.00 cm
CY 15.00 cm
F1P 10.500 ppm
F1 4201.46 Hz
F2P -0.500 ppm
F2 -199.95 Hz
PPMCH 0.45833 ppm/c
HZCH 183.39201 Hz/cr

ppm

Current Data Parameters
NAME PJT-18
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040625
Time 13.18
INSTRUM spect
PROBHD 5 mm QNP 1H/15
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 113
DS 2
SWH 27777.777 Hz
FIDRES 0.847710 Hz
AQ 0.5898920 sec
RG 812.7
DW 18.000 usec
DE 6.00 usec
TE 298.0 K
D1 3.00000000 sec
d11 0.03000000 sec
ICREST 0.00000000 sec
MCMARK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 13C
P0 12.05 usec
PL1 3.00 dB
SFO1 125.7879721 MHz

===== CHANNEL f2 =====
CPOPRG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 0.00 dB
PL12 14.89 dB
SFO2 500.2020008 MHz

F2 - Processing parameters
SI 32768
SF 125.7754035 MHz
WDW EM
SSB 0
LB 3.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 23.00 cm
CY 3.00 cm
F1P 220.000 ppm
F1 27670.59 Hz
F2P -10.000 ppm
F2 -1257.75 Hz
PPMCH 10.00000 ppm/c
HZCH 1257.75403 Hz/cm

Current Data Parameters
NAME FJII- (72-73)
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050525
Time 16.14
INSTRUM spect
PROBHD 5 mm DUL 13C-1
PULPROG zg30
TD 16384
SOLVENT CDCl3
NS 32
DS 2
SWH 4401.409 Hz
FIDRES 0.268641 Hz
AQ 1.8612725 sec
RG 812.7
DM 113.600 usec
DE 6.00 usec
TE 297.0 K
D1 1.00000000 sec
MCREST 0.00000000 sec
MCORR 0.01500000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 10.30 usec
PL1 -2.00 dB
SFO1 400.1320173 MHz

F2 - Processing parameters
SI 16384
SF 400.1300165 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 24.00 cm
CY 13.00 cm
FIP 10.500 ppm
F1 4201.35 Hz
F2 -0.500 ppm
F2 -200.06 Hz
PPMCH 0.45933 ppm/c
HZCM 183.39201 Hz/cir

ppm

143.2085
138.9701
137.1551
129.0518
108.8441
103.1088
77.3149
76.9975
76.6796
66.1881
51.9963
40.9202
35.2193
30.4335
27.2347
27.1932

Current Data Parameters
NAME PUJI-168b
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050527
Time 22.36
INSTRUM spect
PROBHD 5 mm DUL 13C-1
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 6144
DS 4
SWH 22075.055 Hz
FIDRES 0.673677 Hz
AQ 0.7422452 sec
RG 2896.3
DM 22.650 usec
DE 6.00 usec
TE 298.5 K
D1 3.00000000 sec
D11 0.03000000 sec
MCREST 0.00000000 sec
MCWRK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 13C
P0 5.00 usec
PL1 -2.00 dB
SF01 100.6228385 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 95.00 usec
PL2 -2.00 dB
PL12 16.50 dB
SF02 400.1316005 MHz

F2 - Processing parameters
SI 32768
SF 100.6127719 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 24.00 cm
CY 4.00 cm
F1P 210.000 ppm
F1 21128.88 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPMCK 9.16667 ppm/c
HZCM 922.28389 Hz/cm

180 160 140 120 100 80 60 40 20 0

Current Data Parameters
NAME PUJI-168 (13-15)
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20090616
Time 21:39
INSTRUM spect
PROBHD 5 mm DUL 13C-1
PULPROG zgpgdc
TD 32768
SOLVENT CDCl3
NS 122
DS 4
SWH 22075.065 Hz
FIDRES 0.673677 Hz
AQ 0.742262 sec
RG 2896.3
DN 22.650 usec
DE 6.00 usec
TE 299.0 K
D1 3.00000000 sec
D11 0.03000000 sec
HCREST 0.00000000 sec
HCHIRK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 13C
P0 5.00 usec
PL1 -2.00 dB
SFO1 100.6228385 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 95.00 usec
PL2 -2.00 dB
PL12 16.50 dB
SFO2 400.1316005 MHz

F2 - Processing parameters
SI 32768
SF 100.6127712 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 24.00 cm
CY 3.50 cm
F1P 210.000 ppm
F1 21128.66 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPHMM 9.16667 ppm/c
HZCM 922.28369 Hz/cm

26.6573
26.9825
27.4761
30.0636
34.7002
42.9955
50.6217
76.6838
77.0011
77.3487
104.3193
108.8180
127.5042
137.3543
144.1567
146.8206
193.0560

Current Data Parameters
NAME FULL-196
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050329
Time 22.56
INSTRUM spect
PROBHD 5 mm QNP 13C-1
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 5120
DS 4
SWH 23148.148 Hz
FIDRES 0.705425 Hz
AQ 0.7074988 sec
RG 1625.5
DM 21.600 usec
DE 6.00 usec
TE 298.2 K
D1 3.00000000 sec
D11 0.03000000 sec
ICREST 0.00000000 sec
NCHMR 0.01500000 sec

***** CHANNEL f1 *****
NUC1 13C
P0 5.00 usec
PL1 -2.00 dB
SFO1 100.6233416 MHz

***** CHANNEL f2 *****
CPDPRG2 waltz16
NUC2 1H
PCPD2 95.00 usec
PL2 -2.00 dB
PL12 16.50 dB
SFO2 400.1315005 MHz

F2 - Processing parameters
SI 32768
SF 100.6127705 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 24.00 cm
CY 5.00 cm
FIP 220.000 ppm
F1 22134.81 Hz
F2 -10.000 ppm
F2 -1006.13 Hz
PPMCH 9.58333 ppm/c
HZCM 964.20563 Hz/cm

0.7796
0.9438
1.2292
1.3789
1.3941
1.4014
1.4169
1.9269
1.9496
1.9613
2.0563
2.0562
2.0666
2.0679
2.0764
2.2194
2.2226
2.2226
2.2300
2.2382
2.2432
2.2487
2.2534

5.5327
5.5359
5.5411
5.5441
5.5474
5.5524
5.5555
6.2776
6.2799
6.2822
6.2841
7.2147
7.2232
7.2304
7.2402
7.3496
7.3533
7.3555
7.3592
7.4011
7.4053
7.4097

Current Data Parameters
NAME PUJI-129a
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050322
Time 20.52
INSTRUM spect
PROBHD 5 mm DUL 13C-1
PULPROG zg30
TO 16384
SOLVENT COC13
NS 46
DS 2
SWH 4401.409 Hz
FIDRES 0.266641 Hz
AQ 1.8612725 sec
RG 256
DM 113.600 usec
DE 6.00 usec
TE 298.0 K
D1 1.00000000 sec
MCREST 0.00000000 sec
MCWRR 0.01500000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 10.30 usec
PL1 -2.00 dB
SFO1 400.1320173 MHz

F2 - Processing parameters
SI 16384
SF 400.1300165 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 24.00 cm
CY 12.00 cm
F1P 10.500 ppm
F1 4201.35 Hz
F2P -0.500 ppm
F2 -200.05 Hz
PPMCH 0.45833 ppm/c
HZCM 183.39201 Hz/cir

Integral

165.7646
143.8400
142.3461
139.4951
126.9557
125.4604
107.9862
77.2596
77.0057
76.7514
73.0474
38.6432
35.8036
30.7370
29.6532
28.4305
27.7627
23.8381
19.0809

Current Data Parameters
NAME FUL-129a
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050324
Time 22.23
INSTRUM spect
PROBHD 5 mm GNP 1H/15
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 5632
DS 2
SWH 27777.777 Hz
FIDRES 0.847710 Hz
AQ 0.5698920 sec
RG 1625.5
DM 18.000 usec
DE 5.00 usec
TE 298.1 K
D1 3.0000000 sec
d11 0.0300000 sec
MCREST 0.0000000 sec
MCWRT 0.0150000 sec

===== CHANNEL f1 =====
NUC1 13C
P0 12.05 usec
PL1 3.00 dB
SF01 125.7879773 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 0.00 dB
PL12 14.88 dB
SF02 500.2020008 MHz

F2 - Processing parameters
S1 32768
SF 125.7753951 MHz
WDW EM
SSB 0
LB 3.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 23.00 cm
CY 3.50 cm
F1P 210.000 ppm
F1 26412.84 Hz
F2P -10.000 ppm
F2 -1257.75 Hz
PPMCM 9.56522 ppm/c
HZCM 1203.06505 Hz/cm

ppm

nm

Current Data Parameters
NAME PJ1-286
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040529
Time 18.42

INSTRUM spect
PROBHD 5 mm DUL 13C-1
PULPROG zgpg30

TO 32768
SOLVENT CDCl3
NS 1073

DS 4
SWH 22075.055 Hz
FIDRES 0.673677 Hz

AQ 0.7422452 sec
RG 812.7
DW 22.650 usec

DE 6.00 usec
TE 298.0 K
D1 3.0000000 sec

D11 0.03000000 sec
MCREST 0.00000000 sec
MCWRK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 13C
P0 5.00 usec
PL1 -2.00 dB
SF01 100.628372 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 95.00 usec
PL2 -2.00 dB
PL12 16.50 dB
SF02 400.1316005 MHz

F2 - Processing parameters
SI 32768
SF 100.6127705 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 22.00 cm
CY 12.00 cm
F1P 210.000 ppm
F1 21128.68 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPHMC 10.00000 ppm/cm