Supporting Information

An Improved Asymmetric Reformatsky Reaction Mediated by (-)-*N,N*-Dimethylaminoisoborneol

Ralf J. Kloetzing, Tobias Thaler and Paul Knochel*

Department of Chemistry and Biochemistry, Ludwig-Maximilians-University, Butenandtstrasse 5-13, 81377 Munich, Germany

paul.knochel@cup.uni-muenchen.de

1. General

All reactions were carried out under argon using standard Schlenk techniques. Melting points are uncorrected. ^1H and ^{13}C NMR spectra were recorded on a Bruker AMX 300 or AMX 600 or Varian VXR 400 S instrument. Chemical shifts (δ) are given as ppm relative to the residual solvent peak (chloroform-*d*1 7.25 ppm/77.0 ppm; benzene-*d*6 7.16 ppm/128.0 ppm). IR spectra were recorded on a PERKIN ELMER 1420 Infrared Spectrometer. Mass spectra were recorded on a FINNIGAN MAT 95 Q spectrometer. Optical rotations were measured on a PERKIN ELMER 241 polarimeter (temperature: 20 °C). Column chromatography was performed on MERCK silica gel 60 (230-400 mesh ASTM). Enantiomeric excesses were determined by HPLC (Chiralcel OD-H, Daicel Chemical Industries, column was run with *n*-heptane/*i*-propanol as mobile phase (flow rate 0.8 mL/min) and detection was carried out by a diode array UV-VIS detector at the given wavelength) or by chiral GC (Instrument Hewlett Packard HP 6890 GC System equipped with a Varian CP-Chiralsil-Dex-CB column 25m x 0.25 mm x 0.25 μm ; carrier gas H_2 2.0 mL/min; isothermal, temperatures given, FID-detector). Conversions were determined by GC (Hewlett Packard 5890 Series II equipped with a HP Ultra 2 column 12 m x 0.2 mm x 0.35 μm ; FID-detector) using tetradecane as internal standard after a reaction time of 12 h. THF was dried with sodium/benzophenone and distilled. Zinc (dust) and methyl bromoacetate were provided by WACKER Chemie GmbH (München), aldehydes were purchased from Aldrich or Acros. Aliphatic aldehydes were distilled prior to use. 3-Methoxy-3-methylbutanal (Table 2, entry 13) was synthesized from the alcohol by Swern-oxidation. Diethylzinc was provided by WITCO GmbH (Bergkamen).

2. General procedures

2.1. Preparation of the Reformatsky Reagent (1a)

In a 25 mL three-necked flask equipped with an efficient reflux condenser, a dropping funnel, a magnetic stirring bar and a thermometer, zinc (dust, 1.26 g, 19.3 mmol) was suspended in THF (8 mL) and stirred vigorously. The temperature was raised to 40 °C and trimethylsilyl chloride (0.30 mL, 0.26 g, 2.4 mmol) for activation of zinc was added. The temperature was then raised to 55 °C and kept for 15 min. The pressure in the flask was reduced to 330 mbar which assures that the temperature is kept at 37 °C (refluxing THF). Methyl bromoacetate (1.68 mL, 2.70 g, 17.7 mmol) was added during 10 min (CAUTION: exothermic reaction) and stirred for 5 min. The flask was then refilled with argon and the remaining solid material was allowed to settle. The supernatant liquid was decanted with a cannula. This solution of the Reformatsky reagent is stable at ambient temperature for several days. The concentration was determined by titration with iodine (approx. 1.0 mmol) in THF (2 mL) at 0 °C. A typical concentration is 1.7 M.

With *tert*-butyl bromoacetate a less soluble Reformatsky reagent is obtained. It was used as suspension.

2.2. Preparation of the Difluoro Reformatsky Reagent (1c)

Zinc (dust, 358 mg, 5.47 mmol) was suspended in THF (1.5 mL) and trimethylsilyl chloride (0.050 mL, 43 mg, 0.39 mmol) for activation was added. The mixture was heated to 60 °C for 15 min. The heating was stopped and ethyl bromodifluoroacetate (1.11 g, 5.47 mmol) in THF (9.5 mL) was added slowly. The mixture was stirred 25 min at 50 °C.

2.3. General Procedure for the Addition of the Reformatsky Reagents to Aldehydes

(-)-DAIB (4, 109 mg, 0.55 mmol) was dissolved in THF (0.5 mL) and diethylzinc (2M in THF, 0.16 mL, 0.32 mmol) was added at 0 °C. After the mixture had been stirred for 10 min, the temperature was cooled

to $-20\text{ }^{\circ}\text{C}$ (Reformatsky reagent **1a**) or to $-10\text{ }^{\circ}\text{C}$ (difluoro Reformatsky reagent **1c**) and the Reformatsky reagent was added. Then, the mixture was stirred for 20 min. The aldehyde (0.47 mmol) was added as a solution in THF (0.5 mL) together with tetradecane as internal standard. After 2 h, the reaction temperature was allowed to rise to $0\text{ }^{\circ}\text{C}$ (Reformatsky reagent **1a**) or to $+10\text{ }^{\circ}\text{C}$ (difluoro-Reformatsky reagent **1c**). Conc. aq. ammonia (3 mL) and satd. soln. of NH_4Cl (30 mL) were added after 12 h and the mixture was extracted with ethyl acetate. The organic phase was washed with brine and dried over MgSO_4 . The solvent was evaporated and the product was purified by column chromatography (SiO_2 , pentane/diethyl ether 12:1 to 2:1). Caution: the fluorinated products are evaporated on longer drying in high vacuum.

(-)-DAIB could be recovered by extraction of the organic phase with 1M HCl. For the heterocyclic products, (-)-DAIB could be recovered by column chromatography with ethyl acetate/diethyl ether 1:1.

3. Analytical Data

3.1 Methyl (*S*)-3-phenyl-3-hydroxypropanoate (**7a**, Table 1, entry 4)

With benzaldehyde (48.5 mg, 0.457 mmol), alcohol **7a** was obtained (64.1 mg, 0.356 mmol, 75%).

$[\alpha]_D = -17.0^\circ$ (c 1.00, MeOH)

The analytical data is in accordance with the lit.¹

Chiral GC **7a** (Table 1, entry 3)

(120 °C): 43.8 min (*R*), 44.6 min (*S*); 26% *ee*.

Chiral GC **7a** (Scheme 2, add.: thiophene)

(120 °C): 42.8 min (*R*), 43.5 min (*S*); 90% *ee*.

3.2 Methyl (*S*)-3-(4-bromophenyl)-3-hydroxypropanoate (**7b**, Table 2, entry 1)

With 4-bromobenzaldehyde (81.8 mg, 0.442 mmol), alcohol **7b** was obtained (78 mg, 0.30 mmol, 87% based on 79% conv).

$[\alpha]_D = -30.3$ ($c = 2.43$, CHCl_3)

m.p.: 48 °C

IR: (KBr): $\tilde{\nu} = 2996$ (w), 2954 (w), 2906 (w), 1705 (vs), 1489 (m), 1442 (m), 1303 (s), 1266 (s), 1166 (s), 1075 (m), 1062 (s), 1012 (m), 820 (m), 522 (m) cm^{-1} .

EI MS: (70 eV): m/z (%) = 260 (20) [M^+ ; ^{81}Br], 258 (22) [M^+ ; ^{79}Br], 187 (82), 185 (100), 157 (18), 105 (16), 77 (46).

HR EI MS: for $^{12}\text{C}_{10}^{1}\text{H}_{11}^{81}\text{Br}^{16}\text{O}_3$: calcd.: 259.9898
found: 259.9894.
for $^{12}\text{C}_{10}^{1}\text{H}_{11}^{79}\text{Br}^{16}\text{O}_3$: calcd.: 257.9892
found: 257.9910.

^1H -NMR: (300 MHz, CDCl_3): δ 7.48 (d, $^3J = 8.4$ Hz, 2H), 7.25 (d, $^3J = 8.4$ Hz, 2H), 5.13-5.05 (m, 1H), 3.72 (s, 3H), 3.27 (d, $^3J = 3.5$ Hz, 1H), 2.78-2.64 (m, 2H).

^{13}C -NMR: (75 MHz, CDCl_3): δ 172.6, 141.4, 131.7 (CH), 127.4 (CH), 121.6, 69.7 (CH), 52.0 (CH_3), 42.9 (CH_2).

chir GC: (155 °C): 27.3 min (*R*), 28.5 min (*S*); 81% *ee*.

¹H-NMR (300 MHz, CDCl₃)

¹³C-NMR (75 MHz, CDCl₃)

3.3 Methyl (*S*)-3-(2-bromophenyl)-3-hydroxypropanoate (**7c**, Table 2, entry 2)

With 2-bromobenzaldehyde (89.4 mg, 0.483 mmol), alcohol **7c** was obtained (80 mg, 0.31 mmol, 91% based on 71% conv).

$[\alpha]_D = -29.7$ ($c = 1.58$; acetone)

The analytical data is in accordance with the lit.²

chir GC: (150 °C): 23.3 min (*S*), 24.2 min (*R*); 78% *ee*.

^{13}C -NMR (75 MHz, CDCl_3)

3.4 Methyl (*S*)-3-(4-chlorophenyl)-3-hydroxypropanoate (**7d**, Table 2, entry 3)

With 4-chlorobenzaldehyde (80.5 mg, 0.573 mmol), alcohol **7d** was obtained (71 mg, 0.33 mmol, 68% based on 85% conv).

$[\alpha]_D = -12.9$ ($c = 1.21$; CH₃OH)

The analytical data is in accordance with the lit.³

chir GC: (160 °C): 13.9 min (*R*), 14.4 min (*S*); 80% *ee*.

¹³C-NMR (75 MHz, CDCl₃)

3.5 Methyl (*S*)-3-(4-cyanophenyl)-3-hydroxypropanoate (7e, Table 2, entry 4)

With 4-cyanobenzaldehyde (53.6 mg, 0.409 mmol), alcohol 7e was obtained (66 mg, 0.32 mmol, 88% based on 89% conv).

$[\alpha]_D = -33.7$ ($c = 1.18$; CHCl_3)

The analytical data is in accordance with the lit.⁴

chir GC: (155 °C for 30 min, then 5 °C/min for 5 min, then 160 °C): 46.8 min (*R*), 48.5 min (*S*); 72% *ee*.

¹³C-NMR (75 MHz, CDCl₃)

3.6 Methyl (*S*)-3-(4-isopropylphenyl)-3-hydroxypropanoate (**7f**, Table 2, entry 5)

With 4-isopropylbenzaldehyde (84.9 mg, 0.573 mmol), alcohol **7f** was obtained (72 mg, 0.32 mmol, 90% based on 63% conv).

$[\alpha]_D = -14.0$ ($c = 1.04$; CH₃OH)

IR: $\tilde{\nu} = 3456$ (w), 2958 (s), 2871 (m), 1732 (s), 1436 (m), 1362 (w), 1200 (m), 1159 (s), 1054 (m), 1034 (m), 832 (m) cm⁻¹.

EI MS: (70 eV): m/z (%) = 222 (13) [M⁺], 179 (28), 149 (100), 133 (17), 105 (21), 91 (14), 79 (18).

HR EI MS: for ¹²C₁₃¹H₁₈¹⁶O₃:
cald.: 222.1256
found: 222.1234.

¹H-NMR: (300 MHz, C₆D₆): δ 7.25 (dd, ³ $J = 8.2$ Hz, 2H), 7.07 (dd, ³ $J = 8.2$ Hz 2H), 5.05 (ddd, ³ $J = 9.2$ Hz, ³ $J = 3.8$ Hz, 1H), 3.25 (s, 3H), 3.05 (d, ³ $J = 3.8$ Hz, 1H), 2.78-2.59 (m, 2H), 2.46 (dd, ² $J = 16.1$ Hz, ³ $J = 3.8$ Hz, 1H), 1.13 (d, ³ $J = 7.1$ Hz, 6H).

¹³C-NMR: (75 MHz, C₆D₆): δ 172.5, 148.2, 141.1, 126.9 (CH), 126.1 (CH), 70.4 (CH), 51.2 (CH₃), 43.7 (CH₂), 34.1 (CH), 24.1 (CH₃).

chir GC: (160 °C): 14.9 min (*R*), 15.5 min (*S*); 88% *ee*.

¹H-NMR (300 MHz, C₆D₆)

¹³C-NMR (75 MHz, C₆D₆)

3.7 Methyl (*S*)-3-(4-methylthiophenyl)-3-hydroxypropanoate (7g, Table 2, entry 6)

With 4-methylthiobenzaldehyde (94.2 mg, 0.619 mmol), alcohol **7g** was obtained (74 mg, 0.33 mmol, 93% based on 57% conv).

$[\alpha]_D = -14.5$ ($c = 2.06$; CH₃OH)

m.p.: 89-90 °C

IR: $\tilde{\nu} = 3405$ (s), 2996 (w), 2952 (w), 1696 (s), 1437 (m), 1262 (m), 1247 (m), 1164 (m), 816 (m) cm⁻¹.

EI MS: (70 eV): m/z (%) = 226 (22) [M⁺], 153 (100), 109 (21).

HR EI MS: for ¹²C₁₁¹H₁₄³²S¹⁶O₃: cald.: 226.0664
found: 226.0660.

¹H-NMR: (300 MHz, C₆D₆): δ 7.14-7.06 (m, 4H), 4.96 (ddd, ³ $J = 9.2$ Hz, ³ $J = 3.8$ Hz, 1H), 3.26 (s, 3H), 3.06 (d, ³ $J = 3.8$ Hz, 1H), 2.56 (dd, ² $J = 16.0$ Hz, ³ $J = 9.2$ Hz, 1H), 2.40 (dd, ² $J = 16.0$ Hz, ³ $J = 3.8$ Hz, 1H), 1.99 (s, 3H).

¹³C-NMR: (75 MHz, C₆D₆): δ 172.3, 140.4, 138.3, 126.9 (CH), 126.5 (CH), 70.0 (CH), 51.6 (CH₃), 43.9 (CH₂), 15.8 (CH₃).

chir GC: (160 °C): 40.1 min (*R*), 41.6 min (*S*); 93% *ee*.

¹H-NMR (300 MHz, C₆D₆)

¹³C-NMR (75 MHz, C₆D₆)

3.8 Methyl (*S*)-3-(3-benzothieryl)-3-hydroxypropanoate (7h, Table 2, entry 7)

With benzothiophene-3-aldehyde (74.8 mg, 0.461 mmol), alcohol **7h** was obtained (64 mg, 0.27 mmol, 94% based on 62% conv).

$[\alpha]_D = -47.5$ ($c = 1.46$; CHCl_3)

IR: (KBr): $\tilde{\nu} = 3081$ (w), 2952 (w), 1732 (vs), 1460 (w), 1437 (m), 1279 (m), 1167 (s), 1069 (m), 762 (s), 736 (s) cm^{-1} .

EI MS: (70 eV): m/z (%) = 236 (48) $[\text{M}^+]$, 163 (100), 135 (81), 91 (19).

HR EI MS: for $^{12}\text{C}_{12}^{1}\text{H}_{12}^{16}\text{O}_3^{32}\text{S}$:
cald.: 236.0507
found: 236.0489.

$^1\text{H-NMR}$: (300 MHz, CDCl_3): δ 7.90-7.81 (m, 2H), 7.42 (s, 1H), 7.41-7.31 (m, 2H), 5.56-5.49 (td, $^3J = 6.3$ Hz, $^3J = 3.9$ Hz, 1H), 3.75 (s, 3H), 3.36 (d, $^3J = 3.9$ Hz, 1H), 2.91 (d, $^3J = 6.3$ Hz, 2H).

$^{13}\text{C-NMR}$: (75 MHz, CDCl_3): δ 172.7, 140.9, 137.3, 136.9, 124.5 (CH), 124.2 (CH), 122.9 (CH), 122.6 (CH), 122.0 (CH), 66.1 (CH), 52.0 (CH_3), 41.4 (CH_2).

chir GC: (160 °C): 58.6 min (*S*), 60.5 min (*R*); 90% *ee*.

¹H-NMR (300 MHz, CDCl₃)

¹³C-NMR (75 MHz, CDCl₃)

3.9 Methyl (*S*)-3-(2-thienyl)-3-hydroxypropanoate (**7i**, Table 2, entry 8)

With thiophene-2-aldehyde (47.8 mg, 0.426 mmol), alcohol **7i** was obtained (58 mg, 0.31 mmol, 85% based on 86% conv). $[\alpha]_D = -25.0$ ($c = 0.90$; CHCl_3)

The analytical data is in accordance with the lit.³

chir HPLC: (*n*-heptane/isopropanol 84:16; 233 nm): 40.1 min (*S*), 41.6 min (*R*); 93% *ee*.

No.	Ret.Time min	Peak Name	Height mAU	Area mAU*min	Rel.Area %
1	9,99	n.a.	822,539	267,366	50,01
2	26,02	n.a.	214,794	267,237	49,99
Total:			1037,333	534,603	100,00

No.	Ret.Time min	Peak Name	Height mAU	Area mAU*min	Rel.Area %
1	10,05	n.a.	236,247	62,174	95,90
2	28,37	n.a.	4,058	2,656	4,10
Total:			240,306	64,829	100,00

^{13}C -NMR (75 MHz, CDCl_3)

3.10 Methyl (*S*)-3-(3-thienyl)-3-hydroxypropanoate (**7j**, Table 2, entry 9)

With thiophene-3-aldehyde (54.1 mg, 0.482 mmol), alcohol **7j** was obtained (69 mg, 0.37 mmol, 87% based on 88% conv).

$[\alpha]_D = -35.3$ ($c = 1.19$; CHCl_3)

IR: (KBr): $\tilde{\nu} = 3105$ (w), 2953 (w), 1732 (vs), 1438 (m), 1280 (m), 1210 (m), 1167 (s), 1067 (m), 1034 (m), 791 (m) cm^{-1} .

EI MS: (70 eV): m/z (%) = 186 (47) [M^+], 154 (12), 137 (8), 113 (100), 85 (94).

HR EI MS: for $^{12}\text{C}_8^{1}\text{H}_{10}^{16}\text{O}_3^{32}\text{S}$:
cald.: 186.0351
found: 186.0358.

$^1\text{H-NMR}$: (300 MHz, CDCl_3): δ 7.30 (dd, $^3J = 5.0$ Hz, $^4J = 2.8$ Hz, 1H), 7.22 (dd, $^4J = 2.8$ Hz, $^4J = 1.3$ Hz, 1H), 7.06 (dd, $^3J = 5.0$ Hz, $^4J = 1.3$ Hz, 1H), 5.24-5.16 (m, 1H), 3.71 (s, 3H), 3.25 (d, $^3J = 4.0$ Hz, 1H), 2.85-2.72 (m, 2H).

$^{13}\text{C-NMR}$: (75 MHz, CDCl_3): δ 172.6, 143.9, 126.3 (CH), 125.4 (CH), 121.0 (CH), 66.7 (CH), 51.9 (CH₃), 42.4 (CH₂).

chir GC: (120 °C): 50.4 min (*R*), 51.3 min (*S*); 93% *ee*.

¹H-NMR (300 MHz, CDCl₃)

¹³C-NMR: (75 MHz, CDCl₃)

3.11 Methyl (*S*)-3-(2-furyl)-3-hydroxypropanoate (**7k**, Table 2, entry 10)

With furan-2-aldehyde (43.0 mg, 0.448 mmol), alcohol **7k** was obtained (65 mg, 0.38 mmol, 85%).

$[\alpha]_D = -24.1$ ($c = 1.10$; CHCl_3)

The analytical data is in accordance with the lit.³

chir GC: (105 °C): 35.3 min (*R*), 36.0 min (*S*); 84% *ee*.

¹³C-NMR (75 MHz, CDCl₃)

3.12 Methyl (*R*)-3-hydroxy-5,5-dimethylhexanoate (**7I**, Table 2, entry 11)

With 3,3-dimethylbutanal (53.1 mg, 0.530 mmol), alcohol **7I** was obtained (57 mg, 0.33 mmol, 81% based on 77% conv).

$[\alpha]_D = +3.7$ ($c = 0.54$; CH₃OH)

The analytical data is in accordance with the lit.⁵

chir GC: (115 °C): 6.7 min (*S*), 7.1 min (*R*); 92% *ee*.

^{13}C -NMR (75 MHz, CDCl_3)

3.13 Methyl (*R*)-3-hydroxyoctanoate (**7m**, Table 2, entry 12)

With hexanal (56.1 mg, 0.560 mmol), alcohol **7m** was obtained (49 mg, 0.28 mmol, 78% based on 64% conv).

$[\alpha]_D = -0.35$ ($c = 0.80$; CH₃OH)

The analytical data is in accordance with the lit.⁶

chir GC: (120 °C): 9.1 min (*S*), 9.4 min (*R*); 78% *ee*.

^{13}C -NMR (75 MHz, CDCl_3)

3.14 Methyl (*R*)-3-hydroxy-5-methoxy-5-methylhexanoate (**7n**, Table 2, entry 13)

With 3-methoxy-3-methylbutanal (65.0 mg, 0.560 mmol), alcohol **7n** was obtained (52 mg, 0.27 mmol, 80% based on 61% conv).

$[\alpha]_D = +0.72$ ($c = 0.83$; CH₃OH)

IR: (film) $\tilde{\nu} = 3462$ (m), 2973 (m), 2832 (m), 1735 (s), 1437 (w), 1367 (w), 1187 (m), 1160 (m), 1132 (m), 1069 (m), 837 (w) cm⁻¹.

EI MS: (70 eV): m/z (%) = 175 (0.1) [M⁺-CH₃], 157 (18), 103 (14), 73 (100).

HR EI MS: for ¹²C₈¹H₁₅¹⁶O₄ [M-CH₃]⁺: calcd.: 175.0970
found: 175.0988.

¹H-NMR: (300 MHz, C₆D₆): δ 4.57-4.44 (m, 1H), 4.06 (s, 1H), 3.35 (s, 3H), 2.78 (s, 3H), 2.59 (dd, ² $J = 15.5$ Hz, ³ $J = 7.1$ Hz, 1H), 2.32 (dd, ² $J = 15.5$ Hz, ³ $J = 5.8$ Hz, 1H), 1.64 (dd, ² $J = 14.3$ Hz, ³ $J = 10.0$ Hz, 1H), 1.30 (dd, ² $J = 14.3$ Hz, ³ $J = 2.2$ Hz, 1H), 0.99 (s, 3H), 0.87 (s, 3H).

¹³C-NMR: (75 MHz, C₆D₆): δ 172.0, 75.9, 65.9 (CH), 51.0 (CH₃), 48.7 (CH₃), 47.1 (CH₂), 42.8 (CH₂), 25.5 (CH₃), 23.2 (CH₃).

chir GC: (115 °C): 12.9 min (*S*), 13.4 min (*R*); 74% *ee*.

¹H-NMR (300 MHz, C₆D₆)

¹³C-NMR (75 MHz, C₆D₆)

3.15 Methyl (*S*)-4-benzyloxy-3-hydroxybutanoate (**7o**, Table 2, entry 14)

With 2-benzyloxyacetaldehyde (81.5 mg, 0.543 mmol), alcohol **7o** was obtained (84 mg, 0.38 mmol, 87% based on 79% conv).

$[\alpha]_D = -4.8$ ($c = 1.22$; acetone)

The analytical data is in accordance with the lit⁷

chir GC: (135 °C): 54.1 min (*R*), 55.2 min (*S*); 66% *ee*.

¹H-NMR (300 MHz, C₆D₆)

¹³C-NMR (75 MHz, C₆D₆)

3.16 (*E*)-Methyl (*S*)-3-hydroxy-5-phenyl-4-pentenoate (**7p**, Table 2, entry 15)

With cinnamylaldehyde (62.6 mg, 0.474 mmol), alcohol **7p** was obtained (43 mg, 0.21 mmol, 92% based on 48% conv). $[\alpha]_D = +4.5$ ($c = 0.45$; acetone)

The analytical data is in accordance with the lit.¹

chir HPLC: (*n*-heptane/isopropanol 90:10; 250 nm): 18.0 min (*R*), 24.6 min (*S*); 71% *ee*.

No.	Ret.Time min	Peak Name	Height mAU	Area mAU*min	Rel.Area %
1	17,75	n.a.	389,248	249,168	50,35
2	24,24	n.a.	292,590	245,751	49,65
Total:			681,838	494,918	100,00

No.	Ret.Time min	Peak Name	Height mAU	Area mAU*min	Rel.Area %
1	18,00	n.a.	39,019	24,556	14,42
2	24,56	n.a.	170,941	145,749	85,58
Total:			209,960	170,305	100,00

^{13}C -NMR (75 MHz, CDCl_3)

3.17 Ethyl (*R*)-3-phenyl-2,2-difluoro-3-hydroxypropanoate (**8a**, Table 3, entry 1)

With benzaldehyde (77.9 mg, 0.734 mmol), alcohol **8a** was obtained (127 mg, 0.552 mmol, 75% based on 82% conv). $[\alpha]_D = -3.30$ ($c = 1.15$; acetone)

The analytical data is in accordance with the lit.⁸

chir HPLC: (*n*-heptane/isopropanol 96:4; 215 nm): 20.5 min (*S*), 22.1 min (*R*); 88% *ee*.

No.	Ret.Time min	Peak Name	Height mAU	Area mAU*min	Rel.Area %
1	22,14	n.a.	194,847	150,311	49,77
2	23,70	n.a.	177,525	151,704	50,23
Total:			372,373	302,015	100,00

No.	Ret.Time min	Peak Name	Height mAU	Area mAU*min	Rel.Area %
1	20,45	n.a.	11,903	7,311	5,92
2	22,08	n.a.	145,744	116,096	94,08
Total:			157,647	123,407	100,00

3.18 Ethyl (*R*)-3-(4-bromophenyl)-2,2-difluoro-3-hydroxypropanoate (8b**, Table 3, entry 2)**

With 4-bromobenzaldehyde (86.0 mg, 0.465 mmol), alcohol **8b** was obtained (114 mg, 0.369 mmol, 83% based on 96% conv).

$[\alpha]_D = -1.2$ ($c = 0.35$; CHCl_3)

IR: (KBr): $\tilde{\nu} = 2986$ (m), 2940 (w), 1759 (s), 1490 (m), 1402 (w), 1376 (m), 1315 (m), 1192 (m), 1108 (s), 1074 (s), 1012 (m), 855 (m), 830 (m), 779 (m), 740 (m) cm^{-1} .

EI MS: (70 eV): m/z (%) = 310 (4) [M^+ ; ^{81}Br], 308 (5) [M^+ ; ^{79}Br], 187 (97), 185 (100), 159 (11), 157 (14), 136 (6), 77 (32).

HR EI MS: for $^{12}\text{C}_{11}^{1}\text{H}_{11}^{81}\text{Br}^{19}\text{F}_2^{16}\text{O}_3$: calcd.: 309.9849

found: 309.9849.

for $^{12}\text{C}_{11}^{1}\text{H}_{11}^{79}\text{Br}^{19}\text{F}_2^{16}\text{O}_3$: calcd.: 307.9860

found: 307.9828.

^1H -NMR: (600 MHz, CDCl_3): δ 7.52 (d, $^3J = 8.3$ Hz, 2H), 7.32 (d, $^3J = 8.3$ Hz, 2H), 5.14 (dd, $^3J(\text{H},\text{F}) = 15.5$ Hz, $^3J(\text{H},\text{F}) = 7.4$ Hz, 1H), 4.31 (q, $^3J = 7.2$ Hz, 2H), 2.8 (s, br, 1H), 1.30 (t, $^3J = 7.2$ Hz, 3H).

^{13}C -NMR: (151 MHz, CDCl_3): δ 163.3 (t, $^2J(\text{C},\text{F}) = 31.7$ Hz), 133.4, 131.6 (CH), 129.4 (CH), 123.5, 113.3 (dd, $^1J(\text{C},\text{F}) = 261$ Hz, $^1J(\text{C},\text{F}) = 255$ Hz), 73.2 (t, dd, $^2J(\text{C},\text{F}) = 28.2$ Hz, $^2J(\text{C},\text{F}) = 24.7$ Hz, CH), 63.3 (CH_2), 13.9 (CH_3).

chir GC: (160 $^\circ\text{C}$): 28.4 min (*S*), 29.4 min (*R*); 87% *ee*.

¹H-NMR (600 MHz, CDCl₃)

¹³C-NMR (151 MHz, CDCl₃)

3.19 Ethyl (*R*)-3-(4-cyanophenyl)-2,2-difluoro-3-hydroxypropanoate (**8c**, Table 3, entry 3)

With 4-cyanobenzaldehyde (58.7 mg, 0.448 mmol), alcohol **8c** was obtained (92 mg, 0.36 mmol, 88% based on 91% conv).

$[\alpha]_D = -5.7$ ($c = 0.46$; CHCl_3)

The analytical data is in accordance with the lit.⁹

chir GC: (160 °C): 50.5 min (*S*), 51.6 min (*R*); 84% *ee*.

^{13}C -NMR (75 MHz, CDCl_3)

3.20 Ethyl (*S*)-3-(2-thienyl)-2,2-difluoro-3-hydroxypropanoate (**8d**, Table 3, entry 4)

With thiophene-2-aldehyde (52.1 mg, 0.465 mmol), alcohol **8d** was obtained (65 mg, 0.28 mmol, 90% based on 66% conv).

$[\alpha]_D = -11.0$ ($c = 0.58$; CHCl_3)

The analytical data is in accordance with the lit.¹⁰

chir HPLC: (*n*-heptane/isopropanol 92:8; 235 nm): 23.0 min (*S*), 28.1 min (*R*); 90% *ee*.

No.	Ret.Time min	Peak Name	Height mAU	Area mAU*min	Rel.Area %
1	23,09	n.a.	27,542	22,885	50,17
2	27,99	n.a.	23,015	22,727	49,83
Total:			50,557	45,612	100,00

No.	Ret.Time min	Peak Name	Height mAU	Area mAU*min	Rel.Area %
1	23,05	n.a.	84,921	75,656	95,01
2	28,14	n.a.	4,837	3,972	4,99
Total:			89,758	79,627	100,00

¹H-NMR (600 MHz, CDCl₃)

¹³C-NMR (151 MHz, CDCl₃)

3.21 Ethyl (*R*)-3-(3-thienyl)-2,2-difluoro-3-hydroxypropanoate (**8e**, Table 3, entry 5)

With thiophene-3-aldehyde (52.1 mg, 0.465 mmol), alcohol **8e** was obtained (64 mg, 0.27 mmol, 84% based on 69% conv).

$[\alpha]_D = -2.9$ ($c = 0.82$; acetone)

IR: (KBr): $\tilde{\nu} = 3114$ (w), 2986 (m), 1759 (vs), 1375 (m), 1307 (m), 1202 (m), 1105 (s), 1073 (s), 776 (m), 691 (m) cm^{-1} .

EI MS: (70 eV): m/z (%) = 236 (16) $[\text{M}^+]$, 143 (7), 113 (100), 85 (34).

HR EI MS: for $^{12}\text{C}_9^{1}\text{H}_{10}^{19}\text{F}^{16}\text{O}_3^{32}\text{S}$:
calcd.: 236.0319
found: 236.0291.

^1H -NMR: (300 MHz, C_6D_6): δ 7.04-6.96 (m, 2H), 6.80 (dd, $^3J = 4.9$ Hz, $^4J = 3.1$ Hz, 1H), 5.13-5.01 (m, 1H), 3.21 (q, $^3J = 7.1$ Hz, 2H), 2.27 (d, br, $^3J = 4.9$ Hz, 1H), 0.76 (t, $^3J = 7.1$ Hz, 3H).

^{13}C -NMR: (75 MHz, C_6D_6): δ 163.6 (dd, $^2J(\text{C},\text{F}) = 32.6$ Hz, $^2J(\text{C},\text{F}) = 30.8$ Hz), 136.5, 126.9 (CH), 126.0 (CH), 124.7 (CH), 114.4 (dd, $^1J(\text{C},\text{F}) = 260$ Hz, $^1J(\text{C},\text{F}) = 254$ Hz), 70.7 (dd, $^2J(\text{C},\text{F}) = 28.9$ Hz, $^2J(\text{C},\text{F}) = 24.7$ Hz, CH), 62.7 (CH_2), 13.6 (CH_3).

chir GC: (140 °C): 26.5 min (*S*), 26.8 min (*R*); 87% *ee*.

¹H-NMR (300 MHz, C₆D₆)

¹³C-NMR (75 MHz, C₆D₆)

3.22 Ethyl (*R*)-2,2-difluoro-3-hydroxy-5,5-dimethylhexanoate (**8f**, Table 3, entry 6)

With 3,3-dimethylbutanol (47.5 mg, 0.474 mmol), alcohol **8f** was obtained (78 mg, 0.35 mmol, 81% based on 91% conv).

$[\alpha]_D = +17.3$ ($c = 1.07$; acetone)

IR: (KBr): $\tilde{\nu} = 2958$ (s), 2872 (w), 1760 (s), 1718 (m), 1470 (w), 1370 (m), 1325 (m), 1218 (m), 1104 (s), 1077 (s), 856 (w), 700 (w) cm^{-1} .

EI MS: (70 eV): m/z (%) = 225 (1) $[\text{M}+\text{H}]^+$, 191 (25), 163 (19), 124 (32), 101 (51), 57 (100).

HR EI MS: for $^{12}\text{C}_{10}^{1}\text{H}_{19}^{19}\text{F}^{16}\text{O}_3^+$:
calcd.: 225.1302
found: 225.1301 $[\text{M}+\text{H}]^+$.

^1H -NMR: (300 MHz, C_6D_6): δ 4.22-4.05 (m, 1H), 3.88 (q, $^3J = 7.3$ Hz, 2H), 1.64 (d, $^2J = 14.7$ Hz, 2H), 1.55 (s, br, 1H), 1.33 (dd, $^2J = 14.7$ Hz, $^3J = 9.5$ Hz, 1H), 0.84 (t, $^3J = 7.3$ Hz, 3H), 0.83 (s, 9H).

^{13}C -NMR: (75 MHz, C_6D_6): δ 163.9 (dd, $^2J(\text{C},\text{F}) = 32.6$ Hz, $^2J(\text{C},\text{F}) = 31.4$ Hz), 115.7 (dd, $^1J(\text{C},\text{F}) = 257$ Hz, $^1J(\text{C},\text{F}) = 255$ Hz), 70.0 (dd, $^2J(\text{C},\text{F}) = 27.0$ Hz, $^2J(\text{C},\text{F}) = 24.7$ Hz, CH), 62.6 (CH₂), 42.9 (CH₂), 29.8 (3CH₃), 13.7 (CH₃).

chir GC: (130 °C): 5.0 min (*S*), 5.2 min (*R*); 80% *ee*.

¹H-NMR (300 MHz, C₆D₆)

¹³C-NMR (75 MHz, C₆D₆)

-
- ¹ Denmark, S. E.; Beutner, G. L.; Wynn, T.; Eastgate, M. D., *J. Am. Chem. Soc.* **2005**, *127*, 3774-3789.
- ² Ruck, R. T.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2002**, *124*, 2882-2883.
- ³ Ratovelomanana-Vidal, V.; Girard, C.; Touati, R.; Tranchier, J. P.; Hassine, B. B.; Genêt, J. P. *Adv. Synth. Catal.* **2003**, *345*, 261-274.
- ⁴ Suh, Y.; Rieke, R. D. *Tetrahedron Lett.* **2004**, *45*, 1807-1809. (racemic)
- ⁵ Sugimura, T.; Nakagawa, S.; Tai, A. *Bull Chem Soc. Jpn.* **2002**, *75*, 355-363.
- ⁶ Utaka, M.; Watabu, H.; Higashi, H.; Sakai, T.; Tsuboi, S.; Torii, S. *J. Org. Chem.* **1999**, *55*, 3917-3921.
- ⁷ a) Hinterding, K.; Jacobsen, E. R. *J. Org. Chem.* **1999**, *64*, 2164-2165. b) Hoffmann, T.; Seebach, D. *Liebigs Ann.* **1996**, *8*, 1277-1282.
- ⁸ Andrés, J. M.; Martínez, M. M.; Pedrosa, R.; Pérez-Encabo, A. *Synthesis* **1996**, 1070-1072.
- ⁹ Jimenez, O.; Bosch, Maria P.; Guerrero, A. *Synthesis* **2000**, 1917-1924. (racemic)
- ¹⁰ Linderman, R. J.; Graves, D. M. *J. Org. Chem.* **1989**, *54*, 661-668. (racemic)