An Intramolecular Diels-Alder Approach to the Eunicellins: Enantioselective Total Synthesis of Ophirin B

Michael T. Crimmins* and Brandon H. Brown

Venable and Kenan Laboratories of Chemistry

The University of North Carolina at Chapel Hill

Chapel Hill, North Carolina 27599-3290

crimmins@email.unc.edu

Experimental Section

Materials and Methods: General. Infrared (IR) spectra were obtained using a Jasco 460 Plus Fourier transform infrared spectrometer. Proton and carbon nuclear magnetic resonance (^1H and ^{13}C NMR) spectra were recorded on the following instruments: Bruker model Avance 400 (^1H at 400 MHz; ^{13}C at 100 MHz) and Bruker model Avance 500 (^1H at 500 MHz; ^{13}C at 125 MHz). Optical rotations were determined using a Jasco P1010 polarimeter. Thin layer chromatography (TLC) was conducted on silica gel F₂₅₄ TLC plates purchased from Scientific Adsorbents, Inc. Flash column chromatography was carried out using silica gel (32 to 63 μm) purchased from Scientific Adsorbents, Inc. Diethyl ether (Et_2O), tetrahydrofuran (THF), and dichloromethane (CH_2Cl_2) were dried by being passed through a column of neutral alumina under nitrogen immediately prior to use. Alkylamines, benzene, and toluene were distilled from calcium hydride immediately prior to use. Dimethyl sulfoxide (DMSO) was distilled from calcium hydride under reduced pressure and stored over 4 Å molecular sieves. Anhydrous *N,N*-dimethylformamide (DMF) was purchased from Aldrich chemical company in 1L Sure/SealTM bottles. Pivaloyl chloride was distilled and stored over 4 Å molecular sieves. Methallyl iodide was distilled and stored over copper wire at -20°C . Acetic anhydride and methyl iodide were distilled and stored under a blanket of argon. Titanium (IV) isopropoxide and (+)-diethyl tartrate were distilled under reduced pressure and stored in a dark desiccator. All other reagents and solvents were used as received from the manufacturer. All air and water sensitive reactions were preformed in flasks flame dried under positive flow argon and conducted under an argon atmosphere.

(2S)-1-Benzyloxy-but-3-en-2-ol. Into a flask equipped with a Dean-Stark trap was added trimethylsulfonium iodide (37.0g, 183.0 mmol). Then 50 mL of toluene was added and distilled to azeotrope any water. After distillation of the toluene, the salt was taken up in 350 mL of THF and cooled to -8°C (acetone/ice bath). *n*-Butyl lithium (1.56 M in hexanes, 110.0 mL, 165.0 mmol) was then added dropwise via addition funnel over 30 minutes. After stirring the reaction for 30 minutes at -8°C , (*S*)-benzyl glycidyl ether **7** (10.0 g, 61.0 mmol) in 150 mL of THF was added via an addition funnel over 30 minutes at such a rate as to maintain the temperature below -5°C . After the addition was

complete, the cooling bath was removed and the reaction was allowed to warm to ambient temperature over 30 minutes. After stirring at room temperature for 30 minutes, the reaction was quenched by the addition of water. The mixture was diluted in 300 mL of diethyl ether and the organic layer was washed with brine. The aqueous layers were back extracted with diethyl ether. The organic layers were combined, dried over Na₂SO₄, and concentrated in vacuo. Purification by flash column chromatography (10% then 30% EtOAc/hexanes) provided 10.76 g (99%) of the allylic alcohol as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 3.36 (dd, *J* = 9.6, 8 Hz, 1H), 3.53 (dd, *J* = 9.6, 3.2 Hz, 1H), 4.31-4.36 (m, 1H), 4.56 (s, 2H), 5.18 (ddd, *J* = 10.8, 1.6, 1.6 Hz, 1H), 5.35 (ddd, *J* = 17.2, 1.2, 1.2 Hz, 1H), 5.82 (ddd, *J* = 17.2, 10.8, 5.6 Hz, 1H), 7.27-7.37 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ 71.74, 73.60, 74.91, 116.73, 127.99, 128.06, 128.70, 136.72, 138.02; IR (film) 3433 (br), 2861, 1454 cm⁻¹; [α]_D²³ = -2.95 (c = 3.3, CHCl₃); MS (electrospray ionization) calculated for C₁₁H₁₄O₂Na [M + Na]⁺: 201.09, found: 201.03.

(2S)-1-Benzyloxy-2-(4-methoxy-benzyloxy)-but-3-ene (8). Into a flask equipped with a mechanical stirrer and an addition funnel was added sodium hydride (60% dispersion in mineral oil, 16.2 g, 404 mmol). The sodium hydride was rinsed with hexanes three times, diluted in 400 mL of THF and cooled to 0 °C. The allylic alcohol (36.0 g, 202 mmol) in 400 mL of THF was added dropwise via addition funnel. After stirring for 30 minutes at 0 °C, 500 mL of DMF was added. This was followed by *p*-methoxybenzylchloride (63.3 g, 404 mmol) in 200 mL of THF via addition funnel. The orange slurry was then warmed to room temperature and stirred overnight. The reaction was quenched by the slow addition of saturated aqueous NH₄Cl and diluted with diethyl ether. The organic layer was washed with water and brine, dried over Na₂SO₄, and concentrated in vacuo. Purification by flash column chromatography (5% to 20% EtOAc/hexanes) provided 57.3 g (95%) of alkene **8** as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 3.49-3.59 (m, 2H), 3.79 (s, 3H), 3.99-4.04 (m, 1H), 4.49 (AB *J*_{AB} = 11.2 Hz, Δ*v*_{AB} = 70.9 Hz, 2H), 4.55 (AB *J*_{AB} = 12 Hz, Δ*v*_{AB} = 12.3 Hz, 2H), 5.29-5.30 (m, 2H), 5.30 (dddd, *J* = 28, 2.8, 2, 1.2 Hz, 1H), 5.79 (ddd, *J* = 17.2, 10.4, 7.2 Hz, 1H), 6.83-6.87 (m, 2H), 7.24-7.32 (m, 7H); ¹³C NMR (100 MHz, CDCl₃) δ 55.49, 70.43, 73.31, 73.55, 79.26, 113.95, 118.49, 127.74, 127.83, 128.53, 129.54, 130.83, 136.1, 138.59, 159.31; IR (film) 2860, 1613, 1513, 1454, 1274, 1092 cm⁻¹; [α]_D²³ = +20.2 (c = 4, CHCl₃); MS (electrospray ionization) calculated for C₁₉H₂₂O₃Na [M + Na]⁺: 321.15, found: 321.12.

(3R)-4-Benzyloxy-3-(4-methoxy-benzyloxy)-butan-2-one (6). : A solution of alkene **8** (.500g, 1.67 mmol) and mercuric acetate (.640g, 2.01 mmol) in 12 mL of THF and 4 mL of water was stirred at ambient temperature for 12 hours. The reaction was then transferred to a solution of LiCl (.140g, 3.36 mmol), PdCl₂ (.300g, 1.67 mmol) and CuCl₂ (.680g, 5.04 mmol) in 12 mL THF and 4 mL water and stirred at 55 °C for 3 hours. Saturated NaHCO₃ was added and the product was extracted with diethyl ether three times. The combined organic layers were dried over Na₂SO₄ and concentrated in vacuo. Purification by flash column chromatography (10% then 25% EtOAc/hexanes) afforded .467 g (89%) of ketone **6** as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 2.19 (s, 3H), 3.70 (d, *J* = 4.8 Hz, 2H), 3.78 (s, 3H), 3.94 (dd, *J* = 4.4, 4.4 Hz, 1H), 4.52 (s, 2H), 4.54 (AB *J*_{AB} = 11.6 Hz, Δ*v*_{AB} = 28.8 Hz, 2H), 6.85-6.87 (m, 2H), 7.23-7.33 (m, 7H); ¹³C NMR (100 MHz, CDCl₃) δ 27.55, 55.70, 70.71, 72.73, 73.92, 84.03, 114.31, 128.08, 128.14, 128.80, 129.84, 130.06, 138.18, 159.88, 209.89; IR (film) 2863, 1716 (str), 1612, 1514, 1249, 1103, 1033 cm⁻¹; [α]_D²³ = +24.2 (c = 2.35, CHCl₃); MS (electrospray ionization) calculated for C₁₉H₂₂O₄Na [M + Na]⁺: 337.14, found: 337.07.

(2R,3R)-1-benzyloxy-2-(4-methoxy-benzyloxy)-3-methyl-hept-6-en-3-ol (9). A flask equipped with an addition funnel was charged with freshly prepared butenyl magnesium bromide (1.0 M in THF, 244 mL, 244 mmol) in 700 mL of THF. After the Grignard was cooled to -78°C , ketone **6** (35g, 112 mmol) was diluted in 300 mL of THF and added to the Grignard dropwise via addition funnel over 1.5 hours. After the reaction was stirred at -78°C for 2 hours, saturated NH_4Cl was added slowly and the mixture was allowed to warm to room temperature. The mixture was diluted in diethyl ether and washed with water and brine. The aqueous layers were back extracted with diethyl ether and the combined organic layers were dried over Na_2SO_4 and concentrated in vacuo. Purification by flash column chromatography (10% then 25% EtOAc/hexanes) afforded 38 g (92%) of tertiary alcohol **9** as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 1.11 (s, 3H), 1.51-1.65 (m, 2H), 2.01-2.19 (m, 2H), 2.69 (br s, 1H), 3.47 (dd, $J = 5.6$, 4 Hz, 1H), 3.60-3.75 (m, 2H), 3.79 (s, 3H), 4.53 (s, 2H), 4.64 (AB $J_{\text{AB}} = 10.8$ Hz, $\Delta\nu_{\text{AB}} = 103$ Hz, 2H), 4.92 (dddd, $J = 10$, 1.2, 1.2, 1.2 Hz, 1H), 4.98 (dddd, $J = 16.8$, 1.6, 1.6, 1.6 Hz, 1H), 5.79 (dddd, $J = 16.8$, 10, 6.4, 6.4 Hz, 1H), 6.84-6.87 (m, 2H), 7.23-7.36 (m, 7H); ^{13}C NMR (100 MHz, CDCl_3) δ 22.99, 27.97, 37.93, 55.47, 70.98, 73.51, 73.75, 73.90, 82.86, 114.00, 114.40, 127.84, 127.94, 128.65, 129.81, 130.72, 138.14, 139.24, 159.47; IR (film) 3494, 2932, 1612, 1514, 1248, 1086, 1035 cm^{-1} ; $[\alpha]_{\text{D}}^{23} = -16.9$ ($c = 1.5$, CHCl_3); MS (electrospray ionization) calculated for $\text{C}_{23}\text{H}_{30}\text{O}_4\text{Na}$ [$\text{M} + \text{Na}$] $^{+}$: 393.20, found: 393.28.

(2R,3R)-1,3-bis-benzyloxy-2-(4-methoxy-benzyloxy)-3-methyl-hept-6-ene (10). Into a flask equipped an addition funnel was added potassium hydride (30% dispersion in mineral oil, 27.3 g, 204 mmol). The potassium hydride was rinsed with hexanes then diluted in 300 mL of THF and cooled to 0°C . The tertiary alcohol **9** (37.8 g, 102 mmol) in 300 mL of THF was added dropwise via addition funnel. After stirring for 30 minutes at 0°C , 300 mL of DMF was added. This was followed by benzylbromide (15.5 mL, 128 mmol) in 100 mL of THF via addition funnel. The orange slurry was then warmed to room temperature and stirred overnight. The reaction was quenched by the slow addition of saturated aqueous NH_4Cl and diluted with diethyl ether. The organic layer was washed with water and brine, dried over Na_2SO_4 , and concentrated in vacuo. Purification by flash column chromatography (5% to 20% EtOAc/hexanes) provided 43.4 g (92%) of alkene **10** as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 1.33 (s, 3H), 1.67-1.82 (m, 2H), 2.13-2.21 (m, 2H), 3.71-3.81 (m, 2H), 3.82 (s, 3H), 3.98 (dd, $J = 10$, 2 Hz, 1H), 4.52 (AB $J_{\text{AB}} = 8.8$ Hz, $\Delta\nu_{\text{AB}} = 19.4$ Hz, 2H), 4.57 (AB $J_{\text{AB}} = 12$ Hz, $\Delta\nu_{\text{AB}} = 7.2$ Hz, 2H), 4.75 (AB $J_{\text{AB}} = 10.8$ Hz, $\Delta\nu_{\text{AB}} = 94.8$ Hz, 2H), 4.95 (dddd, $J = 10$, 1.2, 1.2, 1.2 Hz, 1H), 5.02 (dddd, $J = 17.2$, 1.6, 1.6, 1.6 Hz, 1H), 5.84 (dddd, $J = 16.8$, 10.4, 6.8, 6.8 Hz, 1H), 6.87-6.91 (m, 2H), 7.23-7.38 (m, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ 21.13, 27.91, 34.40, 55.42, 64.50, 72.34, 73.58, 73.78, 78.88, 83.50, 113.78, 114.33, 127.26, 127.39, 127.71, 127.76, 128.40, 128.56, 129.55, 131.41, 138.63, 139.26, 139.81, 158.17; IR (film) 2932, 1639, 1612, 1513, 1454, 1248 cm^{-1} ; $[\alpha]_{\text{D}}^{23} = -9.30$ ($c = 2.3$, CHCl_3); MS (electrospray ionization) calculated for $\text{C}_{30}\text{H}_{36}\text{O}_4\text{Na}$ [$\text{M} + \text{Na}$] $^{+}$: 483.25, found: 483.26.

(2R,3R)-1,3-Bis-benzyloxy-3-methyl-hept-6-en-2-ol. Into a flask equipped with a reflux condenser was added alkene **10** (43.7g, 94.2 mmol) in 600 mL of methanol. Concentrated HCl (2mL) was added and the reaction was heated to reflux for 2 hours. Once the reaction was complete by TLC, a saturated NaHCO₃ solution was added until the reaction was basic. The mixture was concentrated in vacuo and then diluted in diethyl ether. The organic layer was washed with water and brine. The aqueous layers were back extracted with diethyl ether and the combined organic layers were dried over Na₂SO₄ and concentrated in vacuo. Purification by flash column chromatography (hexanes to 15% EtOAc/hexanes) provided 26.5 g (83%) of the alcohol as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 1.25 (s, 3H), 1.68-1.84 (m, 2H), 2.09-2.26 (m, 2H), 2.69 (br s, 1H), 3.57 (dd, *J* = 10, 8 Hz, 1H), 3.69 (dd, *J* = 10, 3.2 Hz, 1H), 3.99 (dd, *J* = 8, 3.2 Hz, 1H), 4.49 (AB *J*_{AB} = 11.2 Hz, Δ*v*_{AB} = 19.4 Hz, 2H), 4.59 (AB *J*_{AB} = 12 Hz, Δ*v*_{AB} = 22.0 Hz, 2H), 4.97 (dddd, *J* = 10, 1.2, 1.2, 1.2 Hz, 1H), 5.03 (dddd, *J* = 17.2, 1.6, 1.6, 1.6 Hz, 1H), 5.84 (dddd, *J* = 16.8, 10, 6.4, 6.4 Hz, 1H), 7.26-7.38 (m, 10H); ¹³C NMR (100 MHz, CDCl₃) δ 19.60, 27.79, 34.02, 63.92, 71.15, 73.58, 74.37, 78.47, 114.62, 127.49, 127.86, 127.87, 128.51, 128.58, 138.21, 138.79, 138.82, 139.22; IR (film) 3472, 2926, 1640, 1454, 1382 cm⁻¹; [α]_D²³ = +7.30 (c = 3.7, CHCl₃); MS (electrospray ionization) calculated for C₂₂H₂₈O₃Na [M + Na]⁺: 363.19, found: 363.11.

(1R,2R)-(2-Benzyloxy-1-benzyloxymethyl-2-methyl-hex-5-enyloxy)-acetic acid (5). Sodium hydride (60% dispersion in mineral oil 9.31g, 233 mmol) in a flask was washed twice with hexanes, suspended in 40 mL of THF and cooled to 0 °C. Bromoacetic acid (11.9 g, 85.3 mmol) in 40mL of THF was added dropwise, and the mixture was stirred for 15 minutes. The secondary alcohol (26.4 g, 77.5 mmol) in 80 mL of DMF was then added dropwise to the suspension. The reaction mixture was allowed to stir at room temperature for 24 hours. The reaction was quenched by the slow addition of water and diluted with diethyl ether. The aqueous layer was acidified by the addition of 10% H₂SO₄, extracted three times with ethyl acetate and the combined organic extracts were dried over Na₂SO₄ and concentrated in vacuo. Purification by flash column chromatography (5% to 30% EtOAc/hexanes) afforded 29.8 g (97%) of acid **5** as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 1.25 (s, 3H), 1.42 (ddd, *J* = 14.8, 12, 5.2, Hz, 1H), 1.81 (ddd, *J* = 14.8, 12, 4, Hz, 1H), 1.91-2.02 (m, 1H), 2.08-2.20 (m, 1H), 3.50-3.68 (m, 3H), 4.32 (AB *J*_{AB} = 17.6 Hz, Δ*v*_{AB} = 28.2 Hz, 2H), 4.42 (AB *J*_{AB} = 11.2 Hz, Δ*v*_{AB} = 7.84 Hz, 2H), 4.58 (AB *J*_{AB} = 12 Hz, Δ*v*_{AB} = 25.7 Hz, 2H), 4.91-4.94 (m, 1H), 4.96 (dddd, *J* = 10, 1.6, 1.6, 1.6 Hz, 1H), 5.71 (dddd, *J* = 16.4, 10, 6.4, 6.4 Hz, 1H), 7.25-7.38 (m, 10H), 11.32 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 19.42, 27.12, 34.20, 64.22, 69.98, 70.62, 73.98, 79.68, 85.55, 115.24, 128.02, 128.06, 128.13, 128.45, 128.76, 128.88, 134.96, 136.91, 137.86, 172.58; IR (film) 3423 (br), 2929, 1762 (str), 1640, 1454, 1365, 1127 cm⁻¹; [α]_D²³ = +11.9 (c = 1.85, CHCl₃); MS (electrospray ionization) calculated for C₂₄H₃₀O₅Na [M + Na]⁺: 421.20, found: 421.17.

(4S)-3-{2-[(1R,2R)-2-Benzyloxy-1-benzyloxymethyl-2-methyl-hex-5-enyloxy]-acetyl}-4-isopropyl-oxazolidin-2-one (11). A flask was charged with a solution of carboxylic acid **5** (29.8 g, 74.8 mmol) in 450 mL of THF. Triethylamine (11.5 mL, 82.3 mmol) was added via syringe and the mixture was cooled to -78 °C. Pivaloyl chloride (10.1 mL, 82.3 mmol) was added dropwise via syringe over 15 minutes and then the mixture was warmed to 0 °C, where it was stirred for 1 hour and subsequently cooled to -78 °C. In a separate flask, (S)-(-)-4-isopropyl-2-oxazolidinone (12.1 g, 93.5mmol) was dissolved in 500 mL of THF and cooled to -78 °C. *n*-Butyl lithium (1.6M in hexanes, 54 mL, 86 mmol) was added dropwise via syringe and the mixture was stirred for 1 hour. The lithiated

oxazolidinone was added via cannula to the mixed anhydride and the reaction was stirred for an additional 15 minutes before being warmed to 0 °C where stirring continued for 1 hour. The reaction was quenched by the addition of water and extracted twice with ethyl acetate. The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated in vacuo. Purification by flash column chromatography (5% to 30% EtOAc/hexanes) provided 33.9 g (89%) of acyl oxazolidinone **11** as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 0.75 (d, 6.8 Hz, 3H), 0.81 (d, 7.2 Hz, 3H), 1.35 (s, 3H), 1.65-1.85 (m, 2H), 2.12-2.20 (m, 2H), 2.29-2.37 (m, 1H), 3.63 (dd *J* = 8.4, 8.4 Hz, 1H), 3.72-3.81 (m, 2H), 3.91-3.94 (m, 1H), 3.98-4.06 (m, 2H), 4.42 (AB *J*_{AB} = 12.4 Hz, Δ*v*_{AB} = 27.3 Hz, 2H), 4.54 (AB *J*_{AB} = 11.2 Hz, Δ*v*_{AB} = 41.7 Hz, 2H), 4.91 (dddd, *J* = 10.4, 1.6, 1.6, 1.6 Hz, 1H), 4.95 (AB *J*_{AB} = 17.6 Hz, Δ*v*_{AB} = 78.6 Hz, 2H), 5.00 (dddd, *J* = 17.2, 1.6, 1.6, 1.6 Hz, 1H), 5.81 (dddd, *J* = 17.2, 10.4, 6.8, 6.8 Hz, 1H), 7.19-7.36 (m, 10H); ¹³C NMR (100 MHz, CDCl₃) δ 14.50, 17.94, 21.25, 27.94, 28.14, 34.31, 58.20, 64.01, 64.51, 72.11, 72.20, 72.82, 78.52, 85.54, 114.36, 126.73, 127.23, 127.35, 127.43, 128.36, 128.50, 138.70, 139.10, 139.61, 154.09, 170.73; IR (film) 2965, 1781 (str), 1719 (str), 1389, 1258, 1210, 1139 cm⁻¹; [α]_D²³ = +40.0 (c = 2.6, CHCl₃); MS (electrospray ionization) calculated for C₃₀H₃₉NO₆Na [M + Na]⁺: 532.27, found: 532.16.

(4S)-3-((2R)-2-((1R,2R)-2-Benzyloxy-1-benzyloxymethyl-2-methyl-hex-5-enyloxy)-4-methyl-pent-4-enoyl)-4-isopropyl-oxazolidin-2-one (4). A flask equipped with a low-temperature thermometer was charged sodium bis(trimethylsilyl)amide (0.75 M in toluene/THF, 30.0 mL, 22.5 mmol) and 75 mL of THF. After the flask was cooled to -78 °C, acyl oxazolidinone **11** (7.65 g, 15.0 mmol) in 50 mL was added dropwise via an addition funnel at a rate as to maintain the reaction temperature below -65 °C. After stirring for 1 hour at -78 °C, methallyl iodide (13.65 g, 75.0 mmol) in 25 mL of THF was added dropwise via syringe. After 15 minutes, the reaction was warmed to -45 °C and stirred at that temperature for 1 hour. The reaction was quenched by the slow addition of half-saturated NH₄Cl and warmed to room temperature. The solution was extracted twice with 50% ethyl acetate/hexanes. The combined organic extracts were washed with brine, dried over Na₂SO₄, and concentrated in vacuo. Purification by flash column chromatography (5% to 25% EtOAc/hexanes) provided 7.86 g (93%) of diene **4** as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 0.71 (d, *J* = 6.8 Hz, 3H), 0.75 (d, *J* = 7.2 Hz, 3H), 1.33 (s, 3H), 1.68-1.81 (m, 2H), 1.85 (s, 3H), 2.12-2.25 (m, 3H), 2.33-2.50 (m, 2H), 2.83 (dd, *J* = 8.8, 8.8 Hz, 1H), 3.68-3.74 (m, 3H), 3.88-3.94 (m, 2H), 4.34 (AB *J*_{AB} = 12.4 Hz, Δ*v*_{AB} = 27.8 Hz, 2H), 4.53 (AB *J*_{AB} = 11.6 Hz, Δ*v*_{AB} = 65.4 Hz, 2H), 4.80-4.84 (m, 2H), 4.92 (dddd, *J* = 10.4, 1.2, 1.2, 1.2 Hz, 1H), 5.01 (dddd, *J* = 17.2, 1.6, 1.6, 1.6 Hz, 1H), 5.50 (dd, *J* = 9.6, 3.6 Hz, 1H), 5.82 (dddd, *J* = 16.8, 10.4, 6.8, 6.8 Hz, 1H), 7.20-7.36 (m, 10H); ¹³C NMR (100 MHz, CDCl₃) δ 14.70, 17.98, 21.72, 22.27, 28.02, 28.12, 34.19, 42.39, 58.22, 63.07, 64.65, 72.25, 72.46, 77.55, 78.66, 79.14, 86.57, 114.23, 114.29, 126.12, 127.33, 127.47, 128.44, 128.75, 138.84, 139.36, 139.68, 141.51, 153.88, 173.29; IR (film) 2967, 1778 (str), 1711 (str), 1453, 1387, 1120 cm⁻¹; [α]_D²³ = +41.51 (c = 3.25, CHCl₃); MS (electrospray ionization) calculated for C₃₄H₄₅NO₆Na [M + Na]⁺: 586.31, found: 586.26.

(2S)-2-[(1R,2R)-2-benzyloxy-1-benzyloxymethyl-2-methyl-hex-5-enyloxy]-4-methyl-pent-4-en-1-ol (13). Acyl oxazolidinone **4** (.438 g, .777 mmol) and anhydrous methanol (.094 mL, 2.33 mmol) in 10 mL of diethyl ether were added to a flask and cooled to 0 °C. Lithium borohydride (2.0M in THF, 1.17 mL, 2.33mmol) was added dropwise via syringe. After stirring at 0 °C for 1 hour, the reaction was quenched by the slow addition of a saturated Na/K tartrate solution. The mixture was warmed to room temperature and allowed to stir vigorously for 45 minutes. After stirring, the mixture was extracted with 50% ethyl acetate/hexanes. The combined organic layers were dried over Na₂SO₄ and concentrated in vacuo. Purification by flash column chromatography (20% to 50% EtOAc/hexanes) afforded .313g (92%) of alcohol **13** as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 1.28 (s, 3H), 1.48-1.59 (m, 1H), 1.59 (s, 3H), 1.67-1.78 (m, 1H), 1.99-2.23 (m, 3H), 2.39 (dd, *J* = 13.6, 4.4 Hz, 1H), 3.46 (br s, 1H), 3.58 (dd, *J* = 9.2, 9.2 Hz, 1H), 3.63-3.74 (m, 1H), 3.75 (d, *J* = 8.8 Hz, 1H), 3.77-3.84 (m, 1H), 3.92 (d, *J* = 7.6 Hz, 1H), 4.42 (AB *J*_{AB} = 11.2 Hz, Δ*v*_{AB} = 32.9 Hz, 2H), 4.58 (AB *J*_{AB} = 12 Hz, Δ*v*_{AB} = 41.92 Hz, 2H), 4.68 (s, 1H), 4.73 (d, *J* = 0.8 Hz, 1H), 4.92 (dddd, *J* = 10.4, 1.2, 1.2, 1.2 Hz, 1H), 4.97 (dddd, *J* = 16.8, 1.6, 1.6, 1.6 Hz, 1H), 5.77 (dddd, *J* = 16.8, 10.4, 6.8, 6.8 Hz, 1H), 7.22-7.36 (m, 10H); ¹³C NMR (100 MHz, CDCl₃) δ 20.53, 22.80, 27.52, 34.06, 41.01, 64.05, 65.06, 71.06, 73.82, 79.18, 80.96, 83.08, 113.18, 114.58, 127.42, 127.49, 128.12, 128.20, 128.46, 128.73, 137.35, 138.79, 139.29, 142.38; IR (film) 3451 (br), 2935, 1641, 1454, 1092 cm⁻¹; [α]_D²³ = +15.5 (c = 3.0, CHCl₃); MS (electrospray ionization) calculated for C₂₈H₃₈O₄Na [M + Na]⁺: 461.27, found: 461.26.

(2R,8R,9R)-(8-benzyloxy-9-benzyloxymethyl-4,8-dimethyl-2,3,6,7,8,9-hexahydro-oxonin-2-yl)-methanol (14). Into a 2L flask equipped with a reflux condenser was added diene **13** (1.0 g, 2.28 mmol) in 1.2L of benzene. Argon was bubbled through the stirring solution using a needle for 1 ½ hours. The needle was removed and Cl₂(Cy₃P)(sIMES)Ru=CHPh (.194g, .228 mmol) was added in one portion. The solution was heated to reflux at 80 °C for 5 hours and then cooled to room temperature. The solution was stirred vigorously opened to air overnight and then concentrated in vacuo. Purification by flash chromatography provided .832g (89%) of oxonene **14** as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 1.18 (s, 3H), 1.61 (ddd, *J* = 14, 6.8, 5.2 Hz, 1H), 1.72 (s, 3H), 1.91 (dddd, *J* = 14, 8.4, 8.4, 5.2 Hz, 1H), 2.05 (dd, *J* = 14.4, 1.2 Hz, 1H), 2.30 (dd, *J* = 14.4, 6.8 Hz, 1H), 2.40 (ddd, *J* = 13.6, 8.4, 4.8 Hz, 1H), 2.53 (dddd, *J* = 13.6, 6.8, 6.8, 6.8 Hz, 1H), 3.45-3.53 (m, 1H), 3.58 (dd, *J* = 10.4, 10.4 Hz, 1H), 3.71-3.95 (m, 4H), 4.49 (s, 2H), 4.58 (AB *J*_{AB} = 11.6 Hz, Δ*v*_{AB} = 14.3 Hz, 2H), 5.49 (dd, *J* = 7.6, 7.6 Hz, 1H), 7.19- 7.35 (m, 10H); ¹³C NMR (100 MHz, CDCl₃) δ 22.62, 23.41, 26.16, 34.78, 36.00, 64.74, 67.24, 72.44, 73.93, 79.39, 80.29, 83.72, 127.18, 127.21, 127.89, 128.01, 128.19, 128.42, 128.67, 133.13, 138.01, 140.05; IR (film) 3442 (br), 2927, 1453, 1377, 1086 cm⁻¹; [α]_D²³ = -69.5 (c = 1.25, CHCl₃); MS (electrospray ionization) calculated for C₂₆H₃₄O₄Na [M + Na]⁺: 433.24, found: 433.23.

(2R,8R,9R)-8-Benzyloxy-9-benzyloxymethyl-4,8-dimethyl-2,3,6,7,8,9-hexahydro-oxonine-2-carbaldehyde. A flask was charged with alcohol **14** (2.75g, 6.70 mmol) in 50 mL of reagent grade dichloromethane. Pyridine (2.71mL, 33.5 mmol) was added followed by Dess-Martin periodinane (3.12g, 7.37mmol) in one portion. The reaction was stirred at room temperature for 1 hour and then quenched by the addition of a 1:1 mixture of saturated solutions of NaHCO₃ and Na₂S₂O₃. The mixture was diluted in dichloromethane and washed with a 1M solution of NaHSO₄, water, and brine. The organic layer was dried over Na₂SO₄ and concentrated in vacuo. The crude aldehyde was filtered through a small plug of silica gel eluting with 50% EtOAc/hexanes to afford the aldehyde. The aldehyde was immediately used in the following Wittig reaction.

3-[(2R,8R,9R)-8-Benzyloxy-9-benzyloxymethyl-4,8-dimethyl-2,3,6,7,8,9-hexahydro-oxonin-2-yl]-2-methyl-acrylic acid ethyl ester (15**).** A 100 mL flask was charged with aldehyde (“6.70” mmol) in 40 mL of toluene. (1-Ethoxycarbonyl-ethylidene)triphenylphosphorane (7.39g, 20.4 mmol) was added in one portion and the reaction was heated to 80 °C overnight. The following morning the reaction was cooled to room temperature and concentrated in vacuo. Purification by flash column chromatography (10% to 25% EtOAc/hexanes) provided 3.27g (99% from alcohol **14**) of α,β -unsaturated ester **15** as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 1.23 (s, 3H), 1.29 (t, *J* = 7.2 Hz, 1H), 1.66 (ddd, *J* = 13.6, 9.6, 4.4, Hz, 1H), 1.72 (s, 3H), 1.77 (d, *J* = 1.2 Hz, 3H), 1.89 (dddd, *J* = 13.6, 10.8, 6.4, 4.4 Hz, 1H), 2.07 (dd, *J* = 14, 4.8 Hz, 1H), 2.21-2.33 (m, 1H), 2.37 (ddd, *J* = 12.8, 6.4, 3.6, Hz, 1H), 2.61 (dd, *J* = 14, 3.6 Hz, 1H), 3.69 (dd, *J* = 10, 7.2 Hz, 1H), 3.92-4.00 (m, 2H), 4.15-4.24 (m, 2H), 4.48 (s, 2H), 4.46 (s, 1H), 4.57 (s, 2H), 4.73 (ddd, *J* = 8.8, 4.4, 4.4, Hz, 1H), 5.57 (dd, *J* = 7.6, 7.6 Hz, 1H), 6.80 (dddd, *J* = 7.6, 0.8, 0.8, 0.8 Hz, 1H), 7.19-7.35 (m, 10H); ¹³C NMR (100 MHz, CDCl₃) δ 13.01, 14.54, 22.16, 23.71, 26.37, 35.54, 35.57, 60.81, 65.07, 73.44, 73.81, 75.10, 80.30, 80.92, 126.18, 127.11, 127.17, 127.77, 127.83, 128.37, 128.55, 129.33, 131.61, 138.55, 140.27, 143.65, 168.31; IR (film) 2928, 1710 (str), 1453, 1366, 1252, 1075 cm⁻¹; [α]_D²³ = -15.2 (c = 1.25, CHCl₃); MS (electrospray ionization) calculated for C₃₁H₄₀O₅Na [M + Na]⁺: 515.28, found: 515.23.

3-[(2R,8R,9R)-8-Benzyloxy-9-benzyloxymethyl-4,8-dimethyl-2,3,6,7,8,9-hexahydro-oxonin-2-yl]-2-methyl-prop-2-en-1-ol. A 200 mL flask was charged with diisobutyl aluminum hydride (1.0 M in heptane, 19.4 mL, 19.4 mmol) followed by 20 mL of diethyl ether. After the mixture was cooled to -78 °C, α,β -unsaturated ester **15** (3.10g, 6.48 mmol) was diluted in 40 mL of diethyl ether and added to the hydride dropwise over 45 minutes. After the addition was complete, the reaction was stirred at -78 °C for an additional 30 minutes. The reaction was quenched by the careful addition of a saturated sodium/potassium tartrate solution. The mixture was slowly warmed to room temperature and stirred

vigorously until the mixture turned into two clear layers. The layers were separated and the aqueous layer was extracted three times with diethyl ether. The organic layers were dried over Na₂SO₄ and concentrated in vacuo. Purification by flash column chromatography (20% to 50% EtOAc/hexanes) provided 2.67g (92%) of the alcohol as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 1.22 (s, 3H), 1.61 (d, *J* = 1.2 Hz, 3H), 1.65 (ddd, *J* = 13.2, 9.2, 4.4 Hz, 1H), 1.75 (s, 3H), 1.89 (dddd, *J* = 14, 9.6, 6.8, 4.4 Hz, 1H), 2.08 (dd, *J* = 14, 5.2 Hz, 1H), 2.20-2.31 (m, 1H), 2.40 (ddd, *J* = 11.6, 6.8, 4 Hz, 1H), 2.56 (dd, *J* = 14, 3.2 Hz, 1H), 3.69 (dd, *J* = 10, 6.8 Hz, 1H), 3.91-4.01 (m, 4H), 4.50 (s, 2H), 4.55 (AB *J*_{AB} = 12 Hz, Δ*v*_{AB} = 8.7 Hz, 2H), 4.63 (ddd, *J* = 8, 4.8, 3.2 Hz, 1H), 5.55 (dd, *J* = 7.6, 7.6 Hz, 1H), 5.56 (dddd, *J* = 8.4, 1.6, 1.6, 1.6 Hz, 1H), 7.17-7.34 (m, 10H); ¹³C NMR (100 MHz, CDCl₃) δ 14.16, 22.19, 23.40, 26.63, 35.37, 36.46, 64.97, 68.58, 73.35, 73.66, 74.35, 80.45, 80.68, 127.04, 127.16, 127.70, 127.77, 128.26, 128.32, 128.51, 132.64, 134.11, 134.85, 138.72, 140.41; IR (film) 3421 (br), 2921, 1453, 1376, 1252, 1078 cm⁻¹; [α]_D²³ = -24.3 (c = 4.4, CHCl₃); MS (electrospray ionization) calculated for C₂₉H₃₈O₄Na [M + Na]⁺: 473.27, found: 473.27.

(2*R*,8*R*,9*R*)-3-Benzyloxy-2-benzyloxymethyl-3,7-dimethyl-9-[2-methyl-3-(tetrahydro-pyran-2-yloxy)-propenyl]-2,3,4,5,8,9-hexahydro-oxonine (16). A 100 mL flask was charged with the primary alcohol (2.65g, 5.87 mmol) in 60 mL of dichloromethane. Dihydropyran (.589 mL, 6.46 mmol) was added followed by pyridinium *p*-toluenesulfonate (148mg, .587 mmol) and the reaction was stirred at room temperature for 3 hours. The reaction was quenched by the addition of a saturated NaHCO₃ solution and diluted in dichloromethane. The layers were separated and the organic layer was washed with water and brine. The organic layers were dried over Na₂SO₄ and concentrated in vacuo. Purification by flash column chromatography (5% to 20% EtOAc/hexanes) provided 3.16g (quantitative yield) of tetrahydropyranyl acetal **16** as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 1.21 (s, 3H), 1.45-1.94 (m, 8H), [1.61 (d), *J* = 0.8 Hz; 1.63 (d), *J* = 0.8 Hz, 3H], [1.74 (s); 1.76 (s), 3H] 2.06 (dd, *J* = 14, 4.4 Hz, 1H), 2.18-2.29 (m, 1H), 2.35-2.44 (m, 1H), 2.57 (ddd, *J* = 13.6, 3.2, 3.2 Hz, 1H), 3.45-3.51 (m, 1H), 3.63-3.70 (m, 1H), 3.81-3.89 (m, 1H), 3.91-3.98 (m, 2H), 4.06-4.12 (m, 1H), 4.50 (s, 2H), 4.56 (s, 2H), 4.59 (dd, *J* = 4, 4 Hz, 1H), 4.60-4.66 (m, 1H), 5.53 (dd, *J* = 9.2, 9.2 Hz, 1H), 5.56-5.61 (m, 1H), 7.15-7.35 (m, 10H); ¹³C NMR (100 MHz, CDCl₃) δ 14.59, 14.62, 19.73, 22.22, 23.57, 23.62, 25.73, 26.53, 26.59, 30.89, 35.44, 62.36, 62.38, 65.07, 65.11, 72.59, 72.79, 73.39, 73.69, 74.34, 74.40, 77.56, 80.51, 80.67, 97.70, 98.14, 127.03, 127.22, 127.66, 127.78, 127.81, 128.34, 128.48, 128.53, 130.08, 131.45, 131.48, 132.73, 138.77, 139.79, 140.49; IR (film) 2938, 1643, 1453, 1376 cm⁻¹; [α]_D²³ = -30.4 (c = 1.25, CHCl₃); MS (electrospray ionization) calculated for C₃₄H₄₆O₅Na [M + Na]⁺: 557.32, found: 557.27.

(2*R*,3*R*,9*R*)-2-Hydroxymethyl-3,7-dimethyl-9-[2-methyl-3-(tetrahydro-pyran-2-yloxy)-propenyl]-2,3,4,5,8,9-hexahydro-oxonin-3-ol (17). A 2 neck 200 mL round bottom flask was equipped with a dry ice condenser and cooled to -78 °C. Approximately 25 mL of dry ammonia was condensed into the flask. Benzyl ether **16** (1.0g, 1.87 mmol) was added slowly in 30 mL of THF. Sodium (.645g, 28.1 mmol) was cut into small pieces and added to the ammonia / THF mixture. The reaction was stirred at -78 °C until the sodium dissolved and the reaction solution turned a dark

blue/green color (approx. 30 minutes). The reaction was closely monitored by TLC and when the starting material was consumed the reaction was quenched by the careful addition of solid sodium benzoate (1.5g) at -78 °C followed by the careful addition of water. As the mixture was warmed to room temperature the color changed to yellow and vigorous stirring continued to boil off the ammonia. The mixture turned homogenous by the addition of a small amount of water and was extracted by diethyl ether four times. The organic layers were dried over Na₂SO₄ and concentrated in vacuo. Purification by flash column chromatography (20% to 75% EtOAc/hexanes) provided .603g (91%) of diol **17** as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ [1.08 (s); 1.09 (s), 3H], 1.45-1.90 (m, 13H), [1.73 (s); 1.74 (s), 3H], 2.15 (br d, 1H), 2.69-2.82 (m, 2H), 3.32 (br s, 1H), 3.44-3.51 (m, 1H), 3.60-3.69 (m, 3H), 3.79-3.88 m, 2H), 4.06 (dd, *J* = 12.4, 5.2 Hz, 1H), 4.15-4.23 (m, 1H), 4.53-4.59 (m, 1H), 5.25-5.33 (m, 1H), 5.60 (dddd, *J* = 8.8, 2.8, 1.6, 1.6 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 14.36, 14.42, 19.62, 19.77, 23.30, 25.62, 25.69, 25.90, 30.83, 30.85, 39.52, 39.60, 43.64, 62.42, 62.60, 62.84, 62.87, 72.10, 72.51, 73.14, 73.15, 75.27, 75.41, 82.79, 82.87, 98.10, 98.21, 128.91, 129.03, 129.54, 130.81, 130.82, 132.63, 132.66; IR (film) 3447 (br), 2924, 1444, 1021 cm⁻¹; [α]_D²³ = -23.7 (c = 6.4, CHCl₃); MS (electrospray ionization) calculated for C₂₀H₃₄O₅Na [M + Na]⁺: 377.23, found: 377.24.

(2*S*,3*R*,9*R*)-3-Hydroxy-3,7-dimethyl-9-[2-methyl-3-(tetrahydro-pyran-2-yloxy)-propenyl]-2,3,4,5,8,9-hexahydro-oxonine-2-carbaldehyde. A flask was charged with alcohol **17** (.571 g, 1.61 mmol) in 20 mL of reagent grade dichloromethane. Pyridine (.68 mL, 8.45mmol) was added followed by Dess-Martin periodinane (1.79g, 4.23 mmol) in one portion. The reaction was stirred at room temperature for 2 hours and then quenched by the addition of a 1:1 mixture of saturated solutions of NaHCO₃ and Na₂S₂O₃. The mixture was diluted in dichloromethane and washed with a 1M solution of NaHSO₄, water, and brine. The organic layer was dried over Na₂SO₄ and concentrated in vacuo. The crude aldehyde was filtered through a small plug of silica gel eluting with 50% EtOAc/hexanes to afford the aldehyde. The aldehyde was immediately used in the following Wittig reaction.

3-[(2*R*,3*R*,9*R*)-Hydroxy-3,7-dimethyl-9-[2-methyl-3-(tetrahydro-pyran-2-yloxy)-propenyl]-2,3,4,5,8,9-hexahydro-oxonin-2-yl]-acrylic acid methyl ester (18**).** A 100 mL flask was charged with the aldehyde ("1.61" mmol) in 15 mL of dichloromethane. Methoxycarbonylmethylidene triphenylphosphorane (1.13g, 3.22 mmol) was added in one portion and the reaction was heated to reflux. After 45 minutes, a one mL aliquot was taken and the solvent was removed. By ¹H NMR, all of the aldehyde had been consumed. The reaction was then cooled to room temperature and the dichloromethane was removed in vacuo. Purification by flash column chromatography (10% to 20% EtOAc/hexanes) provided .599g (91%) of α,β-unsaturated ester **18** as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 0.99 (s, 3H), 1.46-1.63 (m, 5H), 1.54 (s, 3H), 1.63-1.76 (m, 2H), 1.72 (s, 3H), 1.76-1.94 (m, 3H), 2.65-2.83 (m, 2H), [3.27 (s); 3.28 (s), 1H], 3.45-3.52 (m, 1H), [3.67 (s); 3.68 (s), 3H], 3.80-3.88 (m, 2H), 3.88-3.95 (m, 1H), 4.05 (dd, *J* = 12.4, 9.2 Hz, 1H), 4.23-4.28 (m, 1H), 4.57 (dd, *J* = 3.2, 3.2 Hz, 1H), 5.28-5.35 (m, 1H), 5.49-5.54 (m, 1H), [5.88 (dd) *J* = 5.2, 1.2 Hz; 5.92 (dd) *J* = 5.2, 1.2 Hz, 1H], [6.80 (dd) *J* = 15.6, 6.4 Hz; 6.82 (dd) *J* = 15.6, 6 Hz, 1H]; ¹³C NMR (100 MHz, CDCl₃) δ 14.45, 14.54, 19.62, 19.70, 23.26, 25.68, 25.71, 26.64, 26.68, 30.81, 30.84, 39.34, 39.41, 51.72, 51.74, 62.35,

62.45, 71.80, 71.88, 73.27, 73.46, 73.47, 82.75, 82.76, 97.63, 97.72, 122.47, 122.54, 127.97, 128.42, 130.04, 130.07, 130.77, 130.82, 132.53, 132.61, 145.66, 145.76, 166.71, 166.73; IR (film) 3566 (br), 2938, 1725 (str), 1438, 1278, 1021 cm^{-1} ; $[\alpha]_D^{23} = -43.6$ ($c = 1.7$, CHCl_3); MS (electrospray ionization) calculated for $\text{C}_{23}\text{H}_{36}\text{O}_6\text{Na}$ $[\text{M} + \text{Na}]^+$: 431.24, found: 431.24.

3-[(2R,3R,9R)-3,7-Dimethyl-9-[2-methyl-3-(tetrahydro-pyran-2-yloxy)-propenyl]-3-triethylsilanyloxy-2,3,4,5,8,9-hexahydro-oxonin-2-yl]-acrylic acid methyl ester (19). A 100 mL flask was charged with alcohol **18** (.550g, 1.35 mmol) in 15 mL of dichloromethane. The solution was cooled to -78°C and 2,6-lutidine (.789 mL, 6.75 mmol) was added dropwise via syringe. Triethylsilyl trifluoromethyl sulfonate (.611 mL, 2.70 mmol) was then added slowly via syringe and the reaction was stirred at -78°C for 1 1/2 hours. Once the reaction was complete, a saturated solution of NaHCO_3 was added slowly and the mixture was warmed to room temperature. The layers were separated and the organic layer was washed with water and brine. The organic layers were dried over Na_2SO_4 and concentrated in vacuo. Purification by flash column chromatography (hexanes to 10% EtOAc/hexanes) provided .670g (95%) of the silyl ether **19** as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 0.50 (q, $J = 7.6$ Hz, 6H), 0.88 (t, $J = 7.6$ Hz, 9H), 1.08 (s, 3H), 1.45-1.85 (m, 8H), [1.58 (s), 1.59 (s), 3H], [1.71 (s), 1.72 (s), 3H], 2.02 (ddd, $J = 13.2, 6.4, 2.8$ Hz, 1H), 2.11 (dd, $J = 14, 5.6$ Hz, 1H), 2.30-2.41 (m, 2H), 3.43-3.51 (m, 1H), 3.69 (s, 3H), 3.80-3.89 (m, 2H), 3.94-4.08 (m, 3H), 4.57 (dd, $J = 3.6, 3.6$ Hz, 1H), 5.44 (dd, $J = 8, 8$ Hz, 1H), 5.52 (d, $J = 8.4$ Hz, 1H), [5.86 (dd), $J = 15.6, 1.2$ Hz, 5.87 (dd), $J = 15.6, 1.2$ Hz, 1H], [6.93 (dd), $J = 15.6, 6$ Hz, 6.94 (dd), $J = 15.6, 6$ Hz, 1H]; ^{13}C NMR (100 MHz, CDCl_3) δ 7.09, 7.26, 14.57, 19.68, 19.79, 22.86, 22.88, 25.73, 26.58, 26.64, 27.01, 27.07, 30.86, 30.91, 36.38, 41.46, 41.51, 51.61, 51.64, 62.35, 62.53, 72.08, 72.35, 72.89, 72.96, 78.26, 78.28, 81.83, 81.93, 97.64, 97.79, 121.98, 122.04, 128.61, 129.11, 129.38, 131.50, 131.53, 131.64, 131.69, 147.03, 147.09, 167.08; IR (film) 2950, 1727 (str), 1437, 1274, 1039 cm^{-1} ; $[\alpha]_D^{23} = -47.6$ ($c = 0.96$, CHCl_3); MS (electrospray ionization) calculated for $\text{C}_{29}\text{H}_{50}\text{O}_6\text{SiNa}$ $[\text{M} + \text{Na}]^+$: 545.33, found: 545.30.

3-[(2R,3R,9R)-9-(3-Hydroxy-2-methyl-propenyl)-3,7-dimethyl-3-triethylsilanyloxy-2,3,4,5,8,9-hexahydro-oxonin-2-yl]-acrylic acid methyl ester. A 100 mL flask was charged with the tetrahydropyranyl acetal (.650g, 1.24 mmol) in 15 mL of methanol. Pyridinium *p*-toluenesulfonate (31 mg, .124 mmol) was added in one portion and the reaction was stirred at room temperature for 6 hours. The reaction was quenched by the addition of a saturated NaHCO_3 solution and the mixture was concentrated in vacuo. The mixture was diluted in diethyl ether and the layers were separated and the organic layer was washed with water and brine. The combined aqueous layers were extracted three times with diethyl ether. The organic layers were dried over Na_2SO_4 and concentrated in vacuo. Purification by flash column chromatography (10% to 30% EtOAc/hexanes) provided .528g (97%) of the alcohol as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 0.51 (q, $J = 7.6$ Hz, 6H), 0.89 (t, $J = 8$ Hz, 9H), 1.09 (s, 3H), 1.60 (d, $J = 1.2$ Hz, 3H), 1.60-1.66 (m, 1H), 1.72 (d, $J = 1.2$ Hz, 3H), 1.73-1.80 (m, 1H), 2.03 (ddd, $J = 13.6, 7.2, 2.8$ Hz, 1H), 2.14 (dd, $J = 13.6, 5.6$ Hz, 1H), 2.33 (dd, $J = 14, 3.2$ Hz, 1H), 2.33-2.45 (m, 1H), 3.70 (s, 3H), 3.94-4.00 (m, 3H), 4.03 (dd, $J = 6, 1.2$ Hz, 1H), 5.45 (ddd, $J = 8.4, 8.4, 1.2$ Hz, 1H), 5.50 (dddd, $J = 8.4, 1.2, 1.2, 1.2$ Hz, 1H), 5.88 (dd, $J = 15.6, 1.2$ Hz, 1H), 6.94 (dd, $J = 15.6, 6$ Hz, 1H);

^{13}C NMR (100 MHz, CDCl_3) δ 7.07, 7.26, 14.24, 22.89, 26.68, 27.03, 36.48, 41.56, 51.68, 68.62, 73.08, 78.15, 82.16, 122.05, 127.55, 129.47, 131.39, 134.49, 146.98, 167.07; IR (film) 3433 (br), 2952, 1726 (str), 1655, 1436, 1376, 1277, 1043 cm^{-1} ; $[\alpha]_{\text{D}}^{23} = -80.3$ ($c = 1.25$, CHCl_3); MS (electrospray ionization) calculated for $\text{C}_{24}\text{H}_{42}\text{O}_5\text{SiNa}$ $[\text{M} + \text{Na}]^+$: 461.27, found: 461.27.

3-[(2*R*,3*R*,9*R*)-3,7-Dimethyl-9-(2-methyl-3-oxo-propenyl)-3-triethylsilyloxy-2,3,4,5,8,9-hexahydro-oxonin-2-yl]-acrylic acid methyl ester (20**).** A 25 mL flask was charged with the alcohol (.143g, .327 mmol) in 5 mL of dichloromethane. Pyridine (.079 mL, .98 mmol) was added followed by the addition of the Dess-Martin periodinane (.207g, .490 mmol). The reaction was stirred at room temperature for 30 minutes and quenched by the addition of a 1:1 mixture of saturated solutions of NaHCO_3 and $\text{Na}_2\text{S}_2\text{O}_3$. The mixture was diluted in dichloromethane and washed with a 1M solution of NaHSO_4 , water, and brine. The organic layer was dried over Na_2SO_4 and concentrated in vacuo. The crude aldehyde was filtered through a small plug of silica gel eluting with 20% EtOAc/hexanes to afford aldehyde **20**. The aldehyde was immediately used in the following Wittig reaction.

3-[(2*R*,3*R*,9*R*)-9-(4-Benzyloxy-2-methyl-buta-1,3-dienyl)-3,7-dimethyl-3-triethylsilyloxy-2,3,4,5,8,9-hexahydro-oxonin-2-yl]-acrylic acid methyl ester (3**).** A 25 mL flask was charged with benzyloxymethyltriphenylphosphonium chloride (.410g, .980 mmol) and purged with argon. The salt was taken up in 3 mL of THF to give a white slurry. The suspension was cooled to -78°C and potassium *tert*-butoxide (92 mg, .818 mmol) in 1 mL of THF was added dropwise over 10 minutes. After the addition was complete, the ylide (brown / yellow milky slurry) was stirred at -78°C for one hour. Aldehyde **20** (.327 mmol) was diluted in 2 mL of THF and added to the ylide via syringe over 10 minutes. The reaction was stirred at -78°C for 45 minutes and then quenched with 2 mL of water and then warmed to room temperature. The mixture was diluted in diethyl ether and the organic layer was washed with water and brine. The combined organic layers were dried over Na_2SO_4 and concentrated in vacuo. The crude vinyl ethers (3:1 *E,E:E,Z*) was used directly in the following cycloaddition reaction.

(3*R*,4*R*,7*R*,8*R*,14*R*,15*R*,16*S*)-4-Benzyloxy-6,10,14-trimethyl-14-triethylsilyloxy-15-oxatricyclo [6.6.1.0^{2,7}]pentadeca-5,10-diene-3-carboxylic acid methyl ester (22**).** The crude vinyl ether (0.327 mmol) was taken up in 4 mL of benzene and kept at room temperature. After approximately two hours, the cycloaddition was complete leaving the reaction mixture enriched in tetraene **21** and

cycloadduct **22**. The benzene was removed in vacuo and the resultant mixture was purified by flash column chromatography (hexanes to 25% EtOAc/hexanes) providing a mixture of *Z* vinyl ether **21** and cycloadduct **22**. The mixture was diluted in 5 mL of benzene and phenyl disulfide (4mg, 0.016mmol) was added in one portion. The solution was then irradiated with a 450 watt Hanovia medium pressure mercury vapor lamp equipped with a uranium glass filter sleeve for 15 minutes. By ^1H NMR, a 1:1 mixture of tetraenes **3** and **23** existed. This mixture was allowed to stand at room temperature for two hours after which time the reaction mixture was enriched in tetraene **23** and cycloadduct **22**. The process was repeated (2 more times) until all of the tetraene **23** had been funneled to the cycloadduct **22**. The benzene was removed in vacuo and cycloadduct was purified by flash chromatography (hexanes to 25% ethyl acetate) to afford .142g (80% yield from the allylic alcohol) of cycloadduct **22** as a colorless oil: ^1H NMR (400 MHz, C_6D_6 , 25 °C) δ 0.64 (q, J = 7.6 Hz, 6H), 1.00 (t, J = 7.6 Hz, 9H), 1.23 (s, 3H), 1.53 (s, 3H), 1.59-1.69 (m, 1H), 1.75-1.83 (m, 1H), 1.80 (s, 3H), 2.23 (br s, 2H), 2.48 (br s, 2H), 2.82 (dd J = 6.8, 6.8 Hz, 1H), 3.14 (ddd J = 7.6, 7.6, 3.2 Hz, 1H), 3.20-3.29 (m, 1H), 3.37 (s, 3H), 4.01 (br s, 1H), 4.09 (br s, 1H), 4.48-4.53 (m, 1H), 4.52 (AB J_{AB} = 12 Hz, $\Delta\nu_{\text{AB}}$ = 25.6 Hz, 2H), 5.57 (ddd, J = 2.4, 1.2, 1.2 Hz, 1H), 5.68 (dd, J = 8.4, 8.4 Hz, 1H), 7.05-7.11 (m, 1H), 7.13-7.19 (m, 2H), 7.30-7.36 (m, 2H); ^1H NMR (400 MHz, C_6D_6 , 60 °C) δ 0.66 (q, J = 7.6 Hz, 6H), 1.00 (t, J = 7.6 Hz, 9H), 1.27 (s, 3H), 1.55 (s, 3H), 1.67 (ddd, J = 14, 8.4, 2.8 Hz, 1H), 1.75-1.84 (m, 1H), 1.79 (s, 3H), 2.13-2.29 (m, 2H), 2.43-2.55 (m, 2H), 2.81 (dd, J = 6.8, 6.8 Hz, 1H), 3.14 (ddd, J = 8, 8, 4 Hz, 1H), 3.22 (dd, J = 8.8, 6.8 Hz, 1H), 3.40 (s, 3H), 4.03 (ddd, J = 6.8, 6.8, 3.2 Hz, 1H), 4.10 (d, J = 3.6 Hz, 1H), 4.45-4.50 (m, 1H), 4.52 (AB J_{AB} = 11.6 Hz, $\Delta\nu_{\text{AB}}$ = 21.6 Hz, 2H), 5.58 (ddd, J = 3.2, 1.6, 1.6 Hz, 1H), 5.65 (dd, J = 8.8, 8.8 Hz, 1H), 7.05-7.11 (m, 1H), 7.14-7.19 (m, 2H), 7.30-7.34 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3 , 60 °C) δ 7.13, 7.48, 22.88, 23.45, 25.10, 29.56, 39.05, 40.21, 43.38, 48.24, 51.84, 70.92, 76.18, 77.45, 77.93, 80.01, 90.26, 122.35, 127.75, 127.95, 128.50, 128.95, 132.78, 136.17, 138.66, 175.42; IR (film) 2952, 1736 (str), 1455, 1168, 1089 cm^{-1} ; $[\alpha]_{\text{D}}^{23}$ = -64.2 (c = 1.35, CHCl_3); MS (electrospray ionization) calculated for $\text{C}_{32}\text{H}_{48}\text{O}_5\text{SiNa}$ $[\text{M} + \text{Na}]^+$: 563.32, found: 563.27.

2-((3*R*,4*R*,7*R*,8*R*,14*R*,15*R*,16*S*)-4-Benzoyloxy-6,10,14-trimethyl-14-triethylsilanyloxy-15-oxatricyclo[6.6.1.0^{2,7}]pentadeca-5,10-dien-3-yl)-propan-2-ol (24**).** A 25 mL round bottom flask was charged with methylmagnesium chloride (1.0 M in THF, 3.32 mL, 3.32 mmol) followed by 2 mL of THF. The Grignard was cooled to 0 °C and ester **22** (.120 g, .222 mmol) was added dropwise in 2 mL of THF. The reaction was stirred at 0 °C for 1 hour and then warmed to room temperature. After 12 hours, the reaction was cooled to 0 °C and quenched by the slow addition of a saturated NH_4Cl solution. The mixture was diluted in diethyl ether and washed with water and brine. The water layers were extracted with diethyl ether three times. The combined organic layers were dried over Na_2SO_4 and concentrated in vacuo. Purification by flash column chromatography (5% to 15% EtOAc/hexanes) provided .102g (85%) of alcohol **24** as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 0.58 (q, J = 7.6 Hz, 6H), 0.91 (t, J = 7.6 Hz, 9H), 1.13 (s, 3H), 1.17 (s, 3H), 1.32 (s, 1H), 1.46 (s, 3H), 1.71 (d, J = 0.8 Hz, 3H), 1.76-1.92 (m, 4H), 1.81 (s, 3H), 2.40 (d, J = 8.4 Hz, 1H), 2.50-2.59 (m, 1H), 2.52 (s, 1H), 2.75 (d, J = 14.8 Hz, 1H), 2.95 (dd, J = 9.6, 9.6 Hz, 1H), 3.91 (d, J = 5.2 Hz, 1H), 4.25 (d, J = 10 Hz, 1H), 4.31 (ddd, J = 4, 2, 2 Hz, 1H), 4.52 (AB J_{AB} = 12.4 Hz, $\Delta\nu_{\text{AB}}$ = 31.7 Hz, 2H), 5.52 (dd, J = 10.8, 5.6 Hz, 1H), 5.67 (d, J = 5.2 Hz, 1H), 7.20-7.35 (m, 5H); ^{13}C NMR (100 MHz, CDCl_3) δ 7.24, 7.39, 22.17, 23.53, 28.39, 28.46, 28.70, 29.15, 38.68, 39.03, 39.29, 45.28, 47.34, 69.89, 70.59, 73.08, 78.49, 81.30, 88.48, 122.46, 127.45, 127.86, 128.40, 130.33, 130.68, 138.26, 139.50; IR (film) 3465 (br), 2913, 1455, 1370, 1238, 1133, 1065 cm^{-1} ; $[\alpha]_{\text{D}}^{23}$ = -47.0 (c = 1.0, CHCl_3); MS (electrospray ionization) calculated for $\text{C}_{33}\text{H}_{52}\text{O}_4\text{SiNa}$ $[\text{M} + \text{Na}]^+$: 563.35, found: 563.27.

(5R,6R,7S,8R,9R,15R,16R)-5-Benzyloxy-6-(1-hydroxy-1-methyl-ethyl)-3,9,13-trimethyl-15-oxa-tricyclo[6.6.1.0^{2,7}]pentadeca-3,12-dien-9-ol (27**).** A 25 mL round bottom flask was charged with triethylsilyl ether **24** (80mg, .148 mmol) in 3 mL of THF. Tetrabutylammoniumfluoride (TBAF) (1.0M in THF, .296 mL, .296 mmol) was slowly added dropwise by syringe. The reaction was stirred at room temperature for 30 minutes. After complete by TLC, the reaction was quenched with a solution of saturated NH₄Cl and diluted in ethyl acetate. The organic layer was washed with water and brine. The combined aqueous layers were extracted with ethyl acetate three times. The combined organic layers were dried over Na₂SO₄ and concentrated in vacuo. Purification by flash column chromatography (10% to 30% EtOAc/hexanes) provided 59mg (94%) of diol **27** as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 1.15 (s, 3H), 1.23 (s, 3H), 1.28 (s, 1H), 1.34 (s, 3H), 1.41 (s, 1H), 1.69-1.77 (m, 2H), 1.74 (d, *J* = 1.2 Hz, 3H), 1.86 (d, *J* = 4 Hz, 1H), 1.87-1.97 (m, 1H), 2.45 (d, *J* = 7.6 Hz, 1H), 2.51 (s, 1H), 2.52-2.62 (m, 1H), 2.76 (d, *J* = 14.8 Hz, 1H), 3.05 (dd, *J* = 9.6, 8.4 Hz, 1H), 3.99 (d, *J* = 5.2 Hz, 1H), 4.30 (d, *J* = 10 Hz, 1H), 4.35 (ddd, *J* = 3.6, 1.6, 1.6 Hz, 1H), 4.53 (s, 2H), 5.53 (dd, *J* = 11.2, 6 Hz, 1H), 5.73 (d, *J* = 5.2 Hz, 1H), 7.21-7.34 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ 22.08, 23.22, 28.02, 29.26, 29.42, 29.71, 37.23, 38.83, 39.31, 45.39, 46.89, 70.01, 71.32, 73.02, 74.99, 80.80, 87.70, 122.95, 127.66, 127.73, 128.61, 130.67, 130.69, 138.49, 139.22; IR (film) 3398 (br), 2919, 1443, 1375, 1150, 1062 cm⁻¹; [α]_D²³ = -36.0 (c = 0.45, CHCl₃); MS (electrospray ionization) calculated for C₂₇H₃₈O₄Na [M + Na]⁺: 449.27, found: 449.30.

Acetic acid 1-((3R,4R,7R,8R,14R,15R,16S)-4-benzyloxy-14-hydroxy-6,10,14-trimethyl-15-oxa-tricyclo[6.6.1.0^{2,7}]pentadeca-5,10-dien-3-yl)-1-methyl-ethyl ester (28**).** A 25 mL round bottom flask was charged with KH (30% dispersion in mineral oil (156 mg, 1.17 mmol) and washed with hexanes three times. The dried KH was taken up in 2 mL of THF and cooled to 0 °C. Diol **27** (50mg, .117 mmol) was diluted in 1 mL of THF and slowly added to the KH slurry dropwise via syringe. After the slurry was stirred at 0 °C for 30 minutes, acetic anhydride (.110 mL, 1.17mL) in 0.5 mL of THF was added dropwise. After one hour, the reaction was quenched slowly with water and warmed to room temperature. The solution was diluted in diethyl ether and washed with brine. The aqueous layers were extracted with diethyl ether and the combined organic layers were dried over Na₂SO₄ and concentrated in vacuo. Purification by flash column chromatography (10% to 25% EtOAc/hexanes) provided 30mg (55%, 90%BRSM) of acetate **28** as a white solid: ¹H NMR (400 MHz, CDCl₃) δ 1.34 (s, 3H), 1.40 (s, 3H), 1.46 (s, 3H), 1.62 (dd, *J* = 13.2, 7.6 Hz, 1H), 1.74 (d, *J* = 1.2 Hz, 3H), 1.76-1.85 (m, 2H), 1.82 (s, 3H), 1.87-1.98 (m, 1H), 1.96 (s, 3H), 2.40 (d, *J* = 7.6 Hz, 1H), 2.58 (ddd, *J* = 12, 12, 8.4 Hz, 1H), 2.77 (d, *J* = 14.8 Hz, 1H), 2.98 (dd, *J* = 8.4, 8.4 Hz, 1H), 3.03 (s, 1H), 3.83 (d, *J* = 5.2 Hz, 1H), 4.34-4.37 (m, 1H), 4.37 (d, *J* = 14 Hz, 1H), 4.52 (AB *J*_{AB} = 11.6 Hz, Δ*v*_{AB} = 65.9 Hz, 2H), 5.49 (dd, *J* = 10.8, 6 Hz, 1H), 5.71 (d, *J* = 5.2 Hz, 1H), 7.21-7.36 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ 21.93, 22.98, 23.20, 24.90, 24.98, 29.21, 29.49, 37.58, 38.17, 39.44, 43.88, 45.70, 69.99, 71.06, 74.63, 80.41, 84.75, 86.99, 122.88, 127.70, 127.86, 128.61, 130.65, 130.76, 138.06, 138.99, 170.85; IR (film) 3452 (br),

2919, 1732 (str), 1444, 1367, 1257, 1130, 1061 cm^{-1} ; $[\alpha]^{23}_{\text{D}} = -7.2$ ($c = 0.35$, CHCl_3); MS (electrospray ionization) calcd for $\text{C}_{29}\text{H}_{40}\text{O}_5\text{Na}$ $[\text{M} + \text{Na}]^+$: 491.28, found: 491.2.

(5*R*,6*R*,7*S*,8*R*,9*R*,15*R*,16*R*)-Acetic acid 6-(1-acetoxy-1-methyl-ethyl)-5-benzyloxy-3,9,13-trimethyl-15-oxa-tricyclo[6.6.1.0^{2,7}]pentadeca-3,12-dien-9-yl ester (29). A 2 mL vial was charged with alcohol **28** (15 mg, .032 mmol) in 200 μL of THF and 100 μL of acetic anhydride. The solution was cooled to -78°C and $\text{Bi}(\text{OTf})_3$ (3 g, 4.8 μmol) in 100 μL of THF was added dropwise. The reaction was stirred at -78°C for one hour and then warmed to -45°C for 24 hours. The reaction was cooled to -78°C and slowly quenched with a saturated solution of NaHCO_3 . The mixture was warmed to room temperature and diluted in ethyl acetate. The organic phase was washed with water and brine dried over Na_2SO_4 and concentrated in vacuo. Purification by flash column chromatography (15% to 30% EtOAc /hexanes) provided 12mg (75%) of diacetate **29** as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 1.35 (s, 3H), 1.49 (s, 3H), 1.65 (dd, $J = 14, 8$ Hz, 1H), 1.73 (d, $J = 1.2$ Hz, 3H), 1.77 (s, 3H), 1.82 (s, 3H), 1.83 (dd, $J = 11.2, 4.4$ Hz, 1H), 1.92 (s, 3H), 1.95-1.99 (m, 1H), 1.95 (s, 3H), 2.41 (d, $J = 8.4$ Hz, 1H), 2.58 (ddd, $J = 11.6, 11.6, 7.2$ Hz, 1H), 2.78 (d, $J = 14.8$ Hz, 1H), 2.84-2.97 (m, 2H), 3.21 (s, 1H), 3.85 (d, $J = 5.6$ Hz, 1H), 4.35 (ddd, $J = 3.6, 1.6, 1.6$ Hz, 1H), 4.51 (AB $J_{\text{AB}} = 12.4$ Hz, $\Delta\nu_{\text{AB}} = 92.2$ Hz, 2H), 4.56 (d, $J = 10$ Hz, 1H), 5.50 (dd, $J = 11.2, 6$ Hz, 1H), 5.72 (d, $J = 5.2$ Hz, 1H), 7.20-7.32 (m, 5H); ^{13}C NMR (100 MHz, CDCl_3) δ 21.98, 22.83, 22.86, 22.87, 23.67, 25.25, 25.55, 29.09, 33.13, 38.83, 39.34, 42.62, 45.73, 69.99, 70.64, 80.51, 84.56, 85.63, 86.71, 122.80, 127.39, 127.59, 128.53, 130.59, 130.62, 137.74, 139.08, 170.11, 170.21; IR (film) 2918, 1731(str), 1369, 1254, 1099, 1066 cm^{-1} ; $[\alpha]^{23}_{\text{D}} = -53.9$ ($c = 0.48$, CHCl_3); MS (electrospray ionization) calculated for $\text{C}_{31}\text{H}_{42}\text{O}_6\text{Na}$ $[\text{M} + \text{Na}]^+$: 533.29, found: 533.33.

Acetic acid 1-((3*R*,4*R*,7*R*,8*R*,14*R*,15*R*,16*S*)-14-acetoxy-4-hydroxy-6,10,14-trimethyl-15-oxa-tricyclo[6.6.1.0^{2,7}]pentadeca-5,10-dien-3-yl)-1-methyl-ethyl ester (30). A 10 mL round bottom flask was charged with benzyl ether **29** (8mg, 16 μmol) and taken up in 2 mL of ethyl acetate. Palladium on carbon (10% weight, 3mg) was added to the solution and the reaction atmosphere was evacuated and filled with hydrogen (balloon). The reaction was stirred at room temperature for approximately 1 1/2 hours until less than 15% of the benzyl ether remained by TLC. The hydrogen balloon was removed and the reaction mixture was filtered through a small pad of celite washing thoroughly with 10 mL of ethyl acetate and twice with 10 mL of ethanol. The solvents were removed in vacuo and purification by flash column chromatography (10% to 30% EtOAc /hexanes) provided 4.5mg (70%) of allylic alcohol **30** as a colorless oil: ^1H NMR (500 MHz, CDCl_3) δ 1.35 (s, 3H), 1.50 (s, 3H), 1.67-1.72 (m, 1H), 1.73 (s, 3H), 1.75 (s, 3H), 1.81 (s, 3H), 1.85 (dd, $J = 14.5, 4$ Hz, 1H), 1.96-1.99 (m, 1H), 1.97 (s, 3H), 2.02 (s, 3H), 2.40 (d, $J = 8$ Hz, 1H), 2.49-2.60 (m, 2H), 2.77 (d, $J = 14.5$ Hz, 1H), 2.90 (s, 1H), 2.98 (dd, $J = 8.5, 8.5$ Hz, 1H), 4.08-4.13 (m, 1H), 4.36-4.40 (m, 1H), 4.45 (d, $J = 10.5$ Hz, 1H), 5.51 (dd, $J = 11, 6.5$

Hz, 1H), 5.69 (d, $J = 5$ Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 21.77, 22.75, 22.86, 22.92, 24.00, 25.21, 25.23, 34.33, 38.59, 39.28, 45.57, 47.21, 64.13, 77.43, 80.17, 84.61, 86.31, 124.91, 130.39, 130.96, 136.73, 170.30, 170.76; IR (film) 3423 (br), 2919, 1732 (str), 1369, 1254 cm^{-1} ; $[\alpha]_{\text{D}}^{23} = +9.8$ ($c = 0.16$, CHCl_3); MS (electrospray ionization) calculated for $\text{C}_{24}\text{H}_{36}\text{O}_6\text{Na}$ $[\text{M} + \text{Na}]^+$: 443.24, found: 443.21.

Ophirin B (2). A small vial was charged with allylic alcohol **30** (4mg, 9.5 μmol) in 300 μL of dichloromethane. Pyridine (4 μL , .048mmol) was added to the solution followed by a small crystal of DMAP. Acetic anhydride (4.5 μL , .048mmol) was added and the reaction was stirred at room temperature for 30 minutes. The reaction was quenched with a saturated solution of NaHCO_3 and diluted in dichloromethane. The organic layer was washed with water and brine and dried over Na_2SO_4 . The solvents were removed in vacuo and purification by flash column chromatography (10% to 20% EtOAc/hexanes) provided 4 mg (95%) of ophirin B **2** as a colorless oil: $R_f = 0.55$ (30% EtOAc/hexanes); ^1H NMR (500 MHz, CDCl_3) δ 1.36 (s, 3H), 1.57 (s, 3H), 1.66-1.71 (m, 1H), 1.77 (s, 3H), 1.78 (s, 3H), 1.84 (s, 3H), 1.86 (dd, $J = 14$, 4.5 Hz, 1H), 1.97 (s, 3H), 1.97-1.99 (m, 1H), 2.00 (s, 3H), 2.01 (s, 3H), 2.45 (d, $J = 8$ Hz, 1H), 2.59 (ddd, $J = 12$, 11, 9 Hz, 1H), 2.68 (dd, $J = 12.5$, 12.5 Hz, 1H), 2.82 (d, $J = 14$ Hz, 1H), 2.97 (dd, $J = 8.5$, 8.5 Hz, 1H), 3.04 (s, 1H), 4.39 (dd, $J = 2.5$, 1.5 Hz, 1H), 4.42 (d, $J = 10.5$ Hz, 1H), 5.37 (d, $J = 5$ Hz, 1H), 5.53 (dd, $J = 11.5$, 6.5 Hz, 1H), 5.63 (d, $J = 5.5$ Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 21.33, 21.66, 22.44, 22.51, 22.60, 23.84, 25.43, 25.52, 28.87, 33.63, 38.37, 39.09, 43.29, 45.34, 66.35, 79.90, 83.59, 84.34, 85.97, 120.85, 130.41, 130.45, 139.41, 169.92, 170.00, 170.45; IR (film) 2929, 1735 (str), 1369, 1254, 1130 cm^{-1} ; $[\alpha]_{\text{D}}^{23} = -31.1$ ($c = 0.10$, CHCl_3); MS (electrospray ionization) calculated for $\text{C}_{26}\text{H}_{38}\text{O}_7\text{Na}$ $[\text{M} + \text{Na}]^+$: 485.25, found: 485.20.

(3R,4R,7R,8R,14R,15R,16S)-3-(1-Hydroxy-1-methyl-ethyl)-6,10,14-trimethyl-15-oxatricyclo [6.6.1.0^{2,7}]pentadeca-5,10-diene-4,14-diol (25). ^1H NMR (400 MHz, CDCl_3) δ 1.09 (s, 3H), 1.24 (s, 3H), 1.37 (s, 3H), 1.69-1.76 (m, 2H), 1.72 (d, $J = 0.8$ Hz, 3H), 1.77-1.86 (m, 1H), 1.80 (s, 3H), 1.86-1.96 (m, 1H), 2.49-2.59 (s, 1H), 2.37 (d, $J = 8$ Hz, 1H), 2.54 (m, 1H), 2.75 (d, $J = 14.8$ Hz, 1H), 2.78 (br s, 2H), 2.97 (dd, $J = 9.2$, 9.2 Hz, 1H), 3.02 (br s, 1H), 4.11 (d, $J = 10$ Hz, 1H), 4.30 (d, $J = 4.8$ Hz, 1H), 4.35 (ddd, $J = 3.6$, 1.6, 1.6 Hz, 1H), 5.52 (dd, $J = 11.2$, 6.4 Hz, 1H), 5.65 (d, $J = 4.8$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 21.91, 23.19, 27.03, 29.07, 29.32, 29.47, 37.28, 38.87, 39.28, 45.15, 50.70, 64.64, 73.31, 75.18, 80.77, 88.22, 124.85, 130.55, 130.70, 137.18; IR (film) 3366 (br), 2921, 1654, 1443, 1375, 1151 cm^{-1} ; $[\alpha]_{\text{D}}^{23} = 8.68$ ($c = 0.85$, CHCl_3); MS (electrospray ionization) calcd for $\text{C}_{20}\text{H}_{32}\text{O}_4\text{Na}$ $[\text{M} + \text{Na}]^+$: 359.22, found: 359.10.

Tetracycle (26).

^1H NMR (400 MHz, CDCl_3) δ 0.98 (s, 3H), 1.06 (s, 1H), 1.30 (s, 3H), 1.31 (s, 3H), 1.34 (d, $J = 0.4$ Hz, 3H), 1.51-1.58 (m, 1H), 1.63-1.78 (m, 2H), 1.76 (dd, $J = 1.2, 1.2$ Hz, 3H), 1.81-1.92 (m, 1H), 2.34 (dd, $J = 10, 4.4$ Hz, 1H), 2.39-2.49 (m, 2H), 2.66 (d, $J = 14.8$ Hz, 1H), 2.87 (ddd, $J = 9.6, 5.6, 3.6$ Hz, 1H), 3.59 (d, $J = 5.6$ Hz, 1H), 3.84 (ddd, $J = 4.4, 4.4, 1.6$ Hz, 1H), 5.52 (dd, $J = 10.8, 6.8$ Hz, 1H), 6.14 (d, $J = 8$ Hz, 1H), 6.46 (dd, $J = 7.2, 7.2$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 22.95, 23.21, 28.25, 28.64, 29.01, 29.37, 37.54, 38.03, 43.69, 44.92, 53.34, 73.31, 73.53, 73.98, 83.74, 92.32, 130.16, 131.27, 135.46, 136.39; IR (film) 3430 (br), 2926, 1444, 1376, 1112, 1071 cm^{-1} ; $[\alpha]_D^{23} = -19.7$ (c = 0.5, CHCl_3); MS (electrospray ionization) calculated for $\text{C}_{20}\text{H}_{30}\text{O}_3\text{Na}$ $[\text{M} + \text{Na}]^+$: 341.21, found: 341.14.