Supporting information

for

Speciation and Structure Studies of Europium Phosphotungstate

Cheng Zhang, Robertha C. Howell, Kymora B. Scotland, Frances G. Perez, Louis
Todaro and Lynn C. Francesconi*

Department of Chemistry, Hunter College and the Graduate School of the City University of New
York, New York, New York 10021, E-mail: lfrances@hunter.cuny.edu

Table S1. ^{31}P NMR chemical shift (δ , ppm) of $\text{A-PW}_9\text{O}_{34}^{9-}$ under different pH, buffer and counter cation conditions.

pH	Counter cation	δ , ppm	Assignment	Amount, % ^a
pH 1	Na (0.27 M)	.6410	PO_4^{3-}	36
		-11.8417	PW_{11}^{7-}	30
		-14.5522	PW_{12}^{3-}	36
pH 3	Na (0.27 M)	.5091	PO_4^{3-}	31
		-10.8308	PW_{11}^{7-}	59
		-11.8197	?	1.8
		-12.7501	?	1.8
		-14.8379	PW_{12}^{3-}	6.2
pH 4.75	Cs (0.04 M)	2.8806	PO_4^{3-}	40
		-10.1104	PW_{11}^{7-}	60
	K (0.50 M)	2.8257	PO_4^{3-}	43
		-10.0958	PW_{11}^{7-}	57
	Na (0.50 M)	2.9355	PO_4^{3-}	40
		-10.0958	PW_{11}^{7-}	60
	Li (0.50 M)	2.9941	PO_4^{3-}	40
		-10.3778	PW_{11}^{7-}	60
pH 3	Al (0.27 M)	-10.6049	PW_{11}^{7-}	74
		-10.7074	$\beta\text{-P}_2\text{W}_{18}\text{O}_{62}^{6-}$ or Al complex of $\alpha 2\text{-P}_2\text{W}_{17}\text{O}_{61}^{10-}$, see text	13
		-11.4546	$\beta\text{-P}_2\text{W}_{18}\text{O}_{62}^{6-}$ or Al complex of $\alpha 2\text{-P}_2\text{W}_{17}\text{O}_{61}^{10-}$, see text	13
pH 2	Al (0.27 M)	-10.9260	PW_{11}^{7-}	24
		-12.1128	?	53
		-13.0065	?	5.9
		-13.2995	?	2.9
pH 5	Na (0.27 M)	.7875	PO_4^{3-}	20
		-1.6812	?	8

		-10.1129	PW_{11}^{7-}	72
pH 7.3	Na (0.27 M)	3.0364	PO_4^{3-}	45
		-1.6812	?	2.8
		-8.9994	?	1.7
		-10.1056	PW_{11}^{7-}	50
pH 8	Na (0.27 M)	3.2708	PO_4^{3-}	100
pH 10.35	Na (0.27 M)	3.4467	PO_4^{3-}	100

a. The amount was assessed by the integrated intensity of all ^{31}P NMR peaks and expressed in %.

Figure S1. ^{31}P NMR spectra of $\text{PW}_9\text{O}_{34}^{9-} + \text{Eu}^{3+}$ (1:1 stoichiometry) as a function of differing buffer and pH conditions. Top series of spectra: LiAc buffer (0.5 M), bottom series: NaAc buffer (0.5 M). The species are labeled 1,2,3,4 and correspond to the complexes described in the text.

Figure S2. ^{31}P NMR spectra of $\text{PW}_9\text{O}_{34}^{9-} + \text{Eu}^{3+}$ (1:1 stoichiometry) in KAc buffer (0.5 M) as a function of pH.

Figure S3. ^{31}P NMR spectra of reaction of the Na^+ salt of $\text{A-PW}_9\text{O}_{34}^{9-} + \text{Eu}^{3+}$ (1:1 stoichiometry, pH ca 6-7) a) before heat, b) after heating at 90°C .

Figure S4. ^{31}P NMR spectra of crystalline sample of species **1** as a function of pH

Figure S5. Representation as 50% ellipsoids of the asymmetric unit of **1**. Uncoordinated water molecules and hydrogen atoms removed for clarity.

Table S2. Bond Valence Sum (BVS) for the oxygen atoms in the core
 $[\text{Eu}_2\text{PW}_{10}\text{O}_{38}]_4(\text{W}_3\text{O}_8(\text{H}_2\text{O})_2(\text{OH})_4)]^{22-}$ (Brown, I.D.; Altermatt, D. *Acta Cryst.* **1985**, *B41*, 244-247; Thorp, H.H. *Inorg. Chem.* **1992**, *31*, 1585-1588)

	Oxygen	BVS		Oxygen	BVS
Terminal	W(2)-O(8)	1.707		W(7C)-O(14C)-Eu(5)	2.126
	W(3)-O(11)	1.579		W(8C)-O(16C)-Eu(5)	1.934
	Eu(3)-O(15)	0.335		W(10C)-O(18C)-Eu(6)	1.879
	Eu(7)-O(17)	0.350		W(10C)-O(19C)-Eu(5)	1.942
	W(6A)-O(11A)	1.657		W(10C)-O(20C)-Eu(4)	1.957
	W(7A)-O(13A)	1.657		W(4C)-O(36C)-W(10C)	1.863
	W(8A)-O(15A)	1.600		W(9C)-O(37C)-W(10C)	1.820
	W(9A)-O(17A)	1.721		W(5B)-O(10B)-Eu(4)	1.862
				W(6B)-O(12B)-Eu(4)	1.884
μ_2 -Bridge	W(1)-O(5) –Eu(8)	1.957		W(7B)-O(14B)-Eu(3)	1.820
	W(1)-O(6)-Eu(4)	1.928		W(8B)-O(16B)-Eu(3)	1.765
	W(2)-O(7)-Eu(1)	1.842		W(10B)-O(18B)-Eu(4)	1.992
	W(2)-O(9)-Eu(3)	1.814		W(10B)-O(20B)-Eu(8)	1.987
	W(3)-O(10)-Eu(5)	2.163		W(4B)-O(36B)-W(10B)	1.896
	W(3)-O(12)-Eu(7)	1.839		W(9B)-O(37B)-W(10B)	1.567
	Eu(1)-O(13)-Eu(2)	0.904		W(5D)-O(10D)-Eu(8)	1.851
	Eu(5)-O(14)-Eu(6)	0.927		W(6D)-O(12D)-Eu(8)	1.852
	Eu(3)-O(16)-Eu(4)	1.040		W(7D)-O(14D)-Eu(7)	1.766
	Eu(7)-O(18)-Eu(8)	1.010		W(8D)-O(16D)-Eu(8)	1.980
	W(5A)-O(10A)-Eu(2)	1.902		W(10D)-O(20D)-Eu(4)	1.956
	W(6A)-O(12A)-Eu(2)	1.896		W(4D)-O(36D)-W(10D)	1.931
	W(7A)-O(14A)-Eu(1)	1.845		W(9D)-O(37D)-W(10D)	1.616
	W(8A)-O(16A)-Eu(1)	1.987			
	W(10A)-O(18A)-Eu(2)	1.927	μ_3 - Bridge	μ_3 -O1, Eu1, Eu3, W1	1.894
	W(10A)-O(19A)-Eu(1)	2.062		μ_3 -O2, Eu5, W1, W2	2.060
	W(10A)-O(20A)-Eu(8)	2.006		μ_3 -O3, W3, W1, Eu1	2.094
	W(4A)-O(36A)-W(10A)	1.829		μ_3 -O4, W1, Eu5, Eu7	1.847
	W(9A)-O(37A)-W(10A)	1.789		μ_3 -O19B, W10B, Eu2, Eu3	2.066
	W(5C)-O(10C)-Eu(6)	1.929		μ_3 -O19D, W10D, Eu6, Eu7	2.021
	W(6C)-O(12C)-Eu(6)	1.973			

The BVS equation was shown: $\text{BVS} = \sum \text{Bvi} = \sum \exp[(r_0 - r)/B]$, where $r_0 = 1.917(4)$ Å for W(6+) – O(2-); $r_0 = 2.074(5)$ Å for Eu(2+) – O(2-); B=0.37. The bond valence sums of oxygens in this core Eu8-

W-O unit were given in Table S2. For the terminal oxygen atoms, their values were within the range 1.579-1.707 except for O15 and O17, they had much lower BVS values (0.335 and 0.350), compare with the bond length of Eu(3)-O(15) 2.479(11) Å and Eu(7)-O(17) 2.463(13) Å, we believe that O15 and O17 are terminal oxygens of water molecules. Whereas for the bridging oxygens (μ_2 -O, μ_3 -O) atoms, their values fall within the range 1.80-2.16 except for O13, O14, O16 and O18, we note that the BVS values were rather high for the bridging oxygens atoms in Eu-O-W and W-O-W bonds than that for terminal oxygens W-O. The bridging atoms, O13, O14, O16 and O18, had much lower BVS values (0.904, 0.927, 1.040 and 1.010), were presumed to be hydroxo groups. Overall, the four keggin-like structures are each completed by two Eu^{3+} that are further connected to form the “cluster” through μ -O, μ -OH and tungstate units.

Figure S6 Representation as 50% ellipsoids of **3**. Uncoordinated water molecules and hydrogen atoms removed for clarity.

Figure S7. Position of Cs^+ ions in **3**. The $\text{PW}_{11}\text{O}_{39}$ unit is represented in polyhedral notation. Color code: Cs^+ ion: pink. W: blue; Eu: green.