Supplementary Material

Aryl Annulation of Cyclic Ketones via a Magnesium**Carbometallation-6- π -Electrocyclization Protocol**

**Pierre E. Tessier,[†] Natalie Nguyen, Matthew Clay,
and Alex G. Fallis***

Department of Chemistry, University of Ottawa, 10 Marie Curie,

Ottawa, Ontario, Canada K1N 6N5

afallis@science.uottawa.ca

2-Trifluoromethanesulfonylbicyclo[2.2.1]heptene (7)³: *n*-Butyllithium (6.4 mL, 2.5 M, 16.0 mmol), diisopropylamine (2.10 mL, 16.0 mmol), norcamphor (1.6 g, 14.5 mmol) in THF (5 mL), *N*-phenyltrifluoromethanesulfonimide (5.45 g, 15.3 mmol) were reacted in THF (10 mL). Chromatography (100% petroleum ether) afforded **7** as a colourless oil (2.06 g, 59%). ¹H NMR (CDCl₃, 300 MHz): δ 5.65 (d, J = 3.5 Hz, 1H), 2.98 – 2.96 (m, 2H), 1.82 – 1.69 (m, 2H), 1.67 – 1.61 (m, 1H), 1.49 – 1.34 (m, 1H), 1.25 – 1.13 (m, 3H), 0.89 – 0.79 (m, 2H); ¹³C NMR (CDCl₃, 75 MHz): δ 155.3, 125.3, 121.1, 120.3, 116.8, 112.6, 48.0, 44.1, 41.7, 25.9, 24.6.

Trifluoromethanesulfonyl-1,7,7-trimethylbicyclo[2.2.1]hept-2-ene (10)⁴: Diisopropylamine (8.10 mL, 57.8 mmol) was dissolved in THF (125 mL), cooled to –78 °C and a solution of *n*-BuLi in hexane (2.5 M, 23.1 mL, 57.8 mmol) was added. The solution was stirred for 15 minutes, and then a solution of camphor (8.00 g, 52.6 mmol) in THF (75 mL) was added. After stirring at –78 °C for 90 minutes, a solution of PhNTf₂ (19.7 g, 55.2 mmol) in THF (75 mL) was added and the reaction was stirred at –78 °C for 15 minutes and then at 0 °C for 1 hour. The reaction was concentrated to ~100 mL on the rotary evaporator (T = 40 °C), diluted with water (200 mL), and extracted with ether (3 x 200 mL). The combined organics were dried (MgSO₄), filtered, and carefully concentrated (CAUTION: Product is volatile!). Column chromatography (petroleum ether) afforded the title compound as a clear colorless oil (13.30 g, 89%). ¹H NMR (500 MHz, CDCl₃) δ 5.67 (d, J = 3.8 Hz, 1H), 2.46 (dd, J = 3.8 and 3.8 Hz, 1H), 1.91-1.97 (m, 1H), 1.66 (ddd, J = 12.2, 8.6 and 3.7 Hz, 1H), 1.34 (ddd, J = 12.4, 9.1 and 3.7 Hz, 1H), 1.16 (ddd, J = 12.5, 9.1 and 3.7 Hz, 1H), 1.04 (s, 3H), 0.93 (s, 3H), 0.80 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 155.2, 118.6, 117.7, 56.9, 53.8, 50.1, 30.8, 25.3, 19.6, 18.9, 9.3; IR (neat): ν = 2964, 2880, 1624, 1424, 1321, 1212, 1142, 1111, 1085, 1004, 914, 880, 813, 755 cm⁻¹; MS (EI) m/z 284 (M⁺) (42), 269 (14), 151 (38), 123 (100), 95 (62), 55 (59), 41 (77); HRMS calculated for C₁₁H₁₅O₃F₃S (M⁺) 284.0694, found 284.0703.

Trifluoromethanesulfonyl-7,7-dimethylbicyclo[3.1.1]hept-2-ene (13): *n*-Butyllithium (9.3 mL, 1.6 M, 14.8 mmol), diisopropylamine (1.94 mL, 14.8 mmol), nopinone (1.9 mL, 13.46 mmol), *N*-phenyltrifluoromethanesulfonimide (5.05 g, 14.13 mmol) were reacted in THF (10 mL). Chromatography (100% ether) afforded **13** as a colourless oil (1.94 g, 54%). IR (neat); 2949 (s),

1658 (m), 1414 (s), 1224 (s) cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz): δ 5.51 (q, $J = 3.3$ Hz, 1H), 2.54 (dt, $J = 9.2$ Hz, 5.6 Hz, 1H), 2.22 – 2.40 (m, 3H), 2.09 – 2.13 (m, 1H), 1.35 (d, $J = 9.1$ Hz, 1H), 1.32 (s, 3H), 0.91 (s, 3H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 155.4, 125.2, 121.0, 116.8, 112.6, 111.8, 46.6, 40.5, 40.1, 32.1, 28.6, 25.8, 21.2; MS (EI) m/z 270.1 (M^+ (2.6), 226.0 (16.2), 162.0 (9.0), 77.0 (52.2), 55.0 (100.0); Exact mass calc'd. for $\text{C}_{10}\text{H}_{13}\text{F}_3\text{O}_3\text{S}$: 270.0517, found 270.0496.

2-Trifluoromethanesulfonylbicyclo[3.2.1]oct-2-ene (16): *n*-Butyllithium (7.0 mL, 2.5 M, 17.6 mmol), diisopropylamine (2.30 mL, 17.6 mmol), bicyclo [3.2.1] octanone (1.99 g, 16.0 mmol) in THF (5 mL), *N*-phenyltrifluoromethanesulfonimide (6.0 g, 16.8 mmol) were reacted in THF (10 mL). Chromatography (100% petroleum ether) afforded **16** as a colourless (3.55 g, 87%). IR (neat); 2955 (s), 2865 (s), 1416 (s), 1210 (s) cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz): δ 5.40 – 5.38 (m, 1H), 2.53 (t, $J = 4.7$ Hz, 1H), 2.42 – 2.36 (m, 2H), 2.15 – 2.26 (m, 1H), 1.96 – 1.84 (m, 2H), 1.68 – 1.80 (m, 2H), 1.62 – 1.54 (m, 1H), 1.47 – 1.38 (m, 1H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 154.2, 125.2, 121.0, 116.8, 112.6, 114.1, 40.0, 36.0, 34.8, 34.4, 32.5, 30.3; MS (EI) m/z 256.0 (M^+ (44.2), 228.0 (20.3), 95.1 (100.0), 79.1 (57.4), 67.1 (74.8)); Exact mass calc'd. for $\text{C}_9\text{H}_{11}\text{F}_3\text{O}_3\text{S}$: 256.0379, found 256.0376.

1-Trifluoromethanesulfonyl-5-methoxy-3,4-dihydronaphthalene (34) *n*- Butyllithium (6.0 mL, 2.5 M, 14.6 mmol), diisopropylamine (1.92 mL, 14.6 mmol), 6-methoxy tetralone (2.35 g, 13.3 mmol) in THF (5 mL), *N*-phenyltrifluoromethanesulfonimide (5.05 g, 14.13 mmol) were reacted in THF (10 mL). Chromatography (petroleum ether/ethyl acetate, 19:1) afforded **36** as beige oil (3.30 g, 82%). IR (neat); 2930 (s), 2724 (m), 1693 (s), 1608 (m), 1382 (m), 1230 (m) cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz): δ 7.27 (d, $J = 8.4$ Hz, 1H), 6.78 – 6.69 (m, 2H), 5.85 (t, $J = 4.8$ Hz, 1H), 3.79 (s, 3H), 2.82 (t, $J = 8.0$ Hz, 2H), 2.47 (dt, $J = 4.6$ Hz, 8.3 Hz, 2H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 160.6, 146.7, 138.7, 131.3, 125.5, 123.1, 121.9, 121.1, 116.9, 115.1, 114.6, 112.6, 111.7, 55.6, 27.7, 22.6; MS (EI) m/z 308.0 (M^+ (70.6), 175.1 (41.1), 147.1 (100.0), 115.1 (37.9), 77.0 (28.8)); Exact mass calc'd. for $\text{C}_{12}\text{H}_{11}\text{F}_3\text{O}_4\text{S}$: 308.0328, found 308.0326.

1-Trifluoromethanesulfonyl-3,4-dihydro-naphthalene (37)⁵: *n*-Butyllithium (7.0 mL, 2.4 M, 16.7 mmol), diisopropylamine (2.18 mL, 16.7 mmol), tetralone (2.02 mL, 15.2 mmol) in THF (5 mL), *N*-phenyltrifluoromethanesulfonimide (5.7 g, 16.0 mmol) were reacted in THF (10 mL). Chromatography (petroleum ether) afforded **37** as a colourless oil (6.11 g, 82 %). ^1H NMR (CDCl_3 , 500 MHz): δ 7.37 – 7.35 (m, 1H), 7.27 – 7.25 (m, 2H), 7.18 – 7.17 (m, 1H), 6.02 (t, $J = 4.8$ Hz, 1H), 2.86 (t, $J = 8.0$ Hz, 2H), 2.52 – 2.48 (m, 2H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 146.4, 136.2, 129.1, 128.6, 127.7, 126.9, 122.4, 121.2, 119.9, 117.7, 117.4, 26.8, 22.3.

1-Trifluoromethanesulfonylcyclohex-1-ene (40)⁶: *n*-Butyllithium (5.96 mL, 2.5 M, 14.9 mmol), diisopropylamine (1.95 mL, 14.9 mmol), cyclohexanone (1.4 mL, 13.5 mmol), *N*-phenyltrifluoromethanesulfonimide (5.1 g, 14.2 mmol) in THF (10 mL). Chromatography (100% petroleum ether) afforded **40** as a colourless oil (1.72 g, 55 %). ^1H NMR (CDCl_3 , 200 MHz): δ 5.73 (t, $J = 3.8$ Hz, 1H), 2.29 – 2.20 (m, 2H), 2.17 – 2.13 (m, 2H), 1.82 – 1.70 (m, 2H), 1.63 – 1.54 (m, 2H); ^{13}C NMR (CDCl_3 , 50 MHz): δ 149.3, 128.0, 121.7, 118.3, 115.3, 108.9, 27.3, 23.6, 22.4, 20.7.

Sonogashira Coupling of Propargyl Alcohol to a Vinyl Triflate: $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (0.05 eq) and copper (I) iodide (0.1 eq) were dissolved in a 1:1 solution of THF:DEA (diethylamine). The

solution was degassed with N₂ for 10 minutes. The appropriate vinyl triflate (1 eq) was dissolved in THF and the solution degassed with N₂ for an additional 10 minutes. Propargyl alcohol (1.1 eq) was added, via syringe, and the initial green solution changed from yellow, to orange to red. The solution was refluxed for 16 hours. The THF and DEA were removed under reduced pressure (rotary evaporator). Purification was achieved by flash silica gel chromatography with an appropriate petroleum ether/ethyl acetate solvent mixture.

3-Bicyclo[2.2.1]hept-2-yl-prop-2-yn-1-ol (8): Pd(PPh₃)₂Cl₂ (269 mg, 0.38 mmol), copper (I) iodide (145 mg, 0.76 mmol), THF:DEA (50 mL), (7) (1.85 g, 7.65 mmol) in THF (10 mL), and propargyl alcohol (0.49 mL, 8.42 mmol) were reacted together. Chromatography (petroleum ether/ethyl acetate 5:1) afforded **8** as a viscous red oil (680 mg, 60%). IR (CDCl₃): 3606 (s), 3445 (s), 2968 (s), 2865 (s), 1378 (s) cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 6.17 (d, *J* = 3.0 Hz, 1H), 4.32 (s, 2H), 3.24 (br s, 1H), 2.82 – 2.80 (m, 2H), 1.67 – 1.52 (m, 3H), 1.39 – 1.33 (m, 2H), 1.19 – 0.94 (m, 4H); ¹³C NMR (CDCl₃, 75 MHz): δ 142.1, 129.3, 90.7, 82.4, 51.7, 48.4, 47.3, 43.0, 25.5, 24.7; MS (EI) *m/z* 148.1 (M⁺ (37.6), 120.1 (63.9), 91.1 (100.0), 77.0 (33.8); Exact mass calc'd. for C₁₂H₁₆O: 176.1183, found 176.1164.

(1,7,7-Trimethyl-bicyclo[2.2.1]hept-2-en-2-yl)-prop-2-yn-1-ol (11): Triflate **10** (2.00 g, 7.03 mmol), CuI (0.13 g, 0.70 mmol), Pd(PPh₃)₂Cl₂ (0.25 g, 0.35 mmol), Et₂NH (20 mL) and THF (20 mL) were added to a flask and degassed with argon for 20 minutes. Propargyl alcohol (0.49 mL, 8.44 mmol) was added and the reaction was heated at reflux overnight. The next day the reaction was cooled to room temperature and silica gel (~3 g) was added. After removal of the solvent, dry-pack column chromatography (9:1, petroleum ether:ethyl acetate) afforded the title compound as a yellow oil that solidified upon refrigeration (1.30 g, 97%). mp < 20 °C. ¹H NMR (500 MHz, CDCl₃) δ 6.27 (d, *J* = 3.4 Hz, 1H), 4.43 (s, 2H), 2.34 (dd, *J* = 3.6 and 3.6 Hz, 1H), 1.88 (m, 2H), 1.55 (ddd, *J* = 11.9, 9.1 and 3.4 Hz, 1H), 1.07 (m, 2H), 1.05 (s, 3H), 1.00 (ddd, *J* = 12.4, 9.1 and 3.4 Hz, 1H), 0.79₁ (s, 3H), 0.78₇ (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 141.4, 131.7, 91.1, 82.2, 60.6, 56.7, 55.8, 52.3, 52.0, 31.3, 25.0, 19.8, 19.6, 12.0; IR (neat): ν = 3329, 3062, 2955, 2871, 2214, 1387, 1366, 1291, 1141, 1026, 1005, 825 cm⁻¹; MS (EI) *m/z* 190 (M⁺) (100), 175 (66), 162 (59), 147 (63), 129 (36), 115 (48), 105 (37), 91 (73), 77 (48), 41 (66); HRMS calculated for C₁₃H₁₈O (M⁺) 190.1358, found 190.1349.

(Bicyclo[3.2.1]oct-2-en-2-yl)-prop-2-yn-1-ol (17): Pd(PPh₃)₂Cl₂ (344 mg, 0.49 mmol), copper (I) iodide (187 mg, 0.98 mmol), THF:DEA (80 mL), **16**, (2.50 g, 9.80 mmol) in THF (5 mL), and propargyl alcohol (0.63 mL, 10.8 mmol) were reacted together. Chromatography (neutral alumina gel (petroleum ether/ethyl acetate 7:1) afforded **17** as a viscous orange oil (1.03 g, 65%). IR (neat): 3341 (s), 2942 (s), 2865 (m), 1029 (m) cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 5.72 (t, *J* = 3.1 Hz, 1H), 4.30 (s, 2H), 3.05 (br s, 1H), 2.39 – 2.23 (m, 3H), 1.89 – 1.72 (m, 3H), 1.69 – 1.61 (m, 1H), 1.56 – 1.52 (m, 1H), 1.45 – 1.30 (m, 3H); ¹³C NMR (CDCl₃, 75 MHz): δ 132.2, 128.3, 86.9, 85.8, 51.6, 40.7, 37.7, 35.5, 35.3, 32.8 30.9; MS (EI) *m/z* 162.1 (M⁺ (100.0), 134.1 (26.9), 105.1 (33.8), 91.1 (67.7), 77.0 (45.8)); Exact mass calc'd. for C₁₁H₁₄O: 162.1037, found 162.1030.

(5-Methoxy-3,4-dihydro-naphthalen-1-yl)-prop-2-yn-1-ol (35): Pd(PPh₃)₂Cl₂ (204 mg, 0.29 mmol), copper (I) iodide (110 mg, 0.58 mmol), THF:DEA (50 mL), **(34)** (1.76 mg, 5.8 mmol) in THF (3 mL), and propargyl alcohol (0.37 mL, 6.3 mmol) were reacted together. Chromatography

(petroleum ether/ethyl acetate, 7:1) afforded **35** as a viscous red oil (600 mg, 53%). IR (neat); 3405 (s), 2944 (s), 2842 (s), 1690 (m) cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz): δ 7.45 (d, J = 8.4 Hz, 1H), 6.71 (dd, J = 8.4 Hz, 2.6 Hz, 1H), 6.65 (d, J = 2.6 Hz, 1H), 6.31 (t, J = 4.8 Hz, 1H), 4.48 (s, 2H), 3.78 (s, 3H), 2.73 (t, J = 8.0 Hz, 2H), 2.36 – 2.29 (m, 2H), 2.29 (br s, 1H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 159.5, 137.1, 133.9, 126.6, 126.0, 121.0, 113.9, 111.5, 88.3, 84.1, 55.7, 52.0, 28.0, 23.9; MS (EI) m/z 214.1 (M^+ (100.0), 171.1 (12.9), 152.1 (17.4), 115.1 (17.3), 63.0 (7.6)); Exact mass calc'd. for $\text{C}_{14}\text{H}_{14}\text{O}_2$: 214.0982, found 214.0971.

(3,4-Dihydro-naphthalen-1-yl)-prop-2-yn-1-ol (38): $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (428 mg, 0.61 mmol), copper (I) iodide (232 mg, 1.22 mmol), THF:DEA (50 mL), Compound **37** (3.4 g, 12.2 mmol) in THF (10 mL), and propargyl alcohol (0.71 mL, 12.2 mmol) were reacted together. Chromatography (petroleum ether/ethyl acetate, 7:1) afforded **38** as a viscous red oil (1.19 g, 53%). IR (neat); 3359 (s), 2935 (s), 2830 (s), 1487 (m) cm^{-1} ; ^1H NMR (CDCl_3 , 500 MHz): δ 7.59 (d, J = 7.5 Hz, 1H), 7.23 (td, J = 7.4 Hz, 0.8 Hz, 1H), 7.18 (td, J = 7.4 Hz, 1.4 Hz, 1H), 7.10 (d, J = 7.3 Hz, 1H), 6.47 (t, J = 4.8 Hz, 1H), 4.52 (s, 2H), 2.78 (t, J = 7.9 Hz, 2H), 2.75 (br s, 1H), 2.38 – 2.34 (m, 2H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 135.8, 134.8, 132.3, 127.6, 127.2, 126.5, 124.8, 121.1, 88.1, 83.3, 51.3, 26.9, 23.4; MS (EI) m/z 184.1 (M^+ (100.0), 152.1 (62.8), 141.1 (51.8), 115.1 (41.4)); Exact mass calc'd. for $\text{C}_{13}\text{H}_{12}\text{O}$: 184.0860, found 184.0832.

(Cyclohex-1-enyl)-prop-2-yn-1-ol (41): $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (175 mg, 0.25 mmol), copper (I) iodide (95 mg, 0.5 mmol), THF:DEA (30 mL), triflate **40** (1.15 g, 5.0 mmol) in THF (10 mL), and propargyl alcohol (0.29 mL, 5.5 mmol) were reacted together. Chromatography (petroleum ether/ethyl acetate, 4:1) afforded **41** as a viscous red oil (431 mg, 64%). IR (neat); 3619 (s), 3438 (s), 2929 (s), 2871 (s), 1442 (s) cm^{-1} ; ^1H NMR (CDCl_3 , 500 MHz): δ 6.10 (quintet, J = 2.1 Hz, 1H), 4.35 (s, 2H), 2.10 – 2.05 (m, 4H), 1.63 – 1.54 (m, 4H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 135.5, 120.1, 87.6, 84.5, 51.6, 29.1, 25.6, 22.2, 21.4; MS (EI) m/z 136.1 (M^+ (100.0), 107.0 (51.7), 91.1 (58.1), 79.1 (78.1)); Exact mass calc'd. for $\text{C}_9\text{H}_{12}\text{O}$: 136.0887, found 135.0886.

Carbometallation of Propargyl Alcohols: The appropriate vinyl propargyl alcohol was dissolved in either toluene or cyclohexane. Vinyl magnesium chloride in THF (3.2 eq) was added via syringe and the resulting solution refluxed for 16 hours. The reaction was cooled to 0 °C, the appropriate electrophile added, and stirred for 30 minutes. The reaction was quenched with water (50 mL) or saturated aqueous ammonium chloride solution. The reaction mixture was extracted with diethyl ether (2 x 50 mL) and the concentrated extract purified by silica gel chromatography with an appropriate petroleum ether/ethyl acetate solvent eluent. (More recent experiments have indicated yields are increased for these Grignard additions by addition of 10 mol percent of CuI and an equivalent of MgCl_2)

Tricyclo[6.2.1.0^{2,7}]undeca-2,4,-dien-4-methanol (9): Compound **8** (210 mg, 1.42 mmol) in toluene (5 mL), and vinyl magnesium chloride (2.7 mL, 1.7 M, 4.54 mmol) were reacted together and the reaction quenched with saturated ammonium chloride (30 mL). Chromatography (petroleum ether/ethyl acetate 6:1) afforded **9** as an orange oil (105 mg, 42%). IR (CDCl_3); 3439 (m), 2958 (s), 2867 (s), 1469 (m), 1371 (m) cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz): δ 5.57 (s, 1H), 5.55 – 5.53 (m, 1H), 4.04 (q, J = 12.3 Hz, 2H), 2.71 (s, 1H), 2.15 – 2.02 (m, 3H), 1.82 – 1.59 (m, 4H), 1.47 (br s, 1H), 1.42 – 1.30 (m, 4H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 153.4, 137.8, 120.0, 110.4, 66.1, 44.4, 44.2, 42.1, 39.5, 31.0, 27.2, 26.5; MS (EI) m/z 176.1

(M^+ (26.0), 145.1 (48.3), 117.1 (87.1), 105.1 (96.9), 91.1 (100.0), 39.0 (60.7)); Exact mass calc'd. for $C_{12}H_{16}O$: 176.1183, found 176.1164.

1,11,11-Trimethyltricyclo[6.2.1.0^{2,7}]undeca-2,4,-dien-4-methanol (12): Compound **11** (467 mg, 2.46 mmol) in cyclohexane (10 mL), and vinyl magnesium chloride (6.15 mL, 1.6 M, 9.84 mmol) were reacted together and the reaction quenched with saturated ammonium chloride (30 mL). Chromatography (petroleum ether/ethyl acetate 7:1) afforded **12** as an orange oil (391 mg, 73%). IR (neat); 3355 (s), 2952 (s), 2869 (s), 1670 (m), 1448 (m), 1045 (m) cm^{-1} ; 1H NMR (C_6D_6 , 200 MHz): δ 5.59 (s, 1H), 5.52 – 5.49 (m, 1H), 4.07 (s, 2H), 2.55 – 2.39 (m, 1H), 2.20 – 2.01 (m, 2H), 1.89 – 1.71 (m, 3H), 1.31 – 1.13 (m, 1H), 1.04 (s, 3H), 0.96 (s, 3H), 0.83 (s, 3H); ^{13}C NMR (C_6D_6 , 50 MHz): δ 154.3, 137.5, 118.0, 109.8, 65.5, 50.9, 50.4, 48.1, 45.5, 32.3, 30.5, 27.7, 21.8, 20.7, 11.8; MS (EI) m/z 218.0 (M^+ (17.3), 216.0 (20.9), 171.0 (34.8), 157.0 (83.0), 143.0 (100.0)); Exact mass calc'd. for $C_{15}H_{22}O$: 218.1685, found 218.1700.

(6,6-Dimethylbicyclo[3.1.1]hept-2-en-2-yl)buta-1,3-dien-2-methanol (15): Compound **14** (1.0 g, 5.3 mmol) in toluene (10 mL) and vinyl magnesium chloride (10 mL, 1.7 M, 16.96 mmol) were reacted together and the reaction quenched with saturated ammonium chloride (30 mL). Chromatography (petroleum ether/ethyl acetate 7:1) afforded **15** as an orange oil (830 mg, 77%). IR (neat); 3357 (s), 2990 (s), 2920 (s), 1461 (m), 1008 (m) cm^{-1} ; 1H NMR ($CDCl_3$, 300 MHz): δ 6.75 (ddd, J = 0.9 Hz, 11.2 Hz, 17.9 Hz, 1H), 5.96 (s, 1H), 5.54 – 5.57 (m, 1H), 5.31 (d, J = 17.9 Hz, 1H), 5.10 (dt, J = 11.2 Hz, 1.4 Hz, 1H), 4.28 (s, 2H), 2.34 – 2.40 (m, 3H), 2.24 (dt, J = 1.4 Hz, 5.7 Hz, 1H), 2.05 – 2.11 (m, 1H), 1.73 (br s, 1H), 1.27 (s, 3H), 1.19 (d, J = 8.7 Hz, 1H), 0.85 (s, 3H); ^{13}C NMR ($CDCl_3$, 75 MHz): δ 144.8, 135.3, 133.6, 131.5, 125.5, 114.4, 65.1, 46.9, 40.7, 38.2, 32.4, 31.9, 26.6, 21.5; MS (EI) m/z 204.2 (M^+ (15.2), 161.1 (24.0), 143.1 (38.8), 129.1 (52.8), 117.1 (43.9), 105.1 (100.0), 91.1 (61.3)); Exact mass calc'd. for $C_{14}H_{20}O$: 204.1501, found 204.1487.

(Bicyclo[3.2.1]oct-2-en-2-yl)buta-1,3-dien-2-methanol (18): Compound **17** (1.0 g, 6.17 mmol) in toluene (15 mL) and vinyl magnesium chloride (11.6 mL, 1.7 M, 19.7 mmol) were reacted together and the reaction quenched with saturated ammonium chloride (30 mL). Chromatography (petroleum ether/ethyl acetate 8:1) afforded **18** as a yellow oil (750 mg, 64%). IR (neat); 3329 (s), 2936 (s), 2865 (s), 1442 (m), 1088 (m) cm^{-1} ; 1H NMR ($CDCl_3$, 300 MHz): δ 6.74 (dd, J = 17.9 Hz, 11.3 Hz, 1H), 5.97 (m, 1H), 5.34 (m, 1H), 5.31 (d, J = 17.8 Hz, 1H), 5.09 (d, J = 11.2 Hz, 1H), 4.30 (s, 2H), 2.47 – 2.31 (m, 3H), 2.01 – 1.38 (m, 8H); ^{13}C NMR ($CDCl_3$, 75 MHz): δ 142.5, 135.5, 133.5, 132.3, 126.1, 114.2, 65.2, 41.6, 37.8, 35.6, 35.2, 33.1, 31.1; MS (EI) m/z 190.1 (M^+ (20.8), 159.1 (23.0), 129.1 (29.8), 105.1 (29.0), 91.1 (100.0), 79.1 (43.4)); Exact mass calc'd. for $C_{13}H_{18}O$: 190.1349, found 190.1341.

(Bicyclo[3.2.1]oct-2-en-2-yl)buta-1,3-dien-2-methoxymethyl (22): Compound **18** (680 mg, 3.6 mmol) was dissolved in dry THF (20 mL). $KOtBu$ in THF (17.9 mL, 1M, 18.0 mmol) was added via syringe and stirred for 10 minutes. MeI (1.98 mL, 18.0 mmol) was added via syringe and the reaction stirred at room temperature for 3 hours. The reaction was quenched with water (50 mL) and extracted with diethyl ether (3 x 30 mL) to afford **22** as a yellow oil (728 mg, 99 %). IR (neat); 2943 (s), 2865 (m), 1448 (w), 1094 (m) cm^{-1} ; 1H NMR ($CDCl_3$, 300 MHz): δ 6.75 (dd, J = 17.8 Hz, 11.2 Hz, 1H), 5.93 (m, 1H), 5.34 (m, 1H), 5.31 (d, J = 18.1 Hz, 1H), 5.06 (d, J = 11.2 Hz, 1H), 4.06 (s, 2H), 3.31 (s, 3H), 2.47 – 2.25 (m, 3H), 1.95 – 1.59 (m, 5H), 1.52 – 1.38 (m,

2H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 142.5, 133.6, 132.7, 131.4, 125.9, 114.4, 75.0, 58.2, 41.5, 39.8, 35.6, 35.2, 32.7, 29.2; MS (EI) m/z 204.2 (M^+ (27.8), 159.1 (22.0), 129.1 (41.6), 104.1 (45.1), 91.1 (100.0), 45.0 (66.3)); Exact mass calc'd. for $\text{C}_{14}\text{H}_{20}\text{O}$: 204.1520, found 204.1526.

7,7-Dimethylbicyclo[3.1.1]hept-2,3-benzo-3'-carbaldehyde (23):

Compound **15** (150 mg, 0.74 mmol) was dissolved in toluene (10 mL) in a sealed tube heated at 150 °C for 16 h. The crude product was added to toluene (20 mL), DDQ (500 mg, 1.85 mmol) was added and the reaction mixture refluxed for 3 hours. The DDQ was removed by filtration and the toluene removed with a rotary evaporator under reduced pressure. Chromatography (petroleum ether/ethyl acetate 20:1) afforded **23** as a yellow oil (82 mg, 56%). IR (neat); 2946 (m), 2840 (m), 1609 (m), 1444 (s), 1248 (s) cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz): δ 9.90 (s, 1H), 7.62 (d, J = 7.7 Hz, 1H), 7.41 (s, 1H), 7.25 (d, J = 7.2 Hz, 1H), 3.02 (s, 2H), 2.82 (t, J = 5.5 Hz, 1H), 2.69 – 2.62 (m, 1H), 2.31 – 2.27 (m, 1H), 1.38 (s, 3H), 1.20 (d, J = 9.6 Hz, 1H), 0.60 (s, 3H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 192.8, 148.2, 143.5, 134.2, 128.9, 128.8, 126.7, 48.0, 40.5, 39.5, 34.0, 32.1, 26.4, 21.7; MS (EI) m/z 200.1 (M^+ , (6.8), 157.1 (38.4), 129.1 (100.0), 69.0 (28.7); Exact mass calc'd. for $\text{C}_{14}\text{H}_{16}\text{O}$: 200.1202, found 200.1201.

7,7-Dimethylbicyclo[3.1.1]hept-2,3-benzo-3'-hydroxymethyl (21): Vinyl magnesium bromide (Acros Organics, 15% wt in THF, 2.6 mL, 4.34 mmol) was added dropwise to a solution of 7,7-Dimethylbicyclo[3.1.1]hept-2-en-2-yl)-prop-2-yn-1-ol (**14**) (191 mg, 1.08 mmol) in toluene:THF (5:1, 13 mL). The brown reaction mixture stirred at reflux for 16 h. The reaction was quenched with saturated ammonium chloride solution (10 mL) at 0°C and the mixture stirred for 10 min. Dichloromethane was added followed by water and the aqueous phase was extracted with dichloromethane (3x). The combined organic phases was washed with H_2O (2x) and brine, dried with MgSO_4 and concentrated. Chromatography (petroleum ether/diethyl ether, 3:1) afforded product as a clear oil (115 mg, 53%). IR (neat) ν 3336.6, 2980.6, 2916.2, 2867.7, 2833.5, 1721.4, 1622.1, 1495.4, 1025.1 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.11-7.07 (m, 2H) 6.90 (s, 1H) 4.60 (s, 2H) 2.94 (bs, 2H) 2.71 (t, 5.5 Hz, 1H) 2.60 (dt, 9.3, 5.9 Hz, 1H) 2.25 (m, 1H) 1.35 (s, 3H) 1.23 (bs, 1H) 1.21 (d, 9.3 Hz, 1H) 0.61 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 147.1, 137.3, 134.4, 127.8, 124.7, 124.5, 62.4, 47.7, 40.4, 39.0, 32.6, 31.8, 26.0, 21.2; MS (EI) m/z 202.1 (M^+ , 8.5) 159.1 (14.2) 129.1 (100) 115.1 (7.2); HRMS calcd for $\text{C}_{14}\text{H}_{18}\text{O}$ 202.1356 (M^+), found 202.1353.

Bicyclo[3.2.1]oct-2,3-benzo-3'-carbaldehyde (24): Compound **25** (90.0 mg, 0.44 mmol) was dissolved in toluene (30 mL). DDQ (245 mg, 0.88 mmol) was added and the reaction was refluxed and stirred for 3 hours. The reaction was quenched with water (30 mL) and extracted with diethyl ether (3 x 20 mL). Chromatography (petroleum ether/ethyl acetate 19:1) afforded **24** as a yellow oil (52 mg, 64%). IR (CDCl_3); 2949 (m), 2872 (w), 1693 (s), 1609 (m), 1223 (w) cm^{-1} ; ^1H NMR (CDCl_3 , 500 MHz): δ 9.90 (s, 1H), 7.57 (dd, J = 7.8 Hz, 1.8 Hz, 1H), 7.49 (d, J = 7.17 (d, J = 7.8 Hz, 1H), 3.13 – 3.07 (m, 2H), 2.67 (d, J = 17.8 Hz, 1H), 2.57 – 2.54 (m, 1H), 1.98 – 1.89 (m, 2H), 1.80 – 1.72 (m, 3H), 1.48 – 1.44 (m, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 192.3, 146.3, 142.8, 134.3, 130.1, 128.2, 127.4, 41.1, 40.0, 36.2, 35.3, 33.3, 29.7; MS (EI) m/z 186.1 (M^+ (42.4), 162.0 (9.0), 143.0 (7.2), 129.1 (100.0), 115.1 (20.8); Exact mass calc'd. for $\text{C}_{13}\text{H}_{14}\text{O}$: 186.1012, found 186.0979.

1,11,11,3'-Tetramethyltricyclo[6.2.1.0^{2,7}]undeca-2,4,-dien-4-methanol (26): Propargyl alcohol **11** (465 mg, 2.45 mmol) in cyclohexane (5 mL), vinyl magnesium chloride (4.6 mL, 1.7 M, 7.84 mmol), and MeI (1.5 mL, 24.5 mmol) were reacted together. Chromatography (petroleum ether/ethyl acetate, 6:1) afforded **26** as a yellow oil (520 mg, 92%). IR (neat) 3370 (s), 2954 (s), 2872 (s), 1472 (m), 1387 (m) cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz): δ 5.65 (t, $J = 3.9$ Hz, 1H), 4.22 (d, $J = 12.3$ Hz, 1H), 4.02 (d, $J = 12.3$ Hz, 1H), 2.37 – 2.24 (m, 1H), 2.07 – 1.99 (m, 1H), 1.85 (s, 3H), 1.59 – 1.49 (m, 3H), 1.27 – 1.21 (m, 2H), 1.17 (s, 3H), 1.04 (s, 3H), 0.99 – 0.88 (m, 1H), 0.83 (s, 3H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 146.4, 140.2, 122.9, 118.5, 65.1, 53.0, 52.0, 48.1, 47.8, 34.5, 31.1, 28.2, 22.9, 21.5, 16.5, 13.0; MS (EI) m/z 232.2 (M^+ (53.1), 145.1 (87.7), 133.1 (100.0), 105.1 (67.4), 41.0 (71.9)); Exact mass calc'd. for $\text{C}_{16}\text{H}_{24}\text{O}$: 232.1804, found 232.1781.

4,14,14-Trimethyl-7-boryl-(1-hydroxy)-8-oxa-tetracyclo[9.2.1.0^{5,13}.0^{6,10}]dodeca-5,10- diene (28): Propargyl alcohol **11** (430 mg, 2.26 mmol) and magnesium chloride (323 mg, 3.39 mmol) were dissolved in dry cyclohexane (5 mL). Vinyl magnesium chloride (4.3 mL, 1.7 M, 7.23 mmol) was added via syringe at room temperature and the resulting red solution refluxed for 14 hours. The solution was cooled to 0 °C and quenched with trimethyl borate (0.76 mL, 6.78 mmol) and stirred for 1 hour. The reaction was quenched with water (30 mL) and extracted with diethyl ether (3 x 20 mL). Chromatography (petroleum ether/ethyl acetate, 15:1) afforded **28** as white oily solid (303 mg, 55%). IR (neat); 3612 (m), 3296 (m), 2955 (s), 1667 (s), 1402 (s), 1255 (m) cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz): δ 5.22 (br s, 1H), 5.11 (s, 1H), 4.61 (s, 2H), 2.46 – 2.37 (m, 1H), 2.23 – 2.09 (m, 1H), 1.96 – 1.85 (m, 1H), 1.72 – 1.63 (m, 2H), 1.56 – 1.43 (m, 1H), 1.32 – 1.20 (m, 1H), 1.18 (s, 3H), 0.99 (s, 3H), 0.89 (s, 3H), 0.83 – 0.76 (m, 1H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 167.9, 142.4, 110.8, 70.8, 53.7, 52.7, 48.3, 48.2, 32.9, 30.6, 28.1, 22.6, 21.4, 14.0; MS (EI) m/z 244.2 (M^+ , (11.6), 174.1 (68.8), 162.0 (36.3), 143.0 (31.8), 131.1 (100.0), 100.0 (65.5), 69.0 (78.7)); Exact mass calc'd. for $\text{C}_{13}\text{H}_{21}\text{BO}_2$: 244.1642, found 244.1650.

1,7,7-Trimethyl-2,3-benzo-(5,6-(1-hydroxybora2-2oxacyclopentan)-bicyclo[2.2.1]heptane (30): DDQ (70.4 mg, 0.309 mmol, 1.1 equiv) was added to a solution of **28** (68.8 mg, 0.281 mmol) in toluene (1.87 mL). The mixture was refluxed for 4 hrs. The mixture was diluted with water and the aqueous phase was extracted with diethyl ether (3 x 10 mL). The combined organic layers were washed with water (1 x 10 mL), brine (1 x 10 mL) and dried over MgSO_4 . Chromatography (petroleum ether/ Et_2O , 19:1) afforded **30** (32.6 mg, 48%) as a white solid (mp 256-258). IR (neat) 3358, 3050, 2953, 2872, 1709, 1604, 1490, 1470, 1445, 1430, 1405, 1376, 1278, 1196, 1020, 985, 795 cm^{-1} . ^1H NMR (CDCl_3 , 500 MHz): δ 7.21 (d, $J = 7.4$, 1H), 7.01 (d, $J = 7.4$, 1H), 5.39 (s, 1H) 5.10-5.03 (m, 2H) 2.81 (d, $J = 3.9$, 1H) 2.09-2.03 (m, 1H) 1.86-1.81 (m, 1H) 1.49 (s, 3H) 1.17-1.12 (m, 1H) 1.08-1.03 (m, 1H) 0.95 (s, 3H) 0.53 (s, 3H). ^{13}C NMR (CDCl_3 , 126 MHz): δ 155.3, 151.6, 146.9, 123.8, 117.5, 71.3, 57.6, 54.8, 53.0, 33.2, 33.1, 26.3, 20.1, 19.3, 12.9. Exact mass calc'd. for $\text{C}_{15}\text{H}_{19}\text{BO}$: 242.1478, found 242.1478.

1,11,11-Trimethyltricyclo[6.2.1.0^{2,7}]undeca-2,4,-dien-3-iodo-4-hydroxymethanol (29): Magnesium chloride (10.33 g, 108.5 mmol) was added to a solution of propargyl alcohol **11** (9.38 g, 49.30 mmol) in cyclohexane (100 mL), and a solution of vinyl magnesium chloride in THF (1.6 M, 108 mL, 172.5 mmol) was then added. The mixture was heated at reflux overnight. The reaction was then cooled to 0 °C and a solution of I_2 (37.54 g, 147.9 mmol) in THF (140 mL) was added until the dark brown color of I_2 persisted. After stirring at 0 °C for 2 hours, the

reaction was quenched via the addition of 1 M Na₂S₂O₃ (200 mL) and extracted with ether (3 x 300 mL). The organics were combined, dried (MgSO₄), filtered, and concentrated. Column chromatography (10:1, petroleum ether:ethyl acetate) afforded **29** as a yellow oil that solidified upon refrigeration (13.77 g, 81%). Yields varied from 70-80%. mp < 20 °C. ¹H NMR (500 MHz, CDCl₃) δ 5.68-5.70 (m, 1H), 4.28-4.32 (m, 1H), 4.14 (ddd, *J* = 13.6, 1.4 and 1.4 Hz, 1H), 2.46 (dd, *J* = 18.9 and 8.2 Hz, 1H), 1.95-2.04 (m, 1H), 1.73-1.79 (m, 1H), 1.58-1.68 (m, 2H), 1.48-1.54 (m, 1H), 1.36-1.42 (m, 1H), 1.29 (s, 3H), 1.24 (d, *J* = 3.8 Hz, 1H), 1.00-1.05 (m, 1H), 0.97 (s, 3H), 0.68 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 155.8, 138.6, 123.4, 79.1, 69.1, 55.7, 53.2, 52.5, 48.4, 34.0, 31.2, 28.0, 22.9, 21.7, 16.8; IR (neat): ν = 3329, 2956, 2930, 2870, 1625, 1388, 1448, 1304, 1127, 1053, 987, 786 cm⁻¹; MS (EI) *m/z* 344 (M⁺) (100), 257 (16), 199 (18), 118 (41), 91 (34), 69 (26), 41 (45); HRMS calculated for C₁₅H₂₁OI (M⁺) 344.0637, found 344.0652.

1,7,7-Trimethylbicyclo[2.2.1]hept-2,3-benzo-2'-iodophenol (33): Aldehyde **31** (2.00 g, 5.88 mmol) was dissolved in CH₂Cl₂ (100 mL) and cooled to -8 °C. *m*-Chloroperoxybenzoic acid (purified using a phosphate buffer of pH 7.4) (1.52 g, 8.82 mmol) was added and the mixture was permitted to warm to 0 °C over a period of 30 minutes, and then held at that temperature for a further 60 minutes. After removal of the solvent, the solid was dissolved in 30:1 petroleum ether:ethyl acetate and filtered through a pad of silica gel, which was washed with the same solvent until no further product eluted. After removal of the solvent, the crude formate (mixed with the phenol) was dissolved in MeOH (35 mL) and TsOH (0.2 g) was added. After stirring at rt for one hour, the mixture was diluted with water (30 mL) and extracted with ether (3 x 60 mL). The organics were combined, dried (MgSO₄), filtered, and concentrated. Column chromatography (50:1, petroleum ether:ethyl acetate) afforded **33** as a gummy solid (1.10 g, 57%). ¹H NMR (500 MHz, C₆D₆) δ 6.91 (d, *J* = 7.7 Hz, 1H), 6.73 (d, *J* = 7.7 Hz, 1H), 2.66 (d, *J* = 3.8 Hz, 1H), 1.98-2.04 (m, 1H), 1.73 (ddd, *J* = 12.4, 10.0 and 3.9 Hz, 1H), 1.43 (s, 3H), 1.24 (ddd, *J* = 12.8, 9.5 and 3.3 Hz, 1H), 1.04 (ddd, *J* = 12.7, 9.2 and 3.9 Hz, 1H), 0.90 (s, 3H), 0.57 (s, 3H); ¹³C NMR (125 MHz, C₆D₆) δ 151.9, 150.7, 142.6, 122.2, 111.1, 82.4, 57.5, 57.4, 52.3, 32.5, 26.6, 20.1, 19.4, 14.6; IR (neat): ν = 3475, 3055, 2956, 2871, 1560, 1433, 1226, 1177, 1062, 908, 820, 780 cm⁻¹; MS (EI) *m/z* 328 (M⁺) (18), 285 (15), 186 (4), 158 (46), 59 (34), 31 (100); HRMS calculated for C₂₅H₃₀I₂ (M⁺) 328.0324, found 328.0339.

1,11,11-Trimethylbicyclo[2.2.1]hept-2,3-benzo-2'-iodo-3'-carbaldehyde (31): Compound **29** (5-Iodo-4,9,9-trimethyl-1,2,3,4,8,8a-hexahydro-1,4-methano-naphthalen-6-yl)-methanol (0.59 g, 1.71 mmol) was dissolved in toluene (65 mL) and MnO₂ (1.49 g, 17.14 mmol) was added. The reaction was heated at reflux for two days and then filtered through a silica pad using ether as eluent. Column chromatography (30:1, petroleum ether:ethyl acetate) afforded **31** as a yellow solid (0.43 g, 73%). mp = 75.5 – 77.5 °C; ¹H NMR (500 MHz, CDCl₃) δ 10.24 (s, 1H), 7.66 (d, *J* = 7.4 Hz, 1H), 7.16 (d, *J* = 7.4 Hz, 1H), 2.81 (d, *J* = 3.9 Hz, 1H), 2.10-2.16 (m, 1H), 1.83 (ddd, *J* = 13.8, 10.2 and 4.0 Hz, 1H), 1.56 (s, 3H), 1.34 (ddd, *J* = 12.8, 9.3 and 4.0 Hz, 1H), 1.09 (ddd, *J* = 12.8, 9.3 and 4.0 Hz, 1H), 0.96 (s, 3H), 0.60 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 197.1, 157.8, 151.4, 133.7, 129.6, 121.7, 94.8, 58.0, 57.6, 53.7, 32.2, 26.2, 20.3, 19.4, 15.5; IR (neat): ν = 3054, 2987, 2685, 1684, 1422, 1258 cm⁻¹; MS (EI) *m/z* 584 (M⁺) (51), 527 (10); HRMS calculated for C₁₅H₁₇OI (M⁺) 340.0313, found 340.0302.

1,11,11-Trimethylbicyclo[2.2.1]hept-2,3-benzo-2'-iodo-3'-carboxylic acid (32): Compound **31** (500 mg, 1.47 mmol) was dissolved in an aqueous solution of *t*-BuOH (50 mL, 5:2). NaH₂PO₄ (706 mg, 5.88 mmol) and 2-methyl butene (1.09 mL, 10.29 mmol) were added and the solution allowed to stir for 10 minutes at room temperature. After addition of NaClO₂ (798 mg, 8.82 mmol) the solution was stirred for 3 hours at room temperature. *t*-BuOH was removed on the rotary evaporator as the benzene azeotrope (3 x 20 mL). The resulting white solid was dried under vacuum over night. Chromatography (petroleum ether/ethyl acetate, 19:1) afforded **32** as a yellow oil (520 mg, 98%). ¹H NMR (CDCl₃, 300 MHz): δ 7.34 (d, *J* = 7.3 Hz, 1H), 7.04 (d, *J* = 7.3 Hz, 1H), 2.71 (d, *J* = 3.7 Hz, 1H), 2.03 (m, 1H), 1.72 (dt, *J* = 4.0 Hz, 9.5 Hz, 1H), 1.47 (s, 3H), 1.25 (dt, *J* = 3.7 Hz, 9.2 Hz, 1H), 1.02 (dt, *J* = 3.9 Hz, 12.5 Hz, 1H), 0.89 (s, 3H), 0.52 (s, 3H); ¹³C NMR (CDCl₃, 75 MHz): δ 171.2, 152.6, 151.3, 139.1, 127.4, 121.6, 86.9, 57.9, 57.7, 53.0, 32.6, 26.8, 20.7, 19.8, 16.0; MS (EI) *m/z* 356.0 (M⁺, (100.0), 328.0 (33.6), 295.0 (25.7), 201.1 (32.5), 186.1 (93.2)); Exact mass calc'd. for C₁₅H₁₇IO₂: 356.0286, found 356.0273.

5-Methoxy-3,4-dihydronaphthalen-1-yl)buta-1,3-dien-2-methanol (36a): Compound **35** (510 mg, 2.52 mmol) in toluene (20 mL) and vinyl magnesium chloride (4.9 mL, 1.7 M, 8.1 mmol) were reacted together and the reaction quenched with saturated ammonium chloride (30 mL). Chromatography (petroleum ether/ethyl acetate, 3:1) afforded **36a** as a yellow oil (433 mg, 71%). IR (neat); 3396 (s), 2934 (s), 2834 (s), 1567 (s), 1251 (s), 1042 (s) cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 7.06 (d, *J* = 8.4 Hz, 1H), 6.77 – 6.64 (m, 3H), 6.45 (s, 1H), 5.83 (t, *J* = 4.0 Hz, 1H), 5.32 (t, *J* = 18.0 Hz, 1H), 5.10 (d, *J* = 11.3 Hz, 1H), 4.45 (s, 2H), 3.77 (s, 3H), 2.75 (t, *J* = 7.7 Hz, 2H), 2.56 (br s, 1H), 2.38 – 2.32 (m, 1H); ¹³C NMR (CDCl₃, 125 MHz): δ 159.1, 139.0, 138.2, 133.5, 128.7, 128.3, 127.8, 125.9, 114.5, 114.2, 111.3, 64.3, 55.6, 28.8, 23.6; MS (EI) *m/z* 242.1 (M⁺ (100.0), 211.1 (85.9), 196.1 (17.7), 165.1 (25.7), 115.1 (14.4)); Exact mass calc'd. for C₁₆H₁₈O₂: 242.1325, found 242.1343.

8-Methoxy-4-(2-methoxymethyl-buta-1,3-dienyl)-1,2-dihydronaphthalene (36b): Compound **36a** (165 mg, 0.68 mmol) was dissolved in dry THF (20 mL). KO^tBu in THF (3.4 mL, 1M, 3.4 mmol) was added via syringe and stirred for 10 minutes. MeI (378 μL, 3.4 mmol) was added via syringe and the reaction stirred at room temperature for 3h. The reaction was quenched with water (30 mL) and extracted with diethyl ether (2 x 20 mL). This afforded **36b** as a yellow oil (168 mg, 99 %). IR (neat); 2938 (s), 2836 (s), 1493 (s), 1248 (s), 1092 (s) cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 7.12 (d, *J* = 4.0 Hz, 1H), 6.74 – 6.65 (m, 3H), 6.37 (s, 1H), 5.84 (d, *J* = 4.5 Hz, 1H), 5.34 (t, *J* = 18.8 Hz, 1H), 5.08 (d, *J* = 11.2 Hz, 1H), 4.22 (s, 2H), 3.78 (s, 3H), 3.40 (s, 3H), 2.76 (t, *J* = 7.8 Hz, 2H), 2.35 (m, 2H); ¹³C NMR (CDCl₃, 125 MHz): δ 158.8, 137.9, 135.9, 133.2, 130.0, 127.9, 127.4, 125.5, 114.4, 113.8, 111.0, 74.0, 57.9, 55.2, 28.5, 23.2; MS (EI) *m/z* 256.1 (M⁺ (100.0), 225.1 (56.8), 211.1 (93.0), 171.1 (76.0), 115.1 (38.2), 45.0 (74.8)); Exact mass calc'd. for C₁₇H₂₀O₂: 256.1441, found 256.1413.

3,4-Dihydronaphthal-2-en-1-yl)buta-1,3-dien-2-methanol (39a): Compound **38** (415 mg, 2.26 mmol) in cyclohexane (5 mL) and vinyl magnesium chloride (4.0 mL, 1.7 M, 6.78 mmol) were reacted together and the reaction quenched with NH₄Cl (20 mL). Chromatography (petroleum ether/ethyl acetate, 6:1) afforded **39a** as a yellow oil (370 mg, 77%). IR (neat); 3606 (s), 3448 (m), 3054 (s), 2987 (s), 2306 (m), 1422 (s) cm⁻¹; ¹H NMR (CDCl₃, 500 MHz): δ 7.17 (s, 4H), 6.73 (ddd, *J* = 18.0 Hz, 11.3 Hz, 0.9 Hz, 1H), 6.49 (s, 1H), 5.98 (td, *J* = 4.7 Hz, 1.6 Hz, 1H), 5.39 (dt, *J* = 18.0 Hz, 0.9 Hz, 1H), 5.13 (dt, *J* = 11.3 Hz, 1.4 Hz, 1H), 4.48 (d, *J* = 1.1 Hz, 2H), 2.80 (t,

$J = 7.7$ Hz, 2H), 2.42 – 2.37 (m, 2H), 2.20 (br s, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 138.7, 135.9, 134.6, 133.4, 132.9, 129.8, 128.2, 127.4, 127.1, 126.4, 124.2, 114.2, 64.0, 27.8, 23.2; MS (EI) m/z 212.1 (M^+ (54.5), 181.1 (100.0), 165.1 (53.2), 141.1 (35.2), 128.1 (26.5)); Exact mass calc'd. for $\text{C}_{15}\text{H}_{16}\text{O}$: 212.1200, found 212.1199.

4-(2-Methoxymethyl-buta-1,3-dienyl)-1,2-dihydro-naphthalene (39b): Compound **39a** (160 mg, 0.79 mmol) was dissolved in THF (20 mL). Sodium hydride (60 mg, 1.58 mmol) and methyl iodide (0.24 mL, 3.95 mmol) were added and the solution stirred at room temperature for 14 hours. Water (20 mL) was added followed by diethyl ether (3 x 20 mL) extraction. Chromatography (petroleum ether/ethyl acetate, 20:1) afforded **39b** as a colourless oil (155 mg, 87%). IR (CDCl_3); 3161 (m), 2936 (s), 2865 (s), 2814 (s), 1468 (m) cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz): δ 7.14 (s, 4H), 6.70 (ddd, $J = 17.9$ Hz, 11.4 Hz, 0.9 Hz, 1H), 6.41 (s, 1H), 5.98 (td, $J = 5.7$ Hz, 1.5 Hz, 1H), 5.39 (d, $J = 17.8$ Hz, 1H), 5.10 (dt, $J = 11.2$ Hz, 1.5 Hz, 1H), 2.79 (t, $J = 6.7$ Hz, 2H), 2.42 – 2.35 (m, 2H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 136.4, 135.1, 133.9, 133.4, 130.3, 130.2, 127.9, 127.5, 126.8, 124.7, 115.0, 74.4, 58.4, 28.3, 23.6; MS (EI) m/z 226.1 (M^+ (55.7), 196.1 (38.0), 181.1 (100.0), 165.1 (74.7), 141.1 (39.2), 115.1 (41.0)); Exact mass calc'd. for $\text{C}_{16}\text{H}_{18}\text{O}$: 226.1376, found 226.1394.

(Cyclohex-1-en-1-yl)-buta-1,3-dien-2-methanol (42a): Compound **41** (400 mg, 2.94 mmol) in cyclohexane (5 mL) and vinyl magnesium chloride (5.53 mL, 1.7 M, 8.64 mmol) were reacted together and the reaction quenched with saturated ammonium chloride (20 mL). Chromatography (petroleum ether/ethyl acetate, 6:1) afforded **42a** as a yellow oil (433 mg, 90%). IR (neat); 3338 (s), 2929 (s), 2858 (m), 1436 (m) cm^{-1} ; ^1H NMR (CDCl_3 , 500 MHz): δ 6.79 (ddd, $J = 17.9$ Hz, 11.3 Hz, 0.8 Hz, 1H), 5.98 (s, 1H), 5.65 (sextet, $J = 2.0$ Hz, 1H), 5.29 (d, $J = 17.9$ Hz, 1H), 5.07 (dt, $J = 11.2$ Hz, 1.4 Hz, 1H), 2.12 – 2.05 (m, 4H), 1.90 (br s, 1H), 1.64 – 1.54 (m, 4H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 134.7, 134.4, 133.2, 133.2, 129.7, 113.7, 64.8, 29.1, 25.7, 22.7, 22.0; MS (EI) m/z 164.1 (M^+ (17.0), 133.1 (32.9), 121.1 (25.2), 105.1 (28.1), 91.1 (100.0), 79.1 (37.8)); Exact mass calc'd. for $\text{C}_{11}\text{H}_{16}\text{O}$: 164.1189, found 164.1177.

6-Methoxymethyl-1,2,3,4-tetrahydronaphthalene (43): Compound **42b** (120 mg, 0.67 mmol) was dissolved in toluene (20 mL) and placed in a sealed tube. The solution was heated to 250°C for 16 hours. The toluene was removed under reduced pressure. Chromatography (petroleum ether) afforded **43** as a colourless oil (28 mg, 23%). IR (neat); 2936 (s), 2852 (m), 2247 (m), 1094 (m), cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz): δ 7.24 (s, 3H), 4.36 (s, 2H), 3.35 (s, 3H), 2.74 (m, 4H), 1.09 (s, 4H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 137.5, 137.0, 135.5, 129.5, 129.0, 125.4, 75.1, 58.3, 29.7, 29.5, 23.6; MS (EI) m/z 176.1 (M^+ , 85.2), 145.1 (87.8), 129.1 (100.0), 115.1 (59.9), 91.1 (75.5)); Exact mass calc'd. for $\text{C}_{12}\text{H}_{16}\text{O}$: 176.1184, found 176.1167.

4-(2-Methoxymethyl-buta-1,3-dienyl)-1,2-dihydro-naphthalene (39b) : Alcohol **39a** (160 mg, 0.79 mmol) was dissolved in THF (20 mL). Sodium hydride (60 mg, 1.58 mmol) and methyl iodide (0.24 mL, 3.95 mmol) were added and the solution allowed to stir at room temperature for 14 hours. Water (20 mL) was added and aqueous extraction with diethyl ether (3 x 20 mL) was performed. Purification by silica gel chromatography employing a 20:1 pet ether/ethyl acetate solvent system. This afforded **39b** as a colourless oil (155 mg, 87%). IR (CDCl_3); 3161 (m), 2936 (s), 2865 (s), 2814 (s), 1468 (m) cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz): δ 7.14 (s, 4H), 6.70 (ddd, $J = 17.9$ Hz, 11.4 Hz, 0.9 Hz, 1H), 6.41 (s, 1H), 5.98 (td, $J = 5.7$ Hz,

1.5 Hz, 1H), 5.39 (d, $J = 17.8$ Hz, 1H), 5.10 (dt, $J = 11.2$ Hz, 1.5 Hz, 1H), 2.79 (t, $J = 6.7$ Hz, 2H), 2.42 – 2.35 (m, 2H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 136.4, 135.1, 133.9, 133.4, 130.3, 130.2, 127.9, 127.5, 126.8, 124.7, 115.0, 74.4, 58.4, 28.3, 23.6; MS (EI) m/z 226.1 (M^+ (55.7), 196.1 (38.0), 181.1 (100.0), 165.1 (74.7), 141.1 (39.2), 115.1 (41.0); Exact mass calc'd. For $\text{C}_{16}\text{H}_{18}\text{O}$: 226.1376. Observed: 226.1394.

9,10-Dihydrophenanthrene-3-carbaldehyde (44): 2,3-Dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) (26.0 mg, 0.114 mmol, 1.2 equiv) was added to a solution of **39b** (20.3 mg, 0.095 mmol) in toluene (0.637 mL) and the mixture refluxed for 5 hrs. The mixture was diluted with water and the aqueous phase was extracted with diethyl ether (3 x 10 mL). The combined organic layers were washed with water (1 x 10 mL), brine (1 x 10 mL) and dried over MgSO_4 . Chromatography (petroleum ether/ Et_2O , 9:1) afforded **44** (9.57 mg, 48%) as a colourless liquid. IR (neat) 3058, 2930, 2835, 2728, 1698, 1607, 1565, 1491, 1430, 1296, 1175, 1094, 1029, 826, 770 cm^{-1} ; ^1H NMR (CDCl_3 , 500 MHz): δ 10.03 (s, 1H), 8.24 (d, $J = 1.5$ Hz, 1H), 7.83 (d, $J = 7.7$ Hz, 1H), 7.72 (dd, $J = 7.7, 1.6$, 1H), 7.38 (d, $J = 7.7$, 1H) 7.35-7.32 (m, 1H) 7.29-7.25 (m, 2H) 2.96-2.87 (m, 4H). ^{13}C NMR (CDCl_3 , 126 MHz): δ 192.1, 144.6, 137.2, 135.7, 135.5, 133.3, 128.9, 128.9, 128.3, 128.2, 127.3, 124.8, 124.0, 29.5, 28.4. Exact mass calc'd. for $\text{C}_{16}\text{H}_{16}\text{O}$: 208.0888, found 208.0888.