Assembly of 3-Acyl Oxindoles Via CuI/L-Proline Catalyzed Intramolecular Arylation of β -Keto Amides

Biao Lu¹, Dawei Ma^{2*}

¹*Department of the Chemistry, Fudan University, Shanghai, 20043, China*

²*State key Laboratory of Bioorganic and Natural Products Chemistry, Shanghai Institute Of Organic Chemistry, Chinese Academy of Science, 354 Fenglin Lu, Shanghai 20032, China*

Supporting Information

Table of Contents

Experimental-----	S2-S8
Copies of ¹ H NMR and ¹³ C NMR spectrum of 6a-6q -----	S9-S42

Experimental Section

General procedure for acetoacetylation of aniline with 2,2,6-trimethyl-4-H-1,3-dioxin-4-one. A solution of N-alkylated aniline (5 mmol) and 2,2,6-trimethyl-4-H-1,3-dioxin-4-one (5 mmol) in 2 mL of xylene was placed in a 50 mL Erlenmeyer flask which was immersed in an oil bath that has been preheated to approximate 150 °C. The reaction mixture was vigorously stirred. The evolution of acetone became apparent immediately. After the reaction was completed detected by TLC, the cooled mixture was purified by column chromatography to give acetoacetylated product.

N-Benzyl-N-(2-iodophenyl)-3-oxobutanamide: 87% yield, ^1H NMR (300 MHz, CDCl_3) δ 14.2 (s, 0.2H), 7.95 (d, $J = 7.2$ Hz, 1H), 7.26 (m, 5H), 7.20 (m, 1H), 7.04 (m, 1H), 6.73 (d, $J = 7.2$ Hz, 1H), 5.69 (d, $J = 14.0$ Hz, 1H), 4.38 (s, 0.2H), 3.96 (d, $J = 14.0$ Hz, 1H), 3.28 (d, $J = 15.6$ Hz, 0.8H), 3.10 (d, $J = 15.6$ Hz, 0.8H), 2.15 (s, 2.3H), 1.62 (s, 0.7H); ESI-MS m/z 394 (M-H) $^+$, 416 (M-Na) $^+$.

N-Benzyl-N-(4-fluoro-2-iodophenyl)-3-oxobutanamide: 87% yield, ^1H NMR (300 MHz, CDCl_3) 14.2 (s, 0.2H), 7.66 (m, 1H), 7.24 (m, 5H), 6.91 (m, 1H), 6.68 (m, 1H), 5.69 (m, 1H), 4.36 (s, 0.2H), 3.91 (m, 1H), 3.30 (d, $J = 15.9$ Hz, 0.8H), 3.09 (d, $J = 15.9$ Hz, 0.8H), 2.16 (s, 2.3H), 1.60 (s, 0.7H); ESI-MS m/z 412 (M-H) $^+$, 434 (M-Na) $^+$.

***N*-Benzyl-*N*-(2-iodo-4-methylphenyl)-3-oxobutanamide:** 80% yield, ^1H NMR (300 MHz, CDCl_3) δ 14.2 (s, 0.2H), 7.76 (s, 1H), 7.20 (m, 5H), 6.97 (m, 1H), 6.59 (d, $J = 8.1$ Hz, 1H), 5.65 (m, 1H), 4.40 (s, 0.2H), 3.94 (m, 1H), 3.26 (d, $J = 15.9$ Hz, 0.8H), 3.10 (d, $J = 15.9$ Hz, 0.8H), 2.30 (s, 3H), 2.16 (s, 2.3H), 1.80 (s, 0.7H); ESI-MS m/z 408 (M-H) $^+$, 430 (M-Na) $^+$.

***N*-Benzyl-*N*-(2-iodo-5-methoxyphenyl)-3-oxobutanamide:** 95% yield, ^1H NMR (300 MHz, CDCl_3) δ 14.2 (s, 0.2H), 7.76 (m, 1H), 7.26 (m, 5H), 6.65 (m, 1H), 6.23 (s, 1H), 5.71 (m, 1H), 4.42 (s, 0.2H), 3.92 (m, 1H), 3.51 (s, 3H), 3.31 (d, $J = 15.9$ Hz, 0.8H), 3.12 (d, $J = 15.9$ Hz, 0.8H), 2.17 (s, 3H), 1.82 (s, 0.7H); ESI-MS m/z 424 (M-H) $^+$, 446 (M-Na) $^+$.

General procedure for the benzoacetylation. To a solution of Meldrum's acid (288 mg, 2 mmol) and DMAP (977 mg, 4 mmol) in CH_2Cl_2 (3 mL) was slowly added benzoyl chloride (0.28 mL, 2.4 mmol) in 1 mL of CH_2Cl_2 at 0 $^\circ\text{C}$ under nitrogen atmosphere. The reaction mixture was stirred at 0 $^\circ\text{C}$ for another hour and warmed up to room temperature and kept stirring for 1 h, then the reaction mixture was diluted by adding 20 mL of CH_2Cl_2 . The organic phase was washed with 10% HCl, water and brine, respectively. After the solution was dried and concentrated, the residue was dissolved in 3 mL of dioxane. To this resultant solution was added *N*-benzyl-2-iodoaniline (2 mmol) before it was refluxed for 1 h. The cooled solution

was concentrated, and then purified by chromatography to give product (50-81% yields) as colorless oil.

***N*-Benzyl-*N*-(2-iodophenyl)-3-oxo-3-phenylpropanamide:** ^1H NMR (300 MHz, CDCl_3) δ 14.6 (s, 0.3H), 7.90 (m, 1H), 7.75 (d, $J = 6.8$ Hz, 1H), 7.45 (d, $J = 6.8$ Hz, 1H), 7.26 (m, 8H), 7.05 (m, 1H), 7.00 (m, 1H), 6.74 (m, 1H), 5.67 (m, 1H), 4.98 (s, 0.3H), 4.00 (m, 1H), 3.88 (m, 0.8H), 3.56 (m, 0.8H); ESI-MS m/z 456 (M-H) $^+$, 478 (M-Na) $^+$.

***N*-Benzyl-3-(furan-2-yl)-*N*-(2-iodophenyl)-3-oxopropanamide:** ^1H NMR (300 MHz, CDCl_3) δ 14.3 (s, 0.2H), 7.86 (m, 1H), 7.44 (s, 1H), 7.20 (m, 6H), 7.08 (m, 1H), 6.94 (m, 1H), 6.74 (m, 1H), 6.42 (m, 1H), 5.65 (m, 1H), 4.99 (s, 0.2H), 3.90 (m, 1H), 3.68 (m, 0.8H), 3.41 (m, 0.8H); ESI-MS m/z 446 (M-H) $^+$, 468 (M-Na) $^+$.

***N*-Benzyl-*N*-(4-fluoro-2-iodophenyl)-3-oxo-3-(thiophen-2-yl)propanamide:** ^1H NMR (300 MHz, CDCl_3) δ 7.62 (m, 2H), 7.26 (m, 6H), 7.08 (s, 1H), 6.79 (m, 2H), 5.72 (m, 1H), 3.93 (m, 1H), 3.87 (m, 1H), 3.53 (m, 1H); ESI-MS m/z 480 (M-H) $^+$, 502 (M-Na) $^+$.

3-(Benzo[b]thiophen-2-yl)-N-benzyl-N-(2-iodophenyl)-3-oxopropanamide:

^1H NMR (300 MHz, CDCl_3) δ 14.6 (s, 0.2H), 7.89-7.85 (m, 3H), 7.55 (m, 2H), 7.34 (m, 5H), 7.06 (m, 1H), 6.92 (m, 1H), 6.76 (m, 1H), 5.65 (m, 1H), 4.99 (s, 0.2H), 3.96 (m, 1H), 3.89 (m, 0.8H), 3.58 (m, 0.8H); ESI-MS m/z 512 (M-H^+), 534 (M-Na^+).

N-Benzyl-3-(4-chlorophenyl)-N-(2-iodo-4-methylphenyl)-3-oxopropan-

amide: ^1H NMR (300 MHz, CDCl_3) δ 14.7 (s, 0.4H), 7.70 (m, 2H), 7.39 (m, 1H), 7.32 (m, 1H), 7.20 (m, 6H), 6.61 (m, 1H), 6.58 (m, 1H), 5.62 (m, 1H), 4.97 (s, 0.4H), 3.92 (m, 1H), 3.84 (m, 0.6H), 3.52 (m, 0.6H); ESI-MS m/z 504 (M-H^+), 526 (M-Na^+).

General procedure for CuI/L-proline catalyzed intramolecular arylation.

An oven-dried Schlenk tube was charged with substrate (0.5 mmol), CuI (10 mg, 0.05 mmol), L-proline (12 mg, 0.1 mmol), and cesium carbonate (645 mg, 2 mmol), evacuated and backfilled with argon. DMSO (5 mL) was added into the tube, and then the reaction mixture was stirred at room temperature. After the reaction was completed detected by TLC, the mixture was partitioned between ethyl acetate (40 mL) and saturated NH_4Cl (12 mL). The organic layer was washed with brine, dried over Na_2SO_4 , and concentrated in vacuo. The residue thus obtained was purified by silica gel chromatography to give the corresponding 3-acyl oxindole.

6a: ^1H NMR (400 MHz, CDCl_3) δ 13.68 (br s, 1H), 7.38 (d, $J = 7.5$ Hz, 1H), 7.29 (m, 5H), 7.26 (m, 1H), 7.09 (m, 1H), 6.85 (d, $J = 7.5$ Hz, 1H), 5.05 (s, 2H), 2.49 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 173.3, 171.1, 138.1, 136.1, 128.8, 127.6, 127.2, 125.2, 122.4, 122.2, 119.8, 109.4, 101.7, 43.3, 20.4; ESI-MS m/z 266 ($\text{M}-\text{H}$) $^+$, 288 ($\text{M}-\text{Na}$) $^+$; HRMS calcd for $\text{C}_{17}\text{H}_{15}\text{NO}_2$ (M) $^+$ 265.1103, found 265.1111.

6b: ^1H NMR (400 MHz, CDCl_3): δ 13.64 (br s, 1H), 7.37 (d, $J = 7.5$ Hz, 1H), 7.25 (m, 1H), 7.11 (m, 1H), 6.95 (d, $J = 7.6$ Hz, 1H), 3.36 (s, 3H), 2.46 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 172.9, 171.1, 138.9, 125.2, 122.3, 122.1, 119.7, 108.4, 101.8, 25.7, 20.3; ESI-MS m/z 190 ($\text{M}-\text{H}$) $^+$, 212 ($\text{M}-\text{Na}$) $^+$; HRMS calcd for $\text{C}_{11}\text{H}_{11}\text{NO}_2$ (M) $^+$ 189.0790, found 189.0788.

6c: ^1H NMR (400 MHz, CDCl_3) δ 13.60 (br s, 1H), 7.36 (d, $J = 7.5$ Hz, 1H), 7.26 (s, 1H), 7.11 (m, 1H), 6.97 (m, 2H), 6.46 (s, 1H), 6.36 (d, $J = 7.5$ Hz, 1H), 5.00 (s, 2H), 3.87 (s, 3H), 3.25 (s, 3H), 2.47 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 172.9, 171.3, 160.4, 158.0, 138.4, 129.1, 125.1, 122.3, 121.9, 119.6, 116.6, 109.6, 104.4, 101.8, 98.5, 55.4, 55.3, 37.6, 20.4; ESI-MS m/z 326 ($\text{M}-\text{H}$) $^+$, 348 ($\text{M}-\text{Na}$) $^+$; HRMS calcd for $\text{C}_{19}\text{H}_{19}\text{NO}_4$ (M) $^+$ 325.1314, found 325.1315.

6d: ^1H NMR (400 MHz, CDCl_3) δ 13.70 (br s, 1H), 7.30 (m, 5H), 7.09 (m, 1H), 6.79 (m, 1H), 6.75 (m, 1H), 5.03 (s, 2H), 2.46 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 174.5, 171.1, 159.1 (d, $J = 236.6$ Hz), 135.8, 133.9, 128.8, 127.7, 123.5 (d, $J = 9.4$ Hz), 111.4 (d, $J = 24.0$ Hz), 109.7 (d, $J = 8.9$ Hz), 109.6, 107.2 (d, $J = 26.0$ Hz), 101.6, 43.4, 20.3; ESI-MS m/z 284 (M-H) $^+$, 306 (M-Na) $^+$; HRMS calcd for $\text{C}_{17}\text{H}_{14}\text{NO}_2\text{F}$ (M) $^+$ 283.1009, found 283.1013.

6e: ^1H NMR (400 MHz, CDCl_3) δ 13.64 (br s, 1H), 7.30 (m, 6H), 7.05 (m, 1H), 6.84 (s, 1H), 5.01 (s, 2H), 2.46 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 173.7, 171.2, 138.9, 135.6, 130.8, 128.69, 127.8, 127.2, 122.1, 120.9, 120.4, 109.8, 101.1, 43.4, 20.5; ESI-MS m/z 300 (M-H) $^+$, 322 (M-Na) $^+$; HRMS calcd for $\text{C}_{17}\text{H}_{14}\text{NO}_2\text{Cl}$ (M) $^+$ 299.0713, found 299.0716.

6f: ^1H NMR (500 MHz, CDCl_3) δ 13.71 (br s, 1H), 7.32 (m, 5H), 7.14 (m, 1H), 6.81 (m, 1H), 5.00 (s, 2H), 2.46 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 174.9, 171.0, 154.4 (d, $J = 239.4$ Hz), 135.3, 134.3, 129.0, 127.9, 127.2, 122.0 (d, $J = 8.5$ Hz), 117.0 (d, $J = 19.8$ Hz), 110.6, 108.1 (d, $J = 25.8$ Hz), 101.1, 43.5, 20.4; ESI-MS m/z 318 (M-H) $^+$, 342 (M-Na) $^+$; HRMS calcd for $\text{C}_{17}\text{H}_{13}\text{NO}_2\text{ClF}$ (M) $^+$ 317.0619, found 317.0620.

6g: ^1H NMR (400 MHz, CDCl_3) δ 13.67 (br s, 1H), 7.29 (m, 5H), 7.19 (s, 1H), 6.91 (d, $J = 7.9$ Hz, 1H), 6.73 (d, $J = 7.5$ Hz, 1H), 5.02 (s, 2H), 2.48 (s, 3H), 2.36 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ 173.0, 171.1, 136.2, 136.0, 131.6, 128.7, 127.5,

127.2, 125.7, 122.5, 120.6, 109.1, 101.7, 43.3, 21.4, 20.4; ESI-MS m/z 280 (M-H)⁺, 302 (M-Na)⁺; HRMS calcd for C₁₈H₁₇NO₂ (M)⁺ 279.1259, found 279.1272.

6h: ¹H NMR (500 MHz, CDCl₃) δ 13.26 (br s, 1H), 7.28 (m, 5H), 7.26 (m, 1H), 6.61 (m, 1H), 6.44 (s, 1H), 5.01 (s, 2H), 3.75 (s, 3H), 2.43 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 171.7, 170.7, 158.3, 139.5, 136.0, 128.8, 127.6, 127.3, 120.4, 115.4, 106.8, 101.5, 97.1, 55.6, 43.4, 20.1; ESI-MS m/z 296 (M-H)⁺, 318 (M-Na)⁺; HRMS calcd for C₁₈H₁₇NO₃ (M)⁺ 295.1208, found 295.1219.

6i: ¹H NMR (300 MHz, CDCl₃) δ 7.31 (m, 5H), 7.14 (m, 1H), 6.88 (d, J = 7.5 Hz, 1H), 6.62 (d, J = 7.5 Hz, 1H), 4.97 (d, J = 15.2 Hz, 1H), 4.85 (d, J = 15.2 Hz, 1H), 4.57 (s, 1H), 2.29 (s, 3H), 2.18 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 199.4, 170.8, 143.8, 135.5, 135.4, 129.1, 128.9, 127.8, 127.4, 124.7, 122.4, 107.1, 60.7, 44.2, 28.3, 18.6; ESI-MS m/z 280 (M-H)⁺, 302 (M-Na)⁺; HRMS calcd for C₁₈H₁₇NO₂ (M)⁺ 279.1259, found 279.1246.

6j: ¹H NMR (400 MHz, CDCl₃): δ 13.81 (br s, 1H), 7.80 (m, 2H), 7.14 (m, 2H), 6.87 (m, 1H), 7.22 (m, 1H), 7.17 (m, 1H), 6.86 (m, 2H), 5.10 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 172.1, 171.3, 138.3, 136.0, 134.1, 131.4, 128.8, 128.7, 128.4, 127.7, 127.3, 125.9, 121.9, 121.7, 119.8, 109.3, 101.4, 43.5; ESI-MS m/z 328 (M-H)⁺, 350 (M-Na)⁺; HRMS calcd for C₂₂H₁₇NO₂ (M)⁺ 327.1259, found 327.1265.

6k: ^1H NMR (400 MHz, CDCl_3) δ 14.7 (br s, 1H), 8.29 (d, $J = 7.5$ Hz, 1H), 7.80 (s, 1H), 7.30 (m, 6H), 7.22 (m, 1H), 7.17 (m, 1H), 6.87 (d, $J = 7.7$ Hz, 1H), 6.67 (m, 1H), 5.09 (s, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 172.8, 158.6, 149.0, 145.7, 138.2, 135.9, 128.8, 127.6, 127.2, 125.9, 122.3, 122.2, 121.4, 117.0, 112.6, 109.1, 99.4, 43.5; ESI-MS m/z 318 (M-H) $^+$, 340 (M-Na) $^+$; HRMS calcd for $\text{C}_{20}\text{H}_{15}\text{NO}_3$ (M) $^+$ 317.1052, found 317.1047.

6l: ^1H NMR (400 MHz, CDCl_3) δ 7.90 (s, 1H), 7.70 (d, $J = 7.4$ Hz, 1H), 7.66 (m, 1H), 7.29 (m, 7H), 7.12 (m, 1H), 6.97 (m, 1H), 6.88 (d, $J = 7.4$ Hz, 1H), 5.10 (s, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 172.1, 164.4, 138.2, 136.4, 135.9, 130.9, 130.8, 128.8, 127.7, 127.5, 127.3, 126.0, 122.0, 121.6, 119.8, 109.4, 100.9, 43.5; ESI-MS m/z 334 (M-H) $^+$, 356 (M-Na) $^+$; HRMS calcd for $\text{C}_{20}\text{H}_{15}\text{NO}_2\text{S}$ (M) $^+$ 333.0824, found 333.0827.

6m: ^1H NMR (500 MHz, CDCl_3) δ 7.87 (d, $J = 3.6$ Hz, 1H), 7.69 (d, $J = 4.9$ Hz, 1H), 7.41 (dd, $J = 2.3$ Hz, 7.4 Hz, 1H), 7.33 (m, 7H), 6.79 (m, 2H), 5.08 (s, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 172.1, 165.7, 159.0 (d, $J = 237.5$ Hz), 136.0, 135.6, 134.1, 131.4, 131.0, 128.9, 127.8, 127.7, 122.7 (d, $J = 9.4$ Hz), 112.3 (d, $J = 24.1$ Hz), 109.8 (d, $J = 8.8$ Hz), 107.2 (d, $J = 26.8$ Hz), 100.7, 43.7; ESI-MS m/z 352 (M-H) $^+$, 374 (M-Na) $^+$; HRMS calcd for $\text{C}_{20}\text{H}_{14}\text{NO}_2\text{FS}$ (M) $^+$ 351.0729, found 351.0739.

6n: ^1H NMR (400 MHz, CDCl_3) δ 8.07 (s, 1H), 7.93 (m, 2H), 7.71 (d, $J = 7.6$ Hz, 1H), 7.47 (m, 1H), 7.29 (m, 7H), 7.13 (m, 1H), 6.96 (m, 1H), 6.89 (d, $J = 7.6$ Hz, 1H), 5.11 (s, 2H); ^{13}C NMR (100 MHz, CDCl_3): δ 172.0, 164.4, 141.4, 138.8, 138.5, 136.1, 135.8, 128.9, 127.7, 127.3, 126.5, 126.4, 125.0, 124.9, 122.6, 122.2, 121.4, 120.1, 109.6, 102.1, 43.6; ESI-MS m/z 384 (M-H) $^+$, 406 (M-Na) $^+$; HRMS calcd for $\text{C}_{24}\text{H}_{17}\text{NO}_2\text{S}$ (M) $^+$ 383.0980, found 383.0976.

6o: ^1H NMR (500 MHz, CDCl_3) δ 7.76 (d, $J = 8.2$ Hz, 2H), 7.53 (d, $J = 8.2$ Hz, 2H), 7.47 (m, 1H), 7.28 (m, 5H), 6.99 (s, 1H), 6.89 (d, $J = 7.8$ Hz, 1H), 6.74 (d, $J = 7.9$ Hz, 1H), 5.07 (s, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 172.0, 169.5, 137.5, 136.3, 135.9, 132.7, 131.5, 129.9, 129.0, 128.8, 127.7, 127.3, 126.7, 121.4, 120.4, 109.2, 101.6, 43.5, 21.3; ESI-MS m/z 374 (M-H) $^+$, 396 (M-Na) $^+$; HRMS calcd for $\text{C}_{23}\text{H}_{18}\text{NO}_2\text{Cl}$ (M) $^+$ 375.1026, found 375.1030.

6p: ^1H NMR (500 MHz, CDCl_3) δ 7.73 (d, $J = 8.2$ Hz, 2H), 7.53 (d, $J = 8.2$ Hz, 2H), 7.31 (m, 5H), 7.07 (d, $J = 8.6$ Hz, 1H), 6.85 (m, 2H), 5.06 (s, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 172.1, 170.2, 139.3, 137.8, 135.3, 132.3, 131.7, 129.8, 129.2, 129.0, 127.9, 127.3, 122.0, 120.4, 120.0, 109.9, 101.0, 43.6; ESI-MS m/z 396 (M-H) $^+$, 418

(M-Na)⁺; HRMS calcd for C₂₂H₁₅NO₂Cl₂ (M)⁺ 395.0480, found 395.0478.

6q: ¹H NMR (500 MHz, CDCl₃) δ 8.86 (s, 1H), 8.07 (d, *J* = 6.7 Hz, 1H), 7.53 (d, *J* = 8.4 Hz, 1H), 7.29 (m, 5H), 7.14 (m, 2H), 6.91 (m, 2H), 5.09 (s, 2H); ¹³C NMR (125 MHz, CDCl₃) δ 171.7, 166.2, 154.0, 149.5, 138.6, 135.6, 129.2, 128.9, 127.8, 127.3, 126.8, 124.5, 122.4, 120.8, 119.5, 109.8, 102.8, 43.6; ESI-MS *m/z* 363 (M-H)⁺, 385 (M-Na)⁺; HRMS calcd for C₂₁H₁₅Cl₂N₂O₂Cl (M)⁺ 362.0822, found 362.0813.

28.69

L.B-42-11-2

Current Date Parameters	
NAME	M1
EXPHO	10
FROEND	1

F2 - Acquisition Parameters

Time	E.07
1.10	0.07
1.20	0.07
1.30	0.07
1.40	0.07
1.50	0.07
2.00	0.07
2.10	0.07
2.20	0.07
2.30	0.07
2.40	0.07
2.50	0.07
3.00	0.07
3.10	0.07
3.20	0.07
3.30	0.07
3.40	0.07
3.50	0.07
4.00	0.07
4.10	0.07
4.20	0.07
4.30	0.07
4.40	0.07
4.50	0.07
5.00	0.07
5.10	0.07
5.20	0.07
5.30	0.07
5.40	0.07
5.50	0.07
6.00	0.07
6.10	0.07
6.20	0.07
6.30	0.07
6.40	0.07
6.50	0.07
7.00	0.07
7.10	0.07
7.20	0.07
7.30	0.07
7.40	0.07
7.50	0.07
8.00	0.07
8.10	0.07
8.20	0.07
8.30	0.07
8.40	0.07
8.50	0.07
9.00	0.07
9.10	0.07
9.20	0.07
9.30	0.07
9.40	0.07
9.50	0.07
10.00	0.07
10.10	0.07
10.20	0.07
10.30	0.07
10.40	0.07
10.50	0.07
11.00	0.07
11.10	0.07
11.20	0.07
11.30	0.07
11.40	0.07
11.50	0.07
12.00	0.07
12.10	0.07
12.20	0.07
12.30	0.07
12.40	0.07
12.50	0.07
13.00	0.07
13.10	0.07
13.20	0.07
13.30	0.07
13.40	0.07
13.50	0.07
14.00	0.07
14.10	0.07
14.20	0.07
14.30	0.07
14.40	0.07
14.50	0.07
15.00	0.07
15.10	0.07
15.20	0.07
15.30	0.07
15.40	0.07
15.50	0.07
16.00	0.07
16.10	0.07
16.20	0.07
16.30	0.07
16.40	0.07
16.50	0.07
17.00	0.07
17.10	0.07
17.20	0.07
17.30	0.07
17.40	0.07
17.50	0.07
18.00	0.07
18.10	0.07
18.20	0.07
18.30	0.07
18.40	0.07
18.50	0.07
19.00	0.07
19.10	0.07
19.20	0.07
19.30	0.07
19.40	0.07
19.50	0.07
20.00	0.07
20.10	0.07
20.20	0.07
20.30	0.07
20.40	0.07
20.50	0.07
21.00	0.07
21.10	0.07
21.20	0.07
21.30	0.07
21.40	0.07
21.50	0.07
22.00	0.07
22.10	0.07
22.20	0.07
22.30	0.07
22.40	0.07
22.50	0.07
23.00	0.07
23.10	0.07
23.20	0.07
23.30	0.07
23.40	0.07
23.50	0.07
24.00	0.07
24.10	0.07
24.20	0.07
24.30	0.07
24.40	0.07
24.50	0.07
25.00	0.07
25.10	0.07
25.20	0.07
25.30	0.07

5 m 135

ID	60535
----	-------

३६

SAH	120:9.220
SHH	0.00760

2.7263477

26	41.600
27	39.000

300 0

***** CHENP *****

Multi	2H
12	5.00

P.L.	0.00
SECI	200 1356440

Progressing Patient

51 1033B
5 400 1300156

U	3
1/1	1
1/1	2

0.06
0.07

1.00

13 NEW BLUE PATENTERS
CO 250 000

11-	15,000
11	6001.33

0.000	23
0.00	24

01700M	0.72600
H273M	300.09753

Hz

12-93-2

S-34