Supplementary material

**Assessing Future Habitat Availability for Coastal Lowland Anurans in the
Brazilian Atlantic Rainforest**

**Igor Soares de Oliveira^{1, 2, *}, Dennis Rödder³, César Capinha⁴, Faraham
Ahmadzadeh³, Adriele Karlokoski Cunha de Oliveira⁵, and Luís Felipe Toledo²**

¹ Programa de Pós-Graduação em Ecologia, Instituto de Biologia, Universidade Estadual de Campinas, Avenida Bertrand Russel, s/n, Cidade Universitária Zeferino Vaz, Barão Geraldo, CEP 13083-865, Campinas, São Paulo, Brasil

² Laboratório de História Natural de Anfíbios Brasileiros, Departamento de Biologia Animal, Instituto de Biologia, Unicamp, Rua Monteiro Lobato, 255, CEP 13083-862, Campinas, São Paulo, Brasil

³ Zoologisches Forschungsmuseum Alexander Koenig, Herpetology Section, Adenauerallee 160, 53113 Bonn, Germany.

⁴ Centro de Biologia Ambiental, Faculdade de Ciências da Universidade de Lisboa, 1749-016 Lisboa, Portugal.

⁵ Programa de Pós-Graduação em Zoologia, Departamento de Zoologia, Universidade Federal do Paraná, Rua Cel. Francisco Heráclito dos Santos, 210, Jardim das Américas, CEP 81531-970, Curitiba, Paraná, Brasil.

*Corresponding email address: oliveira.snake@gmail.com

Received:

FIGURE S1: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Aparasphenodon arapapa*. Modeled distributions shown in black.

FIGURE S2: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *A. bokermanni*. Modeled distributions shown in black.

FIGURE S3: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Chiasmocleis alagoana*. Modeled distributions shown in black.

FIGURE S4: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *C. atlantica*. Modeled distributions shown in black.

FIGURE S5: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *C. capixaba*. Modeled distributions shown in black.

FIGURE S6: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *C. lacrimae*. Modeled distributions shown in black.

FIGURE S7: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *C. sapiranga*. Modeled distributions shown in black.

FIGURE S8: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Crossodactylus lutzorum*. Modeled distributions shown in black.

FIGURE S9: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Dasypops schirchi*. Modeled distributions shown in black.

FIGURE S10: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Dendrophryniscus skuki*. Modeled distributions shown in black.

FIGURE S11: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Gastrotheca megacephala*. Modeled distributions shown in black.

FIGURE S12: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Haddadus plicifer*. Modeled distributions shown in black.

FIGURE S13: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Leptodactylus hylodes*. Modeled distributions shown in black.

FIGURE S14: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Melanophryniscus setiba*. Modeled distributions shown in black.

FIGURE S15: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Physalaemus atlanticus*. Modeled distributions shown in black.

FIGURE S16: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Phyllodytes punctatus*. Modeled distributions shown in black.

FIGURE S17: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Scinax cretatus*. Modeled distributions shown in black.

FIGURE S18: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Sphaenorhynchus palustris*. Modeled distributions shown in black.

FIGURE S19: Estimated current (A) and future habitat suitability in the B2a (B) and A2a (C) climatic scenario for the year 2080 in the Brazilian Atlantic forest for *Xenohyla truncata*. Modeled distributions shown in black.