Supporting Information

Asymmetric Addition of Dimethylzinc to *N*-Tosylarylimines Catalyzed by a Rhodium-Diene Complex Toward the Synthesis of Chiral 1-Arylethylamines

Takahiro Nishimura, Yuichi Yasuhara, and Tamio Hayashi*

*Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo-ku, Kyoto
606-8502, Japan*

Supporting Data

General. All anaerobic and moisture-sensitive manipulations were carried out with standard Schlenk techniques under predried nitrogen or glovebox techniques under prepurified argon. NMR spectra were recorded on a JEOL JNM LA-500 spectrometer (500 MHz for ^1H , and 125 MHz for ^{13}C). Chemical shifts are reported in δ ppm referenced to an internal SiMe_4 standard for ^1H NMR and chloroform-*d* (δ 77.16) for ^{13}C NMR. Optical rotations were measured on a JASCO DIP-370 polarimeter.

Materials. 1,4-Dioxane was distilled over benzophenone-ketyl under nitrogen. Dimethylzinc (1M solution in hexane) was purchased from Tosoh-finechem. $[\text{RhCl}(\text{C}_2\text{H}_4)_2]_2$,¹ (1*R*,4*R*)-2,5-dibenzylbicyclo[2.2.2]octa-2,5-diene ((*R,R*)-Bn-bod*),² (1*R*,4*R*)-2,5-diphenylbicyclo[2.2.2]octa-2,5-diene ((*R,R*)-Ph-bod*)² were prepared according to the reported procedures.

Preparation of *N*-Tosylarylimines. *N*-Tosylarylimines (**1a–1j**) were prepared according to the reported procedures.³

***N*-Phenylmethyldene-4-methylbenzenesulfonamide (1a)** [CAS:13707-41-0]: 95% yield. ^1H NMR (CDCl_3) δ 2.44 (s, 3H), 7.35 (d, J = 7.9 Hz, 2H), 7.46–7.52 (m, 2H), 7.59–7.64 (m, 1H), 7.87–7.95 (m, 4H), 9.03 (s, 1H).

***N*-(4-Chlorophenyl)methyldene-4-methylbenzenesulfonamide (1b)** [CAS: 3175-65-1]: 99% yield. ^1H NMR (CDCl_3) δ 2.44 (s, 3H), 7.35 (d, J = 8.3 Hz, 2H), 7.45 (d, J = 8.6 Hz, 2H), 7.86 (d, J = 8.6 Hz, 2H), 7.88 (d, J = 8.3 Hz, 2H), 8.99 (s, 1H).

***N*-(3-Chlorophenyl)methyldene-4-methylbenzenesulfonamide (1c)** [CAS: 169829-11-2]: 79% yield. ^1H NMR (CDCl_3) δ 2.45 (s, 3H), 7.36 (d, J = 8.0 Hz, 2H), 7.43 (t, J = 7.9 Hz, 1H),

7.55–7.59 (m, 1H), 7.76–7.80 (m, 1H), 7.86–7.91 (m, 2H), 7.92–7.95 (m, 1H), 8.98 (s, 1H).

***N*-(4-Fluorophenyl)methylidene-4-methylbenzenesulfonamide (1d)** [CAS: 181041-86-1]: 72% yield. ¹H NMR (CDCl₃) δ 2.44 (s, 3H), 7.14–7.21 (m, 2H), 7.35 (d, *J* = 7.9 Hz, 2H), 7.86–7.91 (m, 2H), 7.92–7.98 (m, 2H), 9.00 (s, 1H).

***N*-(4-Trifluoromethylphenyl)methylidene-4-methylbenzenesulfonamide (1e)** [CAS: 198012-02-1]: 83% yield. ¹H NMR (CDCl₃) δ 2.45 (s, 3H), 7.37 (d, *J* = 7.9 Hz, 2H), 7.75 (d, *J* = 8.2 Hz, 2H), 7.90 (d, *J* = 8.3 Hz, 2H), 8.05 (d, *J* = 8.2 Hz, 2H), 9.08 (s, 1H).

***N*-(3-Methoxyphenyl)methylidene-4-methylbenzenesulfonamide (1f)** [CAS: 160955-96-4]: 64% yield. ¹H NMR (CDCl₃) δ 2.44 (s, 3H), 3.84 (s, 3H), 7.13–7.18 (m, 1H), 7.35 (d, *J* = 8.3 Hz, 2H), 7.37–7.42 (m, 1H), 7.43–7.49 (m, 2H), 7.87–7.91 (m, 2H), 8.99 (s, 1H).

***N*-(4-Methylphenyl)methylidene-4-methylbenzenesulfonamide (1g)** [CAS: 75159-10-3]: 83% yield. ¹H NMR (CDCl₃) δ 2.42 (s, 3H), 2.43 (s, 3H), 7.27–7.31 (m, 2H), 7.31–7.34 (m, 2H), 7.79–7.84 (m, 2H), 7.86–7.91 (m, 2H), 8.98 (s, 1H).

***N*-(3-Methylphenyl)methylidene-4-methylbenzenesulfonamide (1h)**. [CAS: 561319-11-7]: 23% yield. ¹H NMR (CDCl₃) δ 2.39 (s, 3H), 2.44 (s, 3H), 7.32–7.40 (m, 3H), 7.40–7.45 (m, 1H), 7.70 (d, *J* = 7.6 Hz, 1H), 7.76 (s, 1H), 7.86–7.92 (m, 2H), 9.00 (s, 1H).

***N*-(2-Methylphenyl)methylidene-4-methylbenzenesulfonamide (1i)**. [CAS: 342598-64-1]: 63% yield. ¹H NMR (CDCl₃) δ 2.43 (s, 3H), 2.60 (s, 3H), 7.24–7.30 (m, 2H), 7.32–7.36 (m, 2H), 7.44–7.49 (m, 1H), 7.87–7.91 (m, 2H), 8.00 (d, *J* = 7.8 Hz, 1H), 9.34 (s, 1H).

***N*-(2-Naphthyl)methylidene-4-methylbenzenesulfonamide (1j)**. [CAS: 17692-84-1]: 20% yield. ¹H NMR (CDCl₃) δ 2.44 (s, 3H), 7.33–7.39 (m, 2H), 7.55–7.60 (m, 1H), 7.60–7.66 (m, 1H), 7.85–7.90 (m, 2H), 7.90–7.98 (m, 3H), 8.00–8.06 (m, 1H), 8.33 (s, 1H), 9.17 (s, 1H).

General Procedure for the Asymmetric Addition of Dimethylzinc to *N*-Tosylarylimines Catalyzed by a Rhodium-Diene Complex. A solution of [RhCl(C₂H₄)₂]₂ (1.2 mg, 0.0030 mmol, 3 mol % Rh) and (*R,R*)-Ph-bod* (3.1 mg, 0.012 mmol, 2 equiv to Rh) in 1,4-dioxane (0.3 mL) was placed in 20 mL Schlenk tube and stirred for 5 min at room temperature. To this solution were added successively imine (0.20 mmol), additional 1,4-dioxane (0.4 mL), and Me₂Zn (0.30 mL, 0.30 mmol; 1.0 M solution in hexane). After 3 h stirring at 50 °C, the resulting mixture was quenched with 7% aqueous NH₃ and extracted

with Et₂O. The combined organic layer was washed with brine, dried over MgSO₄, filtered, and concentrated on a rotary evaporator. The residue was purified by silica gel column chromatography (hexane/ethyl acetate = 4/1) to afford the desired product. The reaction conditions and results are summarized in Table 2.

2a *N*-[(*1S*)-1-Phenylethyl]-4-methylbenzenesulfonamide (**2a**) [CAS: 66558-04-1]: 82% yield, 97% ee. The ee was determined on Chiralcel OD-H column with hexane/2-propanol = 4/1, flow = 0.3 mL/min, wavelength = 254 nm. Retention times: 22.6 min [(*R*)-enantiomer], 25.9 min [(*S*)-enantiomer]. $[\alpha]_D^{20}$ -64.9 (*c* 1.02, EtOH), $[\alpha]_D^{20}$ -62.9 (*c* 1.02, CHCl₃) for 97% ee. ¹H NMR (CDCl₃) δ 1.40 (d, *J* = 6.8 Hz, 3H), 2.37 (s, 3H), 4.45 (quintet, *J* = 6.8 Hz, 1H), 5.27 (broad s, 1H), 7.08–7.19 (m, 7H), 7.58–7.64 (m, 2H). ¹³C NMR (CDCl₃) δ 21.6, 23.7, 53.7, 126.2, 127.2, 127.5, 128.6, 129.5, 137.8, 142.2, 143.2.

2b *N*-[(*1S*)-1-(4-Chlorophenyl)ethyl]-4-methylbenzenesulfonamide (**2b**): 84% yield (white solid), 96% ee. The ee was determined on Chiralcel OD-H column with hexane/2-propanol = 4/1, flow = 0.3 mL/min, wavelength = 254 nm. Retention times: 23.2 min [(*R*)-enantiomer], 25.4 min [(*S*)-enantiomer]. $[\alpha]_D^{20}$ -70.7 (*c* 1.02, CHCl₃) for 96% ee. ¹H NMR (CDCl₃) δ 1.37 (d, *J* = 6.9 Hz, 3H), 2.39 (s, 3H), 4.43 (dq, *J* = 7.1, 6.9 Hz, 1H), 5.36 (d, *J* = 7.1 Hz, 1H), 7.00–7.05 (m, 2H), 7.08–7.14 (m, 2H), 7.17 (d, *J* = 8.0 Hz, 2H), 7.58 (d, *J* = 8.2 Hz, 2H). ¹³C NMR (CDCl₃) δ 21.6, 23.6, 53.2, 127.2, 127.7, 128.7, 129.6, 133.3, 137.6, 140.7, 143.5. Anal. Calcd for C₁₅H₁₆ClNO₂S: C, 58.15; H, 5.21. Found: C, 58.15; H, 5.17.

2c *N*-[(*1S*)-1-(3-Chlorophenyl)ethyl]-4-methylbenzenesulfonamide (**2c**): 83% yield (white solid), 96% ee. The ee was determined on Chiralcel OD-H column with hexane/2-propanol = 4/1, flow = 0.3 mL/min, wavelength = 254 nm. Retention times: 23.8 min [(*R*)-enantiomer], 27.5 min [(*S*)-enantiomer]. $[\alpha]_D^{20}$ -69.7 (*c* 1.02, CHCl₃) for 96% ee. ¹H NMR (CDCl₃) δ 1.37 (d, *J* = 7.0 Hz, 3H), 2.37 (s, 3H), 4.43 (dq, *J* = 7.3, 7.0 Hz, 1H), 5.51 (d, *J* = 7.3 Hz, 1H), 6.98 (s, 1H), 7.01–7.05 (m, 1H), 7.08–7.12 (m, 2H), 7.16 (d, *J* = 8.0 Hz, 2H), 7.59 (d, *J* = 8.3 Hz, 2H). ¹³C NMR (CDCl₃) δ 21.6, 23.6, 53.3, 124.5, 126.6, 127.1, 127.5, 129.5, 129.8, 134.3, 137.5, 143.4, 144.2. Anal. Calcd for C₁₅H₁₆ClNO₂S: C, 58.15; H, 5.21. Found: C, 58.30; H, 5.21.

2d *N*-[(*1S*)-1-(4-Fluorophenyl)ethyl]-4-methylbenzenesulfonamide (**2d**): 84% yield (white solid), 98% ee. The ee was determined on Chiralcel OD-H column with hexane/2-propanol = 4/1, flow = 0.3 mL/min, wavelength =

254 nm. Retention times: 23.9 min [(*R*)-enantiomer], 27.1 min [(*S*)-enantiomer]. $[\alpha]_D^{20}$ -53.1 (*c* 1.02, CHCl₃) for 98% ee. ¹H NMR (CDCl₃) δ 1.37 (d, *J* = 6.8 Hz, 3H), 2.38 (s, 3H), 4.44 (quintet, *J* = 7.0 Hz, 1H), 5.46 (broad s, 1H), 6.81–6.86 (m, 2H), 7.05–7.10 (m, 2H), 7.17 (d, *J* = 8.1 Hz, 2H), 7.60 (d, *J* = 8.2 Hz, 2H). ¹³C NMR (CDCl₃) δ 21.5, 23.7, 53.1, 115.3 (d, *J*_{F-C} = 21.2 Hz), 127.2, 128.0 (d, *J*_{F-C} = 7.8 Hz), 129.5, 137.7, 138.1 (d, *J*_{F-C} = 3.1 Hz), 143.3, 162.1 (d, *J*_{F-C} = 245 Hz). Anal. Calcd for C₁₅H₁₆FNO₂S: C, 61.41; H, 5.50. Found: C, 61.45; H, 5.50.

2e

***N*-[(*1S*)-1-(4-Trifluoromethylphenyl)ethyl]-4-methylbenzenesulfon-**

amide (2e): 83% yield (white solid), 94% ee. The ee was determined on Chiralcel OD-H column with hexane/2-propanol = 4/1, flow = 0.3 mL/min, wavelength = 254 nm. Retention times: 23.1 min [(*R*)-

enantiomer], 25.6 min [(*S*)-enantiomer]. $[\alpha]_D^{20}$ -45.3 (*c* 1.10, CHCl₃) for 94% ee. ¹H NMR (CDCl₃) δ 1.40 (d, *J* = 7.0 Hz, 3H), 2.35 (s, 3H), 4.53 (dq, *J* = 7.3, 7.0 Hz, 1H), 5.69 (d, *J* = 7.3 Hz, 1H), 7.10 (d, *J* = 8.0 Hz, 2H), 7.20 (d, *J* = 8.1 Hz, 2H), 7.37 (d, *J* = 8.2 Hz, 2H), 7.55 (d, *J* = 8.3 Hz, 2H). ¹³C NMR (CDCl₃) δ 21.4, 23.6, 53.5, 124.1 (q, *J*_{F-C} = 271.7 Hz), 125.5 (q, *J*_{F-C} = 3.7 Hz), 126.8, 127.2, 129.5, 129.7 (q, *J*_{F-C} = 32.6 Hz), 137.5, 143.5, 146.1 (q, *J*_{F-C} = 1.4 Hz). Anal. Calcd for C₁₆H₁₆F₃NO₂S: C, 55.97; H, 4.70. Found: C, 55.97; H, 4.71.

2f

***N*-[(*1S*)-1-(3-Methoxyphenyl)ethyl]-4-methylbenzenesulfonamide**

(2f): 91% yield (colorless liquid), 97% ee. The ee was determined on Chiralcel OD-H column with hexane/2-propanol = 4/1, flow = 0.3 mL/min, wavelength = 254 nm. Retention times: 27.3 min [(*R*)-

enantiomer], 31.1 min [(*S*)-enantiomer]. $[\alpha]_D^{20}$ -64.1 (*c* 1.02, CHCl₃) for 97% ee. ¹H NMR (CDCl₃) δ 1.41 (d, *J* = 6.8 Hz, 3H), 2.37 (s, 3H), 3.68 (s, 3H), 4.42 (quintet, *J* = 6.9 Hz, 1H), 5.14 (broad s, 1H), 6.59 (dd, *J* = 2.0, 2.0 Hz, 1H), 6.66–6.72 (m, 2H), 7.05–7.13 (m, 1H), 7.17 (d, *J* = 7.8 Hz, 2H), 7.58–7.64 (m, 2H). ¹³C NMR (CDCl₃) δ 21.6, 23.7, 53.8, 55.2, 111.8, 113.1, 118.5, 127.2, 129.5, 129.7, 137.8, 143.2, 143.8, 159.8. Anal. Calcd for C₁₆H₁₉NO₃S: C, 62.93; H, 6.27. Found: C, 62.96; H, 6.30.

2g

***N*-[(*1S*)-1-(4-Methylphenyl)ethyl]-4-methylbenzenesulfonamide (2g)**

[CAS: 313949-04-1] for the racemic compound **2g**: 81% yield (white solid), 98% ee. The ee was determined on Chiralcel OD-H column with hexane/2-propanol = 4/1, flow = 0.3 mL/min, wavelength = 254 nm.

Retention times: 21.2 min [(*R*)-enantiomer], 23.5 min [(*S*)-enantiomer]. $[\alpha]_D^{20}$ -72.2 (*c* 1.08, CHCl₃) for 98% ee. ¹H NMR (CDCl₃) δ 1.39 (d, *J* = 6.8 Hz, 3H), 2.27 (s, 3H), 2.38 (s, 3H), 4.40 (dq, *J* = 7.0, 6.8 Hz, 1H), 5.04 (d, *J* = 7.0 Hz, 1H), 6.95–7.03 (m, 4H), 7.17 (d, *J* = 8.3 Hz, 2H), 7.62 (d, *J* = 8.3 Hz, 2H). ¹³C NMR (CDCl₃) δ 21.1, 21.6, 23.6, 53.5, 126.2, 127.3,

129.3, 129.5, 137.2, 137.9, 139.3, 143.1.

***N*-[(1*S*)-1-(3-Methylphenyl)ethyl]-4-methylbenzenesulfonamide (2h)**

[CAS: 355151-82-5]: 82% yield (white solid), 98% ee. The ee was determined on Chiralcel OD-H column with hexane/2-propanol = 4/1, flow = 0.3 mL/min, wavelength = 254 nm. Retention times: 21.0 min [(*R*)-enantiomer], 23.7 min [(*S*)-enantiomer]. $[\alpha]_D^{20}$ -76.6 (*c* 0.78, CHCl₃) for 98% ee. ¹H NMR (CDCl₃) δ 1.40 (d, *J* = 7.0 Hz, 3H), 2.20 (s, 3H), 2.38 (s, 3H), 4.42 (quintet, *J* = 6.9 Hz, 1H), 5.09 (broad s, 1H), 6.82 (s, 1H), 6.90 (d, *J* = 7.7 Hz, 1H), 6.96 (d, *J* = 7.4 Hz, 1H), 7.08 (t, *J* = 7.6 Hz, 1H), 7.16 (d, *J* = 8.3 Hz, 2H), 7.61 (d, *J* = 8.3 Hz, 2H). ¹³C NMR (CDCl₃) δ 21.4, 21.6, 23.7, 53.8, 123.3, 127.0, 127.2, 128.2, 128.5, 129.5, 137.9, 138.2, 142.0, 143.1.

***N*-[(1*S*)-1-(2-Methylphenyl)ethyl]-4-methylbenzenesulfonamide (2i)** [CAS:

101287-22-3] for the racemic compound **2i**: 61% yield (white solid), 96% ee. The ee was determined on Chiralcel OD-H column with hexane/2-propanol = 4/1, flow = 0.3 mL/min, wavelength = 254 nm. Retention times: 20.6 min [(*R*)-enantiomer], 25.4 min [(*S*)-enantiomer]. $[\alpha]_D^{20}$ -30.4 (*c* 1.07, CHCl₃) for 96% ee. ¹H NMR (CDCl₃) δ 1.36 (d, *J* = 6.8 Hz, 3H), 2.18 (s, 3H), 2.34 (s, 3H), 4.71 (quintet, *J* = 6.8 Hz, 1H), 5.27 (broad s, 1H), 6.95–7.06 (m, 3H), 7.08–7.14 (m, 3H), 7.55–7.60 (m, 2H). ¹³C NMR (CDCl₃) δ 19.1, 21.5, 23.2, 49.8, 125.5, 126.5, 127.1, 127.2, 129.4, 130.5, 134.5, 137.8, 140.4, 143.1.

***N*-[(1*S*)-1-(2-Naphthalenyl)ethyl]-4-methylbenzenesulfonamide (2j)**

[CAS: 257940-78-6]: 75% yield (white solid), 98% ee. The ee was determined on Chiralcel OD-H column with hexane/2-propanol = 4/1, flow = 0.3 mL/min, wavelength = 254 nm. Retention times: 30.0 min [(*R*)-enantiomer], 36.2 min [(*S*)-enantiomer]. $[\alpha]_D^{20}$ -67.1 (*c* 1.05, CHCl₃) for 98% ee. ¹H NMR (CDCl₃) δ 1.49 (d, *J* = 6.9 Hz, 3H), 2.21 (s, 3H), 4.62 (dq, *J* = 7.3, 6.9 Hz, 1H), 5.32 (d, *J* = 7.3 Hz, 1H), 6.99 (d, *J* = 8.1 Hz, 2H), 7.21 (dd, *J* = 8.5, 1.9 Hz, 1H), 7.39–7.44 (m, 2H), 7.46 (s, 1H), 7.54–7.59 (m, 2H), 7.61–7.66 (m, 2H), 7.70–7.75 (m, 1H). ¹³C NMR (CDCl₃) δ 21.4, 23.5, 54.0, 124.2, 125.2, 126.0, 126.2, 127.2, 127.6, 128.0, 128.5, 129.4, 132.8, 133.2, 137.7, 139.2, 143.2.

Deprotection of 4-Methylbenzenesulfonyl Group with liq. NH₃/Li. (for sulfonamide **2a**)

Lithium metal (104 mg, 15 mmol) was added to liquid ammonia (~10 mL) at -78 °C. To the mixture was added a solution of sulfonamide **2a** (137.7 mg, 0.50 mmol, >99% ee

obtained after recrystallization) in THF (1.0 mL). After 15 min stirring at the same temperature, the solution was quenched with solid NH_4Cl (802 mg, 15 mmol) and allowed to warm up to room temperature. The reaction mixture was quenched with 10% HCl aq and washed with Et_2O . To the aqueous layer was added 50% NaOH aq until pH 14 and the basic aqueous solution was extracted with Et_2O . The organic layer was dried over K_2CO_3 and concentrated on a rotary evaporator with a bath temperature maintained at 0 °C to give (*S*)-1-phenylethylamine as a pale yellow oil (47.9 mg, 0.395 mmol, 79% yield, >99% ee). The ee was determined on Chiralcel OD-H column with hexane/2-propanol/ Et_2NH = 9/1/0.1, flow = 0.4 mL/min, wavelength = 254 nm. Retention times: 18.8 min [(*R*)-enantiomer], 21.4 min [(*S*)-enantiomer]. $[\alpha]_{\text{D}}^{20} -32.7$ (*c* 1.04, CHCl_3). (lit.⁴ $[\alpha]_{\text{D}}^{20} -35.1$ (*c* 1.0, CHCl_3))

References

- 1) Cramer, R. *Inorg. Synth.* **1974**, 15, 14.
- 2) Otomaru, Y.; Okamoto, K.; Shintani, R.; Hayashi, T. *J. Org. Chem.* **2005**, 70, 2503.
- 3) Love, B. E.; Raje, P. S.; Williams, T. C., II *Synlett* **1994**, 493.
- 4) Atobe, M.; Yamazaki, N.; Kibayashi, C. *J. Org. Chem.* **2004**, 69, 5595.