Asymmetric Annulation towards Pyrroloperazinones: Concise Enantioselective Syntheses of Pyrrole Alkaloid Natural Products

Barry M. Trost,* Guangbin Dong

Department of Chemistry, Stanford University, Stanford, California 94305-5080.

bmtrost@stanford.edu

Supporting Information

Compound 8: (eg. Table 1, entry 3) A solution of $[\text{Pd}(\text{C}_3\text{H}_5)\text{Cl}]_2$ (0.92 mg, 0.0025 mmol) and (*R,R*)-**L** (5.2 mg, 0.0075 mmol) in 0.5 ml degassed DCM which had been stirring at rt for 10 min, was added to a solution of compound **6** (20.4 mg, 0.1 mmol) and compound **7** (22 mg, 0.1 mmol) in DCM (0.5 ml) under Ar. The mixture was stirred at rt for 16 hr, and then the solvent was removed under vacuum. Compound **8** with some uncyclized product were isolated via silica gel flash column chromatography (petroleum ether / ethyl acetate = 4/1, then 3/2) as a light yellow oil. After two days at rt under vacuum, the uncyclized intermediate was automatically converted to compound **8** as a light yellow foam. (28 mg, 72%, 95% ee by HPLC OD column, 90:10 heptane/*i*-propanol, 0.8 ml/min): R_f : 0.35 (petroleum ether / ethyl acetate = 3/2); $[\alpha]_D$: -21.12 (DCM, c 0.1); ^1H NMR (CDCl_3 , 500 MHz) δ 7.25 (m, 1H), 6.95 (d, J = 4.0 Hz, 1H), 6.43 (m, 2H), 6.26 (d, J = 4.0 Hz, 1H), 5.64 (ddd, J = 17, 10.5, 5.0 Hz, 1H), 5.10 (dd, J = 10.5, 1.5 Hz, 1H), 4.75 (m, 2H), 4.60 (d, J = 17 Hz, 1H), 4.50 (d, J = 14.5 Hz, 1H), 3.88 (dd, J = 13, 4.5 Hz, 1H), 3.79 (s, 6H), 3.48 (dd, J = 13, 1.5 Hz, 1H); ^{13}C -NMR (CDCl_3 , 125 MHz) δ 160.5, 158.7, 158.5, 133.8, 131.7, 125.6, 118.0, 117.2, 114.1, 112.4, 104.7, 104.2, 98.3, 55.5, 55.4, 54.6, 49.5, 43.7; IR (film) 2936, 1645, 1614, 1418, 1209 cm^{-1} ; HRMS ($\text{C}_{18}\text{H}_{19}\text{BrN}_2\text{O}_3$): Calc'd. 392.055858 (M^+), Found 392.056840.

Compound 9: 9-BBN (99 mg, 0.40 mmol) was added to a solution of **8** (105 mg, 0.27 mmol) in THF (2 ml) at 0 °C under N_2 . The resulting solution was stirred at rt for 3 h, before it was quenched with $\text{NaBO}_3 \cdot \text{H}_2\text{O}$ (81mg, 0.81 mmol) and H_2O (1 ml). The mixture was stirred at rt for 12h before it was extracted with ethyl acetate (20 ml x 3). The combined organic fraction was dried over MgSO_4 . Compound **9** was purified via

silica gel flash column chromatography (5% MeOH in DCM) to give a white foam (90 mg, with 7.7 mg starting material recovered, 82%, brsm 88%): R_f : 0.40 (10% MeOH in DCM); $[\alpha]_D$: -31.64 (c 0.35, DCM); ^1H NMR (CDCl_3 , 500 MHz): δ 7.28 (m, 1H), 6.92 (d, J = 4.0 Hz, 1H), 6.45 (m, 2H), 6.23 (d, J = 4.0 Hz, 1H), 5.00 (d, J = 14.5 Hz, 1H), 4.42 (m, 1H), 4.35 (d, J = 14 Hz, 1H), 3.81 (s, 3H), 3.81 (s, 3H), 3.77 (m, 1H), 3.56 (dd, J = 13, 1.5 Hz, 1H), 3.47 (m, 1H), 3.33 (m, 1H), 1.72 (m, 2H); ^{13}C -NMR(CDCl_3 , 125 MHz) δ 160.7, 158.8, 158.4, 132.3, 125.2, 117.4, 114.2, 112.3, 104.4, 103.7, 98.5, 59.1, 55.5, 50.2, 47.6, 43.2, 35.0, 30.4; IR (film) 3383 (br), 2923, 2851, 1660, 1615, 1417, 1209 cm^{-1} ; HRMS ($\text{C}_{18}\text{H}_{21}\text{BrN}_2\text{O}_4$): Calc'd. 410.066422 (M^+), Found 410.068069.

Compound 10: TFA (0.6 ml) was added to a solution of **9** (58 mg, 0.14 mmol), tetrahydrothiophene (59.4 mg, 0.68 mmol) in DCM (0.6 ml) at rt. The resulting brown solution was stirred at 50 °C for 20 hr, before poured into pre-cooled sat. NaHCO_3 solution. The mixture was extracted with ethyl acetate (20 ml x 3), and the combined organic fraction was dried over Na_2SO_4 . The products, as a ca. 2:1 mixture (mono-bromo-product and debromo-product) were purified via silica gel flash column chromatography (10% MeOH/DCM). The mixture was then dissolved with THF (1 ml) and MeOH (0.5 ml). At 0 °C, NBS (37 mg, 0.21 mmol) was added portionwisely to the above solution. The resulting solution was stirred at rt for 0.5 hr before the solvents were removed under vacuum. Compound **10** was purified via silica gel flash column chromatography (5% MeOH in DCM) to give a white foam (46.1 mg, 96% over two steps): R_f : 0.50 (10% MeOH in ethyl acetate); $[\alpha]_D$: -27.3 (c 0.6, MeOH); ^1H NMR (CD_3OD , 400 MHz): δ 6.92 (s, 1H), 4.60 (m, 1H), 3.80 (dd, J = 13.6, 4.0 Hz, 1H), 3.69-3.64 (m, 3H), 2.00 (m, 1H), 1.85 (m, 1H); ^{13}C -NMR (CD_3OD , 100 MHz) δ 161.1, 126.1, 116.4, 108.1, 101.3, 59.3, 53.3, 43.4, 35.5; IR (film) 3308 (br), 2927, 2882, 1652, 1550, 1465, 1429, 1338 cm^{-1} ; HRMS ($\text{C}_9\text{H}_{10}\text{Br}_2\text{N}_2\text{O}_2$): Calc'd. 335.910900 (M^+), Found 335.910485.

Longamide B (1)

Method A: To CH_3CN (0.5 ml) was added H_5IO_6 (17.6 mg, 0.077mol) and the mixture was stirred vigorously at rt for 15 min. A solution of compound **10** (8.7 mg, 0.026 mmol) in CH_3CN (0.2 ml) was then added at 0 °C, followed by addition of a PCC solution

(0.0013 mmol, in 0.01 ml CH₃CN). The resulting mixture was stirred at rt overnight before diluted with ethyl acetate. The mixture was washed with brine, sat. aq. NaHSO₃ and brine respectively. Longamide B (**1**) was further purified via a classical acid-base treatment as a white powder (5.7 mg, 63%, brsm 90%; 2.6 mg **10** was recovered).

Method B: TEMPO (0.44 mg, 0.0028 mmol) was added to a mixture of compound **10** (9.5 mg, 0.028 mmol), PhI(OAc)₂ (22.7 mg, 0.0705 mmol) and NaHCO₃ (9.4 mg, 0.112 mmol) in DCM (0.15 ml) at rt under air. The resulting solution was stirred at rt for 3 hr, before quenched with sat. Na₂S₂O₃ and sat. NaHCO₃. The mixture was diluted with ethyl acetate, and longamide B (**1**) was further purified via a classical acid-base treatment as a white powder (9.3 mg, 95%).

mp 203-204 °C (decomposed to black tar); [α]_D: -8.21 (c 0.52, MeOH); ¹H NMR (CD₃OD, 500 MHz): δ 6.94 (s, 1H), 4.81 (m, 1H), 3.89 (ddd, *J* = 13.5, 4.5, 1.5 Hz, 1H), 3.66 (dd, *J* = 13.5, 1.5 Hz, 1H), 2.86 (dd, *J* = 16.5, 11 Hz, 1H), 2.55 (ddd, *J* = 16.5, 3.5, 1.5 Hz, 1H); IR: 3282, 3203, 2924, 1715, 1645, 1558, 1428 cm⁻¹; HRMS (C₉H₈Br₂N₂O₃): Calc'd. 349.890164 (M⁺), Found 349.889306.

Agesamides A and B (**5**)

Dess–Martin periodane (34.3 mg, 0.081 mmol) was added to a mixture of compound **10** (16.5 mg, 0.040 mmol), NaHCO₃ (10.1 mg, 0.12 mmol) in DCM (0.4 ml) at 0 °C. The resulting suspension was stirred at rt for 0.5 hr, before quenched with sat. Na₂S₂O₃ and sat. NaHCO₃. The mixture was extracted with ethyl acetate (10 ml x 3), and the combined organic fraction was dried over Na₂SO₄. The solvent was removed, and the resultant aldehyde with (NH₄)₂CO₃ (8.8 mg, 0.1125 mmol) and KCN (3.0 mg, 0.045 mmol) was then dissolved in EtOH/H₂O (1:1, 0.4 ml). The resulting solution was stirred vigorously at 60 °C overnight. The solvents were removed under vacuum, and agesamides A and B (**5**) was purified via preparative silica gel TLC (10% MeOH in ethyl acetate) as a white solid (1:1 mixture, 17.5 mg, 100 % over two steps): ¹H NMR (DMSO-*d*₆, 400 MHz): δ 10.80 (br, 1H), 8.18 (s, 1H), 8.04 (s, 1H), 7.89-7.88 (2H), 6.85 (s, 2H), 4.65 (m, 1H), 4.53 (d, *J* = 11.2 Hz, 1H), 4.20 (ddd, *J* = 11, 3.0, 1.5 Hz, 1H), 4.13 (m, 1H), 3.78-3.70 (2H), 3.54-3.46 (2H), 2.12 (ddd, *J* = 14, 11.5, 3.5 Hz, 1H), 2.04 (ddd, *J* = 14.0, 9.0, 7.0 Hz, 1H), 1.79 (m, 1H), 1.53 (m, 1H); ¹³C-NMR (DMSO-*d*₆, 100 MHz) δ 175.2, 174.9, 157.53, 157.46, 157.2, 157.1, 125.5, 114.0, 113.9, 105.71, 105.67, 99.3, 54.6, 54.3, 50.6, 41.7, 40.2, 33.8, 33.5; IR: 3228 (br), 2926, 2854, 1724, 1646, 1550, 1429, 1338 cm⁻¹; HRMS (C₁₁H₁₀Br₂N₄O₃): Calc'd. 403.911963 (M⁺), Found 403.910347.

X-94-1A
11-10-2006

expt1 s2pu1

SAMPLE DEC. 8 VT
date Nov 10 2006 dfrq 499.751
solvent CDCl3 dn H1
file ACQUISITION exp dpr 30
sfrq 499.751 dm mn 0
tn H1 dm c
at 4.000 dseq 200
np 14.000 dresq
sb 8000.0 homo 1.0
fb not used
bs 4
tpwr 60 dfrq2 DEC2 0
pw 10.0 dn2
dl 0 dpr2 1
tof 0 dot2 n
nt 16 dm2 n
ct 16 dnm2 c
atlock n dm2 200
gain not used dresq2
FLAGS n dresq2 1.0
11 n homo PROCESSING
1n n vfile
dp y proc
hs n wfile 131072 f
DISPLAY -249.9 math
wp 4747.6
vs 151 werr
sc 0 wexp
wc 250 wds
hzmm 18.99 wnt
ls 33.57 wft
fft 5142.7
tft 3028.2
tn 100.000
nm cdc ph

XI-32A
12-05-2006

exp1 s2pu1

SAMPLE		DEC. & VT	
date	Dec 5 2006	dfrq	499.751
solvent	CDCl3	dn	HL
file	exp	dpu	30
ACQUISITION		dot	0
sfrq	499.751	dm	mm
in	HL	dmq	mm
at	4.000	dsq	208
np	64000	dsq	1.0
sw	8000.0	dsq	1.0
zb	not used	homo	n
ls	4	DEC2	0
tpwr	60	dfrq2	0
pu	10.0	dn2	1
dl	0	dpu2	0
tof	0	dot2	0
nt	16	dm2	n
ct	16	dm2	c
clock	n	dm2	200
gain	not used	dsq2	1.0
flags	n	dsq2	n
11	n	homo2	n
in	y	PROC	ft
dp	mn	wtfile	131072
hs	mn	proc	f
DISPLAY		math	
sp	-249.9	math	
wp	4747.6	math	
vs	140	math	
sc	0	math	
wc	250	math	
h2nm	18.99	math	
h1nm	435.11	math	
f2	5142.2	math	
f1	3628.2	math	
th	7	math	
ins	0.850	math	
nm	cdc ph	math

XI-77A-H
01-24-2007

expi sidh

SAMPLE DEC. & VT
date Jan 24 2007 dfrq 400.115
solvent CDCl3 dn 31
f1 31 d1 31
f2 31 d2 31
ACQUISITION exp 30
sfrq 400.115 dm 0
tn 4.002 H1 dmm nnn
at 4.002 dmf c
np 40000 wtfile 200
sw 4997.5 proc
fb not used fl
bs 4 fn not used
tpwr 61
pw 8.0 werr
d1 0.500 wexp
LOF 0 wds
nr 10 wnt
cs 10
clock n
gain not used
flags n
ij n
in n
dp y
DISPLAY
SD -200.2
WD 3801.0
VS 132
SC 0
WC 256
H2NMR 15.256
F1 321.60
F2 1601.8
F3 1324.4
th 10
ins 1.000
nm cdc ph

XI-86A
 02-13-2007
 exp1 std1h
 SAMPLE DEC. & VT
 date 02-13-2007 dffq 400.115
 solvent dmsd dn H1
 file ACQUISITION exp dpr 30
 sffq 400.115 dm dof 0
 tn H1 dnm nm C
 at 4.002 dmf PROCESSING 200
 mp 48000 wtitle
 sw 4997.5 not used proc ft
 fb not used fn not used
 bs 61
 tpr 8.0 weff
 pw 0.500 wepd
 tof 0 wbs
 nt 16 wnt
 ct 16
 alock n
 gain not used
 flags
 i1 n
 tn n
 dp y
 DISPLAY
 sp 280.6
 wp 4601.2
 vs 0
 sc 0
 wc 250
 hzmm 19.21
 is 281.76
 rfi 1473.2
 rfp 996.3
 lh 20
 ins 1.000
 nm cdc ph

Agesamides A and B (5)

XI-86A
 02-18-2007
 exps std13c
 SAMPLE
 date Feb 13 2007 dfrq 400.115
 solvent DMSO d1
 file /export/home/...
 gudong/XI-86A-C-1
 d
 yyy
 w
 ACQUISITION
 sfrq 100.618 dfrq 8900
 tn C13 1b
 at 1.198 1b
 np 59968 wfile
 sw 25000.0 proc
 fd 13600 tn
 not used
 bs 5
 tpr 5
 pw 4.5 wexp
 sm 0.500 wps
 tof 0 wnt
 nt 1200
 ct 832
 alock n
 gain not used
 flags
 11 n
 1n n
 dp y
 DISPLAY
 sd -3060.5
 wd 24899.2
 vs 453
 sc 0
 wc 250
 hz 100.00
 ts 500.00
 rf 2013.2
 rf1 3953.8
 tn 8
 tns 100.000
 nm no ph

Agesamides A and B (5)