Asymmetric Diels-Alder Reactions of a New Enantiomerically pure Sulfinyl-quinone : A straightforward access to functionalised Wieland-Miescher ketone analogues with (*R*) absolute configuration.

Don Antoine Lanfranchi¹ and Gilles Hanquet^{1 *}

¹Laboratoire de Stéréochimie associé au C.N.R.S., ECPM, ULP, 25, rue Becquerel, 67087 Strasbourg, France

E-mail address: ghanquet@chimie.u-strasbg.fr

SUPPORTING INFORMATION

Table of contents

General Methods, Instrument Details and Materials	S-2
Preparation of and characterization of 8	S-3
Preparation of and characterization of 9	S-3, S-4
Preparation of and characterization of 10	S-4
Preparation of and characterization of 10a	S-5
Preparation of and characterization of 11	S-5, S-6
Preparation of and characterization of 12	S-6
Preparation of and characterization of 13	S-6, S-7
Preparation of and characterization of (+)- 2	S-7, S-8
Preparation of and characterization of (-)- R-4	S-8
Preparation of and characterization of (-)- R-14	S-8, S-9
Preparation of and characterization of (-)- R-3	S-9
Preparation of and characterization of (<i>rac</i>)- 13	S-10
General procedure for Diels-Alder reactions	S-11
Preparation of and characterization of 18a	S-11
Characterization of 18b	S-12
Characterization of 19a	S-12
Characterization of 20	S-12
Characterization of 20	S-13
Mass spectra of compounds 8-13	S-14→S-17
¹ H and ¹³ C NMR of compounds 2-4, 12-14	S-17→S-31
¹ H NMR Eu(tfc) ₃ dosage of compounds (+) 2 , (-) 3 , (-) 4	S-32→S-35

¹H and ¹³C NMR of compounds **18-21**

S-36→S-47

VCD/IR measurement of compounds (–)-**R-3**, (–)-**R-4**

S-48→S-54

Crystallographic data of **18a**

S-55

General. Reagents and solvents were purchased as reagent grade and used without further purification. THF was distilled over sodium benzophenone ketyl. Dichloromethane was distilled over CaH₂. Column chromatography: silica gel 60 (230-400 mesh, 0.040-0.063 mm). Demetallated silica gel was prepared according to published procedure.¹ Thin Layer Chromatography (TLC) was performed on glass sheets coated with silica gel 60 F₂₅₄ (otherwise stated), visualization by UV light. IR (cm⁻¹) measurements using Universal ATR sampling accessories. NMR spectra (¹H and ¹³C) were recorded on a 200 MHz or 300 MHz or 400 MHz. Chemical shifts are reported in ppm with the solvent (CDCl₃) resonance as the δ 7.26 ppm in ¹H spectra, and relative to the central CDCl₃ resonance δ 77.0 ppm in ¹³C spectra. Data reported as (s=singlet, d=doublet, t=triplet, q=quartet, m=multiplet, br=broad ; coupling constant(s) in Hz; integration). The enantiomeric excess (e.e.) of the products (+)-**2**, (–)-**3**, and (–)-**4** were determined by ¹H NMR dosage on a 400 MHz with Eu(tfc)₃ versus racemic compounds. Optical rotations were measured on a polarimeter with a sodium lamp and are reported as follows: $[\alpha]_D^{25}$ (c g/100 mL, solvent). GC-MS analyses were kindly performed with a chromatograph coupled with a mass spectrometer, using the following conditions: fused-silica capillary column (dimethylsiloxane) 50 m×0.2 mm i.d., film thickness 0.5 μm. Injection mode: split 1:30. The oven temperature was programmed from 60 °C to 250 °C at 2 °C/min and then held isothermal for 30 min. The carrier gas was He (0.8 ml/min). Injector and detector temperatures were 250 °C and 230 °C respectively. The mass spectra were performed at 70 eV, mass range 35-400. ZnBr₂ was flamed-dried in the reaction flask under vacuum and under Argon before use.

Materials. Precursor dienoxysilane **5** was prepared from commercial ethylvinylketone (1 step, 61%) as previously reported², 2,5-Dimethoxy-4-methyl-benzaldehyde **7** was prepared³ on 200 g scale (2 steps, 80%) from commercial 2-methyl-hydroquinone.

¹ Hubbard, J.S.; Harris, T.M., *J. Org. Chem.*, **1981**, 46, 2566.

² Cazeau, P. ; Duboudin, F. ; Moulines, F. ; Babot, O. ; Dunogues, J. *Tetrahedron*, **1987**, 43, 2089 ; Ackland, D. J. ; Pinhey, J. T., *J. Chem. Soc., Perkin Trans I*, **1987**, 2689 ; Bell, V. L. ; Holmes, A. B. ; Hsu, S.-Y. ; Mock, G.A. ; Raphael, R. A., *J. Chem. Soc., Perkin Trans I*, **1986**, 1507 ; Danishefsky, S. ; Feng Yan, C. *Synth. Comm.*, **1978**, 8, 211.

³ Standridge, R.T.; Howell, H.G.; Gyls, J.A.; Partyka, R.A.; Shulgin, A.T., *J. Med. Chem.*, **1976**, 19, 1400.

2-Hydroxy-5-methoxy-4-methyl-benzaldehyde (8)

Under an Argon atmosphere, 100.00 g (0.555 mol ; 1 eq) of aldehyde **7** dissolved in 1000 mL of MeCN was added dropwise (~10 mn) to a solution of 1000 mL of MeCN containing 74.83 g (0.560 mol ; 1.01 eq) anhydrous AlCl₃. The resulting mixture was heated at 45°C. Anhydrous NaI 124.85 g (0.833 mol ; 1.5 eq) was added under vigorous stirring. The mixture was refluxed for 2 h. The “curry yellow” solution was allowed to cool down at room temperature. TLC indicated complete conversion (Al₂O₃ neutral ; Hexanes / Et₂O : 3 / 1, R_{f8} : 0.42 vs. R_{f7}: 0.30). The solution was filtered to remove the solids, the filtercake was kept and the filtrate was evaporated to dryness under vacuum to give a yellow solid. The filter cake and the yellow solid were dissolved in 1000 mL of EtOAc and 1500 mL of saturated solution of tartaric acid disodium were added under vigorous stirring until the apparition of two distinct layers (~1 h). The aqueous phase was extracted with 2x600 mL of EtOAc. The brown organic layers were combined and washed first with 2x600 mL of an aqueous solution of saturated Na₂S₂O₃ (the organic layer became orange light), then with 600 ml of brine. The combined organic layers were dried over MgSO₄, and concentrated in vacuum to give 136.90 g (99%) of yellow needles **8** used without purification in the next step.

This salicylaldehyde is stable on storage (R.T.) for more than 2 years.

¹H NMR (300 MHz ; CDCl₃) : δ 2.26 (s, 3H) ; 3.83 (s, 3H) ; 6.80 (s, 1H) ; 6.86 (s, 1H) ; 9.81 (s, 1H,) ; 10.76 (s, 1H). ¹³C NMR (75 MHz ; CDCl₃) : δ 17.5 ; 56.2 ; 112.2 ; 118.52 ; 120.0 ; 139.5 ; 151.7 ; 156.6 ; 195.7. MS (70eV), *m/e* 167 (10), 166 (M+, 100), 165 (11), 152 (7), 151 (79), 123 (22), 95 (6), 77 (9), 67 (7), 66 (5), 55 (7), 53 (7), 39 (5). TLC R_f : 0.42 (Al₂O₃ neutral ; Hexanes / Et₂O : 3 / 1) m.p. : 102-103°C.

3-Bromo-2-hydroxy-5-methoxy-4-methyl-benzaldehyde (9)

A solution of the phenol **8** (75.00 g ; 451.62 mmol ; 1 eq) and anhydrous sodium acetate 55.54 g (677.43 mmol ; 1.50 eq) in acetic acid (1500 mL) was stirred at room temperature. After the complete dissolution of all the solids, a solution of bromine (23.66 mL ; 73.55 g; 406.60 mmol ; 1.02 eq), dissolved in acetic acid (100 mL), was added dropwise (~30 mn) to the phenol solution. After two hours the mixture was filtered to remove the solids, the filtercake was kept and the filtrate was evaporated to dryness under vacuum to give a yellow solid. The resulting yellow solid and the filtercake were dissolved in 1000 mL of CH₂Cl₂. The organic layer was washed with 800 mL of water, saturated Na₂S₂O₃ solution (500 mL), saturated NaHCO₃ solution (500 mL), and brine (500 mL), dried over MgSO₄, and concentrated in vacuum to give 107.10 g (96%) of yellow crystals **9** used without purification in the next step.

The analytical sample was obtained by crystallisation from boiling EtOH .

This bromophenol is stable on storage (R.T.) for more than 2 years.

¹H NMR (200 MHz ; CDCl₃) : δ 2.42 (s, 3H) ; 3.86 (s, 3H) ; 6.92 (s, 1H) ; 9.80 (s, 1H) ; 11.42 (s, 1H). ¹³C NMR (75 MHz ; CDCl₃) : δ 17.0 ; 56.3 ; 111.3 ; 114.9 ; 117.8 ; 138.6 ; 151.3 ; 153.1 ; 195.0. MS (70eV), *m/e* 246 (99), 245 (21), 244 (M+, 100), 243 (12), 231 (47), 229 (51), 203 (19), 201 (21), 67 (10), 65 (15), 53 (12). TLC R_f : 0.47 (Et₂O / hexanes ; 1 / 1) m.p. : 103°C (EtOH).

3-Bromo-2,5-dimethoxy-4-methyl-benzaldehyde (**10**)

Under an Argon atmosphere and at R.T., 100.00 g (404.90 mmol ; 1.00 eq) of aldehyde **9** dissolved in 950 mL of CH₂Cl₂ was added and 950 mL of an aqueous basic solution of NaOH (27.02 g ; 676.18 mmol ; 1.67 eq) followed by, under vigorous stirring, 9.30 mL (8.22 g ; 20.24 mmol ; 0.05 eq) aliquat@336 and 76.65 mL (102.14 g ; 809.80 mmol ; 2 eq) of Me₂SO₄. After two hours, TLC indicated complete conversion. The excess of Me₂SO₄ was destroyed by adding, under vigorous stirring, 16.62 g pellets of NaOH into the solution. The reaction mixture was stirred overnight. The organic layer of the reaction mixture was kept and the aqueous layer was extracted with CH₂Cl₂ (2 x 500 mL), the organic layers were combined and washed with brine, dried over MgSO₄ and concentrated. The crude **10** is filtered through a short pad of SiO₂ by using CH₂Cl₂ to remove traces of ammonium salts, the solvent was evaporated under vacuum to give 101.91 g (98%) of yellow crystals **10** used without purification in the next step. This gentilsaldehyde is stable on storage (R.T.) for more than 2 years.

¹H NMR (300 MHz ; CDCl₃) : δ 2.39 (s, 3H) ; 3.85 (s, 3H) ; 3.92 (s, 3H) ; 7.22 (s, 1H) ; 10.30 (s, 1H). ¹³C NMR (75 MHz ; CDCl₃) : δ 17.3 ; 56.5 ; 64.3 ; 106.7 ; 122.2 ; 128.0 ; 128.1 ; 137.0 ; 155.2 ; 189.5. MS (70eV), *m/e* 260 (97), 258 (M+, 100), 245 (51), 243 (57), 214 (42), 212 (42),

200 (22), 121 (22), 108 (19), 93 (21), 77 (27), 65 (19). **TLC R_f** : 0.64 (Hexanes / Et₂O : 1 / 1). **m.p.** : 92°C (cyclohexane). **Anal.** Calcd for C₁₀H₁₁BrO₃ : C, 46.72 ; H, 3.53. Found : C, 46.54 ; H, 3.87

Formic acid 3-bromo-2,5-dimethoxy-4-methyl-phenyl ester (10a)

A solution of 90.00 g (347.52 mmol ; 1 eq) of **10** dissolved in 200 mL of CH₂Cl₂ was added dropwise (~10 mn) at room temperature to a solution of 94.00 g (435.94 mmol ; 1.25 eq) of *m*-CPBA (80 %) in 1000 mL of CH₂Cl₂. The mixture was stirred for 24 h at room temperature. The mixture was filtered to remove the solid *m*-chlorobenzoic acid. The filtercake was washed with CH₂Cl₂ (2 x 200 mL). The combined organic phases were washed with saturated Na₂S₂O₃ solution (600 mL), saturated NaHCO₃ solution (600 mL), and brine (600 mL), dried over MgSO₄ and concentrated to give 93.74 g (98%) of formate **10a** as a very pale orange solid used without purification in the next step. This formate darkened rapidly (few days) at R.T., it is not very stable on storage.

¹H NMR (200 MHz ; CDCl₃) : δ 2.30 (s, 3H) ; 3.76 (s, 3H) ; 3.78 (s, 3H) ; 6.61 (s, 1H) ; 8.28 (s, 1H). **¹³C NMR** (75 MHz ; CDCl₃) : δ 17.3 ; 55.2 ; 60.7 ; 104.0 ; 117.8 ; 120.4 ; 142.1 ; 145.0 ; 159.2 ; 160.2. **MS** (70eV), *m/e* 276 (15), 274 (16), 248 (15), 246 (15), 233 (M⁺, 100), 232 (12), 231 (99), 205 (44), 203 (48), 124 (18), 109 (23), 77 (12), 69 (21), 67 (14), 53 (12). **m.p.**: 82°C (cyclohexane / pentane).

3-Bromo-2,5-dimethoxy-4-methyl-phenol (11)

The orange formate **10a** (91.86g ; 334.08 mmol ; 1 eq) was dissolved in 700 mL of refluxing MeOH. To the resulting orange solution was added dropwise, at room temperature, a solution of aqueous potassium hydroxide (100 mL of distilled water containing 22.2 g KOH ; 395.86 mmol ; 1.18 eq) : the mixture darkened immediately. Five minutes later TLC (Hexanes / Et₂O ; 4 / 3) indicated the completion of the hydrolyse. Methanol was evaporated in vacuum, leaving a black oil. The crude mixture was poured in 1000 mL of water, acidified to acidic pH (~2) by means of concentrated HCl : yellow solids started to appear. The solution was filtered to remove the solids, the filtercake was washed with cold water (3 x 200 mL) and kept and the aqueous filtrate extracted with CH₂Cl₂ (2 x 250 mL). The filtercake is dissolved in 200 mL of CH₂Cl₂, the combined organic layers were washed with water until neutral pH and with brine (300 mL), dried over MgSO₄, and concentrated in vacuum to give 77.55 g (94%) of yellow crystals **11** used without purification in the next step. The analytical sample was obtained as white needles by crystallisation from boiling cyclohexane / pentane.

¹H NMR (200 MHz ; CDCl₃) : δ 2.21 (s, 3H) ; 3.74 (s, 3H) ; 3.81 (s, 3H) ; 5.57 (s, 1H) ; 6.49 (s, 1H). ¹³C NMR (75 MHz ; CDCl₃) : δ. 15.1 ; 56.0 ; 61.1 ; 98.3 ; 118.9 ; 119.5 ; 137.9 ; 147.5 ; 154.9. MS (70eV), *m/e* : 232 (96), 231 (25), 230 (M⁺, 100), 229. TLC R_f : 0.33 (cyclohexane / Et₂O : 2 / 1). m.p. 91-92°C (cyclohexane / pentane). Anal. Calcd for C₉H₁₁BrO₃ : C, 41.23 ; H, 3.89. Found : C, 41.39 ; H, 3.59

3-Bromo-1,2,5-trimethoxy-4-methyl-benzene (**12**)

The synthetic protocol is the same as for **10** (3-Bromo-2,5-dimethoxy-4-methyl-benzaldehyde). The crude **12** was filtered through a short pad of SiO₂ by using CH₂Cl₂ to remove traces of ammonium salts, the solvent was evaporated under vacuum to give orange solids **12** with 97% yield. It could be purified by crystallisation from boiling cyclohexane or by flash chromatography (eluent : hexanes / Et₂O, 3 / 2). This compound is stable on storage (R.T.) for more than 2 years.

¹H NMR (200 MHz ; CDCl₃) : δ 2.55 (s, 3H) ; 3.79 (s, 3H) ; 3.81 (s, 3H) ; 3.87 (s, 3H) ; 6.47 (s, 1H). ¹³C NMR (75 MHz ; CDCl₃) : δ 15.3 ; 56.3 ; 56.4 ; 60.5 ; 96.7 ; 119.2 ; 121.5 ; 140.4 ; 151.3 ; 154.2. MS (70eV), *m/e* : 262(81), 261 (10), 260 (82), 248 (10), 247 (97), 246 (13), 245 (M⁺, 100), 219 (37), 217 (42), 204 (14), 202 (16), 138 (20), 123 (24), 77 (15). TLC R_f : 0.44 (hexanes / Et₂O : 3 / 2). m.p. : 61°C (cyclohexane). Anal. Calcd for C₁₀H₁₃BrO₃ : C, 46.01 ; H, 5.02. Found : C, 45.89 ; H, 5.04

(-)-(S)S-1,2,5-Trimethoxy-4-methyl-3-(toluene-4-sulfinyl)-benzene (13)

Under an Argon atmosphere, to a suspension of 1.38 g magnesium turnings (56.78 mmol ; 1.05 eq) in 60 mL of anhydrous THF, a solution of **12** (14.53 g ; 55.79 mmol ; 1.00 eq) in 60 mL of anhydrous THF was added. After the initiation of Grignard, the mixture was stirred at room temperature for 4 hours. The Grignard solution was cooled down to 0°C and then transferred through double tipped needle by means of Argon pressure over a cooled (0°C) solution of (-)-(*R*)S-menthyl-*p*-toluenesulfonate⁴ (17.40 g ; 59.14 mmol ; 1.06 eq) in 120 mL of anhydrous THF with vigorous stirring. The resulting mixture was stirred overnight. Hydrolysis was performed, at 0°C, with 250 mL of saturated NH₄Cl solution. The aqueous layer was extracted with Et₂O (3 x 150 mL). The combined organic phases were washed with brine (250 mL), dried over MgSO₄, and concentrated in vacuum to give a waxy solid smelling menthol. Trituration of this oily solid in Et₂O (100 mL) afforded pure sulfoxide **13** (11.61 g; 65%) as a white microcrystalline solid. This sulfoxide is stable on storage (R.T.) for more than 2 years.

¹H NMR (400 MHz ; CDCl₃) : δ 2.26 (s, 3H) ; 2.35 (s, 3H) ; 3.78 (s, 3H) ; 3.82 (s, 3H) ; 3.86 (s, 3H) ; 6.58 (s, 1H) ; 7.21 (d, ³J=8.2 Hz, 2H) ; 7.47 (d, ³J=8.2 Hz, 2H). ¹³C NMR (75 MHz ; CDCl₃) : δ 9.9 ; 21.3 ; 56.2 ; 61.7 ; 100.6 ; 119.9 ; 124.3 ; 129.4 ; 137.4 ; 139.8 ; 141.5 ; 141.8 ; 151.0 ; 154.6. MS (70eV), *m/e* : 320 (43), 304 (46), 303 (M⁺, 100), 288 (67), 273 (33), 272 (75), 258 (32), 257 (33), 228 (60), 167 (28), 166 (34), 151 (35), 136 (34), 121 (31), 105 (62), 91 (42), 77 (28). **m.p.** : 129°C. **Anal.** Calcd for C₁₇H₂₀O₄S : C, 63.73 ; H, 6.29 ; S, 10.01. Found : C, 63.48 ; H, 6.45 ; S, 10.21. $[\alpha]_D^{20} = -154.0$ (c = 1.21 ; Acetone)

(+)-(S)S-5-Methoxy-2-methyl-3-(toluene-4-sulfinyl)-[1,4]benzoquinone (2)

⁴ Andersen, K.K.; Gaffield, W.; Papanikolaou, N.E., *J. Am. Chem. Soc.*, **1964**, 56, 5635. Solladié, G., *Synthesis*, **1981**, 185.

An aqueous solution (500 mL) of ammonium cerium (IV) nitrate (CAN) (47.270 g ; 86.222 mmol ; 2.5 eq) was added rapidly (< 30 s) to a solution of sulfoxide **13** (10.012 g ; 31.268 mmol ; 1.0 eq) in MeCN (500 mL) at room temperature. After stirring for 40 mn, the ^1H NMR indicated that the oxidative demethylation was complete (extended reaction time tends to decrease frankly the yield). The solvent was evaporated and the mixture extracted with CH_2Cl_2 (2x300 mL). The combined extracts were washed with water (300 mL), brine (300 mL), and dried over MgSO_4 . The solvent was evaporated in vacuum leaving the sulfinylquinone as an orange solid. Recrystallization of the sulfinylquinone (+)-**2** from boiling MeOH (70 mL) gave pure (+)-**2** (8.343 g ; 92%) as bright orange needles. This sulfinylquinone is stable on storage (4–8°C, protected from light) for more than 2 years.

^1H NMR (400 MHz ; CDCl_3) : δ 2.39 (s, 3H) ; 2.51 (s, 3H) ; 3.79 (s, 3H) ; 5.96 (s, 1H) ; 7.30 (d, $^3J=8.3$ Hz, 2H) ; 7.64 (d, $^3J=8.1$ Hz, 2H). ^{13}C NMR (75 MHz ; CDCl_3) : δ 9.4 ; 21.4 ; 56.5 ; 107.9 ; 124.9 ; 130.1 ; 139.4 ; 144.5 ; 148.1 ; 157.8 ; 179.1 ; 185.2. IR (cm^{-1}) : 1668 ; 1626 ; 1591 ; 1205. m.p. : 133–134°C (MeOH) (m.p. racemic : 132–133°C from MeOH). Anal. Calcd for $\text{C}_{15}\text{H}_{14}\text{O}_4\text{S}$: C, 62.05 ; H, 4.86 ; S, 11.04. Found : C, 61.89 ; H, 4.80 ; S, 11.06. $[\alpha]_D^{20} = +445.5$ (c = 1.00 ; CH_2Cl_2)

(–)-R-2-Methoxy-4a,7,8-trimethyl-4a,5-dihydro-[1,4]naphthoquinone (4)

To a solution of sulfinylquinone (+)-**2** (290.20 mg ; 1.00 mmol ; 1.0 eq) in 10 mL of dry CH_2Cl_2 , under Argon, piperylene (Aldrich, mixture of isomers; 0.199 mL ; 136.24 mg ; 2.00 mmol; 2.0 eq) was added. After 10 to 12 days at R.T., the solvent was evaporated leaving the crude cycloadduct. Flash chromatography on demetalled silica gel (gradient elution 9 / 1 $\rightarrow$ 3 / 1 cyclohexane / EtOAc) afforded pure (–)-**4** as a yellow oil (200.6 mg, 90%) who crystallised on standing. (Note : chromatography of reaction mixture using standard silica gel and cyclohexane / EtOAc, 2 / 1 as eluent afforded (–)-**4** in 72–75% yield)

^1H NMR (200 MHz ; CDCl_3) : 1.39 (s, 3H) ; 2.35 (s, 3H) ; 2.53 (d, $J=2$ Hz, 1H) ; 2.55 (d, $J=1$ Hz, 1H) ; 3.81 (s, 3H) ; 5.89 (s, 1H) ; 6.09 (m, 1H) ; 6.22 (m, 1H). ^{13}C NMR (75 MHz ; C_6D_6) : δ 21.9 ; 26.0 ; 32.8 ; 45.5 ; 55.5 ; 107.7 ; 129.0 ; 131.0 ; 133.3 ; 145.5 ; 164.1 ; 180.9 ; 199.9. IR (cm^{-1}) : 1650 ; 1602 ; 1201. TLC Rf : 0.26 (cyclohexane / EtOAc : 2 / 1). m.p. : 132°C. Anal. Calcd for $\text{C}_{13}\text{H}_{14}\text{O}_3$: C, 71.58 ; H, 6.42. Found : C, 71.39 ; H, 6.31. $[\alpha]_D^{20} = -114.5$ (c = 0.40 ; CHCl_3)

(-)-R-2-Methoxy-4a,8-dimethyl-7-trimethylsilanyloxy-4a,5-dihydro-[1,4]naphthoquinone (14)

Advertising : The dienoxysilane **5** can be kept for more than a year under an inert atmosphere at – 20~15°C without apparent decomposition (¹H NMR). Nevertheless, before launching the Diels-Alder reaction with sulfinylquinone **2** and a not freshly prepared diene, the latter should be dissolved in Et₂O, washed with a saturated NaHCO₃ solution, then brine and dried over MgSO₄, and distilled (50°C / 30 mbar) as a white oil

To a solution of quinone (+)-**2** (1.000 g ; 3.445 mmol ; 1.00 eq) in 15 mL of dry CH₂Cl₂ under Argon diene **5** (0.108 g ; 6.890 mmol ; 2.01 eq) was added, at temperature inferior at – 20°C. The reaction flask was put into the freezer (– 20°C) for 10 days. Flash chromatography on demetallated silica gel (gradient elution 9 / 1 → 3 / 1 cyclohexane / EtOAc) afforded pure (–)-**3** as a yellow oil (0.960 g, 91%) which crystallised on standing. (Note : chromatography of reaction mixture using standard silica gel afforded (–)-**3** and (–)-**4** in only 15 and 20% yield respectively)

¹H NMR (300 MHz ; CDCl₃) : δ 0.26 (s, 9H) ; 1.29 (s, 3H) . 2.29 (s, 3H) ; 2.53 (d, *J*=2 Hz, 1H) ; 2.55 (d, *J*=1 Hz, 1H) ; 3.82 (s, 3H) ; 5.35 (m, 1H) ; 5.91 (s, 1H). ¹³C NMR (75MHz ; CD₂Cl₂) : δ 0.3 (SiMe₃) ; 15.7 ; 25.3 ; 31.3 ; 53.0 ; 55.0 ; 107.5 ; 108.5 ; 131.4 ; 145.7 ; 149.3 ; 164.0 ; 180.9 ; 199.2. IR (cm⁻¹) : 1652 ; 1604 ; 1536 ; 1213. TLC R_f : 0.30 (cyclohexane / EtOAc : 3 / 1). m.p. : 117-118°C (Et₂O / Pentane). Anal. Calcd for C₁₆H₂₂O₄Si : C, 62.75 ; H, 7.18. Found : C, 62.63 ; H, 7.39. $[\alpha]_D^{20} = -160.7$ (c = 0.42 ; CH₂Cl₂)

(-)-R-3-Methoxy-5,8a-dimethyl-8,8a-dihydro-7H-naphthalene-1,4,6-trione (3)

A solution of TBAF (1M in THF 1.633 mL ; 1.633 mmol; 1 eq) was slowly added to a THF (10 mL) solution of **3** (500 mg; 1.633 mmol ; 1 eq) at 0°C under argon. After 10 mn, aqueous ammonium chloride solution (20 mL) was added. The aqueous layer was extracted with EtOAc (3 x 15 mL). The combined organic phases were washed with water (20 mL) brine (20mL), dried over MgSO₄, and concentrated in vacuum to give crude (–)-**4**. This crude triketone may be purified by flash chromatography on demetalled silica gel (eluent cyclohexane / EtOAc : 4 / 1) or crystallised in Et₂O / pentane at – 20°C to afford pure (–)-**4** (363.15 mg ; 95% for chromatography or 313.46 mg ; 82% for crystallisation).

¹H NMR (300 MHz ; CDCl₃) : 1.51 (s, 3H) ; 2.03 (s, 3H) ; 2.29 (m, 2H) ; 2.61 (m, 2H) ; 3.88 (s, 3H) ; 5.99 (s, 1H). **¹³C NMR** (75MHz ; CDCl₃) : δ 12.8 ; 26.0 ; 29.4 ; 33.4 ; 49.3 ; 56.7 ; 109.4 ; 138.9 ; 146.5 ; 163.1 ; 185.1 ; 198.2 ; 198.4. **IR** (cm⁻¹) : 1656 ; 1638 ; 1603 ; 1154. **TLC Rf** : 0.21 (cyclohexane / CH₂Cl₂ / Acetone : 10 / 10 / 1). **m.p.** : 150°C (Et₂O / pentane) (m.p. racemic : 125°C from Et₂O / pentane) . **Anal.** Calcd for C₁₃H₁₄O₄ : C, 66.66 ; H, 6.02. Found : C, 66.41 ; H, 5.96. $[\alpha]_D^{20} = -63.5$ (c = 0.41 ; Acetone)

1,2,5-Trimethoxy-4-methyl-3-(toluene-4-sulfanyl)-benzene

Under an Argon atmosphere, to a suspension of 1.57 g magnesium turnings (64.59 mmol ; 1.12 eq) in 70 mL of anhydrous THF, a solution of **12** (15.00 g ; 57.48 mmol ; 1.00 eq) in 70 mL of anhydrous THF was added. After the initiation of Grignard, the mixture was stirred at room temperature for 4 hours. The Grignard solution was cooled down to 0°C and then transferred through double tipped needle by means of Argon pressure over a cooled (0°C) solution of (p-TolS)₂ (15.60 g ; 63.31 mmol ; 1.10 eq) in 120 mL of anhydrous THF with vigorous stirring. The resulting mixture was stirred overnight. Hydrolysis was performed, at 0°C, with 250 mL of saturated NH₄Cl solution. The aqueous layer was extracted with Et₂O (3 x 150 mL). The combined organic phases were washed with brine (250 mL), dried over MgSO₄, and concentrated in vacuum to give an orange oil. Flash chromatography on silica gel (cyclohexane / Et₂O ; 35 / 15) afforded 13.11 g (75%) of the title compound as an oil which crystallised on standing.

¹H NMR (300 MHz ; CDCl₃) : δ 2.21 (s, 3H) ; 2.25 (s, 3H) ; 3.7 (s, 3H) ; 3.83 (s, 3H) ; 3.89 (s, 3H) ; 6.59 (s, 1H) ; 6.96 (m, 4H) **¹³C NMR** (75MHz ; CDCl₃) : δ 13.4 ; 20.9 ; 56.1 ; 56.3 ; 61.2 ; 99.0 ; 123.6 ; 126.9 ; 127.8 ; 129.5 ; 134.5 ; 134.7 ; 144.8 ; 151.3 ; 154.3 **IR** (cm⁻¹) : 1589 ; 1316 ; 1218 ; 1195 ; 1025 ; 818. **TLC Rf** : 0.48 (cyclohexane / Et₂O ; 35 / 15). **m.p.** 78°C (hexane): . **Anal.** Calcd for C₁₇H₂₀O₃S. C, 67.12 ; H, 6.57. Found : C, 67.14 ; H, 6.62

(±)-1,2,5-Trimethoxy-4-methyl-3-(toluene-4-sulfinyl)-benzene (*rac*-13)

To a cold (0°C) solution of the above diaryl sulfure (500 mg ; 1.643 mmol ; 1.0 eq) in 6 mL of CH_2Cl_2 was added *m*-CPBA (72 %) (374 mg ; 1.610 mmol ; 0.98 eq of active O_2) in 9 mL of CH_2Cl_2 . After 1 h30 at 0°C , the organic phase was treated with 20 mL of a saturated $\text{Na}_2\text{S}_2\text{O}_3$ solution, saturated NaHCO_3 solution (20 mL), and brine (20 mL), dried over MgSO_4 and concentrated to give a white solid. Trituration of this solid in Et_2O (10 mL) afforded (±)-13 (525 mg ; 98%) as a white microcrystalline solid.

General procedure for Diels-Alder reactions in thermal conditions

To a solution of sulfinylquinone (0.75 mmol ; 1 eq) in 10 mL of dry CH_2Cl_2 at room temperature, -20°C , was added the diene (1.5 mmol ; 2 eq) under Argon. After completion of the reaction (see Table 1 for reaction times), the solvent was evaporated at reduced pressure (For entry 6 and 8, crude mixtures were stirred during several hours at room temperature until the pyrolytic elimination of the sulfinyl group was complete). The resulting cycloadduct was chromatographed on silica gel (see below for the different eluent)

General procedure for catalysed Diels-Alder reactions with Lewis acids

A solution of sulfinylquinone (0.75 mmol ; 1 eq) in 10 mL CH_2Cl_2 was added at -78°C to the appropriate Lewis acid (see Table 2 and 3 for conditions) under Argon, and the mixture was stirred for 10 mn. The appropriate diene (see Table 2 and 3 for conditions) was added and the resulting solution was stored in the fridge at -20°C or -40°C (dry ice / MeCN). After completion of the reaction (see Table 2 and 3 for reaction times) the mixture was quenched with cold aqueous NH_4Cl . The heterogeneous solution CH_2Cl_2 / aqueous NH_4Cl of diene 5 was stirred during several hours at room temperature until the pyrolytic elimination of the sulfinyl group was complete. After work up the CH_2Cl_2 was evaporated at reduced pressure. The resulting cycloadduct was chromatographed on silica gel (see below for the different eluent)

white solid

$^1\text{H NMR}$ (200 MHz ; CDCl_3) : δ 2.38 (s, 3H) ; 2.43 (s, 3H) ; 7.14 (d, $J = 8 \text{ Hz}$, 2H) ; 7.19-7.26 (m, 4H) ; 7.46 (d, $J = 8 \text{ Hz}$, 2H). $^{13}\text{C NMR}$ (75MHz ; CDCl_3) : δ 21.5 ; 21.6 ; 124.6 ; 127.6 ; 129.4 ; 130.2 ; 136.5 ; 140.5 ; 142.0 ; 144.6 **TLC** **Rf** : 0.67 (hexane / EtOAc ; 1 / 1).

Compound 18a

Pale yellow solid

¹H NMR (200 MHz ; CDCl₃) : δ 1.68 (d, *J*= 10.2 Hz, 1H) ; 1.85 (s, 3H) ; 2.25 (d, *J*= 10.2 Hz, 1H) ; 2.38 (s, 3H) ; 3.24 (s br, 1H) ; 3.32 (s, 3H) ; 4.06 (s br, 1H) ; 5.47 (s, 1H) ; 6.08 (dd, *J*= 2.6 Hz, *J*=5.4 Hz, 1H) ; 6.23 (dd, *J*= 2.6 Hz, *J*=5.4 Hz, 1H) ; 7.26 (d, *J*= 8 Hz, 2H) ; 7.50 (d, *J*= 8 Hz, 2H). **¹³C NMR** (75MHz ; CDCl₃) : δ 21.4 ; 25.6 ; 43.3 ; 49.8 ; 54.3 ; 55.9 ; 58.0 ; 81.1 ; 111.4 ; 126.3 (2 x CH) ; 129.5 (2 x CH) ; 136.6 ; 136.8 ; 140.2 ; 142.5 ; 163.8 ; 186.8 ; 199.6 **IR** (cm⁻¹) : 1692 ; 1656 ; 1602 ; 1366 ; 1240

; 1182 ; 1040 ; 818 ; 734. **m.p.** : 148-149°C (decomposition). **TLC Rf** : 0.48 (CH₂Cl₂ / EtOAc / Acetone : 40 / 5 / 1). **Anal.** Calcd for C₂₀H₁₈O₄S : C, 67.81 ; H, 5.08. Found : C, 67.85 ; H, 5.13.

[α]_D²⁰ = - 242 (c = 0.5 ; CH₂Cl₂)

Compound 18b

Pale yellow solid

¹H NMR (200 MHz ; CDCl₃) : δ 1.58 (d, *J*= 10 Hz, 1H) ; 2.07 (d, *J*= 10 Hz, 1H) ; 2.17 (s, 3H) ; 2.48 (s, 3H) ; 2.87 (s br, 1H) ; 3.27 (s br, 1H) ; 3.78 (s, 3H) ; 5.80 (dd, *J*= 3.0 Hz, *J*=5.6 Hz, 1H) ; 5.83 (s, 1H) ; 6.26 (dd, *J*= 3.0 Hz, *J*=5.6 Hz, 1H) ; 7.38 (d, *J*= 8 Hz, 2H) ; 7.44 (d, *J*= 8 Hz, 2H). **¹³C NMR** (75MHz ; CDCl₃) : δ 21.7 ; 24.8 ; 34.4 ; 43.2 ; 48.2 ; 50.9 ; 53.2 ; 56.5 ; 59.4 ; 79.2 ; 111.8 ; 126.7 ; 129.4 ; 135.5 ; 136.6 ; 141.0 ; 143.4 ; 164.8 ; 190.4 ; 200.5 **IR** (cm⁻¹) : 1681 ; 1648 ; 1600 ; 1453 ; 1360 ; 1242 ;

1137 ; 1068 ; 748. **m.p.** : 140-141°C (decomposition). **TLC Rf** : 0.29 (CH₂Cl₂ / EtOAc / Acetone : 40 / 5 / 1). **Anal.** Calcd for C₂₀H₁₈O₄S : C, 67.81 ; H, 5.08. Found : C, 67.86 ; H, 5.11.

[α]_D²⁰ = + 526 (c = 0.98 ; CH₂Cl₂)

Compound 19a

Pale yellow solid

¹H NMR (400 MHz ; C₆D₆) : δ 1.21 (s br, 3H) ; 1.43 (s br, 3H) ; 1.78 (s, 3H) ; 1.82 (d, *J*= 15.6 Hz, 1H) ; 1.88 (s, 3H) ; 2.05 (d, *J*= 17 Hz, 1H) ; 2.20 (d, *J*= 17 Hz, 1H) ; 2.64 (d, *J*= 15.6 Hz, 1H) ; 2.92 (s, 3H) ; 6.06 (s, 1H) ; 6.86 (d, *J*= 8 Hz, 2H) ; 7.20 7.26 (d, *J*= 8 Hz, 2H) **¹³C NMR** (75MHz ; CDCl₃) : δ 16.5 ; 18.6 ; 18.7 ; 21.6 ; 29.6 ; 44.9 ; 49.6 ; 56.2 ; 79.1 ; 112.9 ; 121.6 ; 122.5 ; 125.0 (2 x CH) ; 130.0 (2 x CH) ; 134.8 ; 142.9 ; 161.3 ; 186.5 ; 198.4. **IR** (cm⁻¹)

: 1670 ; 1605 ; 1453 ; 1360 ; 1243 ; 1195 ; 1054 ; 807. **m.p.** : 125-126°C (decomposition). **TLC Rf** : 0.17 (CH₂Cl₂ / Acetone : 40 / 1). **Anal.** Calcd for C₂₁H₂₄O₄S : C, 67.75 ; H, 6.45. Found : C, 67.72 ; H, 6.53. $[\alpha]_D^{20} = -208$ (c = 1.0 ; CH₂Cl₂)

Compound 20

Yellow oil. **¹H NMR** (200 MHz ; CD₂Cl₂) : δ 1.19 (s, 3H) ; 1.89 (s br, 3H) ; 1.92 (s br, 3H) ; 2.4-2.5 (m, 2H) ; 3.81 (s, 3H) ; 5.91 (s, 1H) ; 7.09 (s, 1H) **¹³C NMR** (75MHz ; CD₂Cl₂) : δ 16.7 ; 20.4 ; 25.6 ; 39.2 ; 44.5 ; 56.3 ; 109.3 ; 124.9 ; 131.6 ; 137.9 ; 140.0 ; 163.5 ; 179.3 ; 200.5 **TLC Rf** : 0.34 (CH₂Cl₂ / Acetone : 40 / 1).: **Anal.** Calcd for C₁₄H₁₆O₃ : C, 72.44 ; H, 6.89. Found : C, 72.61 ; H, 6.98. $[\alpha]_D^{20} = +13.0$ (c = 0.40 ; CHCl₃)

Compound 21

Pale yellow solid

¹H NMR (200 MHz ; CD₂Cl₂) : δ 0.28 (s, 9H) ; 1.69 (s, 3H) ; 2.02-2.14 (m, 1H) ; 2.38-2.42 (m, 1H) ; 2.36 (s, 3H) ; 3.35 (s, 3H) ; 5.12 (s br, 1H) ; 5.54-5.62 (m, 1H) ; 5.80 (s, 1H) ; 5.92-5.98 (m, 1H) ; 7.23 (d, *J* = 8 Hz, 2H) ; 7.55 (d, *J* = 8 Hz, 2H). **¹³C NMR** (75MHz ; CDCl₃) : δ 0.6 (SiMe₃) ; 18.4 ; 21.3 ; 39.2 ; 55.7 ; 69.9 ; 82.5 ; 109.1 ; 124.1 ; 126.4 (2 x CH) ; 129.2 (2 x CH) ; 129.9 ; 135.3 ; 141.5 ; 160.6 ; 184.4 ; 196.3. **IR** (cm⁻¹) : 1669 ; 1605 ; 1580 ; 1379 ; 1252 ; 1240 ; 1209 ; 1049 ; 832 ; 813. **m.p.**

: 53°C. **Anal.** Calcd for C₂₂H₂₈O₅SSi : C, 67.81 ; H, 5.08. Found : C, 67.86 ; H, 5.11. $[\alpha]_D^{20} = -131$ (c = 0.42 ; Acetone)

Abundance

m/z→

167 (10), 166 (100), 165 (11), 152 (7), 151 (79), 123 (22), 95 (6), 77 (9), 67 (7), 66 (5), 55 (7), 53 (7), 39 (5).

UD₅ : C₉H₉BrO₃
Mol. Wt.: 245,1
m/e: 244,0 (100,0%), 246,0 (98,3%), 245,0
(10,0%), 247,0 (9,8%), 248,0 (1,0%)

Abundance

Scan 5130 (72.959 min): UD5.D

246 (99), 245 (21), 244 (100), 243 (12), 231 (47), 229 (51), 203 (19), 201 (21), 67 (10), 65 (15), 53 (12).

UD₃ : C₁₀H₁₁BrO₃
Mol. Wt.: 259,1
m/e: 258,0 (100,0%), 260,0 (98,4%), 259,0
(11,1%), 261,0 (10,8%), 262,0 (1,1%)

Abundance

Scan 5123 (72.875 min): UD3.D

260 (97), 258 (100), 245 (51), 243 (57), 214 (42), 212 (42), 200 (22), 121 (22), 108 (19), 93 (21), 77 (27), 65 (19).

UD₈ : C₁₀H₁₁BrO₄
 Mol. Wt.: 275.1
 m/e: 274.0 (100.0%), 276.0 (98.7%),
 275.0 (11.1%), 277.0 (10.9%), 278.0
 (1.3%)

Abundance

276 (15), 274 (16), 248 (15), 246 (15), 233 (100), 232 (12), 231 (99), 205 (44), 203 (48), 124 (18), 109 (23), 77 (12), 69 (21), 67 (14), 53 (12).

UD₁₀ : C₉H₁₁BrO₃
 Mol. Wt.: 247.1
 m/e: 246.0 (100.0%), 248.0 (98.3%), 247.0
 (10.0%), 249.0 (9.8%), 250.0 (1.0%)

Abundance

232 (96), 231 (25), 230 (100), 229 (16), 217 (32), 215 (33), 189 (23), 187 (24), 159 (10), 79 (16), 53 (18), 51 (9).

262 (81), 261 (10), 260 (82), 248 (10), 247 (97), 246 (13), 245 (100), 219 (37), 217 (42), 204 (14), 202 (16), 138 (20), 123 (24), 77 (15).

320 (43), 304 (46), 303 (100), 288 (67), 273 (33), 272 (75), 258 (32), 257 (33), 228 (60), 167 (28), 166 (34), 151 (35), 136 (34), 121 (31), 105 (62), 91 (42), 77 (28).

Current Data Parameters
NAME daig.526
EXPNO 11
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050629
Time 5.37
INSTRUM spect
PROBHD 5 mm Dual 13C/
PULPROG zg30
TD 65536
ID CDCl3
SOLVENT CDCl3
NS 16
DS 2
SWH 6172.839 Hz
FIDRES 0.094190 Hz
AQ 5.3084660 sec
RG 322.5
DM 81.000 usec
DE 6.50 usec
TE 300.0 K
D1 1.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 8.60 usec
PL1 0.00 dB
SFO1 300.1318534 MHz

F2 - Processing parameters
SI 32768
SF 300.1300000 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 12.50 cm
FLP 11.000 ppm
F1 3301.43 Hz
F2P -1.000 ppm
F2 -300.13 Hz
PPMCK 0.60000 ppm/cm
HZCM 180.07800 Hz/cm

```

Current Data Parameters
NAME          da1g-526
EXPNO         10
PROCNO        1

F2 - Acquisition Parameters
Date_         20090629
Time          5.34
INSTRUM       spect
PROBHD        5 mm Dual 13C/
PULPROG       zgpg30
TD            65536
SOLVENT       CDCl3
NS            1024
DS            4
SWH           17985.611 Hz
FIDRES        0.274439 Hz
AQ            1.8219508 sec
RG            496.1
DM            27.800 usec
DE            6.50 usec
TE            300.0 K
D1            2.00000000 sec
d11           0.03000000 sec
d12           0.0002000 sec

===== CHANNEL f1 =====
NUC1          13C
P1            5.50 usec
PL1           -6.00 dB
SFO1          75.4752953 MHz

===== CHANNEL f2 =====
CEPRG2        waltz16
NUC2          1H
PCPD2         80.00 usec
PL2           0.00 dB
PL12          20.50 dB
PL13          20.50 dB
SFO2          300.1312005 MHz

F2 - Processing parameters
SI            32768
SF            75.4677490 MHz
WDW           EM
SSB           0
LB            1.00 Hz
GB            0
PC            1.40

1D NMR plot parameters
CX            20.00 cm
CY            19.50 cm
F1P           235.000 ppm
F1            16225.37 Hz
F2P           -5.000 ppm
F2            -377.34 Hz
FPCYC         11.00000 ppm/cm
HZCM          830.14526 Hz/cm

```

Current Data Parameters
 NAME dal3.526
 EXNO 10
 PROCNO 1

F2 - Acquisition Parameters

Date_ 20030629
 Time 7.06
 INSTRUM spect
 PROBD 5 mm Dual 13C/
 PULPROG zgpg30
 TD 65536
 SOVENT CDCl3
 NS 1024
 DS 4
 SWH 17985.611 Hz
 FIDRES 0.274439 Hz
 AQ 1.8219508 sec
 RG 456.1
 DW 27.800 usec
 DE 6.50 usec
 TE 300.0 K
 D1 2.00000000 sec
 d11 0.03000000 sec
 d12 0.00002000 sec

===== CHANNEL f1 =====
 NUC1 13C
 P1 5.50 usec
 PL1 -6.00 dB
 SFO1 75.4752953 MHz

===== CHANNEL f2 =====
 CPDPRG2 waltz16
 NUC2 1H
 PCPD2 80.00 usec
 PL2 0.00 dB
 PL12 20.50 dB
 PL13 20.50 dB
 SFO2 300.132005 MHz

F2 - Processing parameters
 SI 32768
 SF 75.4677490 MHz
 WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 1.40

1D NMR plot parameters
 CX 20.00 cm
 CY 12.50 cm
 F1P 215.000 ppm
 F1 16225.57 Hz
 F2P -5.000 ppm
 F2 -377.34 Hz
 PPMCK4 11.00000 ppm/cm
 HZCM 830.14326 Hz/cm

Current Data Parameters
NAME dalc45.317
EXPNO 10
PROCNO 1

F2 - Acquisition Parameters
Date_ 20030425
Time 0:58
INSTRUM spect
PROBHD 5 mm Dual 13C/
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 1024
DS 2
SWH 17985.611 Hz
FIDRES 0.274439 Hz
AQ 1.8219508 sec
RG 362
DM 27.800 usec
DE 6.50 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 5.50 usec
PL1 -6.00 dB
SFO1 75.4752953 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 80.00 usec
PL2 0.00 dB
PL12 20.50 dB
PL13 20.50 dB
SFO2 300.1312005 MHz

F2 - Processing parameters
SI 32768
SF 75.4677490 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 12.50 cm
FIP 215.000 ppm
F1 16225.57 Hz
F2 -5.000 ppm
F2 -377.34 Hz
PPMCM 11.00000 ppm/cm
HZCM 830.14526 Hz/cm

Current Data Parameters
 NAME dalc.508
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date 20050225
 Time 14.35
 INS120M spect
 PROCNO 5 mm QNP 1H/1
 PULPROG zg30
 ID 32768
 SOLVENT CDCl3
 NS 16
 DS 2
 SWE 3223.685 Hz
 FIDRES 0.250967 Hz
 AQ 1.9923444 sec
 RG 1024
 DM 60.800 usec
 DE 10.00 usec
 TE 300.0 K
 D1 2.00000000 sec
 D11 8.00 usec
 SFO1 400.1324710 MHz
 NUC1 1H
 P11 -3.00 dB

F2 - Processing parameters
 SI 16384
 SF 400.1300000 MHz
 KW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 21.47 cm
 FIP 9.000 ppm
 FI 9601.17 Hz
 F2P -1.000 ppm
 F2 -400.13 Hz
 PPMX 0.46586 ppm/cm
 HZCX 186.40352 Hz/cm

Current Data Parameters
NAME mko43.423
EXPNO 10
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040602
Time 21.48
INSTRUM spect
PROBHD 5 mm Dual 13C/
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 1024
DS 4
SWH 17985.611 Hz
AQ 0.274439 Hz
RG 1.8219508 sec
DE 322.5
DM 27.800 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 5.50 usec
PL1 -6.00 dB
SFO1 75.4752953 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 80.00 usec
PL2 0.00 dB
PL12 20.50 dB
PL13 20.50 dB
SFO2 300.1312005 MHz

F2 - Processing parameters
SI 32768
SF 75.4677490 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 12.50 cm
F1P 215.000 ppm
F1 16225.57 Hz
F2P -5.000 ppm
F2 -377.34 Hz
FPCOM 11.00000 ppm/cm
HZCM 830.14526 Hz/cm

Current Data Parameters
 NAME mkdrip.423
 EXENO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20040603
 Time 14.11

INSRUM spect
 PROBD 5 mm QNP 1H/1
 PULPROG zg30

TD 32768
 SOLVENT CDCl3
 NS 16
 DS 2

SWH 8223.685 Hz
 FIDRES 0.250967 Hz
 AQ 1.9923444 sec

RG 574.7
 DW 60.800 usec
 DE 10.00 usec
 TE 300.0 K

D1 2.00000000 sec
 P1 8.00 usec
 SFO1 400.1324710 MHz
 NUC1 1H
 PL1 -4.00 dB

F2 - Processing parameters
 SI 16384
 SF 400.1300032 MHz
 WDW EM
 SSB 0

LB 0.30 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 20.00 cm
 F1P 9.000 ppm
 F1 3601.17 Hz

F2P -1.000 ppm
 F2 -400.13 Hz
 PPMCM 0.50000 ppm/cm
 HZCM 200.06500 Hz/cm

Current Data Parameters
NAME dal.528
EXPNO 10
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050714
Time 7.23
INSTRUM spect
PROBHD 5 mm Dual 13C/
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 7168
DS 4
SWH 17985.611 Hz
FIDRES 0.27439 Hz
AQ 1.8219508 sec
RG 4096
DE 27.800 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00020000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 5.50 usec
PL1 -6.00 dB
SFO1 75.4752953 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 80.00 usec
PL2 0.00 dB
PL12 20.50 dB
PL13 20.50 dB
SFO2 300.1312005 MHz

F2 - Processing parameters
SI 32768
SF 75.4677490 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 12.50 cm
F1P 215.000 ppm
F1 16225.57 Hz
F2P -5.000 ppm
F2 -377.34 Hz
PPMCM 11.00000 ppm/cm
HZCM 830.14526 Hz/cm

Current Data Parameters
NAME calq.417
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040422
Time 16.14
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zg30
TD 32768
SOLVENT CDCl3
NS 16
DS 2
SWH 6223.685 Hz
FIDRES 0.250967 Hz
AQ 1.9923444 sec
RG 3149.4
DW 60.800 usec
DE 10.00 usec
TE 300.0 K
D1 2.00000000 sec
P1 8.00 usec
SFO1 400.1324710 MHz
NUC1 1H
PL1 -3.00 dB

F2 - Processing parameters
SI 16384
SF 400.1300000 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CH 20.00 cm
F1P 6.623 ppm
F1 2650.14 Hz
F2P 6.445 ppm
F2 2578.88 Hz
PPMCM 0.00891 ppm/cm
HSCM 3.56323 Hz/cm

Current Data Parameters
NAME dalc.417
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040422
Time 16.38
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zg30
TD 32768
SOLVENT CDCl₃
NS 16
DS 2
SWH 8223.685 Hz
FIDRES 0.250967 Hz
AQ 1.9923444 sec
RG 2298.8
DM 60.800 usec
DE 10.00 usec
TE 300.0 K
D1 2.0000000 sec
F1 8.00 usec
SFO1 400.1324710 MHz
NUC1 1H
PC1 -3.00 dB

F2 - Processing parameters
SI 16384
SF 400.1300000 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
F1P 6.788 ppm
F1 2716.07 Hz
F2P 6.479 ppm
F2 2592.54 Hz
FPMCM 0.01544 ppm/cm
HECM 6.17658 Hz/cm

Current Data Parameters
NAME dalc.508
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050225
Time 14.51
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zg30
TD 32768
SOLVENT CDCl₃
NS 16
DS 2
SWH 8223.685 Hz
FIDRES 0.250967 Hz
AQ 1.9923444 sec
RG 1024
DM 60.800 usec
DE 10.00 usec
TE 300.0 K
D1 2.0000000 sec
F1 8.00 usec
SFO1 400.1324710 MHz
NUC1 1H
PC1 -3.00 dB

F2 - Processing parameters
SI 16384
SF 400.1300000 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 21.47 cm
F1P 2.440 ppm
F1 976.25 Hz
F2P 2.292 ppm
F2 917.02 Hz
FPMCM 0.00690 ppm/cm
HECM 2.75927 Hz/cm

Current Data Parameters
 NAME mkqgrip.424
 EXPNO 6
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20040607
 Time 14.41
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zg30
 TD 32768
 SOLVENT CDCl₃
 NS 16
 DS 2
 SWH 5995.204 Hz
 FIDRES 0.182959 Hz
 AQ 2.7329011 sec
 RG 600
 DW 83.400 usec
 DE 10.00 usec
 TE 300.0 K
 D1 2.00000000 sec
 P1 8.00 usec
 SFO1 400.1323766 MHz
 NUC1 1H
 PL1 -3.00 dB

F2 - Processing parameters
 SI 16384
 SF 400.1300000 MHz
 WDW no
 SSB 0
 LB 0.00 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 20.00 cm
 F1P 2.580 ppm
 F1 1032.34 Hz
 F2P 2.480 ppm
 F2 992.32 Hz
 PPMCM 0.00500 ppm/cm
 HZCM 2.00065 Hz/cm

Current Data Parameters
 NAME mkqgrip.424
 EXPNO 3
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20040607
 Time 15.13
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zg30
 TD 32768
 SOLVENT CDCl₃
 NS 16
 DS 2
 SWH 5995.204 Hz
 FIDRES 0.182959 Hz
 AQ 2.7329011 sec
 RG 574.7
 DW 83.400 usec
 DE 10.00 usec
 TE 300.0 K
 D1 2.00000000 sec
 P1 8.00 usec
 SFO1 400.1323766 MHz
 NUC1 1H
 PL1 -3.00 dB

F2 - Processing parameters
 SI 16384
 SF 400.1300000 MHz
 WDW no
 SSB 0
 LB 0.00 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 20.00 cm
 F1P 2.580 ppm
 F1 1032.34 Hz
 F2P 2.480 ppm
 F2 992.32 Hz
 PPMCM 0.00500 ppm/cm
 HZCM 2.00065 Hz/cm

*** Current Data Parameters ***

NAME : DALTMSI
EXPNO : 547
PROCNO : 2

*** Acquisition Parameters ***

AUNM :
BFI : 200.1300000 MHz
NS : 34
O1 : 3200.00 Hz
PULPROG :

SFO1 : 200.1332000 MHz
SW : 14.9602 ppm
SW_h : 2994.012 Hz
TD : 16384
TE : 297 K

*** Processing Parameters ***

LB : 0.00 Hz
SF : 200.1323404 MHz
SI : 8192
SSB : 0.0000000
SW_p : 2994.019760

*** Aspect 3000 Parameters ***

NM : DALTMSI.547
O1 : 3200.00 Hz
DDATE : 1981 Jan 29
ATIME : 17:58:25

322 mJ
96%

1.5997
1.6947
2.1183
2.1275
2.3685

3.3528

5.1206

5.7159
5.7835
5.9186
5.9734

7.2152
7.5312
7.5732

Current Data Parameters

NAME	dal13.346
EXPNO	10
PROCNO	1

F2 - Acquisition Parameters

Date_	20031113
Time	20.16
INSTRUM	spect
PROBHD	5 mm Dual 13C/
PULPROG	zgpg30
TD	65536
SD	32768
CO213	1024
NS	4
DS	4
SWH	17985.611 Hz
FIDRES	0.274439 Hz
AQ	1.8219508 sec
RG	374.7
DM	27.800 usec
DE	6.50 usec
TE	300.0 K
D1	2.00000000 sec
d11	0.03000000 sec
d12	0.00002000 sec

===== CHANNEL f1 =====

NUC1	13C
P1	5.50 usec
PL1	-6.00 dB
SFO1	75.4752953 MHz

===== CHANNEL f2 =====

CPDPRG2	waltz16
NUC2	1H
PCPD2	80.00 usec
P12	0.00 dB
P112	20.50 dB
P113	20.50 dB
SFO2	300.1312005 MHz

F2 - Processing parameters

SF	32768
SE	75.4677490 MHz
WDM	EX
SSB	0
LB	1.00 Hz
GB	0
PC	1.40

1D NMR plot parameters

CX	20.00 cm
CY	12.50 cm
F1P	215.000 ppm
F1	16225.57 Hz
F2P	-5.000 ppm
F2	-377.34 Hz
PPHCKM	11.00000 ppm/cm
HZCM	830.14526 Hz/cm

data24.346

Current Data Parameters

NAME	dal13.346
EXPNO	10
PROCNO	1

F2 - Acquisition Parameters

Date_	20031113
Time	20.16
INSTRUM	spect
PROBHD	5 mm Dual 13C/
PULPROG	zgpg30
TD	65536
SD	32768
SOLVENT	CDCl3
NS	1024
DS	4
SWH	17985.611 Hz
FIDRES	0.274439 Hz
AQ	1.8219508 sec
RG	374.7
DM	27.800 usec
DE	6.50 usec
TE	300.0 K
D1	2.00000000 sec
d11	0.03000000 sec
d12	0.00002000 sec

===== CHANNEL f1 =====

NUC1	13C
P1	5.50 usec
PL1	-6.00 dB
SFO1	75.4752953 MHz

===== CHANNEL f2 =====

CPDPRG2	waltz16
NUC2	1H
PCPD2	80.00 usec
P2	0.00 dB
P12	20.50 dB
P13	20.50 dB
SFO2	300.1312005 MHz

F2 - Processing parameters

SF	32768
SE	75.4677490 MHz
WDW	EM
SSB	0
LB	1.00 Hz
GB	0
PC	1.40

1D NMR plot parameters

CX	20.00 cm
CY	12.50 cm
F1P	215.000 ppm
F1	16225.57 Hz
F2P	-5.000 ppm
F2	-377.34 Hz
PPHMM	11.00000 ppm/cm
HZCM	830.14526 Hz/cm

*** Current Data Parameters ***
 NAME : DALBUT3
 EXPNO : 605
 PROCNO : 2
 *** Acquisition Parameters ***
 AUNM :
 BFI : 200.1300000 MHz
 NS : 63
 OI : 3700.00 Hz
 PULPROG :
 SFO1 : 200.1337000 MHz
 SW : 14.9601 ppm
 SW_h : 2994.012 Hz
 TD : 16384
 TE : 297 K
 *** Processing Parameters ***
 LB : 0.00 Hz
 SF : 200.1337275 MHz
 SI : 8192
 SSB : 0.0000000
 SW_p : 2994.0119760
 *** Aspect 3000 Parameters ***
 NM : DALBUT3.605
 OI : 3700.00 Hz
 DDATE : 1981/Apr/10
 ATIME : 20:02:51

$$\delta = +0,05 \text{ (c=0,4 / wt\%)} \\ [\alpha]_D = +13^\circ$$

Current Data Parameters
NAME dal.607
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060213
Time 17.26
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zg30
TD 32768
SOLVENT C6D6
NS 16
DS 2
SWH 8223.685 Hz
FIDRES 0.250967 Hz
AQ 1.9923444 sec
RG 456.1
DM 60.800 usec
DE 10.00 usec
TE 300.0 K
D1 2.00000000 sec
P1 8.00 usec
SFO1 400.1324710 MHz
NUC1 1H
PL1 -3.00 dB

F2 - Processing parameters
SI 16384
SF 400.1300000 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
FLP 9.000 ppm
F1 3601.17 Hz
F2P -1.000 ppm
F2 -400.13 Hz
PRCM 0.50000 ppm/cm
HZCM 200.06500 Hz/cm

Current Data Parameters
 NAME dalc2.347
 EXPNO 10
 PROCNO 1

F2 - Acquisition Parameters

Date_ 20031119
 Time 23.31
 INSTRUM spect
 PROBD 5 mm Dual 13C/
 PULPROG zgpg30
 TD 65536
 SOLVENT CDCl3
 NS 1024
 DS 4
 SWH 17985.611 Hz
 FIDRES 0.274439 Hz
 AQ 1.8219508 sec
 RG 362
 DW 27.800 usec
 DE 6.50 usec
 TE 300.0 K
 D1 2.00000000 sec
 d11 0.03000000 sec
 d12 0.0002000 sec

===== CHANNEL f1 =====
 NUC1 13C
 P1 5.50 usec
 PL1 -6.00 dB
 SFO1 75.4752953 MHz

===== CHANNEL f2 =====
 CPDPRG2 waltz16
 NUC2 1H
 PCPD2 80.00 usec
 PL2 0.00 dB
 PL12 20.50 dB
 PL13 20.50 dB
 SFO2 300.1312005 MHz

F2 - Processing parameters
 SI 32768
 SF 75.4677490 MHz
 WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 1.40

1D NMR plot parameters
 CX 20.00 cm
 CY 12.50 cm
 F1P 215.000 ppm
 F1 16225.57 Hz
 F2P -5.000 ppm
 F2 -377.34 Hz
 PPMCM 11.00000 ppm/cm
 HZCM 830.14526 Hz/cm

16.566
 18.617
 18.718
 21.587
 29.621
 44.922
 49.578
 56.160
 76.580
 77.003
 77.427
 79.075
 112.919
 121.583
 122.560
 125.033
 129.984
 134.803
 142.922
 161.335
 186.566
 198.443

Current Data Parameters
 NAME dalsu.609
 EXPNO 13
 PROCNO 1

F2 - Acquisition Parameters

Date_ 20060302
 Time 3.51
 INSTRUM spect
 PROBHD 5 mm Dual 13C/
 PULPROG zg30
 TD 65536
 SOLVENT CDCl3
 NS 16
 DS 2
 SWH 6172.839 Hz
 FIDRES 0.094190 Hz
 AQ 5.3084660 sec
 RG 181
 DW 81.000 usec
 DE 6.50 usec
 TE 300.0 K
 D1 1.00000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 8.60 usec
 PL1 0.00 dB
 SFO1 300.1318534 MHz

F2 - Processing parameters
 SI 32768
 SF 300.1300000 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 20.00 cm
 CY 12.50 cm
 FLIP 11.000 ppm
 F1 3301.43 Hz
 F2 -1.000 ppm
 F2 -300.13 Hz
 PPMCM 0.60000 ppm/cm
 HZCM 180.07800 Hz/cm

Current Data Parameters
NAME dalsu.609
EXPNO 12
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060302
Time 3.48
INSTRUM spect
PROBHD 5 mm Dual 13C/
PULPROG zgpg30
TD 65536
SOLVENT CDC13
NS 1024
DS 4
SWH 17985.611 Hz
FIDRES 0.274439 Hz
AQ 1.8219508 sec
RG 5160.6
DM 27.800 usec
DE 6.50 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 5.50 usec
PL1 -6.00 dB
SFO1 75.4752953 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 80.00 usec
PL2 0.00 dB
PL12 20.50 dB
PL13 20.50 dB
SFO2 300.1312005 MHz

F2 - Processing parameters
SI 32768
SF 75.4677490 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 12.50 cm
F1P 215.000 ppm
F1 16225.57 Hz
F2P -5.000 ppm
F2 -377.34 Hz
PPMCM 11.00000 ppm/cm
HZCM 830.14526 Hz/cm

Report on samples from Dr. Gilles Hanquet and Don Antoine Lanfranchi
Measurements by Dr. Xiaolin Cao. Calculations, analysis and report by Prof. Tess Freedman,
Department of Chemistry, Syracuse University

The structures are represented below:

Sample 1

$[\alpha]_D = -64$ (c=0.4, acetone)

Sample 2

$[\alpha]_D = -114$ (c=0.4, chloroform)

Experimental:

Samples were dissolved in CDCl_3 and placed in a 94- μm pathlength cell with BaF_2 windows. IR and VCD spectra were recorded on a modified ChiralIR VCD spectrometer (BioTools, Inc., Wauconda, IL) at 4- cm^{-1} resolution, with 11 h collection and instrument optimized at 1400 cm^{-1} . The VCD spectrum of the corresponding racemic sample was used for the VCD baseline, and the IR spectrum of the solvent has been subtracted from the sample spectrum. The experimental spectra are displayed in Figures 1 and 2.

Calculations: The *R*-configuration of each sample was generated with HyperChem. The geometries were optimized at the DFT (B3LYP functional/6-31G(d) basis set) level with Gaussian 03, and the vibrational frequencies and IR and VCD intensities were calculated at the same level. The optimized geometries are displayed in Figure 3. For both Sample 1 and Sample 2, the conformation with the methoxy group rotated 180° was calculated to lie 2.5 Kcal/mol higher in energy than the conformer shown in Figure 3, and since these latter conformers would not be significantly populated in solution at room temperature, they were not considered further. The rest of the molecule is rigid. The calculated frequencies were scaled by 0.97 and the calculated intensities were converted to Lorentzian bands with 6- cm^{-1} half-width for comparison to experiment.

The experimental VCD and IR spectra for Samples 1 and 2 are compared with the calculations in Figures 4 and 5, respectively. The excellent agreement between experiment and calculation for the *R*-configurations establishes the configuration of both samples as *R*-(-).

Figure 1. IR (lower frame) and VCD (upper frame) spectra of Sample 1, 2.8 mg/100 μL CDCl_3 , 94- μm pathlength BaF_2 cell, 11-h collection at 4- cm^{-1} resolution, with instrument optimized at 1400 cm^{-1} . IR spectrum is solvent subtracted. VCD of racemic sample has been used for the VCD baseline.

Figure 2. IR (lower frame) and VCD (upper frame) spectra of Sample 2, 1.6 mg/100 μL CDCl_3 , 94- μm pathlength BaF_2 cell, 11-h collection at 4- cm^{-1} resolution, with instrument optimized at 1400 cm^{-1} . IR spectrum is solvent subtracted. VCD of racemic sample has been used for the VCD baseline.

Figure 3. Optimized geometries for Samples 1 and 2 (Gaussian 03, DFT with B3LYP functional and 6-31G(d) basis set. Calculations required ~9h for geometry optimization and ~9h for frequency calculation on Pentium 4 PC.

Figure 4. IR (lower frame) and VCD (upper frame) for Hanquet Sample 1 (right axes) compared to calculated spectra for *R*-enantiomer (left axes).

Figure 5. IR (lower frame) and VCD (upper frame) for Hanquet Sample 2 (right axes) compared to calculated spectra for *R*-enantiomer (left axes).

Crystallographic data of **18a**

Summary of Data CCDC 601300

Authors: D.A.Lanfranchi,G.Hanquet

Journal: J.Org.Chem. (0035)

Formula: C₂₀ H₂₀ O₄ S₁

Unit cell parameters: a 7.54300(10) b 10.5080(2) c 21.9210(4)
space group P212121

Formula ; C₂₀ H₂₀ O₄ S
Formula weight ; 356.42
Crystal system ; orthorhombic
Space group ; P 21 21 21
a (Å) ; 7.54300(10)
b (Å) ; 10.5080(2)
c (Å) ; 21.9210(4)
alpha (°) ; 90.00
beta (°) ; 90.00
gamma (°) ; 90.00
V (Å³) ; 1737.50(5)
Z ; 4
Density (g cm⁻³) ; 1.363
mu (Mo Kalpha) (mm⁻¹) ; 0.208
F(000) ; 752

Data collection
Temperature (K) ; 173(2)
Radiation (Å) ; MoK α - 0.71073
Theta min - max ; 2.69 - 30.02
Dataset[h, k, l] ; -10/10, -14/14, -30/30
Tot., Uniq. Data, R(int) ; 5070, 4385, 0.0000
Observed data ; >2sigma(I)

Refinement
Nreflections, Nparameters ; 5070, 226
R₂, R₁, wR₂, wR₁, Goof ; 0.0480, 0.0372, 0.0937, 0.0880, 1.021
P = (Fo² + 2Fc²)/3 ; w=1/[\s²(Fo²)+(0.0472P)²+0.2687P]
Max. and Av. Shift/Error ; 0.002, 0.000
Flack x ; -0.02(6)
Min, Max. Resd Dens. (e/Å³) ; -0.277, 0.208