Supporting information for

Asymmetric Hetero Diels-Alder Reactions of *N*-Sulfinyl Dienophiles Using Chiral Bis(oxazoline)-Copper(II) and -Zinc(II) Triflates

Annette Bayer[†], Molla Mellese Endeshaw,[‡] Odd R. Gautun^{‡,*}

[†]Department of Chemistry, University of Tromsø, NO-9037 Tromsø, Norway.

[‡] Department of Chemistry, Norwegian University of Science and Technology (NTNU), NO-7491 Trondheim, Norway.

Fax: +47 73 59 42 56

E-mail: odd.gautun@chem.ntnu.no

TABLE OF CONTENTS

General	S2
2-Amino- <i>N</i> -benzyloxycarbonyl-1-propanoic Acid Methyl Ester (15) from <i>cis</i> - 11a .	S2
(<i>S</i>)-2-Amino- <i>N</i> -benzyloxycarbonyl-1-propanoic Acid Methyl Ester (15) from (<i>S</i>)- <i>N</i> -Benzyloxycarbonyl-alanine.	S3
2-Amino- <i>N</i> -tosyl-1-propanol (16) from <i>cis</i> - 11b .	S4
(<i>S</i>)-2-Amino- <i>N</i> -tosyl-1-propanol (16) from (<i>S</i>)- <i>N</i> -Tosylalanine.	S5
(2 <i>E</i>)-5-Amino-1-benzyloxy- <i>N</i> -tosyl-2-hexadiene (17) and (2 <i>E</i>)-5-Amino-6-benzyloxy- <i>N</i> -tosyl-2-hexadiene (18) from <i>cis</i> - 13 and <i>cis</i> - 14 .	S5
References	S7

General. Benzyl *N*-sulfinylcarbamate (**1a**)³ and *N*-Sulfinyl-4-toluenesulfonamide (**1b**)⁴ were prepared according to the literature and stored in the freezer as solutions in dry CH₂Cl₂. (2*E*,4*E*)-1-benzyloxy-2,4-hexadiene (**10**),¹⁴ 2,6-bis[4'-(*R*)-phenyloxazolin-2'-yl]-pyridine (**4a**),⁵ 2,6-bis[4'-(*S*)-*tert*-butyloxazolin-2'-yl]pyridine (**4b**)⁵ and 2,2'-isopropylidenebis[(4*S*)-4-*tert*-butyl-2-oxazoline] (**3b**)⁵ were prepared as described in the literature. (*S*)- (**ent-3a**) and (*R*)-2,2'-isopropylidenebis(4-phenyl-2-oxazoline) (**3a**) were prepared by a modification of the method of Evans *et al.*⁵ Solvents were dried according to standard procedures.³³ Reaction products were purified by flash chromatography (FC) using silica (35–70 micron). TLC was performed on silica gel plates and visualized with UV light and a phosphomolybdic acid in ethanol solution. ¹H and ¹³C NMR spectra of CDCl₃ solutions were recorded at 400 MHz and 100 MHz, respectively, and chemical shifts are in ppm and referenced to TMS via the lock signal of the solvent. ¹H NMR signals were assigned by 2D correlation techniques. EI mass spectra were recorded at 70 eV with a direct inlet. HPLC was performed using a 4.6 mm x 25 cm Daicel Chiralpak AD or Chiracel OJ columns. Melting points are reported uncorrected.

2-Amino-*N*-benzyloxycarbonyl-1-propanoic Acid Methyl Ester (**15**)³⁰ from *cis*-11a.

Thiazine oxide *cis*-**11a** (47.8 mg, 0.18 mmol, 77% ee) was dissolved in THF and cooled to –60 °C, before phenylmagnesium bromide in Et₂O (3M, 80 μl, 0.24 mmol) was added. After 30 min, the reaction mixture was quenched with saturated aqueous NH₄Cl (3 ml), allowed to warm to

room temperature and extracted with Et₂O (2 × 3 ml). The combined organics were washed with brine (1 × 2 ml), dried over MgSO₄ and concentrated. The crude product was dissolved in dry CH₂Cl₂ (7 ml) and methanolic NaOH (2.5 M, 2 ml) was added. The solution was cooled to –78 °C and ozone was passed through for 2 h until a pale blue colour persisted. The ozone was removed by passage of oxygen for 10 min, before the reaction mixture was allowed to warm to room temperature and stirred over night. CH₂Cl₂ (5 ml) and water (5 ml) were added and the aqueous layer was extracted with CH₂Cl₂ (3 × 3 ml). The combined organics were dried over MgSO₄ and concentrated. FC (Et₂O/pentane 30/70 to 60/40) of the crude product yielded a clear oil (13.8 mg, 0.058 mmol, 32%, 75% ee). A similar experiment with thiazine oxide *cis*-**12a** gave 37% of **15**. Analytical data for **15**: HPLC (Chiralcel OJ, *i*PrOH/hexane 20/80, 0.5 ml min^{–1}, 254 nm) *t*_R 26.1 (*S*) and 29.7 (*R*) min; ¹H NMR δ 7.39–7.28 (5H, m, Ph), 7.33 (1H, br d, *J* = 5.9 Hz, NH), 5.11 (1H, AB, *J* = 12.2 Hz, Bn), 5.09 (1H, AB, *J* = 12.2 Hz, Bn), 4.39 (1H, br pent, 2-H), 3.74 (3H, s, Me), 1.40 (3H, d, *J* = 7.0 Hz, 3-H); ¹³C NMR δ 173.6 (C=O), 155.7 (C=O), 136.4 (C), 128.6 (CH), 128.23 (CH), 128.18 (CH), 67.0 (CH₂), 52.5 (CH₃), 49.7 (CH), 18.7 (CH₃); IR (neat) 3337 (s), 3031 (w), 2951 (m), 1719 (s); EIMS *m/z* (relative intensity) 237 (*M*⁺, 0.8), 91 (100).

(*S*)-2-Amino-*N*-benzyloxycarbonyl-1-propanoic Acid Methyl Ester (15**)³⁰ from (*S*)-*N*-Benzyloxycarbonyl-alanine.** (*S*)-*N*-Benzyloxycarbonylalanine (0.46 g, 2.0 mmol) was dissolved in DMF (2 ml) and cooled to 0 °C. K₂CO₃ (0.30 g, 2.2 mmol) was added and the resulting suspension was stirred for 10 min at 0 °C before methyl iodide (249 μl, 4.0 mmol) was

added. The reaction was monitored by TLC (Et₂O/pentane 80/20), and no starting material was detected after 30 min at 0 °C and 1 h at room temperature. The reaction mixture was filtered. The filtrate was diluted with EtOAc (10 ml), washed with H₂O (3 ml) and brine (3 × 3 ml), dried over MgSO₄ and concentrated. FC (Et₂O/pentane 75/25 to 90/10) of the crude product provided a white solid (90%). Mp 46–47 °C (lit.^{30b} 45–46 °C). >98% ee (HPLC, see (*R*)-**15**).

2-Amino-N-tosyl-1-propanol (16**)³² from *cis*-**11b**.** Thiazine oxide *cis*-**11b** (34.0 mg, 0.12 mmol, 78% ee) was dissolved in MeOH (3 ml), stirred at room temperature for 30 min and then cooled to –78 °C. Ozone was passed through the solution for 45 min until a pale blue colour persisted. Excess ozone was removed by passage of oxygen for 10 min, before NaBH₄ (0.30 g, 8 mmol) was added. The resulting solution was allowed to warm to room temperature and three portions of NaBH₄ (3 × 0.30 g, 24 mmol) were added over a period of 2 h. The reaction mixture was concentrated. The residue was stirred with EtOAc (4 ml) and H₂O (4 ml) for 30 min before the layers were separated and the aqueous layer was extracted with EtOAc (3 × 3 ml). The combined organics were dried over MgSO₄ and concentrated. FC (Et₂O/pentane 90/10) of the crude product yielded a clear oil (15.3 mg, 0.067 mmol, 56%, 75% ee). A similar experiment with thiazine oxide *cis*-**12b** (97% ee) gave 82% of **16** (95% ee). Analytical data for **16**: HPLC (Chiralcel AD, *i*PrOH/hexane 20/80, 0.5 ml min^{–1}, 254 nm) *t*_R 14.3 (*S*) and 21.2 (*R*) min; ¹H NMR δ 7.77 (1H, d, *J* = 8.4 Hz, Ts), 7.29 (2H, d, *J* = 8.4 Hz, Ts), 5.30 (1H, d, *J* = 7.3 Hz, NH), 3.56–3.53 (1H, m, 1-H), 3.43–3.32 (2H, m, 1/2-H), 2.65 (1H, br s, OH), 2.41 (3H, s, Me), 0.99

(3H, d, $J = 6.6$ Hz, 3-H); ^{13}C NMR δ 143.6 (C), 137.6 (C), 129.8 (CH), 127.2 (CH), 66.3 (CH₂), 51.5 (CH), 21.6 (CH₃), 17.7 (CH₃); IR (neat) 3491 (s), 3285 (s), 2929 (s), 1598 (m); EIMS m/z (relative intensity) 229 (M^+ , 0.1), 91 (100).

(*S*)-2-Amino-*N*-tosyl-1-propanol (16)³² from (*S*)-*N*-Tosylalanine. (*S*)-*N*-Tosylalanine was reduced with $\text{NaBH}_4\text{-I}_2$ according to a procedure described by McKennon *et al.*^{5d} FC (Et₂O/pentane 90/10) yielded (*S*)-*N*-Ts-alaninol (57%) as a white solid. Mp 61–63 °C (lit. 58–60 °C).^{5d} > 99% ee (HPLC, see (*R*)-**16**).

(*2E*)-5-Amino-1-benzyloxy-*N*-tosyl-2-hexadiene (17) and (*2E*)-5-Amino-6-benzyloxy-*N*-tosyl-2-hexadiene (18) from *cis*-13** and *cis*-**14**.** A mixture (2.2 : 1) of thiazine oxide *cis*-**13** and *cis*-**14** (295 mg, 0.73 mmol) was stirred with aqueous NaOH (5% w/v, 4 ml) at 95 °C for 2 h before aqueous HCl (5% w/v, 12 ml) was added at room temperature. The resulting suspension was extracted with CH_2Cl_2 (3 × 3 ml). The combined organics were dried over MgSO_4 and

concentrated. The crude contained a mixture (2.2 : 1) of the amino alcohols **17** and **18**. FC (EtOAc/pentane 30/70) separated **17** (99 mg, 0.27 mmol, 38%) and **18** (40 mg, 0.11 mmol, 15%), both as colorless, viscous oils. A similar experiment with a mixture (8.3 : 1) of thiazine oxide *trans*-**13** and *trans*-**14** (98 mg, 0.24 mmol) gave **17** (50 mg, 0.14 mmol, 58%) as the major product and **18** (5.2 mg, 0.0144 mmol, 6%) as the minor isomer. Analytical data for **17**: ^1H NMR δ 7.74 (2H, d, J = 8.1 Hz, Ts), 7.36–7.24 (7H, m, Ts/Ph), 5.56 (1H, tAB, J = 15.4, 5.9 Hz, 2-H), 5.45 (1H, tAB, J = 15.4, 7.0 Hz, 3-H), 4.72 (1H, br d, J = 5.1 Hz, NH), 4.46 (2H, s, Bn), 3.88 (2H, d, J = 5.9 Hz, 1-H), 3.35 (1H, sept, J = 6.6 Hz, 5-H), 2.39 (3H, s, Me), 2.11 (2H, t, J = 6.6 Hz, 4-H), 1.05 (3H, d, J = 6.6 Hz, 6-H); ^{13}C NMR δ 143.3, 138.3, 138.1, 130.7, 129.7, 128.6, 128.5, 127.8, 127.7, 127.2, 71.7, 70.0, 49.0, 39.5, 21.1, 20.9; IR (neat) 3500 (w), 3300 (s), 3030 (m), 1593 (m), 1490 (m); EIMS m/z (relative intensity) 359 (M^+ , 0.02), 288 (2), 252 (31), 198 (81), 155 (67), 91 (100); Anal. Calcd for $\text{C}_{20}\text{H}_{25}\text{NO}_3\text{S}$: C, 66.82; H, 7.01; N, 3.90. Found: C, 66.68; H, 7.26; N, 3.71.

Analytical data for **18**: ^1H NMR δ 7.71 (2H, d, J = 8.1 Hz, Ts), 7.35–7.22 (7H, m, Ts/Ph), 5.42–5.33 (1H, m, 2-H), 5.12–5.01 (1H, m, 3-H), 4.75 (1H, br d, J = 7.3 Hz, NH), 4.38 (2H, s, Bn), 3.42 (1H, dd, J = 8.8, 3.7 Hz, 6-H), 3.34–3.25 (2H, m, 5-H/6-H), 2.41 (3H, s, Me), 2.22–2.11 (2H, m, 4-H), 1.54 (3H, d, J = 6.2 Hz, 1-H); ^{13}C NMR δ 143.3, 137.9, 129.6, 129.3, 128.5, 127.8, 127.7, 127.2, 126.0, 73.2, 71.0, 53.2, 35.4, 21.6, 18.0; IR (neat) 3300 (s), 3030 (m), 1593 (w), 1490 (w); EIMS m/z (relative intensity) 359 (M^+ , 0.03), 304 (6), 262 (4), 238 (5), 155 (20), 91 (100); Anal. Calcd for $\text{C}_{20}\text{H}_{25}\text{NO}_3\text{S}$: C, 66.82; H, 7.01; N, 3.90. Found: C, 66.84; H, 7.27; N, 3.73.

References

- (3) Garigipati, R.S.; Freyer, A. J.; Whittle, R. R.; Weinreb, S. M. *J. Am. Chem. Soc.* **1984**, *106*, 7861-7867.
- (4) Kresze, G.; Wucherpfenning, W. *Angew. Chem. Int. Ed. Engl.* **1967**, *6*, 149-167.
- (5) For the preparation of bis(oxazolinyl)pyridines, see: (a) Davies, I.; Gerena, L.; Lu, N.; Larsen, R. D.; Reider, P. *J. Org. Chem.* **1996**, *61*, 9629-9630; For the preparation of the bis(oxazolines), **3a**, see: (b) Corey, E. J.; Imai, N.; Zhang, H.-Y. *J. Am. Chem. Soc.* **1991**, *113*, 728-729; **3b**, see: (c) Evans, D. A.; Peterson, G. S.; Johnson, J. S.; Barnes, D. M.; Campos, K. R.; Woerpel, K. A. *J. Org. Chem.* **1998**, *63*, 4551-4544; For reduction of the amino acids, see: (d) McKennon, M. J.; Meyers, A. I.; Dauz, K.; Schwarm, M. *J. Org. Chem.* **1993**, *58*, 3568-3571.
- (14) Li, Z.-H.; Wang, T.-S.; Yao, E.-Y.; Gao, Z.-H. *Chinese J. Chem.* **1990**, *3*, 265-270.
- (30) (a) Kim, S.; Lee, J. I.; Kim, Y. C. *J. Org. Chem.* **1985**, *50*, 560-565; (b) Dhaon, M.K.; Olsen, R. K.; Ramasamy, K. *J. Org. Chem.* **1982**, *47*, 1962-1965; (c) Huneck, S.; Porzel, A., *Z. Naturforsch. B: Chem. Sci.* **1994**, *49*, 569-575.
- (32) Abiko, A.; Masamune, S. *Tetrahedron Lett.* **1992**, *33*, 5517-5518.
- (33) Armarego, W. L. F.; Perrin, D. D. *Purification of Laboratory Chemicals*, Butterworth Heinemann, 4. Ed., Oxford, 1996.