Online Supplemental Material

Castro-Schilo, Grimm, & Widaman

Demonstration of the Column Rank of Lambda

Using matrix algebra, we can decompose the covariances of multitrait-multimethod (MTMM) manifest variables into the parameters in the model. The covariance matrix, Σ , for the trait-method units, Y_{tm} (i.e., the manifest variables) is the expected value of the second central moment $Y_{tm}Y'_{tm}$, and gives rise to the covariance structure model,

$$\Sigma = \Lambda_t \Psi_t \Lambda'_t + \Lambda_m \Psi_m \Lambda'_m + \Theta$$

We can also express the covariance structure model in a traditional factor analytic way,

$$\Sigma = \Lambda \Psi \Lambda' + \Theta$$

by acknowledging that Λ and Ψ contain trait and method components. This covariance structure model corresponds to that of the Correlated Trait-Correlated Method (CT-CM) model for MTMM data.

The MTMM literature (Grayson & Marsh, 1994; Kenny & Kashy, 1992) has established one condition that leads to empirical underidentification of the CT-CM model: the case in which columns of Λ have equal entries. This is the case of equal within-factor loadings in the trait or method factors. Specifically, we can consider a CT-CM model with 3 Traits and 3 Methods (3x3) measured with 9 manifest variables, for which the Λ matrix would have 9 rows and 6 columns. For example,

$$\Lambda = \begin{bmatrix} 0.50 & 0.00 & 0.00 & 0.60 & 0.00 & 0.00 \\ 0.00 & 0.70 & 0.00 & 0.60 & 0.00 & 0.00 \\ 0.00 & 0.00 & 0.40 & 0.60 & 0.00 & 0.00 \\ 0.50 & 0.00 & 0.00 & 0.00 & 0.30 & 0.00 \\ 0.00 & 0.70 & 0.00 & 0.00 & 0.30 & 0.00 \\ 0.00 & 0.00 & 0.40 & 0.00 & 0.30 & 0.00 \\ 0.50 & 0.00 & 0.00 & 0.00 & 0.00 & 0.40 \\ 0.00 & 0.70 & 0.00 & 0.00 & 0.00 & 0.40 \\ 0.00 & 0.00 & 0.40 & 0.00 & 0.00 & 0.40 \end{bmatrix}$$

where the first 3 columns correspond to Trait factors and the last 3 to Method factors. Notice the non-zero factor loadings are equal for each column. In other words, this matrix of factor loadings conforms to the condition of empirical underidentification that Grayson & Marsh (1994) and Kenny and Kashy (1992) described. Indeed, we can show this matrix is of deficient column rank, which means there is a dependency in its columns and therefore, it is singular and cannot be inverted. To show this, we must recall that the rank of a matrix, in this case Λ , equals the rank of $\Lambda'\Lambda$. We also call to mind that a singular matrix has a determinant of zero. Thus, premultiplying the factor loading matrix by its transpose and getting the determinant of the resulting squared matrix shows its column rank deficiency,

$$\Lambda'\Lambda = \begin{bmatrix} 0.75 & 0.00 & 0.00 & 0.30 & 0.15 & 0.20 \\ 0.00 & 1.47 & 0.00 & 0.42 & 0.21 & 0.28 \\ 0.00 & 0.00 & 0.48 & 0.24 & 0.12 & 0.16 \\ 0.30 & 0.42 & 0.24 & 1.08 & 0.00 & 0.00 \\ 0.15 & 0.21 & 0.12 & 0.00 & 0.27 & 0.00 \\ 0.20 & 0.28 & 0.16 & 0.00 & 0.00 & 0.48 \end{bmatrix}$$

$$|\Lambda'\Lambda| = 0$$

As noted before, Grayson and Marsh (1994) provided a mathematical proof for the general case.

Importantly, the covariance structure of the Fully Augmented Correlated-Trait Correlated-Method (FACT-CM) model is identical to that of the CT-CM model. The key difference across models is in the dimensions of Λ . The factor loading matrix in a 3x3 FACT-CM model would have 15 rows by 6 columns. We can extend the previous Λ matrix with equal factor loadings in each column to this case,

$$\Lambda_{\text{FACT-CM}} = \begin{bmatrix} 0.50 & 0.00 & 0.00 & 0.60 & 0.00 & 0.00 \\ 0.00 & 0.70 & 0.00 & 0.60 & 0.00 & 0.00 \\ 0.00 & 0.00 & 0.40 & 0.60 & 0.00 & 0.00 \\ 0.50 & 0.00 & 0.00 & 0.00 & 0.30 & 0.00 \\ 0.00 & 0.70 & 0.00 & 0.00 & 0.30 & 0.00 \\ 0.00 & 0.00 & 0.40 & 0.00 & 0.30 & 0.00 \\ 0.50 & 0.00 & 0.00 & 0.00 & 0.00 & 0.40 \\ 0.00 & 0.70 & 0.00 & 0.00 & 0.00 & 0.40 \\ 0.00 & 0.00 & 0.40 & 0.00 & 0.00 & 0.40 \\ 0.50 & 0.00 & 0.00 & 0.00 & 0.00 & 0.00 \\ 0.00 & 0.70 & 0.00 & 0.00 & 0.00 & 0.00 \\ 0.00 & 0.00 & 0.40 & 0.00 & 0.00 & 0.00 \\ 0.00 & 0.00 & 0.00 & 0.60 & 0.00 & 0.00 \\ 0.00 & 0.00 & 0.00 & 0.00 & 0.30 & 0.00 \\ 0.00 & 0.00 & 0.00 & 0.00 & 0.00 & 0.40 \end{bmatrix}$$

where, again, the first 3 columns correspond to Trait factors and the last 3 to Method factors, and the additional rows *augment* the matrix.

Using the same approach as before, we can show that premultiplying $\Lambda_{\text{FACT-CM}}$ by its transpose, the determinant of the resulting squared matrix is non-zero,

$$\Lambda'_{\text{FACT-CM}} \Lambda_{\text{FACT-CM}} = \begin{bmatrix} 1.00 & 0.00 & 0.00 & 0.30 & 0.15 & 0.20 \\ 0.00 & 1.96 & 0.00 & 0.42 & 0.21 & 0.28 \\ 0.00 & 0.00 & 0.64 & 0.24 & 0.12 & 0.16 \\ 0.30 & 0.42 & 0.24 & 1.44 & 0.00 & 0.00 \\ 0.15 & 0.21 & 0.12 & 0.00 & 0.36 & 0.00 \\ 0.20 & 0.28 & 0.16 & 0.00 & 0.00 & 0.64 \end{bmatrix}$$

$$|\Lambda'_{\text{FACT-CM}} \Lambda_{\text{FACT-CM}}| = 0.1821$$

Moreover, even when only one single identifying indicator is added to the original CT-CM factor loading matrix – resulting in the Single Indicator Partially Augmented CT-CM model (SIPACT-CM) – the determinant of $\Lambda' \Lambda$ is non-zero,

$$\Lambda_{\text{SIPACT-CM}} = \begin{bmatrix} 0.50 & 0.00 & 0.00 & 0.60 & 0.00 & 0.00 \\ 0.00 & 0.70 & 0.00 & 0.60 & 0.00 & 0.00 \\ 0.00 & 0.00 & 0.40 & 0.60 & 0.00 & 0.00 \\ 0.50 & 0.00 & 0.00 & 0.00 & 0.30 & 0.00 \\ 0.00 & 0.70 & 0.00 & 0.00 & 0.30 & 0.00 \\ 0.00 & 0.00 & 0.40 & 0.00 & 0.30 & 0.00 \\ 0.50 & 0.00 & 0.00 & 0.00 & 0.00 & 0.40 \\ 0.00 & 0.70 & 0.00 & 0.00 & 0.00 & 0.40 \\ 0.00 & 0.00 & 0.40 & 0.00 & 0.00 & 0.40 \\ 0.50 & 0.00 & 0.00 & 0.00 & 0.00 & 0.00 \end{bmatrix}$$

$$\Lambda'_{\text{SIPACT-CM}} \Lambda_{\text{SIPACT-CM}} = \begin{bmatrix} 1.00 & 0.00 & 0.00 & 0.30 & 0.15 & 0.20 \\ 0.00 & 1.47 & 0.00 & 0.42 & 0.21 & 0.28 \\ 0.00 & 0.00 & 0.48 & 0.24 & 0.12 & 0.16 \\ 0.30 & 0.42 & 0.24 & 1.08 & 0.00 & 0.00 \\ 0.15 & 0.21 & 0.12 & 0.00 & 0.27 & 0.00 \\ 0.20 & 0.28 & 0.16 & 0.00 & 0.00 & 0.48 \end{bmatrix}$$

$$|\Lambda'_{SIPACT-CM}\Lambda_{SIPACT-CM}| = 0.008$$

but its small value suggests the SIPACT-CM can still lead to empirical underidentification. Indeed, the condition number for this matrix is 8.8620381, which is rather large and can lead to an ill-conditioned Hessian matrix. We can put the condition number for the lambda matrix of the SIPACT-CM model in context by comparing it with that of the CT-CM model, 3.538017e+16 (i.e., over 35 quadrillions), and that of the FACT-CM model, 3.6830897.

Online Supplemental Material

Figure S1. Boxplots illustrating distribution of mean bias in method-factor loading matrices by model and condition. Mean biases are based on conditions that converged to proper solutions, which can be identified in Table 1.

Figure S2. Boxplots illustrating distribution of mean bias in method-factor correlation matrices by model and condition. Mean biases are based on conditions that converged to proper solutions, which can be identified in Table 1.

Figure S3. Boxplots illustrating distribution of mean bias in error matrices by model and condition. Mean biases are based on conditions that converged to proper solutions, which can be identified in Table 1.