Supporting Information

Bifunctional Organocatalysts for Enantioselective aza-Morita-Baylis-Hillman (aza-MBH) Reaction

Katsuya Matsui, Shinobu Takizawa, and Hiroaki Sasai*

The Institute of Scientific and Industrial Research (ISIR), Osaka University,
Mihogaoka, Ibaraki, Osaka 567-0047, Japan
sasai@sanken.osaka-u.ac.jp

Experimental Section.

General. Commercially available organic and inorganic compounds were used without further purification except for the solvent, which was distilled from sodium/benzophenone or CaH₂. Column chromatography on SiO₂ was performed with Kanto Silica Gel 60 (40-100 μ m).

¹H and ¹³C NMR spectra were recorded with JEOL JNM-EX270 FT NMR (¹H-NMR 270 MHz, ¹³C-NMR 67.7 MHz). All signals are expressed as ppm down field from tetramethylsilane used as an internal standard. FT-IR spectra were recorded on a SHIMADZU FTIR-8300. Optical rotations were measured with JASCO P-1030 polarimeter. HPLC analyses were performed on a JASCO HPLC system (JASCO PU 980 pump and UV-975 UV/Vis detector) using a mixture of hexane and *i*-PrOH as the eluent. Mass spectra were obtained on JEOL JMS-DX300 (for EI-MS), JEOL JMS-700 (for FAB-MS), and JMS-T100LC (for ESI-MS). Elemental analysis was performed on PERKIN-ELMER 2400.

General procedure for synthesis of (*S*)-3-(*N*-isopropyl-*N*-3-pyridinylaminomethyl)BINOL

(*S*)-II

A solution of corresponding 3-*N*-methylaminopyridine¹⁾ (26.0 mg, 0.24 mmol) or 3-*N*-isopropylaminopyridine²⁾ (32.7 mg, 0.24 mmol) in THF (0.7 mL) was added to a heterogeneous solution of NaH (16 mg, 0.24 mmol) in THF (0.3 mL) at 0°C. After stirring for 2 h at 60°C, a solution of (*S*)-I³⁾ (0.20 mmol) in THF (1.0 mL) was combined with the reaction mixture. The mixture was stirred for 15 min at room temperature and then quenched with saturated NH₄Cl aq at 0°C. The organic phase was extracted with CH₂Cl₂, dried over Na₂SO₄, filtered and concentrated. The obtained residue was purified by flash chromatography (SiO₂, CH₂Cl₂/MeOH = 19/1) to afford the titled compound (*S*)-II (**IIa**, R = Me, quant.; **IIb**, R = *i*-Pr, 60% yield)

(*S*)-**IIa** (R = Me) colorless oil; IR (neat): ν 3083, 2912, 2812, 1581, 1494, 1444, 1346, 1222 cm⁻¹; ¹H-NMR (CDCl₃): δ 8.25 (1H, d, *J* = 3.0 Hz), 7.99 (2H, d, *J* = 9.2 Hz), 7.89 (1H, d, *J* = 8.6 Hz), 7.75 (1H, d, *J* = 8.1 Hz), 7.61 (1H, s), 7.60 (1H, d, *J* = 8.4 Hz), 7.41-7.11 (6H, m), 7.19 (2H, d, *J* = 7.0 Hz), 5.16 (1H, d, *J* = 7.3 Hz), 5.06 (1H, d, *J* = 7.3 Hz), 4.91

(2H, s), 4.67 (1H, d, $J = 5.9$ Hz), 4.50 (1H, d, $J = 5.9$ Hz), 3.28 (3H, s), 3.18 (3H, s), 3.10 (3H, s); $^{13}\text{C-NMR}$ (CDCl_3): δ 152.6, 152.4, 145.1, 137.6, 134.5, 133.7, 133.1, 130.7, 130.5, 129.9, 129.6, 127.8, 127.7, 126.8, 125.9, 125.8, 125.5, 125.4, 125.3, 125.0, 124.1, 123.5, 120.4, 118.2, 116.4, 99.3, 94.8, 57.0, 56.1, 52.7, 38.7; HRMS (ESI) calcd for $\text{C}_{31}\text{H}_{32}\text{N}_2\text{NaO}_4$, $m/z = 517.2103$ $[(\text{M}+\text{Na})^+]$; found, $m/z = 517.2093$; $[\alpha]_{\text{D}}^{20} -40.8^\circ$ (c 1.0, CHCl_3).

(*S*)-**IIb** ($\text{R} = i\text{-Pr}$) brown oil; IR (neat): ν 3095, 2945, 2829, 1596, 1514, 1444, 1375, 1221 cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3): δ 8.27 (1H, brs), 7.99 (2H, d, $J = 8.9$ Hz), 7.89 (1H, d, $J = 8.1$ Hz), 7.75 (1H, d, $J = 5.7$ Hz), 7.74 (1H, s), 7.60 (1H, d, $J = 8.4$ Hz), 7.41-7.05 (8H, m), 5.15 (1H, d, $J = 7.0$ Hz), 5.07 (1H, d, $J = 7.0$ Hz), 4.74 (2H, s), 4.70 (1H, d, $J = 5.7$ Hz), 4.53 (1H, d, $J = 5.7$ Hz), 4.48-4.36 (1H, m), 3.18 (3H, s), 3.12 (3H, s), 1.33 (3H, d, $J = 6.7$ Hz), 1.32 (3H, d, $J = 6.7$ Hz); $^{13}\text{C-NMR}$ (CDCl_3): δ 152.7, 152.0, 145.2, 145.1, 137.4, 133.8, 132.0, 130.9, 129.9, 129.6, 127.8, 127.7, 126.8, 126.5, 125.7, 125.5, 125.4, 125.4, 125.1, 124.9, 124.1, 123.6, 120.6, 119.3, 116.6, 99.3, 95.0, 55.9, 56.0, 48.1, 44.7, 19.8, 19.7; HRMS (ESI) calcd for $\text{C}_{33}\text{H}_{35}\text{N}_2\text{O}_4$, $m/z = 523.2597$ $[(\text{M}+\text{H})^+]$; found, $m/z = 523.2585$; $[\alpha]_{\text{D}}^{20} -43.2^\circ$ (c 1.0, CHCl_3).

(*S*)-2a (or 6a)

Trimethylsilyl bromide (53 μL , 0.4 mmol) was added to the mixture of corresponding (*S*)-**II** (0.2 mmol) and MS 4A (10 mg) in CH_2Cl_2 (1 mL) at 0°C . After stirring for 30 min, the reaction mixture was quenched with water. The organic phase was extracted with CH_2Cl_2 . The combined organic extracts were then dried (Na_2SO_4) and evaporated in vacuo. The residue was purified by flash chromatography (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH} = 9/1$) to give the title compound (*S*)-**2a** (92% yield) or **6a** (90% yield).

(*S*)-**2a** ($\text{R} = \text{Me}$) white solid; mp $119\text{-}120^\circ\text{C}$ (hexane/ CH_2Cl_2); IR (neat): ν 3389, 3012, 2933, 1284, 1612, 1588, 1510, 1424, 1412, 1367, 1220, 1012 cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3): δ 8.28-8.18 (1H, br), 7.96 (2H, d, $J = 9.2$ Hz), 7.89 (2H, d, $J = 8.4$ Hz), 7.79 (1H, d, $J = 8.1$ Hz), 7.70 (1H, s), 7.39 (1H, d, $J = 8.9$ Hz), 7.34-7.13 (6H, m), 4.76 (2H, s), 3.18 (3H, s); $^{13}\text{C-NMR}$ (CDCl_3): δ 153.3, 151.4, 145.2, 136.6, 133.8, 133.0, 129.0, 128.8, 128.1, 127.8, 127.1, 126.9, 126.5, 125.6, 124.4, 124.3, 123.7, 123.3, 119.5, 118.3, 113.3, 112.0, 53.0, 38.6; HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{23}\text{N}_2\text{O}_2$, $m/z = 407.1760$ $[(\text{M}+\text{H})^+]$; found, $m/z = 407.1740$; $[\alpha]_{\text{D}}^{20} -17.8^\circ$ (c 0.7, CHCl_3).

(*S*)-**6a** ($\text{R} = i\text{-Pr}$) brown solid; mp $122\text{-}123^\circ\text{C}$ (hexane/ CH_2Cl_2); IR (neat): ν 3440, 3055, 2966, 2823, 1705, 1620, 1585, 1566, 1497, 1431, 1342, 1272, 1184, 1134, 1096, 1061 cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3): δ 8.18 (1H, d, $J = 2.7$ Hz), 7.94 (1H, dd, $J = 4.0$ and 0.8 Hz), 7.85 (1H, d, $J = 8.9$ Hz), 7.79 (1H, d, $J = 7.8$ Hz), 7.79-7.65 (2H, m), 7.28 (1H, d, $J = 8.9$ Hz), 7.28-7.00 (6H, m), 6.96 (1H, d, $J = 8.1$ Hz), 4.55 (2H, s), 4.04-3.43 (1H, m), 1.21 (3H, d, $J = 6.5$ Hz), 1.18 (3H, d, $J = 6.7$ Hz); $^{13}\text{C-NMR}$ (CDCl_3): δ 152.4, 151.8, 143.8, 140.0, 133.5, 132.9, 130.6, 129.2, 128.8, 128.3, 128.2, 128.1, 127.9, 126.9, 126.7, 126.2, 124.3, 124.3, 124.2, 123.8, 123.5, 123.4, 117.8, 112.4, 112.3, 50.8, 47.0, 19.6, 19.4; HRMS (ESI) calcd for $\text{C}_{29}\text{H}_{27}\text{N}_2\text{O}_2$, $m/z = 435.2073$ $[(\text{M}+\text{H})^+]$; found, $m/z = 435.2062$; $[\alpha]_{\text{D}}^{20} -28.5^\circ$ (c 0.4, CHCl_3).

The effects of reaction conditions on enantioselective aza-MBH reaction promoted by (*S*)-2a

The effect of solvent on organocatalyst **2a** mediated reaction is summarized in Table S-1. Ethereal solvents such as diethyl ether, *t*-BuOMe, cyclopentyl methyl ether (CPME), dimethyl ether (DME) and THF (Entries 1-5) were relatively good in terms of enantioselectivity when compared with other less polar, polar, aprotic and protic solvents such as halogenated solvents, DMF, acetonitrile, and methanol (entries 6-12). A mixed solvent system consisting of CPME and toluene (9/1) is afforded the good result (Entries 13).

Table S-1

Entry	Solvent	Time (h)	Yield (%) ^a	Ee (%) ^b
1	Et ₂ O	108	74	72
2	<i>t</i> -BuOMe	72	92	73
3	CPME	72	97	78
4	DME	60	73	68
5	THF	48	71	59
6	toluene	24	81	72
7	CH ₂ Cl ₂	24	Quant	59
8	CHCl ₃	24	97	54
9	ClCH ₂ CH ₂ Cl	24	87	57
10	DMF	48	7	63
11	MeCN	48	60	67
12	MeOH	60	46	18
13	CPME:toluene (9:1)	72	93	83

^aIsolated yield. ^bDetermined by HPLC (DAICEL CHIRALPAK AD-H).

Table S-2 summarises the effect of temperature on the reaction. While the reaction proceeded smoothly at various temperatures, the best result was obtained at -15°C .

Table S-2

$3\text{a} + 4\text{b} \xrightarrow[\text{CPME/toluene (9/1), Temp}]{2\text{a (10 mol \%)}} 5\text{b}$				
Entry	Temperature ($^{\circ}\text{C}$)	Time (h)	Yield (%) ^a	Ee (%) ^b
1	rt	60	93	83
2	0	96	94	86
3	-10	120	95	87
4	-15	144	97	90
5	-20	360	12	90

^aIsolated yield. ^bDetermined by HPLC (DAICEL CHIRALPAK AD-H).

General procedure for enantioselective aza-MBH reaction promoted by (S)-6a

To a solution of organocatalyst (S)-6a (2.2 mg, 0.005 mmol) in CPME/toluene (9/1, 0.1 mL) was added to methyl vinyl ketone (**3**, 12.5 μ L, 0.15 mmol) and known imines⁴⁾ (**4**, 0.05 mmol) at -15°C . The mixture was stirred until the reaction had reached completion by monitoring with TLC analysis. The mixture was directly purified by flash column chromatography (SiO_2 , *n*-hexane/EtOAc = 2/1) to give the corresponding adducts **5a-j** as white solid. The adducts **5a-f** and **5h** were identical in all respects with reported by Shi and co-workers.⁵⁾ Absolute configurations were assigned by comparison of optical rotation reported in the literature.⁵⁾

Table 1. Enantioselective aza-MBH reaction promoted by (S)-6a

Reaction scheme: **3** + **4** $\xrightarrow[\text{toluene:CPME (1:9), } -15^{\circ}\text{C}]{\text{6a (10 mol \%)}}$ **5**

Entry	3 : R ¹	4 : R ²	Time (h)	Yield (%) ^a	Ee (%) ^b
1	Me (3a)	Ph (4a)	168	5a , 93	87
2	Me (3a)	<i>p</i> -Cl-C ₆ H ₄ (4b)	60	5b , 96	95
3	Me (3a)	<i>m</i> -Cl-C ₆ H ₄ (4c)	72	5c , 93	93
4	Me (3a)	<i>p</i> -Br-C ₆ H ₄ (4d)	36	5d , 93	94
5	Me (3a)	<i>p</i> -MeO-C ₆ H ₄ (4e)	132	5e , 93	94
6	Me (3a)	2-furyl (4f)	48	5f , Quant	88
7	Me (3a)	2-naphthyl (4g)	108	5g , 94	91
8	Me (3a)	<i>p</i> -NO ₂ -C ₆ H ₄ (4h)	12	5h , 91	91
9	Et (3b)	<i>p</i> -NO ₂ -C ₆ H ₄ (4h)	96	5i , 88	88
10	H (3c)	<i>p</i> -NO ₂ -C ₆ H ₄ (4h)	36	5j , 95	94

^aIsolated yield. ^bDetermined by HPLC (DAICEL CHIRALPAK AD-H or OD-H).

5a (R¹ = Me, R² = Ph) 93% yield, 87% ee; ¹H-NMR (CDCl₃): δ 7.64 (2H, d, *J* = 8.4 Hz), 7.31-7.20 (5H, m), 7.11 (2H, d, *J* = 8.4 Hz), 6.09 (1H, s), 6.08 (1H, s), 5.59 (1H, d, *J* = 8.6 Hz), 5.25 (1H, d, *J* = 8.6 Hz), 2.41 (3H, s), 2.13 (3H, s); DAICEL CHIRALPAK AD-H column, detection at 254 nm, *i*-PrOH/*n*-hexane = 1/4, flow rate 0.2 mL/min, 43.6 min (minor isomer, *S*) and 49.6 min (major isomer, *R*).

5b (R¹ = Me, R² = *p*-Cl-C₆H₄) 96% yield, 95% ee; ¹H-NMR (CDCl₃): δ 7.59 (2H, d, *J* = 8.4 Hz), 7.18 (2H, d, *J* = 8.4 Hz), 7.12 (2H, d, *J* = 8.6 Hz), 7.00 (2H, d, *J* = 8.6 Hz), 6.05 (1H, s), 6.03 (1H, s), 5.99 (1H, d, *J* = 8.9 Hz), 5.25 (1H, d, *J* = 8.9 Hz), 2.40 (3H, s), 2.13 (3H, s); DAICEL CHIRALPAK AD-H column, detection at 254 nm, *i*-PrOH/*n*-hexane = 1/4, flow rate 0.3 mL/min, 44.7 min (minor isomer, *S*) and 50.8 min (major isomer, *R*).

5c (R¹ = Me, R² = *m*-Cl-C₆H₄) 93% yield, 93% ee; ¹H-NMR (CDCl₃): δ 7.64 (2H, d, *J* = 8.4 Hz), 7.24 (2H, d, *J* = 8.4 Hz), 7.17-7.11 (2H, m), 7.05-7.00 (2H, m), 6.10 (1H, s), 6.05 (1H, s), 5.78 (1H, d, *J* = 8.9 Hz), 5.23 (1H, d, *J* = 8.9 Hz), 2.40 (3H, s), 2.15 (3H, s); DAICEL CHIRALPAK AD-H column, detection at 254 nm, *i*-PrOH/*n*-hexane = 1/4, flow rate 0.2 mL/min, 56.5 min (minor isomer, *S*) and 70.2 min (major isomer, *R*).

5d ($R^1 = \text{Me}$, $R^2 = p\text{-Br-C}_6\text{H}_4\text{-}$) 93% yield, 94% ee; $^1\text{H-NMR}$ (CDCl_3): δ 7.60 (2H, d, $J = 8.4$ Hz), 7.30 (2H, d, $J = 8.4$ Hz), 7.25 (2H, d, $J = 8.6$ Hz), 6.99 (2H, d, $J = 8.6$ Hz), 6.09 (1H, s), 6.06 (1H, s), 5.69 (1H, d, $J = 8.9$ Hz), 5.16 (1H, d, $J = 8.9$ Hz), 2.40 (3H, s), 2.13 (3H, s); DAICEL CHIRALPAK AD-H column, detection at 254 nm, *i*-PrOH/*n*-hexane = 1/4, flow rate 0.2 mL/min, 48.9 min (minor isomer, *S*) and 56.1 min (major isomer, *R*).

5e ($R^1 = \text{Me}$, $R^2 = p\text{-MeO-C}_6\text{H}_4\text{-}$) 93% yield, 94% ee; $^1\text{H-NMR}$ (CDCl_3): δ 7.66 (2H, d, $J = 8.4$ Hz), 7.25 (2H, d, $J = 8.4$ Hz), 7.00 (2H, d, $J = 8.6$ Hz), 6.73 (2H, d, $J = 8.6$ Hz), 6.09 (2H, s), 5.53 (1H, d, $J = 8.9$ Hz), 5.25 (1H, d, $J = 8.9$ Hz), 3.77 (3H, s), 2.45 (3H, s), 2.19 (3H, s); DAICEL CHIRALPAK OD-H column, detection at 254 nm, *i*-PrOH/*n*-hexane = 1/4, flow rate 0.2 mL/min, 58.8 min (major isomer, *R*) and 65.4 min (minor isomer, *S*).

5f ($R^1 = \text{Me}$, $R^2 = 2\text{-Furyl}$) quant, 88% ee; $^1\text{H-NMR}$ (CDCl_3): δ 7.66 (2H, d, $J = 8.4$ Hz), 7.26 (2H, d, $J = 8.4$ Hz), 7.16 (1H, d, $J = 1.8$ Hz), 6.20 (1H, dd, $J = 3.0$ and 1.8 Hz), 6.10 (1H, s), 6.06 (1H, s), 6.00 (1H, d, $J = 3.0$ Hz), 5.70 (1H, d, $J = 8.6$ Hz), 5.36 (1H, d, $J = 8.6$ Hz), 2.40 (3H, s), 2.20 (3H, s); DAICEL CHIRALPAK AD-H column, detection at 254 nm, *i*-PrOH/*n*-hexane = 1/4, flow rate 0.5 mL/min, 24.8 min (minor isomer, *S*) and 29.1 min (major isomer, *R*).

5g ($R^1 = \text{Me}$, $R^2 = 2\text{-naphthyl}$) 94% yield, 91% ee; white solid 105-106°C (hexane/AcOEt); IR (neat): ν 1661 cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3): δ 7.76-7.50 (3H, m), 7.58 (2H, d, $J = 8.4$ Hz), 7.43 (1H, s), 7.39-7.33 (2H, m), 7.12 (3H, d, $J = 7.5$ Hz), 6.12-6.08 (2H, m), 5.65 (1H, d, $J = 8.9$ Hz), 5.36 (1H, d, $J = 8.6$ Hz), 2.29 (3H, s), 2.11 (3H, s); $^{13}\text{C-NMR}$ (CDCl_3): δ 198.6, 146.4, 143.2, 137.4, 136.0, 132.9, 132.5, 129.3, 128.2, 128.1, 127.8, 127.4, 127.2, 126.1, 126.0, 125.4, 124.2, 58.9, 26.4, 21.4; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{21}\text{NnaO}_3\text{S}$, $m/z = 402.1140$ [($\text{M}+\text{Na}$) $^+$]; found, $m/z = 402.1152$; $[\alpha]_{\text{D}}^{20} -47.5^\circ$ (c 0.6, CHCl_3); DAICEL CHIRALPAK AD-H column, detection at 254 nm, *i*-PrOH/*n*-hexane = 1/4, flow rate 0.3 mL/min, 61.7 min (minor isomer, *S*) and 65.6 min (major isomer, *R*).

5h ($R^1 = \text{Me}$, $R^2 = p\text{-NO}_2\text{-C}_6\text{H}_4\text{-}$) 91% yield, 91% ee; $^1\text{H-NMR}$ (CDCl_3): δ 8.08 (2H, d, $J = 8.4$ Hz), 7.67 (2H, d, $J = 8.4$ Hz), 7.34 (2H, d, $J = 8.6$ Hz), 7.26 (2H, d, $J = 8.6$ Hz), 6.16 (1H, s), 6.10 (1H, s), 5.93 (1H, d, $J = 8.9$ Hz), 5.32 (1H, d, $J = 8.9$ Hz), 2.45 (3H, s), 2.16 (3H, s); DAICEL CHIRALPAK AD-H column, detection at 254 nm, *i*-PrOH/*n*-hexane = 1/4, flow rate 0.5 mL/min, 53.2 min (minor isomer, *S*) and 70.8 min (major isomer, *R*).

5i ($R^1 = \text{Et}$, $R^2 = p\text{-NO}_2\text{-C}_6\text{H}_4\text{-}$) 88% yield, 88% ee; white solid; mp 112-113°C (hexane/AcOEt); IR (neat): ν 1670, 1515, 1342 cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3): δ 8.07 (2H, d, $J = 8.9$ Hz), 7.64 (2H, d, $J = 8.4$ Hz), 7.35 (2H, d, $J = 8.9$ Hz), 7.24 (2H, d, $J = 8.4$ Hz), 6.13 (1H, s), 6.04 (1H, s), 5.90 (1H, d, $J = 9.4$ Hz), 5.31 (1H, d, $J = 9.4$ Hz), 2.50 (2H, q, $J = 7.5$ Hz), 2.41 (3H, s), 0.93 (3H, t, $J = 7.5$ Hz); $^{13}\text{C-NMR}$ (CDCl_3): δ 201.2, 147.0, 146.2, 144.8, 143.6, 137.3, 129.8, 129.5, 128.2, 127.1, 127.1, 126.3, 123.5, 59.2, 31.3, 21.6, 7.8; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{20}\text{N}_2\text{NaO}_5\text{S}$, $m/z = 411.0991$ [($\text{M}+\text{Na}$) $^+$]; found, $m/z = 411.0990$; $[\alpha]_{\text{D}}^{20} 5.3^\circ$ (c 0.6, CHCl_3); DAICEL CHIRALPAK AD-H column, detection at 254 nm, *i*-PrOH/*n*-hexane = 1/4, flow rate 0.5 mL/min, 47.7 min (minor isomer) and 52.9 min (major isomer).

5j ($R^1 = \text{H}$, $R^2 = p\text{-NO}_2\text{-C}_6\text{H}_4\text{-}$) 95% yield, 94% ee; $^1\text{H-NMR}$ (CDCl_3): δ 9.33 (1H, s), 8.00 (2H, d, $J = 8.9$ Hz), 7.61 (2H, d, $J = 8.4$ Hz), 7.29 (2H, d, $J = 8.9$ Hz), 7.19 (2H, d, $J = 8.4$ Hz), 6.50 (1H, s), 6.17 (1H, d, $J = 9.4$ Hz), 6.11 (1H, s), 5.37 (1H, d, $J = 9.4$ Hz), 2.38 (3H, s); $[\alpha]_{\text{D}}^{20} -18.0^\circ$ (c 0.6, CHCl_3); DAICEL CHIRALPAK OD-H column, detection at 254 nm, *i*-PrOH/*n*-hexane = 1/4, flow rate 0.5 mL/min, 43.7 min (major isomer) and 77.6 min (minor isomer).

Figure S-1. Molecular orbital calculation of **6a**

Spartan '04, Job type: Geometry optimization, Method: Hartee-Fock, Basis set: 3-21G*. The N-H atomic distance (1.835 Å). The angle between the N-H-O bonds (145.02°).

References

1. Krishnamurthy, S. *Tetrahedron Lett.* **1982**, 23, 3315-3318.
2. a) Ieno, M. Jpn. Kokai Tokkyo Koho (1999), 4pp. JP11035562; b) Morris, J.; Wishka, D. G. *J. Org. Chem.* **1995**, 60, 2642-2644.
3. a) Matsunaga, S.; Das, J.; Roels, J.; Vogl, E. M.; Yamamoto, N.; Iida, T.; Yamaguchi, K.; Shibasaki, M. *J. Am. Chem. Soc.* **2000**, 122, 2252-2260; b) Lee, K. Y.; Lee, C. G.; Kim, J. M. *Tetrahedron Lett.* **2003**, 44, 1231-1234.
4. a) ten Have, R.; Huisman, M.; Meetsma, A.; van Leusen, A. M. *Tetrahedron* **1997**, 53, 11355-11368. b) Lee, K. Y.; Lee, C. G.; Kim, J. M. *Tetrahedron Lett.* **2003**, 44, 1231-1234.
5. a) Shi, M.; Xu, Y.-M. *Angew. Chem. Int. Ed.* **2002**, 41, 4507-4510; b) Shi, M.; Chen, L.-H. *Chem. Commun.* **2003**, 1310-1311; c) Xu, Y.-M.; Shi, M. *J. Org. Chem.* **2004**, 69, 417-425.