Supporting Information

Biosynthesis of 13-Desmethyl Spirolide C by the Dinoflagellate *Alexandrium ostenfeldii*.

Shawna L. MacKinnon, Allan D. Cembella¹, Ian W. Burton, Nancy Lewis,
Patricia LeBlanc and John A. Walter

*Institute for Marine Biosciences, National Research Council of Canada, 1411 Oxford
Street, Halifax NS, Canada B3H 3Z1.*

¹*Now at the Alfred Wegener Institute for Polar and Marine Research, Bremerhaven,
Germany.*

Table of Contents:

- S2 **FIGURE S1.** Structure of 13-desmethyl spirolide C **1** showing ¹³C resonance assignments (solvent CD₃OD, referred to ¹³CD₃OD = 49.0 ppm) in addition to labelling pattern and coupling constants as for Figure 2 and Table 1.
- S3 to S9 **FIGURES S2A to S2G.** Expanded segments of 125.7 MHz ¹³C NMR spectra of **1** showing differences in ¹³C-¹³C satellite resonance intensities for samples enriched from: (A) [1-¹³C] acetate (Experiment 3); (B) [1,2-¹³C₂] acetate (Experiment 1); (C) [1,2-¹³C₂] acetate/NA acetate (molar ratio 1/3, Experiment 2).
- S10 to S21 **FIGURES S3A to S3L.** ¹³C 2D INADEQUATE spectra of **1** enriched from [1-¹³C] acetate (Experiment 3), [1,2-¹³C₂] acetate (Experiment 1) or [1,2-¹³C₂] acetate/NA acetate (molar ratio 1/3, Experiment 2).
- S22 **FIGURE S4.** Structure of **1** showing average % ¹³C at each position retained from [1-¹³C] acetate (Experiment 3) or [2-¹³CD₃] acetate (Experiment 4), and retention of D from [2-¹³CD₃] acetate.

FIGURE S1. Structure of 13-desmethyl spirolide C 1 showing ¹³C resonance assignments (solvent CD₃OD, referred to ¹³CD₃OD = 49.0 ppm) in addition to labelling pattern and coupling constants as for Figure 2 and Table 1.

FIGURE S2A.

Expanded segments of 125.7 MHz ^{13}C NMR spectra of **1** showing differences in ^{13}C - ^{13}C satellite resonance intensities for samples enriched from:

- (A) $[1-^{13}\text{C}]$ acetate (Experiment 3);
- (B) $[1,2-^{13}\text{C}_2]$ acetate (Experiment 1);
- (C) $[1,2-^{13}\text{C}_2]$ acetate/NA acetate (molar ratio 1/3, Experiment 2).

FIGURE S2B.

Expanded segments of 125.7 MHz ^{13}C NMR spectra of **1** showing differences in ^{13}C - ^{13}C satellite resonance intensities for samples enriched from:

- (C) $[1-^{13}\text{C}]$ acetate (Experiment 3);
- (D) (B) $[1,2-^{13}\text{C}_2]$ acetate (Experiment 1);
- (C) $[1,2-^{13}\text{C}_2]$ acetate/NA acetate (molar ratio 1/3, Experiment 2).

FIGURE S2C.

Expanded segments of 125.7 MHz ^{13}C NMR spectra of **1** showing differences in ^{13}C - ^{13}C satellite resonance intensities for samples enriched from:

- (E) $[1-^{13}\text{C}]$ acetate (Experiment 3);
- (F) (B) $[1,2-^{13}\text{C}_2]$ acetate (Experiment 1);
- (C) $[1,2-^{13}\text{C}_2]$ acetate/NA acetate (molar ratio 1/3, Experiment 2).

FIGURE S2D.

Expanded segments of 125.7 MHz ^{13}C NMR spectra of **1** showing differences in ^{13}C - ^{13}C satellite resonance intensities for samples enriched from:

- (G) $[1-^{13}\text{C}]$ acetate (Experiment 3);
- (H) (B) $[1,2-^{13}\text{C}_2]$ acetate (Experiment 1);
- (C) $[1,2-^{13}\text{C}_2]$ acetate/NA acetate (molar ratio 1/3, Experiment 2).

FIGURE S2E.

Expanded segments of 125.7 MHz ^{13}C NMR spectra of **1** showing differences in ^{13}C - ^{13}C satellite resonance intensities for samples enriched from:

- (I) $[1-^{13}\text{C}]$ acetate (Experiment 3);
- (J) (B) $[1,2-^{13}\text{C}_2]$ acetate (Experiment 1);
- (C) $[1,2-^{13}\text{C}_2]$ acetate/NA acetate (molar ratio 1/3, Experiment 2).

FIGURE S2F.

Expanded segments of 125.7 MHz ^{13}C NMR spectra of **1** showing differences in ^{13}C - ^{13}C satellite resonance intensities for samples enriched from:

- (K) $[1-^{13}\text{C}]$ acetate (Experiment 3);
- (L) (B) $[1,2-^{13}\text{C}]_2$ acetate (Experiment 1);
- (C) $[1,2-^{13}\text{C}]_2$ acetate/NA acetate (molar ratio 1/3, Experiment 2).

FIGURE S2G.

Expanded segments of 125.7 MHz ^{13}C NMR spectra of **1** showing differences in ^{13}C - ^{13}C satellite resonance intensities for samples enriched from:

- (M) $[1\text{-}^{13}\text{C}]$ acetate (Experiment 3);
- (N) (B) $[1,2\text{-}^{13}\text{C}_2]$ acetate (Experiment 1);
- (C) $[1,2\text{-}^{13}\text{C}_2]$ acetate/NA acetate (molar ratio 1/3, Experiment 2).

FIGURE S3A. ¹³C 2D INADEQUATE spectra of **1** enriched from [1-¹³C] acetate, (Experiment 3) optimized for $J_{CC} = 40$ Hz. For expansion of lower right hand region of plane see Figure S3B. Note similarity of peak intensities indicating uniform ¹³C enrichment from scrambled label.

FIGURE S3B. ¹³C 2D INADEQUATE spectra of **1** enriched from [1-¹³C] acetate, (Experiment 3) optimized for $J_{CC} = 40$ Hz (expansion of lower right hand region of Figure S3A). The similarity of peak intensities indicates uniform ¹³C enrichment from scrambled label.

FIGURE S3C. ^{13}C 2D INADEQUATE spectra of **1** enriched from $[1,2-^{13}\text{C}_2]$ acetate (Experiment 1), optimized for $J_{\text{CC}} = 40$ Hz. For expansion of lower right hand region of plane see Figure S3D. Peaks for some coupled pairs have higher intensity, corresponding to incorporation of intact ^{13}C - ^{13}C units in addition to scrambled ^{13}C label (see Figure 1 or Figure S1A).

FIGURE S3D. ^{13}C 2D INADEQUATE spectra of **1** enriched from $[1,2-^{13}\text{C}_2]$ acetate (Experiment 1), optimized for $J_{\text{CC}} = 40$ Hz (expansion of lower right hand region of Figure S3C). Peaks for some coupled pairs have higher intensity, corresponding to incorporation of intact ^{13}C - ^{13}C units in addition to scrambled ^{13}C label (see Figure 1 or Figure S1A).

FIGURE S3E. ^{13}C 2D INADEQUATE spectra of **1** enriched from $[1,2-^{13}\text{C}_2]$ acetate/NA acetate (3/1) (Experiment 2), optimized for $J_{\text{CC}} = 30$ Hz. For expansion of lower right hand region of plane see Figure S3F. Peaks for some coupled pairs have higher intensity, corresponding to incorporation of intact ^{13}C - ^{13}C units in addition to scrambled ^{13}C label (see Figure 1 or Figure S1A).

FIGURE S3F. ^{13}C 2D INADEQUATE spectra of **1** enriched from [1,2- $^{13}\text{C}_2$] acetate/NA acetate (3/1) (Experiment 2), optimized for $J_{\text{CC}} = 30$ Hz (expansion of lower right hand region of Figure S3E). Peaks for some coupled pairs have higher intensity, corresponding to incorporation of intact ^{13}C - ^{13}C units in addition to scrambled ^{13}C label (see Figure 1 or Figure S1A).

FIGURE S3G. ^{13}C 2D INADEQUATE spectra of **1** enriched from $[1,2-^{13}\text{C}_2]$ acetate/NA acetate (3/1) (Experiment 2), optimized for $J_{\text{CC}} = 40$ Hz. For expansion of lower right hand region of plane see Figure S3H. Peaks for some coupled pairs have higher intensity, corresponding to incorporation of intact ^{13}C - ^{13}C units in addition to scrambled ^{13}C label (see Figure 1 or Figure S1A).

FIGURE S3H. ¹³C 2D INADEQUATE spectra of **1** enriched from [1,2-¹³C₂] acetate/NA acetate (3/1) (Experiment 2), optimized for J_{CC} = 40 Hz (expansion of lower right hand region of Figure S3G). Peaks for some coupled pairs have higher intensity, corresponding to incorporation of intact ¹³C-¹³C units in addition to scrambled ¹³C label (see Figure 1 or Figure S1A).

FIGURE S3I. ^{13}C 2D INADEQUATE spectra of **1** enriched from $[1,2\text{-}^{13}\text{C}_2]$ acetate/NA acetate (3/1) (Experiment 2), optimized for $J_{\text{CC}} = 60$ Hz. For expansion of lower right hand region of plane see Figure S3J. Peaks for some coupled pairs have higher intensity, corresponding to incorporation of intact ^{13}C - ^{13}C units in addition to scrambled ^{13}C label (see Figure 1 or Figure S1A).

FIGURE S3J. ^{13}C 2D INADEQUATE spectra of **1** enriched from [1,2- $^{13}\text{C}_2$] acetate/NA acetate (3/1) (Experiment 2), optimized for $J_{\text{CC}} = 60$ Hz (expansion of lower right hand region of Figure S3I). Peaks for some coupled pairs have higher intensity, corresponding to incorporation of intact ^{13}C - ^{13}C units in addition to scrambled ^{13}C label (see Figure 1 or Figure S1A).

FIGURE S3K. ^{13}C 2D INADEQUATE spectra of **1** enriched from $[1-^{13}\text{C}]$ acetate (Experiment 3), optimized for $J_{\text{CC}} = 60$ Hz. For expansion of lower right hand region of plane see Figure S3L. Note similarity of peak intensities indicating uniform ^{13}C enrichment from scrambled label.

FIGURE S3L. ¹³C 2D INADEQUATE spectra of **1** enriched from [1-¹³C] acetate (Experiment 3), optimized for $J_{CC} = 60$ Hz (expansion of lower right hand region of Figure S3K). Note similarity of peak intensities indicating uniform ¹³C enrichment from scrambled label.

FIGURE S4. Structure of **1** showing average % ¹³C at each position retained from [1-¹³C] acetate (Experiment 3) or [2-¹³CD₃] acetate (Experiment 4), and retention of D from [2-¹³CD₃] acetate.