Supporting Information

**Can Trimethylamine-N-Oxide Act to Influence the
Self-Aggregation of tert-Butyl Alcohol?**

Sandip Paul*

Department of Chemistry, Indian Institute of Technology, Guwahati Assam, India-781039

(Dated: February 23, 2016)

FIG. S1: Water-water site-site radial distribution functions for different systems considered.

FIG. S2: TBA-water and TMAO-water selected site-site radial distribution functions. CC_{tb} and C_{tb} represent central carbon atom and methyl carbon of TBA and N_{tm} and C_{tm} are nitrogen atom and methyl carbon atom of TMAO respectively. O_w refers to water oxygen.

FIG. S3: TBA-TMAO site-site radial distribution functions for different systems considered. O_{tm} , O_{tb} and H_{tb} refer to TMAO oxygen, TBA oxygen and TBA hydroxyl hydrogen respectively.

FIG. S4: TBA-TMAO methyl carbon-methyl carbon ($C_{tb} - C_{tm}$) site-site radial distribution functions for all systems considered.

FIG. S5: (a) The average number of different types of hydrogen bonds vs. mole fraction of TBA. W, TB, and TM refer to water, TBA, and TMAO and the subscripts D and A represent donor and acceptor respectively. $Total^{TB}$ is the total number of hydrogen bonds formed by a single TBA molecule. (b) The average energy (in kJ/mol of solution) of different types of hydrogen bonds vs. mole fraction of TBA.

FIG. S6: The time dependence of the correlation function $S_{HB}(t)$ of water-water hydrogen bonds for the lowest ($x_{tba} = 0.002$) and for the highest ($x_{tba} = 0.10$) mole fraction of TBA.

* Electronic address: `sandipp@iitg.ernet.in`