Catalytic Asymmetric Acyl Halide-Aldehyde Cyclocondensation Reactions of Substituted Ketenes

Scott G. Nelson,* Cheng Zhu and Xiaoqiang Shen

Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260

Supporting Information

General Information: Optical rotations were measured on a Perkin-Elmer 241 digital polarimeter with a sodium lamp at ambient temperature and are reported as follows: $[\alpha]_D^{25}$ (c g/100mL). Infrared spectra were recorded on a Nicolet Avatar 360 FT-IR spectrometer. NMR spectra were recorded on a Bruker Avance-300 (300 MHz) spectrometer with chemical shifts reported relative to residual CHCl_3 (7.27 ppm) for ^1H and CDCl_3 (77.0 ppm) for ^{13}C NMR spectra. Analytical high performance liquid chromatography (HPLC) was performed on a Hewlett Packard 1100 liquid chromatograph equipped with a variable wavelength UV detector (deuterium lamp, 190-600 nm) using a Daicel ChiracelTM OD-H column (250 x 4.6 mm) (Daicel Inc.) or Daicel ChiralpakTM AD column (250 x 4.6 mm) (Daicel Inc.) and HPLC-grade isopropanol and hexanes as the eluting solvents. Analytical gas-liquid chromatography (GLC) was performed on a Varian 3900 gas chromatography equipped with a flame ionization detector and split mode capillary injection system using a ChiraldexTM G-TA column (20 m x 0.25 mm) (Advanced Separation Technologies Inc.). Helium was used as the carrier gas at the indicated pressures. Unless otherwise stated, all reactions were carried out in dry glassware under a nitrogen atmosphere using standard inert atmosphere techniques for the manipulation of solvents and reagents. Anhydrous solvents (CH_2Cl_2 , THF, diethyl ether and toluene) were obtained by passage through successive alumina and Q5 reactant-packed columns on a solvent purification system. *N*, *N*-Diisopropylethylamine and triethylamine were distilled under nitrogen from CaH_2 . Trimethylaluminum (2.0M in hexane) and benzotrifluoride (BTF) were purchased from Aldrich in Sure/SealTM bottles. 3-Benzyloxypionyaldehyde¹ and trimethylsilylpropynal² were prepared according to published procedures. All commercially available aldehydes were redistilled under N_2 . Commercially available propionyl bromide and valeryl bromide were distilled from P_2O_5 . Butyryl bromide and isovaleryl bromide were prepared according to literature procedures and were distilled from P_2O_5 .³ Flash chromatography was performed as previously described on EM silica gel 60 (230-240 mesh).⁴

General procedure for asymmetric AAC reaction (Table 1): To a solution of ligand **6** (0.10-0.20 mmol) in 2.0 ml of BTF at ambient temperature was slowly added 0.050 ml of a 2.0 M hexane solution of AlMe_3 (0.10 mmol). After stirring for 2 hours, the resulting catalyst solution was cooled to -25°C and 0.348 ml of *N*, *N*-diisopropylethylamine (2.0 mmol) and 2-4 equiv of acid bromide were added consecutively (see table below). The resulting heterogeneous mixture was stirred 10-30 min at -25°C (10 min: propionyl bromide; 15 min: butyryl bromide; 30 min: valeryl bromide; 20 min: isovaleryl bromide) whereupon aldehyde (1.0 mmol) was added dropwise and the reaction was stirred 14 h at -25°C . The reaction was quenched at -25°C by adding 10 ml of a 3% Et_3N - Et_2O solution (v/v) and the resulting mixture was filtered through silica gel (pretreated with 3% Et_3N -hexanes) eluting with 60 ml of Et_2O . The filtrate was concentrated *in vacuo* and the crude product mixture was purified by flash chromatography (for benzaldehyde-derived lactones, the silica gel was pretreated with 2% Et_3N -hexanes).

Table 1. Acid bromide and catalyst stoichiometry used for enolizable and non-enolizable aldehydes

Ketene \ Aldehyde	Enolizable (catalyst, acid bromide)	Non-enolizable (catalyst, acid bromide)
Methyl ketene	20 mol %, 4 equiv	10 mol %, 2 equiv
Ethyl ketene	10 mol %, 4 equiv	10 mol %, 2 equiv
ⁿ Propyl ketene	10 mol %, 4 equiv	10 mol %, 2 equiv
ⁱ Propyl ketene	NA	10 mol %, 2 equiv

¹ Kusakabe, M.; Kato, H.; Sato, F. *Chem. Lett.* **1987**, 2163-2166.

² Journet, M.; Cai, D.; Dimichele, L. M.; Larsen, R. D. *Tetrahedron Lett.* **1998**, 39, 6427-6428.

³ Burton, T. M.; Degering, E. F.; *J. Am. Chem. Soc.* **1940**, 62, 227.

⁴ Still, W. C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, 43, 2923-2925.

(S)-N-[3,5-bis(trifluoromethyl)benzenesulfonyl]-2-isopropylaziridine: To a solution of 2.05 g of (*S*)-valinol (19.9 mmol, 1.0 eq) and 120 mg of DMAP (1.0 mmol, 0.05 equiv) in 40 ml of CH₂Cl₂ at -25 °C was added 11.0 ml of Et₃N (80 mmol, 4.0 equiv). To the resulting solution was added 6.84 g of 3,5-bis(trifluoromethyl)benzenesulfonyl chloride (21.9 mmol, 1.1 equiv) in portions. After addition was complete, the reaction mixture was warmed to 0 °C and stirred for 1 h whereupon 1.85 ml of MsCl (23.9 mmol, 1.2 equiv) was added dropwise and the resulting reaction mixture was stirred an additional 2 h at 0 °C. The reaction mixture was poured into saturated aqueous NaHCO₃ (50 ml) and the layers were shaken and separated. The organic portion was washed with brine (50 ml), dried over Na₂SO₄ and concentrated *in vacuo*. The crude product mixture was purified by flash chromatography (10% ethyl acetate in hexanes) to afford 4.96 g (69%) of the title compound as colorless oil. [α]_D +14.8° (*c* 2.24, CHCl₃); IR (thin film): 1626, 1470, 1362, 1344, 1281, 1166 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 8.43 (s, 2H), 8.15 (s, 1H), 3.03-2.71 (m, 2H), 2.25 (d, *J* = 4.2 Hz, 1H), 1.51 (m, 1H), 0.96 (d, *J* = 6.8 Hz, 3H), 0.86 (d, *J* = 6.7 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 141.5, 133.0 (q, *J* = 34.4 Hz), 128.3, 126.9 (m), 122.4 (q, *J* = 271.7 Hz), 73.4, 47.3, 33.8, 30.1, 19.4, 18.9; HRMS *m/z* calcd for C₁₃H₁₄NO₂F₆S (M⁺ + H): 362.0649; found: 362.0640.

N-[(S)-N'-trifluoromethylsulfonyl-2-amino-3-methylbutyl]-N'-[3,5-bis(trifluoromethyl)benzenesulfonyl]-2-amino-3-methylbutyl-2,2,2-trifluoroethyl amine (6):

A mixture of 4.96 g of (*S*)-[3,5-bis(trifluoromethyl)benzenesulfonyl]-2-isopropylaziridine **10** (13.7 mmol, 1.0 equiv) and 3.29 ml of 2,2,2-trifluoroethylamine (41.2 mmol, 3.0 equiv) was placed in a sealed Pyrex[®] tube and heated at 50 °C for 12 h. The reaction was then cooled to ambient temperature and the remaining 2,2,2-trifluoroethylamine was evaporated *in vacuo*. The resulting crude (*S*)-N-[3,5-bis(trifluoromethyl)benzenesulfonyl]-N'-(2,2,2-trifluoroethyl)-2-isopropylethylene diamine **11** was combined with 3.43 g of (*S*)-N-trifluoromethylsulfonyl-2-isopropylaziridine (15.8 mmol, 1.2 equiv)⁵ and the mixture stirred at ambient temperature for 2 h then heated at 80 °C for 3 h. After cooling to ambient temperature, the crude product mixture was purified by flash chromatography (10% ethyl acetate in hexane) to afford 9.0 g (97%) of the title compound; crystallization of purified **6** from CH₂Cl₂/hexane afforded 7.68 g of **6** (83 %) as colorless needles: mp 99-100 °C. [α]_D -8.5° (*c* 2.02, CHCl₃); IR (thin film): 3302, 1361, 1281 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 8.33 (s, 2H), 8.09 (s, 1H), 6.08 (d, *J* = 8.9 Hz, 1H), 5.32 (d, *J* = 7.4 Hz, 1H), 3.73-3.48 (m, 3H), 3.21-2.99 (m, 3H), 2.93 (dd, *J* = 14.3, 8.0 Hz, 1H), 2.79 (dd, *J* = 14.0, 8.6 Hz, 1H), 2.07-1.87 (m, 2H), 1.05 (d, *J* = 6.8 Hz, 3H), 1.01 (d, *J* = 6.9 Hz, 3H), 0.81 (d, *J* = 6.9 Hz, 6H); ¹³C NMR (75 MHz, CDCl₃) δ 144.0, 133.0 (q, *J* = 34.4 Hz), 126.0 (q, *J* = 284.0 Hz), 127.2 (m), 126.2 (m), 122.5 (q, *J* = 271.7 Hz), 119.4 (q, *J* = 318.1 Hz), 59.3, 57.6, 55.5, 55.2, 53.7 (q, *J* = 29.0 Hz), 25.9, 29.7, 18.43, 18.39, 17.6, 17.3; HRMS *m/z* calcd for C₂₁H₂₈N₃O₄F₁₂S₂: 678.1330; found: 678.1339.

(3R, 4S)-3-Methyl-4-phenethyloxetan-2-one (9a): The General Procedure was followed employing 136 mg of ligand **6** (0.20 mmol, 20 mol%) and 132 μ L of hydrocinnamaldehyde (1.0 mmol). Successive purification by flash chromatography (2:1 CH₂Cl₂/hexane then 10% ethyl acetate in hexane) gave 135 mg (71%) of the title compound as a colorless oil. Separation of enantiomers by Chiral HPLC [Daicel ChiracelTM OD-H column, flow rate 1.0 mL/min, 5% *i*PrOH, 95% hexane, T_r: 13.7 min (3*S*, 4*R*), 14.5 min (3*R*, 4*S*) provided the enantiomer ratio (3*R*, 4*S*):(3*S*, 4*R*) = 94.5:5.5 (89% ee). [α]_D -42.3 ° (*c* 1.89, CHCl₃); IR (thin film): 1821, 1455cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.34-7.21 (m, 5H), 4.59 (ddd, *J* = 10.5, 6.4, 4.1 Hz, 1H), 3.76 (qd, *J* = 7.8, 6.5 Hz, 1H), 2.90 (ddd, *J* = 13.8, 9.4, 5.3 Hz, 1H), 2.72 (ddd, *J* = 13.8, 8.8, 7.5 Hz, 1H), 2.13-1.97 (m, 2H), 1.29 (d, *J* = 7.8 Hz, 3 H); ¹³C NMR (75 MHz, CDCl₃) δ 172.3, 140.4, 128.6, 128.4, 126.4, 74.6, 47.2, 31.9, 31.5, 8.1; HRMS *m/z* calcd for C₁₂H₁₄O₂: 190.0994; found: 190.0990.

⁵ Cernerud, M.; Adolfsson, H.; Moberg C. *Tetrahedron: Asymmetry* **1997**, 8, 2655-2662

(3R, 4S)-4-(9-Decenyl)-3-methyloxetan-2-one (9b): The General Procedure was followed employing 136 mg of ligand **6** (0.20 mmol, 20 mol%) and 208 μ L (1.0 mmol) of undecylenic aldehyde. Purification by flash chromatography (5% ethyl acetate in hexane) gave 173 mg (77%) of the title compound. Separation of the crude mixture of stereoisomers by chiral GLC [Chiraldex™ G-TA column 20 m x 0.25 mm, flow rate 1.0 mL/min, method: 100 °C for 10 min, ramp @ 5.0 °C/min to 130 °C for 12.0 min, ramp @ 10.0 °C/min to 160 °C for 20.0 min, T_r : 40.1 min and 41.0 (*trans*), T_r : 43.5 min (3*S*, 4*R*), T_r : 43.9 min (3*R*, 4*S*)] provided the enantiomer ratio (3*R*, 4*S*):(3*S*, 4*R*) = 93.5:6.5 (87% ee) and diastereomer ratio *syn:anti* = 94:6. $[\alpha]_D -15.1^\circ$ (*c* 1.61, CHCl₃); IR (thin film): 1824, 1640, 1460 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 5.78 (ddt, *J* = 16.9, 10.2, 6.7 Hz, 1H), 4.99 (d, *J* = 17.2 Hz, 1H), 4.93 (d, *J* = 10.2 Hz, 1H), 4.55 (ddd, *J* = 8.9, 6.4, 4.9 Hz, 1H), 3.73 (qd, *J* = 7.7, 6.5 Hz, 1H), 2.04 (q, *J* = 6.9 Hz, 2H), 1.72-1.65 (m, 2H), 1.55-1.45 (m, 1H), 1.45-1.27 (m, 14H); ¹³C NMR (75 MHz, CDCl₃) δ 172.6, 139.1, 114.1, 75.7, 47.2, 33.7, 30.0, 29.30, 29.27, 29.24, 29.0, 28.9, 25.4, 8.0; HRMS *m/z* calcd for C₁₄H₂₄O₂: 224.1776; found: 224.1780.

(3R, 4S)-4-(2-Benzyloxyethyl)-3-methyloxetan-2-one (9c): The General Procedure was followed employing 136 mg of ligand **6** (0.20 mmol, 20 mol%) and 164 mg (1.0 mmol) of 3-benzyloxypropionaldehyde. Purification by flash chromatography (15% ethyl acetate in hexane) gave 165 mg (75%) of the title compound. Separation of the enantiomers by chiral HPLC (Daicel Chiracel™ OD-H column, flow rate 1.0 mL/min, 10% ⁱPrOH, 90% hexane, T_r 10.8 (4*S*, 3*R*) and 16.5 (4*R*, 3*S*) min) provided the enantiomer ratio: (4*S*, 3*R*):(4*R*, 3*S*) = 95.5:4.5 (91% ee). $[\alpha]_D -40.4^\circ$ (*c* 1.88, CHCl₃); IR (thin film): 2866, 1824, 1455, 1111 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.39-7.30 (m, 5H), 4.81 (ddd, *J* = 8.4, 6.5, 5.1 Hz, 1H), 4.54 (s, 2H), 3.78 (qd, *J* = 7.7, 6.7 Hz, 1H), 3.67-3.62 (m, 2H), 2.10-2.00 (m, 2H), 1.29 (d, *J* = 7.7 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 172.4, 137.8, 128.3, 127.6, 127.5, 73.1, 72.7, 65.7, 47.2, 30.3, 8.1; HRMS *m/z* calcd for (M⁺-CO) C₁₂H₁₆O₂: 192.1150; found: 192.1156.

(3R, 4R)-3-Methyl-4-phenyloxetan-2-one (9d): The General Procedure was followed employing 68 mg of ligand **6** (0.10 mmol, 10 mol%) and 102 μ L (1.0 mmol) of benzaldehyde. Purification by flash chromatography (10% ethyl acetate in hexane) gave 130 mg (80%) of the title compound. Separation of the enantiomers by chiral HPLC (Daicel Chiracel™ OD-H column, flow rate 1.0 mL/min, 3% ⁱPrOH, 97% hexane, T_r 10.9 (3*R*, 4*R*) and 11.7 (3*S*, 4*S*) min) provided the enantiomer ratio: (3*R*, 4*R*):(3*S*, 4*S*) = 98:2 (96% ee). $[\alpha]_D +138^\circ$ (*c* 1.44, CHCl₃); IR (thin film): 2979, 1830, 1497, 1455, 1143, 1108 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.46-7.38 (m, 3H), 7.32-7.29 (m, 2H), 5.66 (d, *J* = 6.5 Hz, 1H), 4.06 (qd, *J* = 7.8, 6.5 Hz, 1H), 0.94 (d, *J* = 7.8 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 172.2, 134.6, 128.6, 128.5, 125.7, 75.2, 50.1, 9.5; HRMS *m/z* calcd for C₁₀H₁₀O₂: 162.0681; found: 162.0677.

(3R, 4R)-3-Methyl-4-(trimethylsilylethynyl)oxetan-2-one (9e): The General Procedure was followed employing 68 mg of ligand **6** (0.10 mmol, 10 mol%) and 143 μ L of trimethylsilylpropynal (1.0 mmol). Separation of the crude mixture of stereoisomers by chiral GC [Chiraldex™ G-TA column 20 m x 0.25 mm, flow rate 0.5 mL/min, method: 100 °C for 10 min, ramp @ 5.0 °C/min to 130 °C for 12.0 min, ramp @ 10.0 °C/min to 160 °C for 15.0 min, T_r : 14.0 min and 14.2 (*trans*), T_r : 16.5 min (3*R*, 4*R*), T_r : 16.7 min (3*S*, 4*S*)] provided the enantiomer ratio (3*R*, 4*R*):(3*S*, 4*S*) = 97.5:2.5 (95% ee) and diastereomer ratio *syn:anti* = 98:2. Due to the volatility of **9e** and the resulting difficulty in isolating the β -lactone, the crude product mixture was treated directly with morpholine in CH₂Cl₂ according to the literature procedure to afford 205 mg (76%) of morpholino-2-(*R*)-methyl-3-(*R*)-hydroxy-5-trimethylsilyl-4-pentynamide.⁶

(3R, 4S)-3-Ethyl-4-phenethyloxetan-2-one (9f): The General Procedure was followed employing 68 mg of ligand **6** (0.10 mmol, 10 mol%) and 132 μ L (1.0 mmol) of hydrocinnamaldehyde. Purification by flash chromatography (9% ethyl acetate in hexane) gave 165 mg (81%) of the title compound. Separation of the enantiomers by chiral HPLC

⁶ Nelson, S. G.; Wan, Z. *Org. Lett.* **2000**, 2, 1883-1886

(Daicel Chiralpak™ AD column, flow rate 0.7 mL/min, 10% *i*PrOH, 90% hexane, *T_r* 8.4 (3*S*, 4*R*) and 9.1(3*R*, 4*S*) min) provided the enantiomer ratio: (3*S*, 4*R*):(3*R*, 4*S*) = 4.5:95.5 (91% ee). [α]_D –48.8° (*c* 1.37, CHCl₃); IR (thin film): 2967, 1821, 1497, 1455, 1106 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.35-7.20 (m, 5H), 4.56 (ddd, *J* = 10.1, 6.4, 3.6 Hz, 1H), 3.55 (ddd, *J* = 8.8, 7.7, 6.4 Hz, 1H), 2.91 (ddd, *J* = 14.0, 9.4, 5.1 Hz, 1H), 2.71 (ddd, *J* = 14.1, 8.7, 7.7 Hz, 1H), 2.16-1.78 (m, 3H), 1.74-1.60 (m, 1H), 1.07 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 171.7, 140.4, 128.5, 128.4, 126.2, 74.4, 54.1, 31.9, 31.5, 17.3, 12.0; HRMS *m/z* calcd for C₁₃H₁₆O₂: 204.1150; found: 204.1158.

(3*R*, 4*S*)-4-(2-Benzyloxyethyl)-3-ethyloxetan-2-one (9g): The General Procedure was followed employing 68 mg of ligand **6** (0.10 mmol, 10 mol%) and 164 mg (1.0 mmol) of 3-benzyloxypropionaldehyde. Purification by flash chromatography (14% ethyl acetate in hexane) gave 194 mg (83%) of the title compound as an 88:12 *syn:anti* mixture. Separation of the enantiomers by chiral HPLC (Daicel Chiralpak™ OD-H column, flow rate 1.0 mL/min, 10% *i*PrOH, 90% hexane, *T_r* 8.5 (3*R*, 4*S*) and 13.3(3*S*, 4*R*) min) provided the enantiomer ratio: (3*R*, 4*S*):(3*S*, 4*R*) = 95.5:4.5 (91% ee). [α]_D –41.7° (*c* 2.06, CHCl₃); IR (thin film): 2969, 2877, 1824, 1496, 1455, 1116 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.39-7.29 (m, 5H), 4.80 (ddd, *J* = 9.8, 6.5, 4.0 Hz, 1H), 4.54 (s, 2H), 3.69-3.55 (m, 3H), 2.14-1.95 (m, 2H), 1.87-1.77 (m, 1H), 1.73-1.63 (m, 1H), 1.07(t, *J* = 7.4 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 171.7, 137.8, 128.2, 127.5, 127.4, 73.0, 72.4, 65.7, 54.0, 30.3, 17.3, 11.8; HRMS *m/z* calcd for (M⁺+H) C₁₄H₁₉O₃: 235.1334; found: 235.1329.

(3*R*, 4*R*)-4-(Benzyloxymethyl)-3-ethyloxetan-2-one (9h): The General Procedure was followed employing 68 mg of ligand **6** (0.10 mmol, 10 mol%) and 140 μ L (1.0 mmol) of benzyloxyacetaldehyde. Purification by flash chromatography (10% ethyl acetate in hexane) gave 172mg (78%) of the title compound. Separation of the enantiomers by chiral HPLC (Daicel Chiralpak™ OD-H column, flow rate 1.0 mL/min, 10% *i*PrOH, 90% hexane, *T_r* 9.8 (3*R*, 4*R*) and 19.3(3*S*, 4*S*) min) provided the enantiomer ratio: (3*R*, 4*R*):(3*S*, 4*S*) = 96.5:3.5 (93% ee). [α]_D –8.0° (*c* 2.0, CHCl₃); IR (thin film): 2970, 2878, 1828, 1455, 1372, 1112 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.40-7.29 (m, 5H), 4.71 (dd, *J* = 11.3, 5.5 Hz, 1H), 4.60 (s, 2H), 3.78 (d, *J* = 5.5 Hz, 2H), 3.64 (ddd, *J* = 8.8, 6.9, 6.9 Hz, 1H), 1.94-1.65 (m, 2H), 1.07 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 171.2, 137.3, 128.4, 127.8, 127.6, 73.6, 72.9, 68.0, 54.2, 17.3, 12.1; HRMS *m/z* calcd for C₁₃H₁₆O₃: 220.1099; found: 220.1109.

(3*R*, 4*R*)-3-Ethyl-4-phenyloxetan-2-one (9i): The General Procedure was followed employing 68 mg of ligand **6** (0.10 mmol, 10 mol%) and 102 μ L (1.0 mmol) of benzaldehyde. Purification by flash chromatography (9% ethyl acetate in hexane) gave 146 mg (83%) of the title compound. Separation of the enantiomers by chiral HPLC (Daicel Chiralpak™ OD-H column, flow rate 0.5 mL/min, 10% *i*PrOH, 90% hexane, *T_r* 13.3 (3*R*, 4*R*) and 16.1(3*S*, 4*S*) min) provided the enantiomer ratio: (3*R*, 4*R*):(3*S*, 4*S*) = 97:3 (94% ee). [α]_D +128° (*c* 2.08, CHCl₃); IR (thin film): 2970, 2938, 1825, 1497, 1455, 1111cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.45-7.32 (m, 5H), 5.64 (d, *J* = 6.4 Hz, 1H), 3.85 (ddd, *J* = 8.3, 8.3, 6.4, 1H), 1.57-1.41 (m, 1H), 1.35-1.21 (m, 1H), 0.85 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 171.6, 134.7, 128.6, 128.4, 125.8, 75.1, 56.8, 18.6, 11.1; HRMS *m/z* calcd for C₁₁H₁₂O₂: 176.0837; found: 176.0837.

(3*R*, 4*S*)-4-(2-Benzyloxyethyl)-3-propyloxetan-2-one (9j): The General Procedure was followed employing 68 mg of ligand **6** (0.10 mmol, 10 mol%) and 164 mg (1.0 mmol) of 3-benzyloxypropionaldehyde. Purification by flash chromatography (12% ethyl acetate in hexane) gave 218 mg (88%) of the title compound as a 91:9 *syn:anti* mixture. Separation of the enantiomers by chiral HPLC (Daicel Chiralpak™ OD-H column, flow rate 1.0 mL/min, 10% *i*PrOH, 90% hexane, *T_r* 8.0 (3*R*, 4*S*) and 12.4(3*S*, 4*R*) min) provided the enantiomer ratio: (3*R*, 4*S*):(3*S*, 4*R*) = 95.5:4.5 (91% ee). [α]_D –37.5° (*c* 1.78, CHCl₃); IR (thin film): 2961, 2873, 1817, 1455, 1115cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.39-7.29 (m, 5H), 4.80 (ddd, *J* = 9.5, 6.4, 4.0 Hz, 1H), 4.54 (s, 2H), 3.71-3.60 (m, 3H), 2.13-1.96 (m, 2H), 1.84-1.33(m, 4H), 0.95 (t, *J* = 7.3 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 171.7, 137.8, 128.1, 127.4, 127.3, 72.9, 72.3, 65.6, 52.1, 30.4, 25.7, 20.4, 13.5; HRMS *m/z* calcd for (M⁺–CO) C₁₄H₂₀O₂: 220.1463; found: 220.1466.

(3R, 4R)-4-Phenyl-3-propyloxetan-2-one (9k): The General Procedure was followed employing 68 mg of ligand **6** (0.10 mmol, 10 mol%) and 102 μ L (1.0 mmol) of benzaldehyde. Purification by flash chromatography (8% ethyl acetate in hexane) gave 161 mg (85%) of the title compound.

Separation of the enantiomers by chiral HPLC (Daicel Chiracel™ OD-H column, flow rate 0.5 mL/min, 10% *i*PrOH, 90% hexane, T_r 11.7 (3*R*, 4*R*) and 13.6(3*S*, 4*S*) min) provided the enantiomer ratio: (3*R*, 4*R*): (3*S*, 4*S*) = 98:2 (96% ee). $[\alpha]_D^{+116^\circ}$ (*c* 1.73, CHCl₃); IR (thin film): 2961, 2874, 1825, 1455, 1114 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.45-7.31 (m, 5H), 5.63 (d, *J* = 6.5 Hz, 1H), 3.93 (ddd, *J* = 8.5, 8.5, 6.5 Hz, 1H), 1.43-1.34 (m, 2H), 1.25-1.18 (m, 2H), 0.77 (t, *J* = 7.0 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 171.7, 134.7, 128.6, 128.4, 125.8, 75.1, 55.1, 27.0, 19.9, 13.5; HRMS *m/z* calcd for C₁₂H₁₄O₂: 190.0994; found: 190.0998.

(2R,3R)-3-Hydroxy-2-isopropyl-5-trimethylsilanylpent-4-ynoic acid, benzylamide [from (3*R*, 4*R*)-3-isopropyl-4-trimethylsilanylethynyl-oxetan-2-one (**9i**)]: The General Procedure was followed employing 68 mg of ligand **6**

(0.10 mmol, 10 mol%) and 134 μ L (1.0 mmol) of trimethylsilanylpropynal. Due to the volatility of **9i** and the resulting difficulty in isolating the β -lactone, the crude product mixture was dissolved in 1 mL of CH₃CN and 109 μ L of benzylamine (1.0 mmol) and DMAP (~1 mg) were added at ambient temperature. The resulting solution was stirred at ambient temperature for 15 h prior to evaporating the volatiles *in vacuo*. The crude product mixture was purified by flash chromatography (20% ethyl acetate in hexane) to afford 225 mg of (2*R*,3*R*)-3-hydroxy-2-isopropyl-5-trimethylsilanylpent-4-ynoic acid, benzylamide (71%). Separation of the enantiomers by chiral HPLC (Daicel Chiracel™ OD-H column, flow rate 1.0 mL/min, 5% *i*PrOH, 95% hexane, T_r 7.4 (2*S*, 3*S*) and 8.9(2*R*, 3*R*) min) provided the enantiomer ratio: (2*R*, 3*R*):(2*S*, 3*S*) = 97:3 (94% ee). $[\alpha]_D^{+50.3^\circ}$ (*c* 1.58, CHCl₃); IR (thin film): 3379, 3312, 2960, 2932, 2171, 1650, 1539, 1250, 1045, 845 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.36-7.30 (m, 5H), 6.38 (br s, 1H), 4.73 (t, *J* = 4.9 Hz, 1H), 4.48 (d, *J* = 5.5 Hz, 2H), 3.19 (d, *J* = 4.6 Hz, 1H), 3.32 (dd, *J* = 8.5, 5.4 Hz, 1H), 2.15 (m, 1H), 1.04 (t, *J* = 7.0 Hz, 6H), 0.10 (s, 9H); ¹³C NMR (75 MHz, CDCl₃) δ 172.7, 138.0, 128.5, 127.7, 127.3, 104.4, 91.6, 62.3, 59.1, 43.4, 28.1, 21.1, 20.6, -0.5; HRMS *m/z* calcd for C₁₈H₂₇NO₂Si: 317.1811; found: 317.1808.

(3R, 4R)-3-Isopropyl-4-phenyloxetan-2-one (9m): The General Procedure was followed employing 68 mg of ligand **6** (0.10 mmol, 20 mol%) and 51 μ L (0.50 mmol) of benzaldehyde.

Purification by flash chromatography (7% ethyl acetate in hexane) gave 80 mg (84%) of the title compound. Separation of the enantiomers by chiral HPLC (Daicel Chiracel™ OD-H column, flow rate 0.5 mL/min, 10% *i*PrOH, 90% hexane, T_r 11.1 (3*R*, 4*R*) and 14.1(3*S*, 4*S*) min) provided the enantiomer ratio: (3*R*, 4*R*): (3*S*, 4*S*) = 98:2 (96% ee). $[\alpha]_D^{+89.2^\circ}$ (*c* 1.50, CH₃Cl₃); IR (thin film): 2960, 1808, 1454, 1368, 1120, 881 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.42-7.39 (m, 5H), 5.60 (d, *J* = 6.4 Hz, 1H), 3.60 (dd, *J* = 11.4, 6.4 Hz, 1H), 1.82 (m, 1H), 1.07 (d, *J* = 6.6 Hz, 3H), 0.53 (d, *J* = 6.5 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 171.1, 134.6, 128.9, 128.4, 126.7, 75.5, 62.5, 25.2, 20.2, 19.9; HRMS *m/z* calcd for C₁₂H₁₄O₂: 190.0994; found: 190.0987.