Catalytic Asymmetric Assembly of Stereodefined Propionate Units: An Enantioselective Total Synthesis of (–)-Pironetin

Supporting Information

Xiaoqiang Shen, Andrew S. Wasmuth, Junping Zhao, Cheng Zhu, and Scott G. Nelson*

Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260

TABLE OF CONTENTS

Compounds 8-11 , 13a , 13b , 14a , 14c , 16 , 18 , 19-26 , 28 , 29	page S1-S10
Stereochemical proofs (X-ray data) for compounds 14c and 16	page S11-S12
¹ H and ¹³ C Spectra for compounds 8 , 9-11 , 13a , 13b , 14a , 14c , 16 , 18 , 19-26 , 28 , 29	page S13-S60

General Information: Unless otherwise stated, all reactions were carried out in dry glassware under a nitrogen atmosphere using standard inert atmosphere (N₂) techniques for the manipulation of solvents and reagents. Proton NMR chemical shifts are reported relative to residual CHCl₃ (7.27 ppm) for ¹H and CDCl₃ (77.00 ppm) for ¹³C NMR spectra. Analytical high performance liquid chromatography (HPLC) was performed using a variable wavelength UV detector (deuterium lamp, 190-600 nm) using a Daicel ChiracelTM OD-H column (250 x 4.6 mm) (Daicel Inc.) and HPLC-grade isopropanol and hexanes as the eluting solvents. Anhydrous solvents (CH₂Cl₂, THF, diethyl ether, pentane, *N*, *N*-dimethylformamide and toluene) were obtained by passage through successive alumina- and Q5 reactant-packed columns on a solvent purification system. *N*, *N*-Diisopropylethylamine and triethylamine were distilled under nitrogen from CaH₂. The enantioenriched aldehydes **12a-c** and **17** were prepared from the corresponding β-lactones using the three-step sequence described for converting β-lactone **8** to aldehyde **9**.^{1,2} Aldehydes **12a-c**, **15**, and **17** were prepared with the indicated stereochemical purities: **12a** (99% ee), **12b** (99% ee), **12c** (99% ee), **15** (95% ee), **17** (>97% de).³ Catalyst **27** was prepared according to the literature procedure.⁴ Anhydrous LiClO₄ and LiI were purchased from Aldrich Chemical Co. and weighed out in a N₂-filled glovebox. *O*-Trimethylsilylquinidine (**3a**, TMSQd)⁵ and *O*-trimethylsilylquinine (**4a**, TMSQn),⁵ [(C₈H₁₄)₂IrCl]₂⁶ and 3-*p*-methoxybenzyloxypropanal⁷ were prepared according to literature procedures. Commercially available propionyl chloride was redistilled under N₂. Butyryl bromide was prepared according to literature procedure and was distilled from P₂O₅.⁸ Flash chromatography was performed as previously described on EM silica gel 60 (230-240 mesh).⁹

(3R, 4S)-4-[(S)-1-(4-Methoxybenzyloxy)propan-2-yl]-3-methyloxetan-2-one (8**):** To a –78 °C solution of *O*-trimethylsilylquinine (**4a**) (40 mg, 0.1 mmol) and LiClO₄ (108 mg, 1.0 mmol) in 2:1 CH₂Cl₂:Et₂O (3 mL) was added *N*, *N*-diisopropylethylamine (0.44 mL, 2.5 mmol) and (S)-3-(4-methoxybenzyloxy)-2-methylpropanal (210 mg, 1.0 mmol). To the resulting mixture

¹ Nelson, S. G.; Cheung, W. S.; Kassick, A. J.; Hilfiker, M. A. *J. Am. Chem. Soc.* **2002**, 124, 13654.

² For aldehyde **12c**, the *p*-methoxybenzyl ether was introduced according to the published procedure: Clark, D. L.; Heathcock, C. H. *J. Org. Chem.* **1993**, 58, 5878.

³ Optical purities for aldehydes **12a-c** and **15** are based on those for the β-lactone precursors assayed by chiral GC, see: Zhu, C.; Shen, X.; Nelson, S. G. *J. Am. Chem. Soc.* **2004**, 126, 5352. The stereochemical purity of aldehyde **17** was assayed by integrating the appropriate resonances in the 500 MHz ¹H NMR.

⁴ Nelson, S. G.; Zhu, C.; Shen, X. *J. Am. Chem. Soc.* **2004**, 126, 14.

⁵ Calter, M. A. *J. Org. Chem.* **1996**, 61, 8006.

⁶ Onderdelinden, A. L.; Van der Ent, A. *Inorg. Chim. Acta* **1972**, 6, 420.

⁷ Zampella, A.; Sepe, V.; D'Orsi, R.; Bifulco, G.; Bassarello, C.; D'Auria, V. *Tetrahedron Asymm.* **2003**, 14, 1787.

⁸ Burton, T. M.; Degering, E. F. *J. Am. Chem. Soc.* **1940**, 62, 227.

⁹ Still, W. C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, 43, 2923.

was added a solution of propionyl chloride (0.17 mL, 2.0 mmol) in 0.5 mL CH₂Cl₂ over 2 h by syringe pump. The reaction was stirred 16 h at -78 °C whereupon Et₂O (10 mL) was added and the reaction was warmed to ambient temperature. The resulting mixture was filtered through a silica gel plug eluting with Et₂O (100 mL). ¹H NMR analysis of the crude product revealed the diastereomer ratio to be 92:8 *syn*, *anti*: *syn*:*syn*. The filtrate was concentrated *in vacuo* and the crude product mixture was purified by flash chromatography (SiO₂, 21% ethyl acetate in hexanes) giving 195 mg (74%) of the title compound as light yellow oil. [α]_D -29 (c 0.31, CHCl₃); IR (thin film): 1823, 1613, 1513, 1463, 1247, 1117, 986 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 1.09 (d, J=6.7 Hz, 3H), 1.42 (d, J=7.7 Hz, 3H), 2.13-2.21 (m, 1H), 3.56 (dd, J=9.0, 5.7 Hz, 1H), 3.64 (dd, J=9.0, 3.1 Hz, 1H), 3.80-3.89 (m, 1H), 3.89 (s, 3H), 4.51 (dd, J=11.0, 6.3 Hz, 1H), 4.54 (s, 2H), 6.96 (d, J=8.5 Hz, 2H), 7.53 (d, J=8.4 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃): δ 8.4, 13.0, 34.2, 46.9, 55.2, 70.8, 72.9, 76.2, 113.7, 129.1, 130.3, 159.1, 172.4; MS (EI, 70 eV): *m/z* 264, 152, 137, 121, 78; HRMS *m/z* calcd for C₁₅H₂₀O₄ (M⁺): 264.1362; found: 264.1360.

(2R, 3S, 4S)-N-Methoxy-N-methyl-3-hydroxy-5-(4-methoxybenzyloxy)- 2, 4-dimethylpentanamide (8'): To a 0 °C solution of *N*, *O*-dimethylhydroxyamine hydrochloride (1.64 g, 16.8 mmol) in 30 mL of CH₂Cl₂ was added Me₂AlCl (1.0 M in

hexane, 16.8 mL, 16.8 mmol) added dropwise. Once addition was complete, the reaction was warmed to ambient temperature and stirred for 60 min. The reaction was cooled to -30 °C and a solution of lactone **7** (2.23 g, 8.40 mmol) in 10 mL of CH₂Cl₂ was added over 60 min via syringe pump. Once addition was complete, the reaction was stirred for 60 min then 100 mL of saturated aqueous sodium tartrate was added. The resulting gel like mixture was warmed to ambient temperature and stirred vigorously until two homogeneous phases resulted. The mixture was diluted with 100 mL of CH₂Cl₂ and the layers were shaken and separated. The aqueous portion was extracted with CH₂Cl₂ (3×50 mL) and the combined organic extracts were dried (Na₂SO₄) and concentrated. The crude product mixture was purified by column chromatography (SiO₂, gradient 10→50% EtOAc in hexanes) to afford 2.66 g (80%) of the title compound as colorless oil. [α]_D = -13 (c 1.8, CHCl₃); IR ν (thin film) 3458, 1636, 1586, 1513, 1463, 1247, 1032, 820 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 0.98 (d, J=6.9, 3H), 1.18 (d, J=7.0 Hz, 3H), 1.82-1.94 (m, 1H), 3.04(broad s, 1H), 3.19 (s, 3H), 3.53 (dd, J=9.0, 5.8 Hz, 1H), 3.63 (dd, J=9.1, 4.0 Hz, 1H), 3.67 (s, 3H), 3.68-3.73(m, 2H), 3.80 (s, 3H), 4.45 (s, 2H), 6.87 (d, J=8.4 Hz, 2H), 7.26 (d, J=8.4 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃): δ 10.4, 14.2, 32.0, 36.0, 36.4, 55.2, 61.4, 72.7, 72.9, 73.8, 113.7, 129.2, 130.6, 159.0, 176.6; MS (ESI): *m/z* 348 (M⁺+Na); HRMS *m/z* calcd for C₁₇H₂₇NO₅Na (M⁺+Na): 348.1787; found: 348.1777.

(2R, 3S, 4S)-N-Methoxy-N-methyl-5-(4-methoxybenzyloxy)-3-trimethylsiloxy-2, 4-dimethylpentanamide (8''). To a -78 °C solution of alcohol **8'** (665 mg, 2.00 mmol) in 18 mL of CH₂Cl₂ containing 2,6-lutidine (0.35 mL, 3.0 mmol) was added TMSOTf

(0.47 mL, 2.6 mmol). The reaction was stirred 30 min then warmed to ambient temperature and stirred an additional 2.5 h. The reaction was quenched by adding aqueous 1 N NaHSO₄ (40 mL) and the resulting mixture was extracted with CH₂Cl₂ (50 mL). The aqueous portion was extracted with CH₂Cl₂. (2×30 mL) and the combined organic extracts dried (Na₂SO₄) and concentrated. The crude product mixture was purified by column chromatography (SiO₂, 20% EtOAc in hexanes) to afford 689 mg (86%) of the title compound as a colorless oil. [α]_D -11 (c 1.2, CHCl₃); IR ν (thin film) 1660, 1613, 1513, 1461, 1249, 1082, 839 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 0.07 (s, 9H), 0.95 (d, J=6.9 Hz, 3H), 1.06 (d, J=6.9 Hz, 3H), 1.79-1.90 (m, 1H), 3.08 (s, 3H), 3.08-3.18 (m, 2H), 3.53 (s, 3H), 3.53-3.57 (m, 1H), 3.84 (dd, J=7.8, 3.5 Hz, 1H), 4.35 (s, 2H), 6.81 (d, J=8.6 Hz, 2H), 7.19 (d, J=8.6 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃): δ 0.7, 14.3, 16.0, 32.2, 37.8, 39.4, 55.3, 61.3, 71.8, 72.6, 77.1, 113.7, 129.2, 131.0, 159.0, 176.6; MS (ESI): *m/z* 420 (M⁺+Na); HRMS *m/z* calcd for C₂₀H₃₅NO₅NaSi (M⁺+Na): 420.2182; found: 420.2172.

(2R, 3S, 4S)-5-(4-Methoxybenzyloxy)-3-trimethylsiloxy-2,4-dimethylpentanal (9): A hexanes solution of *i*Bu₂AlH (1.0 M, 2 mL, 2.0 mmol) was added to a -78 °C solution of Weinreb amide **8''** (588 mg, 1.5 mmol) in 10 mL THF. After being stirred 3 h, the reaction

was quenched at -78 °C by adding 40 mL of an aqueous, saturated sodium tartrate solution. The reaction was warmed to ambient temperature affording a gel-like mixture that was stirred until two homogeneous layers resulted (~30 min). The mixture was diluted with Et₂O (50 mL) and the layers were separated. The aqueous

portion was extracted with Et₂O (3 x 20 mL) and the combined organic portions were dried and concentrated. The crude product mixture was purified by flash chromatography (SiO₂, 10% ethyl acetate in hexane) giving 438 mg (80%) of the title compound as colorless oil. [α]_D -32 (c 0.34, CHCl₃); IR (thin film): 1725, 1613, 1513, 1463, 1249, 1032, 841 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 0.07 (s, 9H), 0.96 (d, J=6.9 Hz, 3H), 1.08 (d, J=7.0 Hz, 3H), 1.87-1.95 (m, 1H), 2.52 (qd, J=6.9, 3.4 Hz, 1H), 3.37 (dd, J=9.0, 5.7 Hz, 1H), 3.46 (dd, J=9.0, 4.5 Hz, 1H), 3.81 (s, 3H), 4.16 (dd, J=7.6, 3.4 Hz, 1H), 4.38 (d, J=11.5 Hz, 1H), 4.45 (d, J=11.5 Hz, 1H), 6.88 (d, J=8.5 Hz, 2H), 7.25 (d, J=8.5 Hz, 2H), 9.69 (s, 1H); ¹³C NMR (75 MHz, CDCl₃): δ 0.5, 7.5, 14.6, 37.6, 49.8, 55.2, 71.6, 72.7, 72.8, 113.7, 129.2, 130.6, 159.1, 205.0; MS (ESI): *m/z* 361 (M⁺+Na); HRMS (ESI) *m/z* calcd for C₁₈H₃₀O₄NaSi (M⁺+Na): 361.1811; found: 361.1823

(3S, 4R)-4-[(3S, 4S)-5-(4-Methoxybenzyloxy)-3-trimethylsiloxy-4-methylpentan-2-yl]-3-methyloxetan-2-one (10): To a solution of *O*-trimethylsilylquinidine (**3a**) (20 mg, 0.05 mmol) and LiI (134 mg, 1.0 mmol) in 0.13 mL of Et₂O and 1.0 mL CH₂Cl₂ was added *N*, *N*-diisopropylethylamine (0.22 mL, 1.25 mmol) and the solution was cooled to -40 °C. A

solution of aldehyde **9** (170 mg, 0.5 mmol) in 0.3 mL of CH₂Cl₂ was added to the reaction whereupon a solution of propionyl chloride (0.09 mL, 1.0 mmol) in 0.4 mL CH₂Cl₂ was added via syringe pump over 90 min. Once addition was complete, the reaction was stirred at -40 °C for 16 h at which time the reaction was diluted with 5 mL of Et₂O. The resulting mixture was diluted with 0.1N HCl (20 mL) and CH₂Cl₂ (20 mL) and the layers were shaken and separated. The aqueous portion was extracted with CH₂Cl₂ (2x15 mL) and the combined organic portions were washed with saturated NaHCO₃ and brine, dried and concentrated. The crude product mixture was purified by flash chromatography (SiO₂, 12% ethyl acetate in hexane) to afford 166 mg (84%) of the title compound as colorless oil. ¹H NMR analysis of the crude product revealed the diastereomer ratio to be 97:3 *syn, anti, syn, anti: anti, anti, syn, anti*. [α]_D +10 (c 0.84, CHCl₃); IR (thin film): 1827, 1613, 1513, 1463, 1249, 1048, 734 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 0.10 (s, 9H), 0.80 (d, J=6.7 Hz, 3H), 0.94 (d, J=6.8 Hz, 3H), 1.31 (d, J=7.7 Hz, 3H), 1.83-1.91 (m, 1H), 1.95-2.00 (m, 1H), 3.31 (dd, J=8.6, 6.9 Hz, 1H), 3.50 (dd, J=8.7, 3.5 Hz, 1H), 3.80-3.74 (m, 2H), 3.80 (s, 3H), 4.35 (dd, J=11.1, 6.3 Hz, 1H), 4.38 (d, J=11.5 Hz, 1H), 4.48 (d, J=11.5 Hz, 1H), 6.88 (d, J=8.5 Hz, 2H), 7.26 (d, J=8.5 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃): δ 0.7, 7.8, 8.5, 14.4, 35.3, 37.6, 46.6, 55.3, 72.2, 72.6, 72.8, 76.2, 113.8, 129.3, 130.8, 159.2, 172.7; MS (ESI): *m/z* 417 (M⁺+Na); HRMS (ESI) *m/z* calcd for C₂₁H₃₄O₅NaSi (M⁺+Na): 417.2073; found: 417.2058

(3R, 4R)-4-[(3S, 4S)-5-(4-Methoxybenzyloxy)-3-trimethylsiloxy-4-methylpentan-2-yl]-3-methyloxetan-2-one (11): To a solution of *O*-methylquinidine (**4b**) (17 mg, 0.05 mmol) and LiI (167 mg, 1.25 mmol) in 0.13 mL of Et₂O and 1.0 mL CH₂Cl₂ was added *N*, *N*-diisopropylethylamine (0.22 mL, 1.25 mmol) and the solution was cooled to -40 °C. A

solution of aldehyde **9** (170 mg, 0.5 mmol) in 0.3 mL of CH₂Cl₂ was added to the reaction whereupon a solution of propionyl chloride (0.9 mL, 1.0 mmol) in 0.4 mL CH₂Cl₂ was added via syringe pump over 90 min. Once addition was complete, the reaction was stirred at -40 °C for 16 h at which time the reaction was diluted with 5 mL of Et₂O. The resulting mixture was diluted with 0.1 N HCl (20 mL) and CH₂Cl₂ (20 mL) and the layers were shaken and separated. The aqueous portion was extracted with CH₂Cl₂ (2x15 mL) and the combined organic portions were washed with saturated NaHCO₃ and brine, dried and concentrated. The crude product mixture was purified by flash chromatography (SiO₂, 12% ethyl acetate in hexane) to afford 142 mg (72%) of the title compound as colorless oil. ¹H NMR analysis of the crude product revealed the diastereomer ratio to be 96:4 *anti, anti, syn, anti: syn, anti, syn, anti*. [α]_D +28 (c 0.73, CHCl₃); IR (thin film): 1827, 1613, 1513, 1463, 1249, 1048, 734 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 0.12 (s, 9H), 0.84 (d, J=6.9 Hz, 3H), 0.93 (d, J=6.9 Hz, 3H), 1.41 (d, J=7.5 Hz, 3H), 1.79-1.90 (m, 2H), 3.22 (qd, J=7.5, 4.0 Hz, 1H), 3.33 (dd, J=8.7, 6.1 Hz, 1H), 3.49 (dd, J=8.7, 3.6 Hz, 1H), 3.79-3.81 (m, 1H), 3.81 (s, 3H), 3.97 (dd, J=10.3, 4.0 Hz, 1H), 4.39 (d, J=11.6 Hz, 1H), 4.49 (d, J=11.6 Hz, 1H), 6.88 (d, J=8.6 Hz, 2H), 7.26 (d, J=8.6 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃): δ 0.7, 7.4, 13.1, 14.4, 37.6, 40.6, 50.3, 55.3, 72.1, 72.5, 72.8, 80.3, 113.8, 129.2, 130.8, 159.2, 172.0; MS (ESI): *m/z* 417 (M⁺+Na); HRMS (ESI) *m/z* calcd for C₂₁H₃₄O₅NaSi (M⁺+Na): 417.2073; found: 417.2063.

(3S, 4R)-4-[(2R, 3S)-5-Phenyl-3-trimethylsilyloxypentan-2-yl]-3-methyloxetan-2-one (13a): A flame-dried 25 mL round bottom flask was charged with *O*-trimethylsilylquinidine (**3a**) (45 mg, 0.11 mmol) and LiI (0.456 g, 3.41 mmol). To the

mixture was added CH_2Cl_2 (3.3 mL), DMF (0.3 mL) and $^i\text{Pr}_2\text{NEt}$ (0.50 mL, 2.84 mmol). The resulting mixture was stirred at ambient temperature for 5 min then cooled to $-50\text{ }^\circ\text{C}$. To the reaction mixture was added aldehyde **12a** (300 mg, 1.14 mmol) whereupon a solution of propionyl chloride (0.20 mL, 2.27 mmol) in CH_2Cl_2 (0.76 mL) was added via syringe pump over 2 h. The heterogeneous reaction mixture was stirred vigorously overnight at $-50\text{ }^\circ\text{C}$. The resulting mixture was partitioned between ether (15 mL) and H_2O (15 mL) and the layers were shaken and separated. The aqueous portion was extracted with ether (2 $\times$) and the combined organic portions were washed with water (2 $\times$) and brine. The combined organics were dried (Na_2SO_4), filtered and concentrated and the crude residue was purified by column chromatography (SiO_2 , 4% ethyl acetate in hexanes) to yield 0.300 g (83%) of β -lactone **13a**. ^1H NMR (300 MHz) analysis of the crude product revealed **13a** to be a single diastereomer (*syn, anti, syn*) within the limits of detection. $[\alpha]_D^{25} = +28$ (*c* 1.17, CHCl_3); IR (thin film) 2952, 1827, 1250, 1148, 1118, 1044, 1001 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 7.33–7.18 (m, 5H), 4.40 (dd, *J* = 11.0, 6.3 Hz, 1H), 4.01 (ddd, *J* = 6.6, 6.6, 0.9 Hz, 1H), 3.77 (dddd, *J* = 7.6, 7.6, 7.6, 7.6 Hz, 1H), 2.67 (ddd, *J* = 13.5, 10.8, 6.0 Hz, 1H), 2.54 (ddd, *J* = 13.5, 10.7, 6.0 Hz, 1H), 1.91–1.70 (m, 3H), 1.33 (d, *J* = 7.7 Hz, 3H), 0.85 (d, *J* = 6.6 Hz, 3H), 0.14 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): δ 172.4, 141.7, 128.3 (2C), 128.1 (2C), 125.7, 75.9, 70.1, 46.3, 37.2, 37.0, 32.5, 8.3, 7.4, 0.3 (3C); EI-MS *m/z* 320 (M^+), 305, 249, 230, 207; HRMS *m/z* calcd for $\text{C}_{18}\text{H}_{28}\text{O}_3\text{Si}$: 320.1808, found: 320.1804.

(3S, 4R)-4-[(2R, 3S)-3-(Benzyloxy)-5-phenylpentan-2-yl]-3-methyl-oxetan-2-one (13b):

A flame-dried 25 mL round bottom flask was charged with *O*-trimethylsilylquinidine (**3a**) (41 mg, 0.10 mmol) and LiClO_4 (0.320 g, 3.00 mmol). To the mixture was added CH_2Cl_2 (2.8 mL), THF (0.3 mL) and $^i\text{Pr}_2\text{NEt}$ (0.44 mL, 2.5 mmol). The resulting mixture was stirred at ambient temperature for 5 min then cooled to $-40\text{ }^\circ\text{C}$. To the reaction mixture was added aldehyde **12b** (282 mg, 1.00 mmol) whereupon a solution of propionyl chloride (0.176 mL, 2.00 mmol) in CH_2Cl_2 (0.70 mL) was added via syringe pump over 4 h. The heterogeneous reaction mixture was stirred vigorously overnight at $-40\text{ }^\circ\text{C}$. The reaction was diluted ether (20 mL) and the resulting mixture was filtered through a silica gel plug eluting with additional ether. The filtrate was concentrated and the crude residue was purified by column chromatography (SiO_2 , 5:1 hexanes:ethyl acetate) to yield 0.257 g (76%) of β -lactone **13b** as a colorless oil. IR (thin film): 2943, 1824, 1496, 1454, 1382, 1120 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.46–7.25 (m, 10 H), 4.74 (d, *J* = 11.5 Hz, 1 H), 4.66–4.60 (m, 2 H), 3.90–3.75 (m, 2 H), 2.87 (m, 2 H), 2.23–2.03 (m, 2 H), 1.94–1.82 (m, 1 H), 1.39 (d, *J* = 7.8 Hz, 3 H), 1.01 (d, *J* = 6.8 Hz, 3 H); ^{13}C NMR (75 MHz, CDCl_3) δ 172.2, 141.6, 138.6, 128.2, 128.1, 128.0, 127.3, 127.2, 125.7, 77.2, 76.0, 72.7, 46.3, 36.7, 33.7, 32.2, 8.1 (note: one carbon missing; most likely hidden under CDCl_3); HRMS *m/z* calcd for $\text{C}_{22}\text{H}_{26}\text{O}_3$: 338.1882; found: 338.1882.

(3R, 4S)-4-[(2R, 3S)-5-Phenyl-3-trimethylsilyloxypentan-2-yl]-3-methyloxetan-2-one (14a):

To a $-78\text{ }^\circ\text{C}$ mixture of 40 mg (0.10 mmol, 0.1 equiv) of *O*-trimethylsilylquinine (**4a**) and 201 mg (1.5 mmol, 1.5 equiv) of LiI in 2.0 mL of CH_2Cl_2 and 0.25 mL of Et_2O was added 0.44 mL (2.5 mmol, 2.5 equiv) of *N, N*-diisopropylethylamine followed by a solution of 264 mg (1.0 mmol) of aldehyde **12a** in 0.75 mL of CH_2Cl_2 . A solution of 0.175 mL (2.0 mmol, 2.0 eq) of propionyl chloride in 0.5 mL of CH_2Cl_2 was added to the reaction mixture over 1 h via syringe pump. The reaction mixture was stirred at $-78\text{ }^\circ\text{C}$ for 16 h and then diluted with 10 mL of ether. The resulting mixture was filtered through a silica gel plug eluting with Et_2O (3 $\times$ 20 mL) and the filtrate was concentrated. ^1H NMR (300 MHz) analysis of the crude product to be a 95.5:4.5 *anti, anti, syn:syn, anti, syn* mixture of diastereomers. Purifying the crude residue by flash chromatography (SiO_2 , 20% ethyl acetate in hexane) afforded 0.259 g (81%) of the title compound as colorless oil. $[\alpha]_D^{25} +42.9$ (*c* 1.06, CHCl_3); IR (thin film): 2954, 1828, 1455, 1251, 1125, 1067, 1023 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.35–7.20 (m, 5 H), 4.07 (dd, *J* = 9.7, 4.1 Hz, 1 H), 4.00 (apparent dt, *J* = 6.6, 2.2 Hz, 1 H), 3.28 (dq, *J* = 7.5, 4.1 Hz, 1 H), 2.73–2.50 (m, 2 H), 1.93–1.75 (m, 3 H), 1.43 (d, *J* = 7.5 Hz, 3 H), 0.92 (d, *J* = 6.9 Hz, 3 H), 0.18 (s, 9 H); ^{13}C NMR (75 MHz, CDCl_3) δ 171.8, 141.7, 128.3, 128.1, 125.8, 80.0, 70.3, 49.7, 42.1, 36.8, 32.3, 12.8, 7.3, 0.4; HRMS *m/z* calcd for $\text{C}_{18}\text{H}_{28}\text{O}_3\text{Si}$: 320.1808; found: 320.1797.

(3R, 4R)-4-[(2R, 3S)-3-(4-Methoxybenzyloxy)-5-phenylpentan-2-yl]-3-methyloxetan-2-one (14c):

To a $-78\text{ }^\circ\text{C}$ mixture of 40 mg of *O*-trimethylsilylquinine (**4a**) (0.10 mmol, 0.1

eq) and 201 mg (1.5 mmol, 1.5 eq) of LiI in 2.0 mL of CH₂Cl₂ and 0.25 mL of Et₂O was added 0.44 mL (2.5 mmol, 2.5eq) of *N,N*-diisopropylethylamine followed by a solution of 312 mg (1.0 mmol) of aldehyde **12c** in 0.75 mL of CH₂Cl₂. A solution of 0.175 mL (2.0 mmol, 2.0 eq) of propionyl chloride in 0.5 mL of CH₂Cl₂ was added to the reaction mixture over 1 h via syringe pump. The reaction mixture was stirred at -78 °C for 16 h and then diluted with 10 mL of ether. The resulting mixture was filtered through a silica gel plug eluting with Et₂O (3×20 mL) and the filtrate was concentrated. Separation of the diastereomers by HPLC [Zorbex column, flow rate 1.0 mL/min, 20% EtOAc, 80% hexane, T_r 6.4 (*trans*) and 7.2 (*cis*) min] provided the diastereomer ratio: (*trans*):(*cis*) = 23 : 1. Purification by flash chromatography (20% ethyl acetate in hexane) afforded 0.300 g (81%) of the title compound as colorless oil: [α]_D +53.1 (*c* 1.68, CHCl₃); IR (thin film): 1824, 1612, 1514, 1456, 1248 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.44-7.30 (m, 7H), 7.03 (app. d, *J* = 8.7 Hz, 2H), 4.69 (d, *J* = 11.1 Hz, 1H), 4.57 (d, *J* = 11.1 Hz, 1H), 4.31 (dd, *J* = 9.2, 4.1 Hz, 1H), 3.96 (s, 3H), 3.81 (td, *J* = 6.6, 2.6 Hz, 1H), 3.41 (qd, *J* = 7.5, 4.1 Hz, 1H), 2.83 (ddd, *J* = 13.7, 10.5, 5.9 Hz, 1H), 2.73 (ddd, *J* = 13.3, 10.3, 6.0 Hz, 1H), 2.18-2.06 (m, 2H), 1.96-1.84 (m, 1H), 1.53 (d, *J* = 7.5 Hz, 3H), 1.10 (d, *J* = 7.0 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 171.9, 159.3, 141.8, 130.8, 129.3, 128.5, 128.3, 126.0, 113.9, 80.3, 77.4, 72.3, 55.3, 49.7, 41.1, 33.5, 32.3, 12.9, 8.1; HRMS *m/z* calcd for C₂₃H₂₈O₄: 368.1988; found: 368.1972.

(3R, 4S)-4-[(2S, 3S)-5-phenyl-3-trimethylsilyloxypentan-2-yl]-3-methyloxetan-2-one

(16): A flame-dried 25 mL round bottom flask was charged with *O*-trimethylsilylquinine (**4a**) (30 mg, 0.076 mmol) and LiI (304 mg, 2.27 mmol). To the mixture was added CH₂Cl₂ (2.1 mL), THF (0.20 mL) and ^tPr₂NEt (0.330 mL, 1.90 mmol). The resulting

mixture was stirred at ambient temperature for 5 min then cooled to -78 °C. To the reaction mixture was added aldehyde **15** (200 mg, 0.76 mmol) whereupon a solution of propionyl chloride (0.130 mL, 1.51 mmol) in CH₂Cl₂ (0.5 mL) was added via syringe pump over 4 h. The heterogeneous reaction mixture was stirred vigorously overnight at -78 °C. The reaction mixture was diluted with ether (4 mL) and the resulting white precipitate was removed by filtration through a silica gel plug eluting with ether (80 mL). The filtrate was concentrated and the crude residue was purified column chromatography (SiO₂, 24:1/hexanes:ethyl acetate) to yield 197 mg (81%) of β-lactone **16** as a colorless solid. ¹H NMR (300 MHz) analysis of the crude product revealed the product lactone to be a single diastereomer (*syn*, *anti*, *anti*) within the limits of detection. m.p. = 76-77 °C; [α]_D = -49 (*c* 1.3, CHCl₃); IR (CH₂Cl₂) 2954, 1826, 1250, 1118, 1087, 1066 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 7.32-7.17 (m, 5H), 4.36 (dd, *J* = 11.3, 6.3 Hz, 1H), 3.97-3.92 (m, 1H), 3.78 (dddd, *J* = 7.7, 7.7, 7.7, 7.7 Hz, 1H), 2.80 (ddd, *J* = 13.6, 10.0, 6.6 Hz, 1H), 2.52 (ddd, *J* = 13.5, 9.6, 7.0 Hz, 1H), 2.10 (dddd, *J* = 10.3, 6.8, 6.8, 6.8, 3.3 Hz, 1H), 1.80-1.70 (m, 2H), 1.35 (d, *J* = 7.7 Hz, 3H), 0.89 (d, *J* = 6.8 Hz, 3H), 0.16 (s, 9H); ¹³C NMR (75 MHz, CDCl₃): δ 171.7, 141.9, 128.0 (4C), 125.4, 75.8, 72.0, 47.1, 38.5, 33.5, 32.6, 9.5, 8.2, 0.1 (3C); EI-MS *m/z* 320 (M)⁺, 261, 249, 230, 207, 174; HRMS *m/z* calcd for C₁₈H₂₈O₃Si: 320.1808, found 320.1825.

(3R, 4S)-4-[(2S, 3R, 4R, 5S)-3, 5-bis(trimethylsilyloxy)-4-methyl-7-phenylheptan-2-yl]-3-methyloxetan-2-one

(18): A flame-dried 25 mL round bottom flask was charged with *O*-trimethylsilylquinine (**4a**) (29 mg, 72.0 μmol) and LiI (289 mg, 2.16 mmol). To the mixture of solids was added CH₂Cl₂ (2.1 mL), Et₂O (0.20 mL) and ^tPr₂NEt (0.31 mL, 1.80 mmol). The resulting mixture was stirred at ambient temperature for 5 min then cooled to -50 °C.

To the reaction mixture was added aldehyde **17** (284 mg, 0.72 mmol) whereupon a solution of propionyl chloride (0.130 mL, 1.44 mmol) in CH₂Cl₂ (0.48 mL) was added via syringe pump over 2h. The heterogeneous reaction mixture was stirred vigorously overnight at -50 °C. The reaction mixture was diluted with ether and filtered through a silica gel plug eluting with ether. The filtrate was concentrated and the crude residue was purified by column chromatography (SiO₂, 24:1/hexanes:ethyl acetate) to yield 258 mg (79%) of β-lactone **18**: ¹H NMR (300 MHz) analysis of the crude product revealed **18** to be a single diastereomer (*syn*, *anti*, *syn*, *anti*, *syn*) within the limits of detection. [α]_D = -4.0 (*c* 1.0, CHCl₃); IR (thin film) 2954, 1828, 1250, 1148, 1117, 1080, 1050, 1002 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 7.32-7.18 (m, 5H), 4.37 (dd, *J* = 11.1, 6.3 Hz, 1H), 3.95-3.91 (m, 2H), 3.77 (dddd, *J* = 14.1, 7.7, 7.7, 7.7 Hz, 1H), 2.59 (dd, *J* = 10.1, 6.4 Hz, 2H), 2.00-1.82 (m, 3H), 1.74-1.65 (m, 1H), 1.34 (d, *J* = 7.8 Hz, 3H), 0.83 (d, *J* = 7.0 Hz, 3H), 0.82 (d, *J* = 6.6 Hz, 3H), 0.14 (s, 18H); ¹³C NMR (75 MHz, CDCl₃): δ 172.1, 141.8, 128.2 (2C), 128.0 (2C), 125.6, 75.9, 72.3, 71.7, 46.3, 41.2, 37.3, 35.2,

32.0, 9.6, 8.2, 7.9, 1.2 (3C), 0.6 (3C); ESI-MS m/z 473 ($M+Na$)⁺; HRMS m/z calcd for $C_{24}H_{42}O_4NaSi_2$ ($M+Na$)⁺: 473.2519, found 473.2532.

(3S, 4R)-4-[2-(4-Methoxybenzyloxy)ethyl]-3-methyloxetan-2-one (20): To a solution of *O*-trimethylsilylquinidine (**3a**) (0.926 g, 2.32 mmol) and $LiClO_4$ (0.737 g, 6.95 mmol) in 23 mL of diethyl ether was added 46 mL of CH_2Cl_2 and the reaction mixture was cooled to $-78^\circ C$. To the resulting mixture was added *N,N*-diisopropylethylamine (10.0 mL, 57.9 mmol) followed by 3-(4-methoxybenzyloxy)propionaldehyde (4.49 g, 23.2 mmol). A solution of propionyl chloride (4.0 mL, 46.4 mmol) in 6.0 mL CH_2Cl_2 was then added over 1 h by syringe pump. The reaction mixture was stirred for 18 h then quenched at $-78^\circ C$ by adding 100 mL of Et_2O and 100 mL saturated aqueous NH_4Cl . The layers were shaken and separated and the aqueous portion was extracted with Et_2O (3×100 mL). The combined organic portions were washed with brine, dried (Na_2SO_4) and concentrated. The crude product mixture was purified by flash chromatography (SiO_2 , 15% ethyl acetate in hexane) to give 4.06 g (70%) of the title compound as a colorless oil (*syn: anti* = 89: 11). Separating the enantiomers by chiral HPLC [Daicel ChiracelTM OD-H column, flow rate 1.0 mL/min, 10% $iPrOH$, 90% hexane, T_r : 13.3 min (3R, 4S), 18.7 min (3S, 4R)] provided only the (3S, 4R) enantiomer (99% ee). $[\alpha]_D +38.8$ (*c* 1.81, $CHCl_3$); IR (thin film): 1817, 1613, 1514, 1248 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ 7.26 (d, *J* = 8.6 Hz, 2H), 6.89 (d, *J* = 8.7 Hz, 2H), 4.79 (ddd, *J* = 8.2, 6.5, 5.2 Hz, 1H), 4.46 (s, 2H), 3.82 (s, 3H), 3.77 (dq, *J* = 7.7, 6.5 Hz, 1H), 3.67-3.56 (m, 2H), 2.05-1.96 (m, 2H), 1.28 (d, *J* = 7.8 Hz, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 172.1, 158.8, 129.7, 128.8, 113.3, 72.5, 72.3, 65.2, 54.7, 46.8, 30.0, 7.7; HRMS m/z calcd for $C_{14}H_{18}O_4$: 250.1205; found: 250.1203.

(2S, 3R)-N-Methoxy-N-methyl-3-hydroxy-5-(4-methoxybenzyloxy)-2-methylpentanamide (20'): To a suspension of *N,N*-dimethylhydroxylamine hydrochloride (2.26 g, 23.2 mmol) in 40 mL CH_2Cl_2 at $0^\circ C$ was added dimethylaluminum chloride (1.0 M solution in hexanes, 23.2 mL, 23.2 mmol) dropwise. The resulting mixture was warmed to ambient temperature and stirred for 2 h, then cooled to $-25^\circ C$ and a solution of **20** (2.90 g, 11.6 mmol) in 30 mL CH_2Cl_2 was added dropwise. The reaction mixture was stirred 2 h then was quenched at $-25^\circ C$ by adding 50 mL of aqueous Rochelle's salt solution. The resulting mixture was extracted with CH_2Cl_2 (3×100 mL). The combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated. The crude product was used without further purification. $[\alpha]_D +10.1$ (*c* 1.92, $CHCl_3$); IR (thin film): 3459 (br), 1655, 1248 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ 7.26 (d, *J* = 8.6 Hz, 2H), 6.87 (d, *J* = 8.7 Hz, 2H), 4.45 (s, 2H), 4.05 (dt, *J* = 9.1, 3.7 Hz, 1H), 3.93 (s, 1H), 3.81 (s, 3H), 3.67 (s, 3H), 3.67-3.63 (m, 2H), 3.19 (s, 3H), 2.92 (br, 1H), 1.89-1.78 (m, 1H), 1.73-1.63 (m, 1H), 1.20 (d, *J* = 7.0 Hz, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 177.1, 158.8, 130.0, 128.8, 113.3, 72.4, 70.1, 67.6, 61.1, 54.8, 39.3, 33.7, 31.5, 11.1; HRMS m/z calcd for $C_{16}H_{25}NO_5$: 311.1733; found: 311.1733.

(2S, 3R)-N-Methoxy-N-methyl-3-(tert-butyldimethylsilyloxy)-5-(4-methoxybenzyloxy)-2-methylpentanamide (20''): To a solution of **20'** (11.6 mmol crude) in 35 mL of DMF at ambient temperature was added imidazole (1.71 g, 25.1 mmol) followed by TBSCl (3.15 g, 20.9 mmol) and DMAP (142 mg, 1.16 mmol). The reaction mixture was stirred for 20 h then diluted with 100 mL saturated aqueous NH_4Cl . The mixture was extracted with CH_2Cl_2 (3×150 mL). The combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated. The crude product mixture was purified by flash chromatography (SiO_2 , 10% ethyl acetate in hexane) to give 4.80 g (97% over 2 steps) of the title compound as a colorless oil. $[\alpha]_D +0.30$ (*c* 1.76, $CHCl_3$); IR (thin film): 1661, 1514, 1250, 837, 776 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ 7.25 (d, *J* = 8.8 Hz, 2H), 6.87 (d, *J* = 8.7 Hz, 2H), 4.41 (s, 2H), 4.03 (dt, *J* = 7.9, 4.9 Hz, 1H), 3.81 (s, 3H), 3.61 (s, 3H), 3.58-3.49 (m, 2H), 3.15 (s, 3H), 3.04-2.96 (m, 1H), 1.86-1.80 (m, 2H), 1.14 (d, *J* = 6.9, 3H), 0.89 (s, 9H), 0.062 (s, 3H), 0.050 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 176.1, 159.1, 130.6, 129.2, 113.6, 72.4, 71.3, 66.2, 61.1, 55.2, 41.2, 35.4, 32.1, 25.9, 18.0, 14.3, -4.4, -4.5; HRMS m/z calcd for (M^+Na) $C_{22}H_{39}NO_5NaSi$: 448.2495; found: 448.2452.

(2S, 3R)-3-(tert-Butyldimethylsilyloxy)-5-(4-methoxybenzyloxy)-2-methylpentanal (21): To a solution of **20''** (4.45 g, 10.45 mmol) in 50 mL of THF at $-50^\circ C$ was added tBu_2AlH (1.0 M in hexanes, 13.0 mL, 13.0 mmol) over 30 min by syringe pump. The reaction mixture was stirred for 30 min then was quenched by adding 100 mL aqueous Rochelle's salt solution.

The mixture was extracted with Et₂O (3 × 100 mL). The combined organic extracts were washed with brine, dried (Na₂SO₄) and concentrated. The crude product mixture was purified by flash chromatography (SiO₂, 11% ethyl acetate in hexane) to give 3.68 g (96%) of the title compound as a colorless oil. [α]_D +47.7 (*c* 1.46, CHCl₃); IR (thin film): 1726, 1613, 1514, 1250, 837, 776 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 9.78 (s, 1H), 7.26 (d, *J* = 8.7 Hz, 2H), 6.89 (d, *J* = 8.6 Hz, 2H), 4.45 (d, *J* = 11.5 Hz, 1H), 4.39 (d, *J* = 11.5 Hz, 1H), 4.31 (ddd, *J* = 7.2, 5.7, 3.7 Hz, 1H), 3.82 (s, 3H), 3.49 (t, *J* = 5.7 Hz, 2H), 2.53-2.44 (m, 1H), 1.82-1.77 (m, 2H), 1.06 (d, *J* = 7.0 Hz, 3H), 0.87 (s, 9H), 0.072 (s, 3H), 0.047 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 204.3, 159.0, 130.1, 129.0, 113.5, 72.4, 69.0, 65.9, 54.8, 51.3, 34.4, 25.5, 17.7, 7.5, -4.7, -4.9; HRMS *m/z* calcd for (M⁺+Na) C₂₀H₃₄O₄SiNa: 389.2124; found: 389.2114.

(3R, 4S)-4-[(2S, 3R)-2-(tert-Butyldimethylsilanyloxy)-4-(4-methoxybenzyloxy)-1-methylbutyl]-3-methyloxetan-2-one (22): To a suspension of *O*-trimethylsilylquinine (**4a**) (397 mg, 0.99 mmol) and LiI (6.64 g, 49.59 mmol) in 4.5 mL of diethyl ether was added 40.0 mL of CH₂Cl₂ and the reaction mixture was cooled to -78 °C. To the

resulting mixture was added *N,N*-diisopropylethylamine (4.3 mL, 24.9 mmol) followed by a solution of **21** (3.43 g, 9.92 mmol) in 8.0 mL CH₂Cl₂. A solution of propionyl chloride (1.73 mL, 20.0 mmol) in 6.0 mL CH₂Cl₂ was then added over 3 h by syringe pump. After the addition of propionyl chloride was complete, the reaction was quenched at the -78 °C by adding 100 mL of Et₂O and 100 mL saturated aqueous NH₄Cl. The resulting mixture was extracted with Et₂O (3 × 100 mL) and the combined organic extracts were washed with brine, dried (Na₂SO₄) and concentrated. The crude product mixture was purified by flash chromatography (SiO₂, 9% ethyl acetate in hexane) to give 3.60 g (91%) of the title compound as a single diastereomer. [α]_D -20.3 (*c* 2.28, CHCl₃); IR (thin film): 1824, 1613, 1250 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.25 (d, *J* = 8.7 Hz, 2H), 6.88 (d, *J* = 8.6 Hz, 2H), 4.47 (d, *J* = 11.6 Hz, 1H), 4.41 (dd, *J* = 11.1, 6.3 Hz, 1H), 4.36 (d, *J* = 11.4 Hz, 1H), 4.10 (t, *J* = 6.2 Hz, 1H), 3.82 (s, 3H), 3.75 (m, 1H), 3.43 (t, *J* = 6.4 Hz, 2H), 1.88-1.78 (m, 3H), 1.26 (d, *J* = 7.7 Hz, 3H), 0.89 (s, 9H), 0.81 (d, *J* = 6.7 Hz, 3H), 0.065 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ 172.6, 159.0, 130.2, 129.1, 113.7, 75.9, 72.5, 67.5, 66.4, 55.1, 46.3, 37.0, 35.0, 25.8, 18.0, 8.2, 7.3, -4.5, -5.0; HRMS *m/z* calcd for (M⁺-*t*Bu) C₁₉H₂₉O₅Si: 365.1784; found: 365.1786.

(2S, 3R, 4S, 5R)-5-(tert-Butyldimethylsilanyloxy)-7-(4-methoxybenzyloxy)-2, 4-dimethylheptane-1,3-diol (22'): To a solution of **22** (3.60 g, 8.52 mmol) in 60 mL of THF at -50 °C was added ^tBu₂AlH (1.0 M in hexanes, 25.6 mL, 25.6 mmol) over 30

min by syringe pump. The reaction mixture was stirred for 10 min then 50 mL of Et₂O and 50 mL aqueous Rochelle's salt solution were added. The reaction mixture was extracted with Et₂O (3 × 100 mL) and the combined organic extracts were washed with brine, dried (Na₂SO₄) and concentrated. The crude product was used without further purification. [α]_D +33.0 (*c* 1.99, CHCl₃); IR (thin film): 3465 (br), 1514, 1250, 837, 778 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.26 (d, *J* = 8.6 Hz, 2H), 6.89 (d, *J* = 8.7 Hz, 2H), 4.46 (d, *J* = 11.5 Hz, 1H), 4.40 (d, *J* = 11.4 Hz, 1H), 4.00-3.96 (m, 1H), 3.94 (dd, *J* = 10.3, 2.0 Hz, 1H), 3.82 (s, 3H), 3.80 (dd, *J* = 10.5, 3.6 Hz, 1H), 3.68 (dd, *J* = 10.6, 5.4 Hz, 1H), 3.60-3.48 (m, 2H), 1.94-1.83 (m, 3H), 1.66-1.63 (m, 1H), 0.96 (d, *J* = 7.0 Hz, 3H), 0.90 (s, 9H), 0.75 (d, *J* = 7.1 Hz, 3H), 0.133 (s, 3H), 0.074 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 159.1, 130.4, 129.2, 113.7, 76.0, 74.9, 72.5, 67.8, 66.6, 55.2, 39.7, 36.2, 31.1, 25.6, 17.8, 13.4, 8.3, -4.5, -5.1.

(2S, 3R, 4S, 5R)-5-(tert-Butyldimethylsilanyloxy)-3-hydroxy-7-(4-methoxybenzyloxy)-2, 4-dimethylheptyl 4-methylbenzenesulfonate (22''): To a solution of **22'** (1.81 g, 4.24 mmol) in 40 mL of CH₂Cl₂ at ambient temperature was added pyridine

(840 mg, 10.6 mmol) followed by DMAP (104 mg, 0.85 mmol) and TsCl (1.21 g, 6.37 mmol). The reaction mixture was stirred for 22 h then was quenched by adding 40 mL CH₂Cl₂ and 50 mL saturated aqueous NH₄Cl. The mixture was extracted with CH₂Cl₂ (3 × 80 mL) and the combined organic extracts were washed with brine, dried (Na₂SO₄) and concentrated. The crude product mixture was purified by flash chromatography (SiO₂, 7% ethyl acetate in hexane) to give 2.05 g (83% over 2 steps) of the title compound as a colorless oil. [α]_D +18.6 (*c* 1.86, CHCl₃); IR (thin film): 3469 (br), 1613, 1514, 1361, 1249, 964, 838 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.80 (d, *J* = 8.3 Hz, 2H), 7.33 (d, *J* = 8.0 Hz, 2H), 7.26 (d, *J* = 8.7 Hz, 2H), 6.89 (d, *J* = 8.7 Hz, 2H), 4.45 (d, *J* = 11.5 Hz, 1H), 4.39 (d, *J* = 11.5 Hz, 1H), 4.12 (d, *J* = 7.3 Hz, 1H), 4.09 (d, *J* = 7.2 Hz, 1H), 3.98-3.91 (m, 2H),

3.82 (s, 3H), 3.70 (dd, $J = 10.2, 1.7$ Hz, 1H), 3.56-3.47 (m, 2H), 2.44 (s, 3H), 1.92-1.70 (m, 4H), 0.87 (s, 9H), 0.84 (d, $J = 6.8$ Hz, 3H), 0.69 (d, $J = 7.1$ Hz, 3H), 0.094 (s, 3H), 0.057 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 159.0, 144.3, 133.2, 130.3, 129.6, 129.1, 127.7, 113.6, 74.6, 73.5, 72.4, 71.7, 66.4, 55.1, 39.3, 35.2, 31.0, 25.6, 21.4, 17.7, 13.0, 8.1, -4.6, -5.2; HRMS m/z calcd for ($\text{M}^+ + \text{Na}$) $\text{C}_{30}\text{H}_{48}\text{O}_7\text{SSiNa}$: 603.2788; found: 603.2778.

(2S, 3R, 4S, 5R)-5-(tert-Butyldimethylsilanyloxy)-3-methoxy-7-(4-methoxybenzyloxy)-2, 4-dimethylheptyl 4-methylbenzenesulfonate (23): To a solution of **22''** (2.09 g, 3.60 mmol) in 40 mL of CH_2Cl_2 at ambient temperature was added proton sponge

(3.20 g, 21.6 mmol) followed by Me_3OBF_4 (4.626 g, 21.6 mmol). The reaction flask was covered with aluminum foil and the reaction stirred for 8 h. The reaction was quenched by adding 100 mL of CH_2Cl_2 and 150 mL of saturated aqueous NH_4Cl . The mixture was extracted with CH_2Cl_2 (3×100 mL) and the combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated. The crude product mixture was purified by flash chromatography (SiO_2 , 8% ethyl acetate in hexane) to give 1.73 g (81%) of the title compound as a colorless oil. $[\alpha]_D -0.67$ (c 1.79, CHCl_3); IR (thin film): 1514, 1365, 1249, 1189, 1178, 835 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.81 (d, $J = 8.3$ Hz, 2H), 7.35 (d, $J = 8.0$ Hz, 2H), 7.25 (d, $J = 8.7$ Hz, 2H), 6.87 (d, $J = 8.7$ Hz, 2H), 4.44 (d, $J = 11.6$ Hz, 1H), 4.38 (d, $J = 11.6$ Hz, 1H), 4.09-4.04 (m, 2H), 4.02-3.94 (m, 1H), 3.81 (s, 3H), 3.41 (td, $J = 6.8, 1.0$ Hz, 2H), 3.33 (s, 3H), 3.26 (dd, $J = 9.4, 1.7$ Hz, 1H), 2.46 (s, 3H), 2.02-1.95 (m, 1H), 1.86-1.75 (m, 2H), 1.56-1.50 (m, 1H), 0.87 (s, 9H), 0.77 (d, $J = 6.8$ Hz, 3H), 0.67 (d, $J = 6.9$ Hz, 3H), 0.068 (s, 3H), 0.062 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 158.7, 144.3, 133.0, 130.2, 129.4, 128.6, 127.5, 113.3, 80.0, 72.3, 72.2, 69.0, 66.5, 60.3, 54.7, 39.8, 35.4, 34.7, 25.7, 21.1, 17.9, 9.0, 8.8, -3.7, -4.6; HRMS m/z calcd for ($\text{M}^+ - \text{tBu}$) $\text{C}_{27}\text{H}_{41}\text{O}_7\text{SSi}$: 537.2342; found: 537.2316.

(3R, 4S, 5R, 6S)-3-(tert-Butyldimethylsilanyloxy)-5-methoxy-1-(4-methoxybenzyloxy)-4, 6-dimethyl-9-decene (24): To a suspension of CuBr (825 mg, 5.75 mmol) in 2.5 mL of Et_2O at -20 $^\circ\text{C}$ was added allylmagnesium bromide (1.0 M in

Et_2O , 11.5 mL, 11.5 mmol) dropwise and the reaction mixture was stirred for 30 min. To the resulting mixture was added a solution of **23** (683 mg, 1.15 mmol) in 4.0 mL Et_2O and the reaction was stirred 10 min at -20 $^\circ\text{C}$ then warmed to ambient temperature and stirred for 4.5 h. The reaction was quenched by adding 20 mL of saturated aqueous NH_4Cl . The mixture was extracted by Et_2O (3×40 mL) and the combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated. The crude product mixture was purified by flash chromatography (SiO_2 , 6% ethyl acetate in hexane) to give 454 mg (85%) of the title compound as an oil. $[\alpha]_D -12.4$ (c 1.65, CHCl_3); IR (thin film): 1640, 1614, 1514, 1249, 1097, 835 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.27 (d, $J = 8.7$ Hz, 2H), 6.88 (d, $J = 8.7$ Hz, 2H), 5.83 (ddt, 17.0, 10.2, 6.7 Hz, 1H), 5.0 (m, 2H), 4.45 (d, $J = 11.5$ Hz, 1H), 4.40 (d, $J = 11.5$ Hz, 1H), 4.13 (td, $J = 6.9, 1.4$ Hz, 1H), 3.81 (s, 3H), 3.47 (s, 3H), 3.43 (td, $J = 6.8, 1.7$ Hz, 2H), 3.13 (dd, $J = 9.4, 1.7$ Hz, 1H), 2.23-2.02 (m, 2H), 1.89-1.82 (m, 2H), 1.69-1.40 (m, 4H), 0.89 (s, 9H), 0.83 (d, $J = 6.7$ Hz, 3H), 0.73 (d, $J = 6.9$ Hz, 3H), 0.091 (s, 3H), 0.083 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 159.0, 139.0, 130.6, 129.1, 114.3, 113.7, 84.7, 72.6, 69.3, 67.2, 60.8, 55.2, 40.7, 35.9, 34.4, 32.0, 26.0, 18.3, 12.8, 9.5, -3.4, -4.3; HRMS m/z calcd for $\text{C}_{27}\text{H}_{48}\text{O}_4\text{Si}$: 464.3322; found: 464.3304.

(E, 3R, 4S, 5R, 6S)-3-(tert-Butyldimethylsilanyloxy)-5-methoxy-1-(4-methoxybenzyloxy)-4, 6-dimethyl-8-decene (25): To a solution of NaBPh_4 (7.7 mg, 0.0225 mmol) in 0.2 mL of acetone was added a solution of $[(\text{C}_8\text{H}_{14})_2\text{IrCl}]_2$

(10.2 mg, 0.0113 mmol) and PCy_3 (19.0 mg, 0.0678 mmol) in 10.8 mL of CH_2Cl_2 . The reaction mixture was stirred for 5 min, then this catalyst solution was added to **24** (550 mg, 1.185 mmol). The reaction mixture was stirred for 3 h and then filtered through silica gel eluting with 120 mL of Et_2O . The filtrate was concentrated and the crude product mixture was purified by flash chromatography (SiO_2 , 6% ethyl acetate in hexane) to give 539 mg (98%) of the title compound as a colorless oil. $[\alpha]_D -8.5$ (c 1.02, CHCl_3); IR (thin film): 1614, 1249, 1097, 835 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.27 (d, $J = 8.7$ Hz, 2H), 6.88 (d, $J = 8.7$ Hz, 2H), 5.51-5.36 (m, 2H), 4.45 (d, $J = 11.6$ Hz, 1H), 4.40 (d, $J = 11.6$ Hz, 1H), 4.12 (t, $J = 6.2$ Hz, 1H), 3.81 (s, 3H), 3.48 (s, 3H), 3.43 (td, $J = 7.7, 2.4$ Hz, 2H), 3.18 (dd, $J = 9.4, 1.6$ Hz, 1H), 2.15 (m, 1H), 2.01 (m, 1H), 1.87 (dd, $J = 6.8, 2.2$ Hz, 1H), 1.82 (dd, $J = 7.3, 3.6$ Hz, 1H), 1.67 (d, $J = 5.0$ Hz, 3H), 1.58 (br, 1H), 1.54-1.46 (m, 1H), 0.89 (s, 9H), 0.81 (d, $J = 6.8$ Hz, 3H), 0.73 (d, $J = 6.9$ Hz, 3H), 0.079 (s, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ 158.9, 130.5, 128.9,

126.0, 113.5, 83.6, 72.4, 69.3, 67.0, 60.6, 54.9, 40.7, 38.2, 35.9, 35.5, 25.9, 18.2, 17.8, 12.7, 9.4, -3.5, -4.5; HRMS m/z calcd for $C_{27}H_{48}O_4Si$: 464.3322; found: 464.3345.

(E, 3R, 4S, 5R, 6S)-3-(tert-Butyldimethylsilanyloxy)-5-methoxy-4, 6-dimethyldec-8-en-1-ol (25'): To a 0 °C solution of DDQ (282 mg, 1.24 mmol) in 2.2 mL of CH_2Cl_2 and 0.45 mL of pH 7 buffer was added a solution of **25** (486 mg, 1.05 mmol) in 7.0 mL of CH_2Cl_2 . The resulting mixture was stirred for 40 min and then quenched by adding 10 mL of H_2O . The mixture was extracted with Et_2O (3×40 mL) and the combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated. The crude product mixture was purified by flash chromatography (SiO_2 , gradient 6%→10%→20% ethyl acetate in hexane) to give 291 mg (81%) of the title compound as an oil. $[\alpha]_D -9.3$ (c 1.40, $CHCl_3$); IR (thin film): 3401(br), 1253, 1077, 835, 774 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ 5.51-5.35 (m, 2H), 4.15-4.10 (m, 1H), 3.69 (td, J = 6.5, 1.4 Hz, 2H), 3.49 (s, 3H), 3.16 (dd, J = 9.2, 1.8 Hz, 1H), 2.15 (m, 1H), 2.01 (m, 1H), 1.84-1.76 (m, 2H), 1.67 (d, J = 5.3 Hz, 3H), 1.64-1.59 (m, 2H), 0.91 (s, 9H), 0.83 (d, J = 6.8 Hz, 3H), 0.77 (d, J = 7.0 Hz, 3H), 0.113 (s, 3H), 0.107 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 130.3, 126.3, 84.2, 69.2, 60.8, 59.4, 40.2, 38.5, 38.2, 35.5, 25.9, 18.2, 17.9, 12.6, 9.6, -3.4, -4.4; HRMS m/z calcd for ($M^+ - tBu$) $C_{15}H_{31}O_3Si$: 287.2042; found: 287.2050.

(E, 3R, 4S, 5R, 6S)-3-(tert-Butyldimethylsilanyloxy)-5-methoxy-4, 6-dimethyldec-8-enal (26): To a solution of DMSO (139 μ L, 1.96 mmol) in 5.0 mL of CH_2Cl_2 at -78 °C was added a solution of $(COCl)_2$ (1.25 equiv, 490 μ L, 0.98 mmol) in CH_2Cl_2 (0.5 mL) dropwise. The reaction mixture was stirred for 40 min whereupon a solution of **25'** (270 mg, 0.785 mmol) in 2.5 mL of CH_2Cl_2 was added. The resulting mixture was stirred for 30 min then Et_3N (548 μ L, 3.92 mmol) was added dropwise and the reaction mixture was stirred for another 30 min then warmed to ambient temperature and quenched by adding 10 mL of saturated aqueous NH_4Cl . The mixture was extracted by CH_2Cl_2 (3×40 mL) and the combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated. The crude product mixture was purified by flash chromatography (SiO_2 , 6% ethyl acetate in hexane) to give 235 mg (88%) of the title compound as an oil. $[\alpha]_D -6.2$ (c 1.62, $CHCl_3$); IR (thin film): 2718, 1728, 1472, 1134, 1089, 836, 775 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ 9.79 (t, J = 2.4 Hz, 1H), 5.53-5.35 (m, 2H), 4.63 (td, J = 6.2, 1.8 Hz, 1H), 3.50 (s, 3H), 3.17 (dd, J = 9.2, 1.8 Hz, 1H), 2.67 (ddd, J = 16.1, 6.2, 2.5 Hz, 1H), 2.55 (ddd, J = 16.1, 6.2, 2.3 Hz, 1H), 2.15 (m, 1H), 2.05 (m, 1H), 1.76-1.66 (m, 1H), 1.67 (d, J = 4.9 Hz, 3H), 1.60-1.50 (m, 1H), 0.89 (s, 9H), 0.82 (d, J = 6.9 Hz, 3H), 0.80 (d, J = 7.0 Hz, 3H), 0.13 (s, 3H), 0.07 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 201.1, 130.3, 126.3, 83.9, 66.9, 60.6, 50.8, 42.6, 38.2, 35.5, 25.8, 18.2, 17.9, 12.8, 10.1, -3.8, -4.5; HRMS (electrospray) m/z calcd for $C_{19}H_{38}O_3NaSi$ ($M+Na$) $^+$: 365.2488; found: 365.2473.

(3S, 4R)-4-[(E, 2R, 3S, 4R, 5S)-2-(tert-Butyldimethylsilanyloxy)-4-methoxy-3, 5-dimethylnon-7-enyl]-3-ethyloxetan-2-one (28): To a solution of triamine ligand (169.5 mg, 0.25 mmol)⁴ in 2.0 mL of benzotrifluoride at ambient temperature was slowly added a solution of $AlMe_3$ (2.0 M in hexane, 0.125 mL, 0.25 mmol). After stirring for 2 h, the resulting catalyst solution was cooled to -25 °C and *N, N*-diisopropylethylamine (0.173 mL, 1.0 mmol) and butyryl bromide (0.212 mL, 2.0 mmol) were added consecutively. The resulting heterogeneous mixture was stirred 20 min at -25 °C whereupon **26** (170 mg, 0.50 mmol) in 1.0 mL of benzotrifluoride was added dropwise and the reaction was stirred 21 h at -25 °C. The reaction was quenched at -25 °C by adding 10 mL of a 3% Et_3N - Et_2O solution (v/v) and the resulting mixture was filtered through silica gel eluting with 60 mL of Et_2O . The filtrate was concentrated and the crude product mixture was purified by flash chromatography (gradient 4%→6% ethyl acetate in hexane) to give 135 mg (65%) of the title compound. $[\alpha]_D +20.2$ (c 1.06, $CHCl_3$); IR (thin film): 1828, 1462, 1254, 836, 775 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ 5.53-5.35 (m, 2H), 4.64 (dt, J = 6.5, 6.5 Hz, 1H), 4.20 (td, J = 6.9, 1.1 Hz, 1H), 3.57 (dt, J = 9.1, 7.4 Hz, 1H), 3.48 (s, 3H), 3.16 (dd, J = 9.5, 1.5 Hz, 1H), 2.15 (m, 1H), 2.02 (m, 1H), 1.89 (t, J = 6.5 Hz, 2H), 1.85-1.79 (m, 1H), 1.74-1.66 (m, 1H), 1.67 (d, J = 4.7 Hz, 3H), 1.62-1.60 (m, 2H), 1.09 (t, J = 7.4 Hz, 3H), 0.91 (s, 9H), 0.82 (d, J = 6.8 Hz, 3H), 0.76 (d, J = 6.9 Hz, 3H), 0.11 (s, 3H), 0.10 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 172.0, 130.4, 126.4, 83.8, 72.6, 69.2, 60.8, 54.2, 41.6, 38.3, 36.6, 35.6, 25.9, 18.3, 18.0, 17.6, 12.7, 12.1, 9.9, -3.7, -3.9; HRMS (electrospray) m/z calcd for $C_{23}H_{44}O_4NaSi$ ($M+Na$) $^+$: 435.2907; found: 435.2897.

(*E,4S,5R,7R,8S,9R,10S*)-tert-Butyl 7-(tert-butyldimethylsilanyloxy)-4-ethyl-5-hydroxy-9-methoxy-8, 10-dimethyl-3-oxotetradec-12-enoate (29**):** To a solution of KHMDS (0.5 M in toluene, 1.14 mL, 0.57 mmol) in 1.0 mL of THF at $-78\text{ }^{\circ}\text{C}$ was added tert-butyl acetate (66.2 mg, 0.57 mmol).

The reaction mixture was stirred for 1h then transferred via cannula to a suspension of $\text{MgBr}_2 \cdot \text{OEt}_2$ (147 mg, 0.57 mmol) in 1.0 mL of THF. The resulting mixture was stirred for 1h whereupon **28** (47 mg, 0.114 mmol) was added and the reaction was stirred for an additional 30 min. The reaction was quenched by adding 10 mL of saturated aqueous NH_4Cl and the mixture was extracted with Et_2O ($3 \times 30\text{ mL}$). The combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated. The crude product mixture was purified by flash chromatography (SiO_2 , 15% ethyl acetate in hexanes) to give 40 mg (66%) of the title compound as an inseparable mixture keto-enol tautomers. IR (thin film): 3467 (br), 1733, 1709 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 5.50-5.33 (m, 2H), 4.14-4.10 (m, 1H), 4.04 (q, $J = 4.9\text{ Hz}$, 1H), 3.48 (d, $J = 1.6\text{ Hz}$, 2H), 3.44 (s, 3H), 3.06 (dd, $J = 8.2, 1.9\text{ Hz}$, 1H), 2.75 (dt, $J = 8.9, 4.3\text{ Hz}$, 1H), 2.16-2.07 (m, 1H), 2.02-1.93 (m, 1H), 1.88-1.73 (m, 3H), 1.67 (d, $J = 5.2\text{ Hz}$, 3H), 1.63-1.58 (m, 4H), 1.47 (s, 9H), 0.92 (t, $J = 7.4\text{ Hz}$, 3H), 0.90 (s, 9H), 0.85 (d, $J = 6.8\text{ Hz}$, 3H), 0.84 (d, $J = 6.9\text{ Hz}$, 3H), 0.11 (s, 3H), 0.10 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 206.6, 166.5, 130.2, 126.5, 84.7, 81.7, 71.6, 69.3, 60.2, 59.0, 52.0, 40.7, 39.2, 38.6, 35.5, 28.3, 28.0, 26.0, 19.8, 17.9, 13.4, 12.2, 11.5, -3.8 , -4.3 ; HRMS m/z calcd for $\text{C}_{29}\text{H}_{56}\text{O}_6\text{Si}$: 528.3846; found: 528.3842.

(*E,4S,5R,7R,8S,9R,10S*)-tert-Butyl 4-ethyl-3,5,7-trihydroxy-9-methoxy-8, 10-dimethyltetradec-12-enoate (29'**):** To a solution of **29** (27 mg, 0.05 mmol) in 1.0 mL of EtOH at $0\text{ }^{\circ}\text{C}$ was added NaBH_4 (6 mg, 0.16 mmol). The resulting mixture was stirred for 40 min and then quenched by adding

50 μL acetic acid. The mixture was extracted with ethyl acetate ($3 \times 30\text{ mL}$) and the combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated. The crude product used without further purification.

(-)-Pironetin (19**):** A solution of **29'** (15 mg crude product from previous transformation) and $\text{TsOH} \cdot \text{H}_2\text{O}$ (5 mg) in 1.0 mL of toluene was heated at $110\text{ }^{\circ}\text{C}$ for 30 min. The mixture was then cooled to ambient temperature and diluted with 50 mL of ethyl acetate. The resulting mixture was washed successively with 10 mL of aqueous NaHCO_3 solution and brine and the organic layer was dried (Na_2SO_4) and concentrated. The crude product mixture was purified by flash chromatography (SiO_2 , 40% ethyl acetate in hexanes) to give 4.6 mg (56% over two steps) of the title compound as a white solid. $[\alpha]_{\text{D}} -135$ (c 0.35, CHCl_3) {lit.¹⁰ $[\alpha]_{\text{D}} -144$ (c 0.50, CHCl_3)}; ^1H NMR (300 MHz, CDCl_3) δ 7.02 (dd, $J = 9.8, 6.0\text{ Hz}$, 1H), 6.03 (d, $J = 9.8\text{ Hz}$, 1H), 5.51-5.32 (m, 2H), 4.74 (m, 1H), 4.21 (brd, 1H), 3.48 (s, 3H), 3.45 (d, $J = 2.3\text{ Hz}$, 1H), 2.99 (dd, $J = 6.1, 4.4\text{ Hz}$, 1H), 2.33-2.25 (m, 1H), 2.13-2.06 (m, 1H), 1.92-1.84 (m, 2H), 1.81-1.76 (m, 1H), 1.74-1.65 (m, 3H), 1.67 (d, $J = 5.3\text{ Hz}$, 3H), 1.58-1.46 (m, 1H), 1.01 (d, $J = 7.1\text{ Hz}$, 3H), 0.97 (t, $J = 7.4\text{ Hz}$, 3H), 0.96 (d, $J = 6.9\text{ Hz}$, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 164.6, 150.6, 128.8, 126.9, 120.9, 91.1, 77.8, 67.5, 61.5, 39.1, 39.0, 37.3, 36.7, 36.2, 20.8, 17.9, 15.2, 12.2, 11.0; HRMS m/z calcd for $(\text{M}^+ - \text{H}_2\text{O}) \text{C}_{19}\text{H}_{30}\text{O}_3$: 306.2195; found: 306.2185.

¹⁰ Kobayashi, S.; Tsochia, K.; Harada, T.; Nishide, M.; Kurokawa, T.; Nakagawa, T.; Shimada, N.; Kobayashi, K. *J. Antibiot.* **1994**, *47*, 697.

Stereochemical Proofs

(3*R*, 4*R*)-4-[(2*R*, 3*S*)-3-(4-Methoxybenzyloxy)-5-phenylpentan-2-yl]-3-methyloxetan-2-one (14*c*): Lactone **14*c*** was converted to the crystalline derivative **14*c*'** whose structure was determined by X-ray diffraction analysis.

Figure S1. X-ray Crystal Structure of (3*S*,4*S*,5*R*,6*R*)-6-(Benzylcarbamoyl)-5-hydroxy-4-methyl-1-phenylheptan-3-yl 4-bromobenzoate (**14*c*'**)

Table S1. Crystal data and structure refinement for **14*c*'**

Empirical formula	C ₂₉ H ₃₂ BrNO ₄	
Formula weight	538.47	
Temperature	295(2) °K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	P2(1)	
Unit cell dimensions	a = 7.3752(8) Å	α = 90°.
	b = 7.6119(8) Å	β = 92.553(2)°.
	c = 24.195(3) Å	γ = 90°.
Volume	1356.9(2) Å ³	
Z	2	
Density (calculated)	1.318 Mg/m ³	
Absorption coefficient	1.548 mm ⁻¹	
F(000)	560	
Theta range for data collection	2.53 to 25.00°.	
Index ranges	-8 ≤ h ≤ 8, 0 ≤ k ≤ 9, 0 ≤ l ≤ 28	
Reflections collected	2579	
Independent reflections	2579 [R(int) = 0.0000]	
Completeness to theta = 25.00°	99.9 %	
Absorption correction	None	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	2579 / 1 / 287	
Goodness-of-fit on F ²	0.851	
Final R indices [I > 2σ(I)]	R1 = 0.0604, wR2 = 0.1269	
R indices (all data)	R1 = 0.1123, wR2 = 0.1383	
Absolute structure parameter	0.39(5)	
Largest diff. peak and hole	0.980 and -0.228 e.Å ⁻³	

(3*R*, 4*S*)-4-[(2*S*, 3*S*)-5-Phenyl-3-trimethylsilyloxypentan-2-yl]-3-methyloxetan-2-one (16): Lactone **16** was converted to the crystalline derivative **16'** whose structure was determined by X-ray diffraction analysis.

Figure S2. X-ray Crystal Structure of (3*S*,4*S*)-4-[(2*S*,3*R*)-3-methyl-4-oxo-2-oxetanyl]-1-phenylpentan-3-yl 3,5-dinitrobenzoate (**16'**)

Table S2. Crystal data and structure refinement for **16'**

Empirical formula	C ₂₂ H ₂₂ N ₂ O ₈	
Formula weight	442.42	
Temperature	295(2) K	
Wavelength	0.71073 Å	
Crystal system	Orthorhombic	
Space group	P2(1)2(1)2(1)	
Unit cell dimensions	a = 7.3533(5) Å	α = 90°.
	b = 10.3188(8) Å	β = 90°.
	c = 27.986(2) Å	γ = 90°.
Volume	2123.5(3) Å ³	
Z	4	
Density (calculated)	1.384 Mg/m ³	
Absorption coefficient	0.107 mm ⁻¹	
F(000)	928	
Theta range for data collection	1.46 to 32.56°.	
Index ranges	-11 ≤ h ≤ 10, -15 ≤ k ≤ 15, -41 ≤ l ≤ 42	
Reflections collected	27932	
Independent reflections	7574 [R(int) = 0.0356]	
Completeness to theta = 32.56°	99.3 %	
Absorption correction	None	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	7574 / 0 / 377	
Goodness-of-fit on F ²	1.056	
Final R indices [I > 2σ(I)]	R1 = 0.0613, wR2 = 0.1365	
R indices (all data)	R1 = 0.0874, wR2 = 0.1499	
Absolute structure parameter	0.2(10)	
Largest diff. peak and hole	0.336 and -0.198 e.Å ⁻³